

v. 11, n. 3, May-June 2020



http://www.ijmp.jor.br ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083

Multi objective fuzzy inventory model with deterioration, price and time dependent demand and time dependent holding cost

Satya Kumar Das Govt. General Degree College at Gopiballavpur-II, India E-mail: satyakrdasmath75@gmail.com

> Sahidul Islam University of Kalyani, India E-mail: sahidul.math@gmail.com

> > Submission: 6/25/2019 Revision: 9/18/2019 Accept: 10/2/2019

ABSTRACT

In this paper, we have formulated an inventory model with time dependent holding cost, selling price as well as time dependent demand. Multi-item inventory model has been considered under limitation on storage space. Due to uncertainty all the require cost parameters are taken as generalized trapezoidal fuzzy number. Our proposed multi-objective inventory model has been solved by using fuzzy programming techniques which are FNLP, FAGP, WFNLP and WFAGP methods. A numerical example is provided to demonstrate the application of the model. Finally to illustrate the model and sensitivity analysis and graphical representation have been shown.

Keywords: Inventory, Deterioration, Multi-item, Generalized trapezoidal fuzzy number, Fuzzy programming technique.





http://www.ijmp.jor.br ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083

v. 11, n. 3, May-June 2020

1. INTRODUCTION

Inventory model deals with decisions that minimize the total average cost or maximize the total average profit. In that way to construct a real life mathematical inventory model on base on various assumptions, notations and approximations.

In ordinary inventory system inventory costs i.e set-up cost, holding cost, deterioration cost, etc. are taken fixed amount but in real life inventory system these costs are not always fixed. So consideration of fuzzy variable is more realistic and interesting.

The inventory problems for deteriorating item such as fashionable items, electronics products, fruits, and green vegetables, and many others and the deterioration is defined as the spoilage, damage, dryness, vaporization etc. This results in decrease of usefulness of the commodity. Inventory problem for deteriorating items have been widely studied by many researchers. The economic order quantity model was first introduced in February 1913 by Harris.

Ghare and Schrader (1963) was the first to establish an economic order quantity (EOQ) model for deteriorating items. Then Covert and Philip (1973) extended their research work by presenting a variable rate of deterioration. Later, there are many papers presented on the deteriorating inventory, such as Sridevi et al. (2010), Bhunia and Shaikh (2014), and Ghosh, Sarkar and Chaudhuri (2015) etc. Kumar, et. al (2016) presented on Optimization of Weibull deteriorating items inventory model under the effect of price and time dependent demand with partial backlogging. Yang, H.L discussed on two warehouse partial backlogging inventory model for deteriorating items under inflation.

Yu-Chung Tsao, Gwo-Ji-Sheen (2008) studied on dynamic pricing promotion and replenishment policies for a deteriorating item under permissible delay in payments. Yang (2016) studied on two warehouse partial backlogging inventory model for deteriorating items under inflation. Liang and Zhou (2011) discussed on a two warehouse inventory model for deteriorating items under conditionally permissible delay in Payment.

The demand of an inventory item depends on the price is the most important in real life. Therefore the inventory system should incorporate the selling price as a decision variable. Bhunia and Shaikh (2014) presented a paper on deterministic inventory model for deteriorating items with selling price dependent demand and three-parameter Weibull distributed deterioration. Alfares and Ghaithan (2016) formulated on inventory and pricing model with price-dependent demand, time-varying holding cost, and quantity discounts.



v. 11, n. 3, May-June 2020

ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083

http://www.ijmp.jor.br

Shah et. al (2009) studied on a lot size inventory model for the Weibull distributed deterioration rate with discounted selling price and stock-dependent demand. Sridevi et. al (2010) discussed on Inventory model for deteriorating items with Weibull rate of replenishment and selling price dependent demand. The limitation of the space of the inventory item is the most important factor in the business management system.

Ghosh (2015) presented a paper on a multi-item inventory model for deteriorating items in limited storage space with stock-dependent demand. Islam and Mandal (2017), discussed fuzzy E.O.Q model with constant demand and shortages in a fuzzy signomial geometric programming (FSGP) approach. Mondal et. al (2003) formulated on an inventory system of ameliorating items for price dependent demand rate.

The concept of fuzzy set theory was first introduced by Zadeh in 1965. Afterward Zimmermann (1985) applied the fuzzy set theory concept with some useful membership functions to solve the linear programming problem with some objective functions. Then the various ordinary inventory model transformed to fuzzy versions model by various authors such as Roy and Maity (1995) presented on fuzzy inventory model with constraints.

Islam and Roy (2006) studied on a fuzzy EPQ model with flexibility and reliability consideration and demand depended unit production cost under a space constraint. Islam and Mandal (2017) discussed on fuzzy inventory model (EOQ model) with unit production cost, time depended holding cost, without shortages under a space constrain in a fuzzy parametric geometric programming (FPGP) approach. Maity (2008) developed a paper on fuzzy inventory model with two ware house under possibility measure in fuzzy goal. Roy (2014) presented on fuzzy inventory model for deteriorating items with price dependent demand.

In this paper, we have considered demand rate is depended on selling price as well as time and holding cost is time dependent. Multi-item inventory has been considered under limitation on storage space. Due to uncertainty all the required cost parameters are taken as generalized trapezoidal fuzzy number. The formulated multi objective inventory model has been solved by using FNLP, FAGP, WFNLP and WFAGP methods. A numerical example is considered to illustrate the model. Finally sensitivity analysis and graphical figures have been shown.

2. MATHEMATICAL MODEL

2.1. Notations

 c_i : Ordering cost per order for ith item.



http://www.ijmp.jor.br ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083

- $H_i(=\alpha_i t)$: Holding cost per unit and per unit time for ith item.
- d_i : Deteriorating cost per unit and per unit time for ith item.
- θ_i : Constant deterioration rate per unit time for the ith item.
- T_i : The length of cycle time for ith item ($T_i > 0$).
- $D(S_i, t)$: Demand rate per unit time for the ith item.
- S_i : Selling price for the ith item.
- P_i : Purchasing cost for the ith item.
- $I_i(t)$: Inventory level of the ith items at time t.
- Q_i : The order quantity for the duration of a cycle of length T_i for ith item.
- $TAP_i(T_i, S_i)$: Total average profit per unit for the ith item.
- w_i : Storage space per unit time for the ith item.
- W: Total area of space.
- $\widetilde{c_{\iota}}$: Fuzzy ordering cost per order for the ith item.
- $\tilde{\theta}_i$: Fuzzy deterioration rate for the ith item.
- \widetilde{P}_i : Purchasing cost for the ith item in fuzziness.
- $\widetilde{w_i}$: Storage space per unit time for the ith item in fuzziness.
- $\widetilde{H}_i(=\widetilde{\alpha}_i t)$: Fuzzy holding cost per unit per unit time for the ith item
- \tilde{d}_i : Fuzzy deteriorating cost per unit per unit time for the ith item.
- $TA\widetilde{P_{l}(T_{l},S_{l})}$: Fuzzy total average profit per unit for the ith item.
- \hat{c}_i : Defuzzyfication of fuzzy ordering cost per order for the ith item.
- $\widehat{\theta}_i$: Defuzzy fication of fuzzy deterioration rate for the ith item.
- \widehat{P}_i : Defuzzyfication of fuzzy purchasing cost for the ith item.
- $\widehat{w_i}$: Defuzzy fication of fuzzy storage space per unit time for the ith item.
- $\widehat{H}_{i}(=\widehat{\alpha}_{i}t)$: Defuzzy fication of fuzzy holding cost per unit per unit time for the ith item
- \hat{d}_i : Defuzzyfication of fuzzy deteriorating cost per unit per unit time for the ith item.



http://www.ijmp.jor.br ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083

 $TAP_{i}(T_{i}S_{i})$: Defuzzy fication of fuzzy total average profit per unit for the ith item.

2.2. Assumptions

- 1. The inventory system involves multi-item.
- 2. The replenishment occurs instantaneously at infinite rate.
- 3. The lead time is negligible.
- 4. Shortages are not allowed.

5. Demand rate is time depended as well as selling price. It is taken as $D(S_i, t) = (a_i - b_i S_i)e^{\lambda_i t}$ where $a_i, b_i > 0$, $(a_i - b_i S_i) > 0$ and $S_i \ge P_i$.

6. Deterioration rate per unit time per cycle is θ_i for the ith item. $0 \le \theta_i \le 1$, θ_i is constant.

2.3. Model Formulation

The inventory model for ith item is illustrated in Figure-1. During the period $[0, T_i]$ the inventory level reduces due to demand rate and deterioration rate for ith item. In this time period, the inventory level is described by the differential equation-

$$\frac{dI_i(t)}{dt} + I_i(t)\theta_i = -D(S_i, t) = -(a_i - b_i S_i)e^{\lambda_i t}, \quad 0 \le t \le T_i$$

$$\tag{1}$$

With boundary condition, $I_i(0) = Q_i$, $I_i(T_i) = 0$.

Solving (1) we have,

$$I_{i}(t) = e^{-\theta_{i}t} \frac{(a_{i}-b_{i}S_{i})}{(\theta_{i}+\lambda_{i})} \left[e^{(\theta_{i}+\lambda_{i})T_{i}} - e^{(\theta_{i}+\lambda_{i})t} \right]$$
(2)

$$Q_i = \frac{(a_i - b_i S_i)}{(\theta_i + \lambda_i)} \left(e^{(\theta_i + \lambda_i)T_i} - 1 \right)$$
(3)







http://www.ijmp.jor.br ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083

Now we are calculating various costs for ith item as following

i) Sales revenue $(SR_i) = S_i \int_0^{T_i} D(S_i, t) dt$

$$=\frac{S_i(a_i-b_iS_i)}{\lambda_i}(e^{\lambda_iT_i}-1)$$

ii) Purchasing cost $(PC_i) = P_i Q_i = \frac{P_i(a_i - b_i S_i)}{(\theta_i + \lambda)} (e^{(\theta_i + \lambda)T_i} - 1)$

iii) Inventory holding cost $(HC_i) = \int_0^{T_i} \alpha_i t I_i(t) dt$

$$=\frac{\alpha_i(a_i-b_iS_i)}{(\theta_i+\lambda_i)}\left[-e^{(\theta_i+\lambda_i)T_i}\left\{\frac{T_ie^{-\theta_iT_i}}{\theta_i}+\frac{1}{\theta_i^2}\left(e^{-\theta_iT_i}-1\right)\right\}-\frac{T_ie^{\lambda T_i}}{\lambda_i}+\frac{1}{\lambda_i^2}\left(e^{\lambda_iT_i}-1\right)\right]$$

iv) Deterioration cost $(DC_i) = d_i \theta_i T_i$

v) Ordering cost $(OC_i) = c_i$

Total average profit per unit time for *ith* item

$$TAP_{i}(T_{i}, S_{i}) = \frac{1}{T_{i}} [SR_{i} - (PC_{i} + HC_{i} + DC_{i} + OC_{i})]$$

$$= \frac{S_{i}(a_{i}-b_{i}S_{i})}{\lambda_{i}T_{i}} (e^{\lambda_{i}T_{i}} - 1) - \frac{P_{i}(a_{i}-b_{i}S_{i})}{(\theta_{i}+\lambda_{i})T_{i}} (e^{(\theta_{i}+\lambda_{i})T_{i}} - 1) - \frac{\alpha_{i}(a_{i}-b_{i}S_{i})}{(\theta_{i}+\lambda_{i})T_{i}} \Big[e^{(\theta_{i}+\lambda_{i})T_{i}} \Big\{ \frac{T_{i}e^{-\theta_{i}T_{i}}}{\theta_{i}} + \frac{1}{\theta_{i}^{2}} (e^{-\theta_{i}T_{i}} - 1) \Big] - d_{i}\theta_{i} - \frac{c_{i}}{T_{i}}$$
(4)

Multi-objective inventory model (MOIM) can be written as:

 $\begin{aligned} & \max \left\{ TAP_{1}(T_{1},S_{1}), TAP_{2}(T_{2},S_{2}), \dots, TAP_{n}(T_{n},S_{n}) \right\} \\ & \text{Subject to, } \sum_{i=1}^{n} w_{i}Q_{i} \leq W, P_{i} \leq S_{i} \leq \frac{a_{i}}{b_{i}} \\ & \text{where } TAP_{i}(T_{i},S_{i}) = \frac{S_{i}(a_{i}-b_{i}S_{i})}{\lambda_{i}T_{i}} \left(e^{\lambda_{i}T_{i}} - 1 \right) - \frac{P_{i}(a_{i}-b_{i}S_{i})}{(\theta_{i}+\lambda_{i})T_{i}} \left(e^{(\theta_{i}+\lambda_{i})T_{i}} - 1 \right) - \frac{a_{i}(a_{i}-b_{i}S_{i})}{(\theta_{i}+\lambda_{i})T_{i}} \left[e^{(\theta_{i}+\lambda_{i})T_{i}} \left\{ \frac{T_{i}e^{-\theta_{i}T_{i}}}{\theta_{i}} + \frac{1}{\theta_{i}^{2}} \left(e^{-\theta_{i}T_{i}} - 1 \right) \right\} - \frac{T_{i}e^{\lambda T_{i}}}{\lambda_{i}} + \frac{1}{\lambda_{i}^{2}} \left(e^{\lambda_{i}T_{i}} - 1 \right) \right] - d_{i}\theta_{i} - \frac{c_{i}}{T_{i}} \end{aligned}$ $\tag{5}$

and
$$Q_i = \frac{(a_i - b_i S_i)}{(\theta_i + \lambda_i)} \left(e^{(\theta_i + \lambda_i)T_i} - 1 \right)$$
 for $i = 1, 2, \dots, n$ (6)

2.4. Fuzzy Model

Due to uncertainty, we assume all parameters as generalized trapezoidal fuzzy number (GTrFN). Let us assume,



http://www.ijmp.jor.br ISSN: 2236-269X

DOI: 10.14807/ijmp.v11i3.1083

$$\begin{split} \widetilde{a_{i}} &= \left(a_{i}^{1}, a_{i}^{2}, a_{i}^{3}, a_{i}^{4}; w_{a_{i}}\right), 0 < w_{a_{i}} \leq 1; \ \widetilde{b_{i}} = \left(b_{i}^{1}, b_{i}^{2}, b_{i}^{3}, b_{i}^{4}; w_{b_{i}}\right), 0 < w_{b_{i}} \leq 1; \\ \widetilde{\theta_{i}} &= \left(\theta_{i}^{1}, \theta_{i}^{2}, \theta_{i}^{3}, \theta_{i}^{4}; w_{\theta_{i}}\right), 0 < w_{\theta_{i}} \leq 1; \ \widetilde{\alpha_{i}} = \left(\alpha_{i}^{1}, \alpha_{i}^{2}, \alpha_{i}^{3}, \alpha_{i}^{4}; w_{\alpha_{i}}\right), 0 < w_{\alpha_{i}} \leq 1; \\ \widetilde{d_{i}} &= \left(d_{i}^{1}, d_{i}^{2}, d_{i}^{3}, d_{i}^{4}; w_{d_{i}}\right), 0 < w_{d_{i}} \leq 1; \ \widetilde{P_{i}} = \left(P_{i}^{1}, P_{i}^{2}, P_{i}^{3}, P_{i}^{4}; w_{P_{i}}\right), 0 < w_{P_{i}} \leq 1; \\ \widetilde{w_{i}} &= \left(w_{i}^{1}, w_{i}^{2}, w_{i}^{3}, w_{i}^{4}; w_{w_{i}}\right), 0 < w_{w_{i}} \leq 1; \ \widetilde{c_{i}} = \left(c_{i}^{1}, c_{i}^{2}, c_{i}^{3}, c_{i}^{4}; w_{c_{i}}\right), 0 < w_{c_{i}} \leq 1 \end{split}$$

Then due to uncertainty, the total average profit for the ith item is given by,

$$TA\widetilde{P_{l}(T_{l}, S_{l})} = \frac{\widetilde{P_{l}(\tilde{a}_{l} - \tilde{b}_{l}S_{l})}}{(\widetilde{\theta}_{l} + \lambda_{i})T_{i}} \left(1 - e^{(\widetilde{\theta}_{l} + \lambda_{i})T_{i}}\right) + \frac{S_{i}(\widetilde{a}_{l} - \widetilde{b}_{l}S_{l})}{\lambda_{i}T_{i}} \left(e^{\lambda_{i}T_{i}} - 1\right) - \frac{\widetilde{\alpha}_{i}(\widetilde{a}_{i} - \widetilde{b}_{i}S_{l})}{(\widetilde{\theta}_{i} + \lambda_{i})T_{i}} \left[e^{(\widetilde{\theta}_{i} + \lambda_{i})T_{i}} \left\{\frac{T_{i}e^{-\widetilde{\theta}_{i}T_{i}}}{\widetilde{\theta}_{i}} + \frac{1}{\widetilde{\theta}_{i}^{2}} \left(e^{-\widetilde{\theta}_{i}T_{i}} - 1\right)\right\} - \frac{T_{i}e^{\lambda_{i}T_{i}}}{\lambda_{i}} + \frac{1}{\lambda_{i}^{2}} \left(e^{\lambda_{i}T_{i}} - 1\right)\right] - \widetilde{d}_{i}\widetilde{\theta}_{i} - \frac{\widetilde{c}_{i}}{T_{i}}(7)$$

and $Q_{i} = \frac{(\widetilde{a}_{i} - \widetilde{b}_{i}S_{i})}{(\widetilde{\theta}_{i} + \lambda_{i})} \left(e^{(\widetilde{\theta}_{i} + \lambda_{i})T_{i}} - 1\right)$ for $i = 1, 2, ..., n$.

And our MOIM problem becomes fuzzy model as

$$\begin{aligned} &\operatorname{Max} \left\{ TA\widetilde{P_{1}(T_{1},S_{1})}, TA\widetilde{P_{2}(T_{2},S_{2})}, TA\widetilde{P_{3}(T_{3},S_{3})}, \dots, TA\widetilde{P_{n}(T_{n},S_{n})} \right\} \\ &\operatorname{Subject to}, \sum_{i=1}^{n} \widetilde{w_{i}} Q_{i} \leq W, \widetilde{P_{i}} \leq S_{i} \leq \frac{\widetilde{a_{i}}}{\widetilde{b_{i}}} \\ &\operatorname{where} TA\widetilde{P_{i}(T_{i},S_{i})} = \frac{\widetilde{P_{i}(\widetilde{a}_{i}-\widetilde{b_{i}}S_{i})}}{(\widetilde{\theta_{i}}+\lambda_{i})T_{i}} \left(1-e^{(\widetilde{\theta_{i}}+\lambda_{i})T_{i}}\right) + \frac{S_{i}(\widetilde{a}_{i}-\widetilde{b_{i}}S_{i})}{\lambda_{i}T_{i}} \left(e^{\lambda_{i}T_{i}}-1\right) - \frac{\widetilde{\alpha_{i}(\widetilde{a}_{i}-\widetilde{b_{i}}S_{i})}}{(\widetilde{\theta_{i}}+\lambda_{i})T_{i}} \left[e^{(\widetilde{\theta_{i}}+\lambda_{i})T_{i}} \left\{\frac{T_{i}e^{-\widetilde{\theta_{i}}T_{i}}}{\widetilde{\theta_{i}}} + \frac{1}{\widetilde{\theta_{i}}^{2}} \left(e^{-\widetilde{\theta_{i}}T_{i}}-1\right)\right\} - \frac{T_{i}e^{\lambda_{i}T_{i}}}{\lambda_{i}} + \frac{1}{\lambda_{i}^{2}} \left(e^{\lambda_{i}T_{i}}-1\right) \right] - \widetilde{d_{i}}\widetilde{\theta_{i}} - \frac{\widetilde{c_{i}}}{T_{i}} \\ &\operatorname{and} \quad Q_{i} = \frac{(\widetilde{a}_{i}-\widetilde{b_{i}}S_{i})}{(\widetilde{\theta_{i}}+\lambda_{i})} \left(e^{(\widetilde{\theta_{i}}+\lambda_{i})T_{i}}-1\right) \operatorname{for} i = 1, 2, \dots, n. \end{aligned}$$

3. MATHEMATICAL ANALYSIS

3.1. Ranking fuzzy numbers with respect to their total integral value.

Ranking fuzzy number is an important aspect of decision making in a fuzzy environment. In reality, decision-makers having different viewpoints will give different ranking outcomes under the same situation. Bortolan and Degani (1985) studied the number of methods for ranking fuzzy numbers. Liou and Wang (1992) proposed a relatively simple computation and easily understood method.

Let $\lambda \in [0,1]$ be a pre-assigned parameter which is called the degree of optimism. The graded mean value (or, total λ -integer value) of generalized trapezoidal fuzzy number (GTrFN) is discussed below.



934

v. 11, n. 3, May-June 2020

ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083

http://www.iimp.ior.br

A fuzzy number $\tilde{A} = (a, b, c, d; w)$ is called generalized trapezoidal fuzzy number (GTrFN) if its membership function is given by

$$\mu_{\tilde{A}}^{w}(x) = \begin{cases} 0 & \text{if } x \leq a \\ w \frac{x-a}{b-a} & \text{if } a \leq x \leq b \\ w & \text{if } b \leq x \leq c \\ w \frac{d-x}{d-c} & \text{if } c \leq x \leq d \\ 0 & \text{if } d \leq x \end{cases}$$

The total λ integer value of $\tilde{A} = (a, b, c, d; w)$ is given as

 $I_{\lambda}^{w}(\tilde{A}) = \lambda I_{R}^{w}(\tilde{A}) + (1-\lambda)I_{L}^{w}(\tilde{A}), \text{ where } I_{L}^{w}(\tilde{A}), I_{R}^{w}(\tilde{A}) \text{ are the left and right interval values}$ of \tilde{A} respectively. Where $I_{L}^{w}(\tilde{A}) = \int_{0}^{1} (\mu_{L\tilde{A}}^{w})^{-1} \alpha \ d\alpha, \ I_{R}^{w}(\tilde{A}) = \int_{0}^{1} (\mu_{R\tilde{A}}^{w})^{-1} \alpha \ d\alpha$ Now, $(\mu_{L\tilde{A}}^{w})^{-1} \alpha = a + \frac{\alpha}{w} (b-a) \& (\mu_{R\tilde{A}}^{w})^{-1} \alpha = d - \frac{\alpha}{w} (d-c)$

So the left & right interval values \widetilde{A} are

$$I_L^w(\tilde{A}) = w\left(\frac{a+b}{2}\right) \text{ and } I_R^w(\tilde{A}) = w\left(\frac{c+d}{2}\right)$$

Hence the total λ - integral value of \tilde{A} is

 $I_{\lambda}^{w}(\tilde{A}) = \lambda w(\frac{c+d}{2}) + (1-\lambda)w(\frac{a+b}{2})$

When $\lambda = 1$, we obtain $I_1^w(\tilde{A}) = I_R^w(\tilde{A}) = w\left(\frac{c+d}{2}\right)$ which reflects an optimistic viewpoint. When $\lambda = 0$, we get $I_0^w(\tilde{A}) = I_L^w(\tilde{A}) = w\left(\frac{a+b}{2}\right)$ which reflects a pessimistic point of view. When $\lambda = 0.5$ the total λ -integral value is

$$I_{0.5}^{w}(\tilde{A}) = \frac{1}{2} [I_{L}^{w}(\tilde{A}) + I_{R}^{w}(\tilde{A})] = \frac{1}{2} \left[w\left(\frac{a+b}{2}\right) + w\left(\frac{c+d}{2}\right) \right] = w\left(\frac{a+b+c+d}{4}\right).$$
(9)

Which reflects a reasonably optimistic decision-makers perspective and is treated as the defuzzification of \tilde{A} .

3.2. Fuzzy Model Using Defuzzification of Fuzzy Number.

Using defuzzification of fuzzy number technique (9), we have the approximated values $(\hat{c}_{l}, \hat{a}_{l}, \hat{b}_{l}, \hat{\theta}_{l}, \hat{\alpha}_{l}, \hat{d}_{l}, \hat{P}_{l}, \widehat{w}_{l})$ of the GTrFN parameters $(\tilde{c}_{l}, \tilde{a}_{l}, \tilde{b}_{l}, \tilde{\theta}_{l}, \tilde{a}_{l}, \tilde{d}_{l}, \tilde{P}_{l}, \widetilde{w}_{l})$.

So the above problem (8) reduces to

 $\operatorname{Max}\left\{TA\widehat{P_1(T_1,S_1)}, TA\widehat{P_2(T_2,S_2)}, TA\widehat{P_3(T_3,S_3)}, \dots, \dots, TA\widehat{P_n(T_n,S_n)}\right\}$





http://www.ijmp.jor.br ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083 v. 11, n. 3, May-June 2020

Subject to, $\sum_{i=1}^{n} \widehat{w_i} Q_i \leq W, \widehat{P}_i \leq S_i \leq \frac{\widehat{a_i}}{\widehat{h_i}}$

where
$$TA\widehat{P_{l}(T_{l}, S_{l})} = \frac{\widehat{P_{l}(\hat{a}_{l} - \widehat{b}_{l}S_{l})}{(\widehat{\theta}_{l} + \lambda_{i})T_{i}} \left(1 - e^{(\widehat{\theta}_{l} + \lambda_{i})T_{i}}\right) + \frac{S_{i}(\widehat{a}_{i} - \widehat{b}_{l}S_{i})}{\lambda_{i}T_{i}} \left(e^{\lambda_{i}T_{i}} - 1\right) - \frac{\widehat{a_{i}}(\widehat{a}_{i} - \widehat{b}_{l}S_{i})}{(\widehat{\theta}_{i} + \lambda_{i})T_{i}} \left[e^{(\widehat{\theta}_{i} + \lambda_{i})T_{i}} \left\{\frac{T_{i}e^{-\widehat{\theta}_{i}T_{i}}}{\widehat{\theta}_{i}} + \frac{1}{\widehat{\theta}_{l}^{2}}\left(e^{-\widehat{\theta}_{l}T_{i}} - 1\right)\right\} - \frac{T_{i}e^{\lambda_{i}T_{i}}}{\lambda_{i}} + \frac{1}{\lambda_{i}^{2}}\left(e^{\lambda_{i}T_{i}} - 1\right)\right] - \widehat{d}_{i}\widehat{\theta}_{l} - \frac{\widehat{c}_{l}}{T_{i}}$$

and $Q_{i} = \frac{(\widehat{a}_{i} - \widehat{b}_{i}S_{i})}{(\widehat{\theta}_{i} + \lambda_{i})} \left(e^{(\widehat{\theta}_{i} + \lambda_{i})T_{i}} - 1\right)$ for $i = 1, 2, ..., n.$ (10)

4. FUZZY PROGRAMMING TECHNIQUE (BASED ON MAX-MIN AND MAX-ADDITIVE OPERATORS) TO SOLVE MOIM

Solve the MOIM (10) as a single objective using only one objective at a time and we ignoring the others. So we get the ideal solutions.

From the above results, we find out the corresponding values of every objective function at each solution obtained. With these values the pay-off matrix can be prepared as follows:

Let $L^k = \min\{TAP_k(T_i^i, S_i^i), i = 1, 2, ..., n\}$ for k = 1, 2, ..., n and

 $U^{k} = TAP_{k}^{*}(T_{k}^{k}, S_{k}^{k}) for k = 1, 2, ..., n.$

Hence U^k, L^k are identified, $L^k \leq TAP_k(T_i^i, S_i^i) \leq U^k$, for i = 1, 2, ..., n; k = 1, 2, ..., n.

Therefore fuzzy linear membership functions $\mu_{TAP_k}(TAP_k(T_k, S_k))$ for the kth objective functions $TAP_k(T_k, S_k)$ respectively for k = 1, 2, ..., n are defined as follows:

$$\mu_{TAP_k}(TAP_k(T_k, S_k)) = \begin{cases} 0 & for \, TAP_k(T_k, S_k) \le L^k \\ \frac{TAP_k(T_k, S_k) - L^k}{U^k - L^k} & for \, L^k \le TAP_k(T_k, S_k) \le U^k \\ 1 & for \, TAP_k(T_k, S_k) \ge U^k \end{cases}$$



INDEPENDENT JOURNAL OF MANAGEMENT & PRODUCTION (IJM&P) http://www.ijmp.jor.br v. 11, n. 3, May-June 2020 ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083

for k = 1, 2, ..., n.

(11)

4.1. 4.1 Fuzzy non-linear programming problem (FNLP) method based on max-min operator

Fuzzy non-linear programming problem (FNLP) is formulated as follows

Max α

Subject to

$$TAP_{k}(T_{k}, S_{k}) - \alpha (U^{k} - L^{k}) \le L^{k}, for \ k = 1, 2, ..., n.$$
(12)
$$0 \le \alpha \le 1,$$

and same constraints and restrictions as the problem (10).

4.2. Fuzzy additive goal programming problem (FAGP) method based on maxadditive operator

Fuzzy additive goal programming problem (FAGP) is formulated as follows

Max
$$\sum_{k=1}^{n} \frac{TAP_k(T_k, S_k) - L^k}{U^k - L^k}$$

Subject to

$$0 \le \mu_{TAP_k} (TAP_k(T_k, S_k)) \le 1, for \ k = 1, 2, \dots, n.$$
(13)

and same constraints and restrictions as the problem (10).

The non-linear programming problems (12) and (13) can be solved by suitable mathematical programming algorithm.

5. WEIGHTED FUZZY PROGRAMMING TECHNIQUE (BASED ON MAX-MIN AND MAX-ADDITIVE OPERATORS) TO SOLVE MOIM

Let us consider positive weights ω_k for each objective $TAP_k(T_k, S_k)$, k =

1,2, ..., *n* where $\sum_{k=1}^{n} \omega_k = 1$.

Using the above membership functions (11) and weights ω_k (k = 1, 2, ..., n), we formulate the following programming problems.

5.1. Weighted fuzzy non-linear programming problem (WFNLP) method based on max-min operator

Weighted fuzzy non-linear programming problem (WFNLP) is formulated as follows



http://www.ijmp.jor.br ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083 v. 11, n. 3, May-June 2020

Max α

Subject to,

$$\omega_k \mu_{TAP_k} (TAP_k(T_k, S_k)) \ge \alpha , for \ k = 1, 2, \dots, n$$
(14)

 $0 \le \alpha \le 1$,

$$\sum_{k=1}^{n} \omega_k = 1$$

and same constraints and restrictions as the problem (10)

5.2. Weighted Fuzzy additive goal programming problem (WFAGP) method based on max-additive operator

Weighted fuzzy additive goal programming problem (WFAGP) is formulated as follows

 $\operatorname{Max} \sum_{k=1}^{n} \omega_k \mu_{TAP_k} (TAP_k(T_k, S_k))$

Subject to,

$$0 \le \mu_{TAP_k} (TAP_k(T_k, S_k)) \le 1, for \ k = 1, 2, \dots, n.$$
(15)

$$\sum_{k=1}^{n} \omega_k = 1$$

and same constraints and restrictions as the problem (10)

The non-linear programming problems (14) and (15) can be solved by suitable mathematical programming algorithm.

6. Numerical Example

Let us consider an inventory model which consist two items with following parameter values in proper units. Also consider the total storage area is W = 600 square unit and $\lambda_1 = 0.01$, $\lambda_2 = 0.02$.

Item I		Item II		
Parametric Value in Fuzzy Number	Defuzzification	Parametric Value in Fuzzy Number	Defuzzification	
	of fuzzy number		of fuzzy number	
$\begin{split} \widetilde{c_1} &= (400, 450, 500, 550; 0.9) \\ \widetilde{a_1} &= (550, 600, 650, 700; 0.96) \\ \widetilde{b_1} &= (2.5, 3, 3.5, 4; 0.92) \\ \widetilde{a_1} &= (2, 3, 4, 5; 0.7) \\ \widetilde{d_1} &= (38, 40, 42, 44; 0.8) \\ \widetilde{\theta_1} &= (0.01, 0.02, 0.03, 0.04; 0.8) \\ \widetilde{P_1} &= (40, 42, 44, 46; 0.9) \end{split}$	$ \hat{c}_1 = 427.5 \hat{a}_1 = 600 \hat{b}_1 = 2.99 \hat{a}_1 = 2.45 \hat{d}_1 = 32.8 \hat{\theta}_1 = 0.02 \hat{P}_1 = 38.7 $	$\begin{split} \widetilde{c_2} &= (450,500;550,600;0.8)\\ \widetilde{a_2} &= (560,610,660,710;0.96)\\ \widetilde{b_2} &= (3,3.5,4,4.5;0.93)\\ \widetilde{a_2} &= (1,3,4,5;0.8)\\ \widetilde{d_2} &= (40,41,42,43;0.8)\\ \widetilde{\theta_2} &= (0.01,0.02,0.03,0.04;0.8))\\ \widetilde{P_2} &= (41,42,43,44;0.9) \end{split}$		

Table 1: Input imprecise data for shape parameters



http://www.ijmp.jor.br ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083 v. 11, n. 3, May-June 2020

$\widetilde{w_1} = (4,5,$	6,7; 0.91)	$\widehat{w_1} = 5.005$	$\widetilde{w_2} = (3,4,5,6;0.89)$	$\widehat{w_2} = 4.005$
	Т	Table 2: Optimal solutions of	MOIM (10)	
	Methods	$TAP_1(T_1^*, S_1^*)$	$TAP_2(T_2^*, S_2^*)$	
	FNLP	18085.52	14482.83	
	FAGP	18081.19	14487.72]

The above Table 2 shows that both FNLP and FAGP methods give almost same results.

The optimal solutions of the MOIM by WFNLP method with different weights are shown in table 3.

Table 3: Optimal solutions of MOIM (10) with different weights by WFNLP method

Weights	$TAP_{1}(T_{1}^{*}, S_{1}^{*})$	$TAP_2(T_2^*, S_2^*)$
$(\omega_1 = 0.5, \omega_2 = 0.5)$	18085.53	14482.84
$(\omega_1 = 0.8, \omega_2 = 0.2)$	17794.61	14640.87
$(\omega_1 = 0.2, \omega_2 = 0.8)$	18214.18	14149.66





Figure 2: Profit of 1st item with respect to the different weights (WFNLP method)

Figure 3: Profit of 2nd item with respect to the different weights (WFNLP method)

Table 4: Optimal solutions of MOIM (10) with different weights by WFAGP method

Weights	$TAP_{1}(T_{1}^{*}, S_{1}^{*})$	$TAP_{2}(T_{2}^{*}, S_{2}^{*})$
$(\omega_1 = 0.5, \omega_2 = 0.5)$	18089.58	14497.67
$(\omega_1 = 0.8, \omega_2 = 0.2)$	18175.47	14205.92
$(\omega_1 = 0.2, \omega_2 = 0.8)$	17846.26	14603.66



Figure 4: Profit of 1st item with respect to the different weights (WFAGP method)



Figure 5: Profit of 2nd item with respect to the different weights (WFAGP method)



http://www.ijmp.jor.br v. 11, n. 3, May-June 2020 ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083

From the above table 3 and table 4 shows that total profit for 1st and 2nd items in all types are almost same.

7. SENSITIVITY ANALYSIS

In the sensitivity analysis the MOIM (10) is solved by using FNLP and FAGP methods for different values of θ , *a*, *b*, λ are given in Table 5, 6,7 and 8 respectively.

Table 5: Optimal so	lutions of MOIM by	y FNLP & FAGP	methods for	different values of θ

Methods	θ	$TAP_{1}(T_{1}^{*}, S_{1}^{*})$	$TAP_{2}(T_{2}^{*}, S_{2}^{*})$
	0.02	18085.52	14482.83
	0.04	18062.70	14466.73
FNLP	0.06	18009.48	14399.27
	0.08	17955.12	14257.89
	0.02	18081.19	14487.72
	0.04	18060.12	14474.77
FAGP	0.06	18043.72	14448.96
	0.08	18006.95	14410.01







Figure 7: Sensitivity analysis for profit of 2^{nd} item w.r.t. θ by FNLP & FAGP methods

From the above figures 6 & 7 shows that profit of the both items is decreased when θ

is increased in both methods.

Table 6: Optimal solutions of MOIM (10) by FNLP & FAGP methods for different values of

a_1, a_2 .						
Methods	<i>a</i> ₁ , <i>a</i> ₂	$TAP_1(T_1^*, S_1^*)$	$TAP_{2}(T_{2}^{*}, S_{2}^{*})$			
	$a_1 = 600,$	18085.52	14482.83			
	$a_2 = 609.6$					
	$a_1 = 650$,	22341.45	18069.17			
	$a_2 = 659.6$					
FNLP	$a_1 = 700$,	22987.17	21854.66			
	$a_2 = 709.6$					
	$a_1 = 750$,	23513.51	25998.57			
	$a_2 = 759.6$					
	$a_1 = 600$,	18081.19	14487.72			
	$a_2 = 609.6$					
	$a_1 = 650$,	22371.25	18129.17			
	$a_2 = 659.6$					
FAGP	$a_1 = 700$,	22967.80	21865.45			
	$a_2 = 709.6$					



http://www.ijmp.jor.br ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083

 $\begin{array}{c|c} a_1 = 750, \\ a_2 = 759.6 \end{array}$ 23589.23 25984.34





v. 11, n. 3, May-June 2020

Figure 8: Sensitivity analysis for profit of 1^{st} item w.r.t. a_1, a_2 by FNLP & FAGP methods

Figure 9: Sensitivity analysis for profit of 2^{nd} item w.r.t. a_1 , a_2 by FNLP & FAGP methods

From the above figures 8 & 9 shows that profit of the both items is increased when values of a_1, a_2 are increased in all methods.

Table 7: Optimal solutions of MOIM (10) b	ŊУ	FNLP	& FAGP	methods	for	different	values	of
	1.	1.						

$b_1, b_2.$						
Methods	<i>b</i> ₁ , <i>b</i> ₂	$TAP_1(T_1^*, S_1^*)$	$TAP_{2}(T_{2}^{*}, S_{2}^{*})$			
	$b_1 = 2.99$	18085.02	14482.64			
	$b_2 = 3.49$					
	$b_1 = 3.09$	17300.56	13934.35			
ENIL D	$b_2 = 3.59$					
FNLP	$b_1 = 3.19$	16432.94	13285.35			
	$b_2 = 3.69$					
	$b_1 = 3.29$	15621.16	12673.22			
	$b_2 = 3.79$					
FAGP	$b_1 = 2.99$	18081.15	14487.65			
	$b_2 = 3.49$					
	$b_1 = 3.09$	17312.43	13954.35			
	$b_2 = 3.59$					
	$b_1 = 3.19$	16467.87	13285.47			
	$b_2 = 3.69$					
	$b_1 = 3.29$	15643.45	12689.28			
	$b_2 = 3.79$					



Figure 10: Sensitivity analysis for profit of 1^{st} item w.r.t. b_1 , b_2 by FNLP & FAGP methods



Figure 11: Sensitivity analysis for profit of 2^{nd} item w.r.t. b_1, b_2 by FNLP & FAGP methods



v. 11, n. 3, May-June 2020

http://www.ijmp.jor.br ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083

From the above figures 10 & 11 shows that profit of the both items is decreased when

 b_1, b_2 are increased in all methods.



Figure 12: Sensitivity analysis for profit of 1st item w.r.t. λ_1 , λ_2 by FNLP & FAGP methods Figure 13: Sensitivity analysis for profit of2nd item w.r.t. λ_1 , λ_2 by FNLP & FAGP methods

From the above figures 12 & 13 shows that profit of the both items is increased when λ_1, λ_2 are increased in all methods.

8. CONCLUSIONS

In this paper, we have developed a real life inventory model in which time dependent holding cost, selling price as well as time dependent demand. Multi-item inventory has been considered under limitation on storage space. First crisp model is formed then it transferred to the fuzzy model due to uncertainty of cost parameters. All fuzzy cost parameters are taken as generalized trapezoidal fuzzy number. The proposed multi-objective inventory model is solved by using FNLP, FAGP, WFNLP and WFAGP methods.

In the future study, it is hoped to further incorporate the proposed model into more realistic assumption, such as probabilistic demand, introduce shortages etc. Also other type of



http://www.ijmp.jor.br ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083

membership functions like as triangular fuzzy number, Parabolic flat Fuzzy Number (*PfFN*), Parabolic Fuzzy Number (*pFN*) etc. can be used to form the fuzzy model.

9. ACKNOWLEDGEMENTS:

The authors are thankful to University of Kalyani for providing financial assistance through DST-PURSE (Phase-II) Programme. The authors are grateful to the reviewers for their comments and suggestions.

REFERENCES:

ALFARES, H. K.; GHAITHAN, A. M. (2016) Inventory and Pricing Model with Price-Dependent Demand, Time-Varying Holding Cost, and Quantity Discounts, **Computers & Industrial Engineering**, doi: http://dx.doi.org/10.1016/j.cie.2016.02.009

BHUNIA, A. K.; SHAIKH, A. A. (2014) A deterministic inventory model for deteriorating items with selling price dependent demand and three-parameter Weibull distributed deterioration, **International Journal of Industrial Engineering Computations**, n. 5, p. 497–510

BORTOLAN, G.; DEGANI R. (1985) A review of some methods for ranking fuzzy numbers, **Fuzzy Sets and Systems**, n. 15, p. 1–19.

COVERT, R. P.; PHILIP, G. C. (1973) An EOQ model for items with Weibull distribution deterioration. **AIIE Transaction**, n. 5, p. 323–326.

GHARE, P. M.; SCHRADER, G. H. (1963) A model for exponentially decaying inventory system. **International Journal of Production Research**, n. 21, p. 449–460.

GHOSH, S. K.; SARKAR, T.; CHAUDHURI, K. (2015) A Multi-Item Inventory Model for Deteriorating Items in Limited Storage Space with Stock-Dependent Demand, American Journal of Mathematical and Management Sciences, v. 34, n. 2, p. 147-161, DOI: 10.1080/01966324.2014.980870.

HARRI, F. W. (1913) How many parts to make at once factory, **Mag. Mannage**., n. 10, p.135-136.

ISLAMM, S.; ROY, T. K. (2006) A fuzzy EPQ model with flexibility and reliability consideration and depended unit production cost under a space constraint: A fuzzy geometric programming approach, **Applied Mathematics and Computation**, v. 176, n. 2, p. 531-544.

ISLAMM, S.; MANDAL, W. A. (2017) A Fuzzy Inventory Model (EOQ Model) with Unit Production Cost, Time Depended Holding Cost, Without Shortages Under a Space Constrain: A Fuzzy Parametric Geometric Programming (FPGP) Approach, **Independent Journal of Management Production**, v. 8, n. 2, p. 299–318. DOI: dx.doi.org/10.14807/ijmp.v8i2.535.

ISLAMM, S.; MONDAL, W. A. (2017) Fuzzy E.O.Q Model with Constant Demand and Shortages: A Fuzzy Signomial Geometric Programming (FSGP) Apprach, **Independent Journal Of Management & Production**, v. 8, n. 4, p. 1191-1209.

KUMAR, S.; SINGH, A. K.; PATEL, M. K. (2016) Optimization of Weibull deteriorating items inventory model under the effect of price and time dependent demand with partial backlogging, **Sadhana**, v. 41, DOI 10.1007/s12046-016-0533-4



http://www.ijmp.jor.br ISSN: 2236-269X DOI: 10.14807/ijmp.v11i3.1083 v. 11, n. 3, May-June 2020

LIANG, Y.; ZHOU, F. (2011) A two warehouse inventory model for deteriorating items under conditionally permissible delay in Payment, **Appl. Math. Model**, n. 35, p. 2221-2231.

LIOU T. S.; WANG, M. J. (1992) Ranking fuzzy numbers with integral value, **Fuzzy Sets** and **Systems**, n. 50, p. 247–255.

MAITY, M. K. (2008) Fuzzy inventory model with two ware house under possibility measure in fuzzy goal, **Euro. J Oper. Res**, n. 188, p. 746-774.

MOMDAL, B.; BHUNIA, A. K.; MAITI, M. (2003) An inventory system of ameliorating items for price dependent demand rate. **Computers and Industrial Engineering**, n. 45, p. 443–456.

ROY, A. (2014) Fuzzy inventory model for deteriorating items with price dependent demand, **International Journal of Management Science and Engineering Management**, DOI: 10.1080/17509653.2014.959086.

ROY, T. K.; MAITY, M. (1995) A fuzzy inventory model with constraints, **Operation** research, v. 32, n. 4, p.287-298.

SHAH, N. H.; SHAH, B. J.; WEE, H. M. (2009) A lot size inventory model for the Weibull distributed deterioration rate with discounted selling price and stock-dependent demand, **Int. J. Data Analysis Techniques and Strategies**, v. 1, n. 4, p. 355–363.

SRIDEVI, G.; NIRUPAMA DEVI, K.; SRINIVASA RAO, K. (2010) Inventory model for deteriorating items with Weibull rate of replenishment and selling price dependent demand, **Int. J. Operational Research**, v. 9, n. 3, p. 329–349.

YANG, H. L. (2016) Two Warehouse Partial Backlogging inventory model for deteriorating items under inflation, **International Journal of Production Economics**, n. 103, p. 362-370.

YU-CHUNG TSAO, GWO-JI-SHEEN (2008) Dynamic pricing promotion and replenishment policies for a deteriorating item under permissible delay in payments, **Computer & operation research**, n. 35, p. 3562-3580.

ZADEH, L. A. (1965) Fuzzy sets, Information and Control, n. 8, p. 338-353.

ZIMMERMANN, H. J. (1985) Application of fuzzy set theory to mathematical programming, **Information Science**, n. 36, p. 29-58.

ZIMMERMANN, H. J. (1992) **Methods and applications of Fuzzy Mathematical programming**, in An introduction to Fuzzy Logic Application in Intelligent Systems, p. 97-120, Kluwer publishers, Boston.

