



This document is a postprint version of an article published in Diseases Of Aquatic Organisms © Inter-Research after peer review. To access the final edited and published work see <https://doi.org/10.3354/dao03388>

Document downloaded from:



1 Fatal *Photobacterium damselae*-induced enteritis in a leatherback turtle (*Dermochelys*
2 *coriacea*)

3 Arturo Oliver-Guimerá¹, María Lourdes Abarca^{2,3}, María Cuvertoret-Sanz⁴, Mariano
4 Domingo^{2,4,5*}

5 ¹Department of Pathology, Microbiology and Immunology, School of Veterinary
6 Medicine, University of California-Davis, Davis, California 95616, USA

7 ²Departament de Sanitat i Anatomia Animals, Universitat Autònoma de Barcelona,
8 08193 Bellaterra, 08193, Spain.

9 ³Veterinary Mycology Group

10 ⁴Servei de Diagnòstic de Patologia Veterinària, Departament de Sanitat i Anatomia
11 Animals, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain.

12 ⁵Centre de Recerca en Sanitat Animal (CRESA), IRTA, Campus de la Universitat
13 Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain.

14
15 Running page head: PDD enteritis in leatherback turtle

16 ABSTRACT: Stranded leatherback turtle (*Dermochelys coriacea*) complete pathology
17 reports are rare and the cause of mortality is difficult to determine in many cases. We
18 conducted a complete pathological study of a stranded leatherback turtle from the
19 western Mediterranean. The main finding was a fibrino-necrotizing enteritis with
20 associated bacteria which were identified as *Photobacterium damselae* subsp. *damselae*
21 according to their biochemical and phenotypical characteristics. This report provides
22 evidence of the pathogenic effect of this bacterium in wild sea turtles.

- 1 KEY WORDS: *Photobacterium damsela*, *Dermochelys coriacea*, necrotizing enteritis,
- 2 intestinal bacteria, Vibrionaceae, stranded animal.

1. INTRODUCTION

Leatherback turtle (*Dermochelys coriacea*, Vandelli, 1761) is a worldwide distributed sea turtle species that can be seen occasionally in the Mediterranean Sea. Despite their occurrence being much lower than that of loggerhead turtle (*Caretta caretta*) and green turtle (*Chelonia mydas*) (Gómez de Segura et al. 2006), records of this species are not infrequent. Over a period of 20 years, there were 411 reported leatherback turtle strandings distributed throughout the whole Mediterranean Sea. Incidental capture by commercial fisheries seems to be the main threat for this species, followed by other anthropogenic lesions, such as head and carapace injuries, due to impacts with boats or propellers (Casale et al. 2003). There is little information about other causes of mortality in leatherback turtles and very few complete pathologic investigations have been carried out. Occasionally, *Photobacterium damsela* has been reported in stranded leatherback turtles from the Mediterranean (Poppi et al. 2012) as well as from other parts of the world (Obendorf et al. 1987). In this report, we present a pathological study of a stranded leatherback turtle from the western Mediterranean (Catalonian coast), with a severe necrotizing enteritis associated to *P. damsela*, showing that this bacterium may be a relevant pathogen for this species.

2. MATERIALS AND METHODS

On November 5th, 2016, an adult female leatherback turtle was found dead on the beach of Montroig (41°02'36"N 1°00'07"E), at the Catalonian Coast. The turtle was slightly autolyzed, with organs intact, and then given a D3 scoring for preservation (Flint et al. 2009). It was transported to the Veterinary School of the Autonomous University of Barcelona, where it was necropsied. Representative tissue samples were collected and preserved in 10% neutral buffered formalin for histopathological examination. After

1 48 hours of fixation, tissues were embedded in paraffin and routinely stained with
2 hematoxylin and eosin. Additional stains (GRAM and Grocott) were performed to detect
3 infectious agents in lesions. Microphotographs were taken with a Leica DM 6000 B
4 Photomicroscope and a Leica camera DFC480, that added a scale bar automatically.
5 Samples from intestinal mucosal were submitted for microbiological studies. Scrapped
6 intestinal mucosa was inoculated onto Columbia agar with 5% sheep blood (Difco) and
7 MacConkey agar (Oxoid) and incubated overnight at 37 °C in 5% CO₂. Investigation for
8 *Salmonella* spp. was also performed following ISO guideline 6579-1:2017 (ISO 2017),
9 with pre-enrichment culture of the sample in buffered peptone water (Oxoid) followed
10 by selective enrichment in modified semi-solid Rappaport Vassiliadis medium (Oxoid).
11 *Salmonella-Shigella* agar (Oxoid) and Xylose lysine deoxycholate agar (Oxoid) were used
12 as solid selective media. Isolates were phenotypically identified using the API 20E and
13 API20NE identification system (bioMérieux).

14

15 **3. RESULTS**

16 The turtle's curved carapace length was 126 cm and its curved carapace width
17 was 91 cm, and its weight was 156 Kg. Upon external examination, a few small barnacles
18 of the species *Chelonibia testudinaria* were detected (Epibiont Research Cooperative
19 2007). A large subcutaneous hemorrhagic and edematous area of about 15 x 10 cm was
20 observed on the left flank of the animal, approximately 20 cm cranial to the hind limb.
21 There were no other remarkable external traumatic lesions.

22 Approximately 3.5 liters of unclotted, dense, serosanguinous fluid were collected
23 from the coelomic cavity, and two or three small filamentous blood clots were seen on
24 the liver surface. There was a small amount of sand inside the larynx, trachea, and main

1 bronchi, apparently without causing occlusion (Fig.1A). The stomach contained a plastic
2 refreshment straw and two short fragments (3-5 cm long) of thin fishing line, without
3 apparent damage to the gastric mucosa. The serosal surface of small intestine was
4 congested, and intestinal loops were turgid and slightly distended with gray-reddish
5 dense fluid contents, up to the ileocecal sphincter (Fig. 1B). At this location there was a
6 protruding firm diverticulum of about 3 cm in diameter. Distal to this ileocecal
7 diverticulum, in the large intestine, there was almost no digesta, with few amounts of
8 gas, indicating an obstructive effect at the level of the ileocecal diverticulum. The
9 mucosa of the distal half of small intestine was uniformly thickened and there was a fine
10 granular brown-orange material (identified as fibrin) diffusely attached to the mucosal
11 surface, (Fig. 1C) effacing the normal honeycomb folds of the small intestinal mucosa of
12 leatherback turtles (Fig. 1D). The ileocecal diverticulum was filled with a 3 cm diameter
13 firm whitish dry mass, which was easily separated from the intestinal mucosa, that on
14 cut surface was shown to contain concentric stratified material (Fig. 1E). No internal
15 parasites were detected macroscopically.

16 Tissues were moderately autolyzed based on poor preservation of cellular morphologic
17 details. Sections of small intestine revealed diffuse necrosis of the upper part of the
18 mucosa (Fig. 2A), which was replaced by a thick band of hypereosinophilic cellular
19 debris and fibrin (Fig. 2B). Large amounts of small rods were found adhered to and
20 admixed with the necrotic cell debris (Fig. 2C). Most of these bacteria were Gram-
21 negative, although some Gram-positive rods were also present. Moderate numbers of
22 these Gram-positive rods were found in all organs investigated and were interpreted
23 as overgrowth of saprophytic bacteria due to autolysis. The mass found in the ileocecal
24 diverticulum corresponded to a granulomatous-necrotizing exudate, disposed in

1 concentric layers with bands of hypereosinophilic necrotic cells (consistent with
2 degenerated macrophages and heterophils) along with fibrin, necrotic material, and
3 bacteria (Fig. 2D). No significant lesions were observed in the other organs investigated
4 (liver, lung, spleen, brain). No fungal structures were detected with Grocott stain.

5 After 24h, a pure culture of β -hemolytic Gram-negative rods was obtained from
6 cultured scrapped mucosa of affected small intestine. The isolated bacterium was
7 identified as *P. damsela* with a percentage of identity of 99.9% using the API 20E
8 (profile 6015004) and API20NE (profile 5300044) identification system. *P. damsela*
9 includes two subspecies: *P. damsela* subsp. *damsela* and *P. damsela* subsp. *piscicida*,
10 which exhibit discriminating biochemical/phenotypical features. According to the
11 positive results obtained for β -hemolysis, nitrate reduction, urease, and lysine
12 decarboxylase, our isolate was identified as *P. damsela* subsp. *damsela*. No
13 *Salmonella* spp. were recovered.

14 **4. DISCUSSION**

15 Infections with bacteria from the family *Vibrionaceae* are frequent in sea turtles,
16 with most of these infections being caused by bacteria of the genus *Vibrio* (Glazebrook
17 et al. 1981, Work et al. 2003, Orós et al. 2005, Zavala-Norzagaray et al. 2015). However,
18 bacteria from the genus *Photobacterium*, another of the eight genera included in the
19 family, are not so frequently isolated from sea turtles. The taxonomy of the genus
20 *Photobacterium* has been recently reviewed (Labella et al. 2017). It contains 27 species
21 with valid names, with *P. damsela* being the most pathogenic for marine vertebrates.
22 Two subspecies of *P. damsela* are described, and both have been implicated in enteritis
23 in leatherback turtles. *Vibrio damsela* (*P. damsela* subsp. *damsela* under the actual
24 taxonomy) caused diphteric enteritis, valvular thrombotic endocarditis, and septicemia

1 in a female leatherback turtle stranded in Tasmania, Australia (Obendorf et al. 1987). *P.*
2 *damselae* subsp. *piscicida* (Gauthier et al. 1995) was determined to be the cause of
3 death of a stranded leatherback turtle with acute hemorrhagic gastroenteritis in the
4 Adriatic Sea (Poppi et al. 2012). Recently, *P. damsela* subsp. *damsela* has been isolated
5 from intestine and other organs from three stranded loggerhead turtles (*Caretta*
6 *caretta*) in the Tuscany coast of Italy (Fichi et al. 2016). Virulence of *P. damsela* subsp.
7 *damsela* is based on the secretion of toxins like phospholipase-D Dly (damselysin) and
8 the pore-forming toxins HlyApl and HlyAch (Rivas et al. 2013). The stranded leatherback
9 turtle presented in this report had a diffuse fibrino-necrotizing enteritis and obstructive
10 ileocecal diverticulitis, and *P. damsela* subsp. *damsela* was isolated in pure culture
11 from a mucosal scrapping, corroborating the pathogenicity of this bacteria for
12 leatherback turtles. The relationship between the observed fibrino-necrotizing enteritis
13 and the ileocecal diverticulitis is uncertain, but probably both lesions are unrelated to
14 each other. The existence of a diverticulum at the ileocecal junction appears to be an
15 anatomical feature of leatherback turtles, not present in other sea turtle species
16 (Magalhães et al. 2012). The diverticulum appears to be well developed in adult
17 individuals, but not in juveniles, and a high number of turtles showed inflammatory
18 changes of the diverticulum, which become distended by exudate. However, in none of
19 their cases the diverticulitis appeared to be associated to other lesions in small or large
20 intestine (Stacy et al. 2015). In our case there was intestinal obstruction at the ileocecal
21 junction, a complication not recorded previously in association with diverticulitis. The
22 intestinal lumen becomes narrowed in cases of diverticulitis (Stacy et al. 2015),
23 therefore we suggest that the thickening of the intestinal mucosa due to the fibrino-
24 necrotizing exudation in concurrence with diverticulitis have led both together to

1 obstruction. This view is supported by the fact that another leatherback turtle
2 necropsied in our laboratory in 2016 (data not shown) had also diverticulitis in absence
3 of other enteric lesions (Fig.1F).

4 *P. damselae* subsp. *damselae* is a water transmitted (Fouz et al. 2000) pathogen
5 of marine fishes, including blacksmith (*Chromis punctipinnis*), sharks, turbot
6 (*Scophthalmus maximus*), yellowtail (*Seriola quinqueradiata*), red-banded sea bream
7 (*Pagrus auriga*), sea bream (*Sparus aurata*) and sea bass (*Dicentrarchus labrax*), in which
8 it causes granulomatous ulcerative dermatitis (Austin 2010, Khouadja et al. 2014). It has
9 been also isolated from healthy skin and wounds from cetaceans, including bottlenose
10 dolphin (*Tursiops truncatus*) and Bryde's whale (*Balaenoptera brydei*). Its role as a
11 human pathogen is limited to causation of wound infections (Rivas et al. 2013).

12 In addition to the enteric bacterial disease, there was a bruise on the turtle's left
13 side, with subcutaneous hematoma, caused most likely by a blunt trauma to this side of
14 the body, possibly by a vessel strike. The excessive accumulation of serosanguinous fluid
15 in the coelomic cavity was probably a consequence of intestinal serositis. Hepatic
16 rupture may have increased the amount of serosanguinous fluid, although liver rupture
17 and hemorrhage were not clearly seen, and the liver was autolyzed and friable upon
18 removal, making it very difficult to distinguish lesions from postmortem alterations.
19 Parasitism is a frequent cause of morbidity and mortality in sea turtles (Greiner 2013)
20 but no significant parasitic diseases were observed in this case.

21 Sand is thought to have been introduced into the airways peri- or post-mortem
22 as the animal was tossed onto the beach by wave action. Despite many studies
23 establishing the ingestion of marine debris as a major threat to marine turtles (Bugoni

1 et al. 2001, Mrosovsky et al. 2009), in this case few foreign bodies were present in the
2 gastrointestinal tract, and those found were not thought to be associated with disease.

3 In view of the severity of intestinal lesions observed in our case, as well as of
4 previous reports of disease by others, the capability of *P. damselae* subsp. *damselae* to
5 act as a primary agent of disease in stranded or sick leatherback turtles should be
6 considered.

7

8 ACKNOWLEDGEMENTS

9 We thank Blanca Pérez, Aida Neira and Lola Perez, from the Servei de Diagnòstic de
10 Patologia Veterinària, Universitat Autònoma de Barcelona, and Carolina Gómez from
11 the Veterinary Mycology Group of the Universitat Autònoma de Barcelona for their
12 valuable technical assistance. Financial support came from Direcció General de Medi
13 Natural i Biodiversitat, Generalitat de Catalunya, and Servei de Diagnòstic de Patologia
14 Veterinària and Servei Veterinari de Bacteriologia i Micologia (Universitat Autònoma
15 de Barcelona).

16

17 REFERENCES

- 18 Austin B (2010) Vibrios as causal agents of zoonoses. *Vet Microbiol* 140:310–317
- 19 Bugoni L, Krause L, Petry M V. (2001) Marine debris and human impacts on sea turtles
20 in Southern Brazil. *Mar Pollut Bull* 42:1330–1334
- 21 Casale P, Nicolosi P, Freggi D, Turchetto M, Argano R (2003) Leatherback turtles
22 (*Dermochelys coriacea*) in Italy and in the Mediterranean basin. *Herpetol J*
23 13:135–139
- 24 Epibiont Research Cooperative (2007) A Synopsis of the Literature on the Turtle
25 Barnacles (Cirripedia: Balanomorpha: Coronuloidea) 1758-2007. Epibiont
26 Research Cooperative Special Publication Number 1
- 27 Fichi G, Cardeti G, Cersini A, Mancusi C, Guarducci M, Guardo G Di, Terracciano G
28 (2016) Bacterial and viral pathogens detected in sea turtles stranded along the
29 coast of Tuscany, Italy. *Vet Microbiol* 185:56–61
- 30 Flint M, Patterson-kane J, Mills P, Limpus C (2009) A veterinarian’s guide for sea turtle
31 post mortem examination and histological investigation. ISBN 9781864999594
32 <http://www.uq.edu.au/vetschool/content/vet-marti/PM.Guide.MSF.pdf>.
- 33 Fouz B, Toranzo AE, Milan M, Amaro C (2000) Evidence that water transmits the

- 1 disease caused by the fish pathogen *Photobacterium damsela* subsp. *damsela*. J
2 Appl Microbiol 88:531–535
- 3 Gauthier G, Lafay B, Ruimy R, Breittmayer V, Nicolas JL, Gauthier M, Christen R (1995)
4 Small-subunit rRNA sequences and whole DNA relatedness concur for the
5 reassignment of *Pasteurella piscicida* (Snieszko et al.) Janssen and Surgalla to the
6 genus *Photobacterium* as *Photobacterium damsela* subsp. *piscicida* comb. nov. Int
7 J Syst Bacteriol 45:139–144
- 8 Glazebrook J, Campbell R, Johnson R (1981) Traumatic ulcerative dermatitis: A disease
9 of captive sea turtles *Chelonia mydas* (L) in north-east Australia. In: Wildlife
10 Diseases of the Pacific Basin and Other Countries, 4th International Conference of
11 the Wildlife Diseases Association, ME Fowler, Sydney, NSW.p 160
- 12 Gómez de Segura A, Tomás J, Pedraza SN, Crespo EA, Raga JA (2006) Abundance and
13 distribution of the endangered loggerhead turtle in Spanish Mediterranean
14 waters and the conservation implications. Anim Conserv 9:199–206
- 15 Greiner EC (2013) Parasites of marine turtles. In: Wyneken J (ed) The Biology of Sea
16 Turtles, Vol III. CRC Press, p 427–446
- 17 Khouadja S, Lamari F, Bakhrouf A, Gaddour K (2014) Virulence properties, biofilm
18 formation and random amplified polymorphic DNA analysis of *Photobacterium*
19 *damsela* subsp. *damsela* isolates from cultured sea bream (*Sparus aurata*) and
20 sea bass (*Dicentrarchus labrax*). Microb Pathog 69–70:13–19
- 21 Labella AM, Arahal DR, Castro D, Lemos ML, Borrego JJ (2017) Revisiting the genus
22 *Photobacterium*: taxonomy, ecology and pathogenesis. Int Microbiol 20:1–10
- 23 Magalhães M dos S, Santos AJB, Silva NB da, Moura CEB de (2012) Anatomy of the
24 digestive tube of sea turtles (Reptilia: Testudines). Zool 29:70–76
- 25 Mrosovsky N, Ryan GD, James MC (2009) Leatherback turtles: The menace of plastic.
26 Mar Pollut Bull 58:287–289
- 27 Obendorf DL, Carson J, McManus TJ (1987) *Vibrio damsela* infection in a stranded
28 leatherback turtle (*Dermochelys coriacea*). J Wildl Dis 23:666–8
- 29 Orós J, Torrent A, Calabuig P, Déniz S (2005) Diseases and causes of mortality among
30 sea turtles stranded in the Canary Islands, Spain (1998-2001). Dis Aquat Organ
31 63:13–24
- 32 Poppi L, Zaccaroni A, Pasotto D, Dotto G, Marcer F, Scaravelli D, Mazzariol S (2012)
33 Post-mortem investigations on a leatherback turtle *Dermochelys coriacea*
34 stranded along the Northern Adriatic coastline. Dis Aquat Organ 100:71–76
- 35 Rivas AJ, Lemos ML, Osorio CR (2013) *Photobacterium damsela* subsp. *damsela*, a
36 bacterium pathogenic for marine animals and humans. Front Microbiol 4:1–6
- 37 Stacy BA, Innis CJ, Daoust PY, Wyneken J, Miller M, Harris H, James MC, Christiansen

- 1 EF, Foley A (2015) Solitary Large Intestinal Diverticulitis in Leatherback Turtles
2 (*Dermochelys coriacea*). Vet Pathol 52:712–715
- 3 Work TM, Balazs GH, Wolcott M, Morris R (2003) Bacteraemia in free-ranging Hawaiian
4 green turtles *Chelonia mydas* with fibropapillomatosis. Dis Aquat Organ 53:41–46
- 5 Zavala-Norzagaray AA, Aguirre AA, Velazquez-Roman J, Flores-Villaseñor H, León-
6 Sicaio N, Ley-Quiñonez CP, Hernández-Díaz LDJ, Canizalez-Roman A (2015)
7 Isolation, characterization, and antibiotic resistance of *Vibrio* spp. in sea turtles
8 from northwestern Mexico. Front Microbiol 6:1–10
- 9


1 FIGURES



2
3 Fig. 1. Gross findings of stranded leatherback turtle necropsy. (A) Deposits of sand in tracheal
4 lumen (B) Distention of small intestine and reddening of the intestinal serosa (esophagus and
5 stomach on the left side). Arrowhead is showing the transition of small intestine to large
6 intestine. Large intestine was distended with gas. (C) Mucosal surface of small intestine
7 showing fibrino-necrotizing enteritis. The surface has been rinsed with water (D) Normal small
8 intestine mucosa of another leatherback turtle showing the characteristic honeycomb pattern
9 (E) Section of the luminal nodulation found in the ileocecal diverticulum, composed of white
10 stratified material. (F) Normal leatherback turtle showing transition of small intestine (left) to
11 large intestine (right), and ileocecal diverticulum (arrowhead) with diverticulitis (presence of a
12 spherical stratified mass composed of fibrin and cellular exudate, see Stacy et al., 2014).

13

1
2
3



4

5 Fig. 2. Small intestine, leatherback turtle, Haematoxylin & Eosin staining. (A) Intestinal
6 mucosa and submucosa, low magnification, showing a fibrino-necrotizing enteritis,
7 with a superficial layer (arrowhead) of necrotic tissue and fibrin. Necrosis extends
8 through almost whole mucosa. Submucosal vessels are distended by gas from autolysis
9 (B) Small intestine, mucosa, closer view of fibrino-necrotizing enteritis (arrowhead).
10 There is a layer of bright eosinophilic necrotic tissue underneath a loose layer of fibrin
11 and cell debris mixed with clumps of basophilic rods. Basophilic, pyknotic cell nuclei
12 are still visible in the necrotic layer. The muscularis mucosae is visible underneath the
13 necrotic mucosa (arrow). (C) Detail of fibrino-necrotic mucosa (upper-left side of the
14 photograph), with pyknotic cell nuclei, and numerous bacteria overlying the necrotic
15 enteric mucosa. Most of these bacteria were GRAM-negative, but also some large
16 GRAM-positive rods were present. (D) Diverticular mass showing stratified layers of
17 inflammatory infiltrates, cellular debris and intestinal contents with bacteria.

18
19
20
21