





الجامعة الإسلامية العالمية ماليزيا
INTERNATIONAL ISLAMIC UNIVERSITY MALAYSIA
يُونِيسْكَوِي اِسْلَامِي اِنْتَارَا اَبْحَثِيَا مَلِيسِيَا

Garden of Knowledge and Virtue

IJOHS

*International Journal
of Orofacial and
Health Sciences*

A scientific journal
published by Kulliyah of
Dentistry, IIUM, Malaysia



ISSN 2716-5434



9 772716 543003

VOL. 1 ISSUE 1

2020



International Journal of Orofacial and Health Sciences

Editorial Board

Chief Editor	: Professor Dr. Zainul Ahmad Rajion
Editor	: Dr. Salwana Supa'at
	: Dr. Basma Ezzat Mustafa Alahmad
	: Assoc. Prof. Dr. Khairani Idah Mokhtar@Makhtar
	: Assoc. Prof. Dr. Solachuddin J.A. Ichwan
	: Assoc. Prof. Dr. Muhannad Ali Kashmoola
	: Dr. Widya Lestari
	: Dr. Azlini Ismail
	: Dr. Mohd Hafiz Arzmi
	: Dr. Mohamad Shafiq Mohd Ibrahim

Aims and Scope:

International Journal of Orofacial and Health Sciences (IJOHS) is a peer reviewed biannual international journal dedicated to publish high quality of scientific research in the field of orofacial sciences, health sciences and interdisciplinary fields, including basic, applied and clinical research. The journal welcomes review articles, original research, case reports and letters to the editor. Areas that are covered include but are not limited to dental sciences, oral microbiology and immunology, oral maxillofacial and craniofacial surgery and imaging, dental stem cells and regenerative medicine, dental biomaterial, oral maxillofacial genetic and craniofacial deformities.

Published by:

Kulliyah of Dentistry,
International Islamic University Malaysia (IIUM),
Indera Mahkota,
25200 Kuantan, Pahang, Malaysia.

Printed and distributed by:

Kulliyah of Dentistry,
International Islamic University Malaysia (IIUM),
Indera Mahkota,
25200 Kuantan, Pahang, Malaysia.

All rights reserved; No part of this publication maybe reproduced, stored in a retrieval system, or transmitted in any form, or by any means, electronic or otherwise, without prior permission of the publisher.

CONTENTS

EDITORIAL

Introduction to IJOHS 3

REVIEW ARTICLES

Genetics of malocclusion: A review 4

ORIGINAL ARTICLES

Potential antibacterial effects of flaxseed and *Nigella sativa* extracts on *Streptococcus pyogenes* 11

Dental treatment needs among patients undergoing screening at a university-based dental institution in Kuantan, Pahang, Malaysia 18

Analysis of the anti-cancer effect of ethyl-p-methoxycinnamate extracted cekur (*Kaempferia galanga*) on cancer cell lines with wild-type and null p53 28

Radiographic findings in panoramic radiographs of patients attending Kulliyah of Dentistry, IIUM 34

Isolation of *Candida* species in children and their biofilm-forming ability on nano-composite surfaces 40

Radiographic findings in panoramic radiographs of patients attending Kulliyah of Dentistry, IIUM

Muhannad Ali Kashmoola^{1*}, Nazih Shaaban Mustafa¹, Azrul Safuan Mohd Ali², Kamal Ammar Dahlan³

¹Department of Oral Maxillofacial Surgery and Oral Diagnosis, Kulliyah of Dentistry, International Islamic University Malaysia

²Department of Orthodontics, Kulliyah of Dentistry, International Islamic University Malaysia

³Ministry of Health, Malaysia

Abstract

This research was done to study the radiographic finding of the jaws amongst the patients attending at the polyclinic using Orthopantomograph (OPG). The OPG was taken using the Planmeca Promax 3D and the Planmeca Romexis software (Version 2.1.1.R). The first step was collecting all the OPG images from 1st April 2009 until 31st January 2011. Then, the abnormal radiographs were further divided into 3 groups which were radiolucent, radiopaque and mixed. This classification includes site, size, border, and possible diagnosis as part of the lesion's appearance. One thousand four hundred and five OPG images were retrieved, 96 images were discarded because of poor quality. The data collected were analyzed statistically by using SPSS Version 16.0. Among 77 abnormal radiographic images, 41 images were radiopaque, 30 images were radiolucent and 6 images were mixed. Out of 77 abnormal images, 34 images that showed bone lesion were from male patients while the rest which was 43 images from female. In conclusion, most of the pathological lesion occur in the mandible.

Keywords: Panoramic, radiograph, jawbone, lesion

*Corresponding Author

Email: drmuhannadkashmoola@iium.edu.my

Tel: +6019-252 9418

Introduction

Radiograph as a method of investigation plays an important role in making diagnostic decision in determining the diseases. Each individual that need radiographic examination are based on findings from dental history and clinical examination and customized by age and general health of the patient (*Sirisha et al.*, 2013). Thus, radiographs mostly referred to as the clinician's key of diagnostic support (*Lee et al.*, 2013). Besides that radiographs are ordered when dentist sought that the radiographs will provide helpful diagnostic information which will affect the treatment plan (*Kapila et al.*, 2011). However, lesions inside the jaw are sometimes accidentally discovered by the dentist during routine examination.

Oral lesions is always a challenge for clinician and based on the experience, constructing a final diagnosis can be complicated and easily misinterpreted.

Many of the oral pathological lesions cannot be viewed clinically, and the indication for dental radiograph can provide important information on a patient's oral health and general health. (*Diz et al.*, 2013).

Panoramic imaging is a two dimensional radiographic technique for producing a single 2D tomographic image of the facial structures that include both maxillary and mandibular dental arches and their supporting structures (*Subbulakshmi, et al.*, 2016). Besides that, the panoramic imaging produces a

single image of a complete representation of the jaws, teeth, temporomandibular joint, alveolar lobes of the maxillary sinuses. This image forms when two adjacent disks are rotating at the same speed in opposite directions as an X-rays been passed through their centers of rotation.

Lesion is defined as a pathologic disturbance of a tissue, with loss of continuity, enlargement, or function (Regezi, Sciubba, & Jordan, 2015). The most common jaw lesions are cyst and tumours. A cyst is defined as an epithelium lined pathologic cavity (Deepthi *et al.*, 2016). On the other hand, tumour of the jaw can be divided by 2 categories; odontogenic and non-odontogenic tumour. As the name implies, odontogenic tumour is the tumour that derived from the tooth structure related to the tooth while non-odontogenic tumours are the most common neoplasm of the jaw (Zegalie, Speight, & Martin, 2015).

The radiographic jaw lesions can be describe as having either a radiolucent, radiopaque or mixed appearance, relative to density of the adjacent bone which more than 80% of them is radiolucent (Ed, 2017). It can be multilocular or unilocular with well-defined or poor defined border. In general, lesions with well-defined borders are usually benign whereas poor defined borders invariably represent aggressive, inflammatory or neoplastic process (Hall, 2017).

This research is primarily concerned with the radiographic study of bone lesions found in patients coming to the polyclinic of the Kulliyah of Dentistry, IIUM. The prevalence of bone lesions in patient will be analyzed based on the radiograph that had been taken beginning from the patient registered 1st April 2009 until 31st January 2011.

Thus, the objectives of this study were to find out the prevalence of bone lesions in patients attending polyclinic, Kulliyah of Dentistry clinics through OPG, its radiographic findings and to assess the most common radiographic lesions and location.

In addition the study was done in order to set a potential data base on bone lesions in patients attending KOD, IIUM.

Material and Methods

A retrospective cross-sectional study was done to assess the prevalence of radiographic bone lesion in the patient coming to the Kulliyah of Dentistry, IIUM. The approach of the study is quantitative, by collecting data (radiographic images) taken in this Kulliyah from 1st April 2009 until 31st January 2011. It is appropriate to use cross sectional study because this study involve different group of people coming to the polyclinic, Kulliyah of Dentistry.

The total number of collected radiographs (OPG) in the polyclinic is 1405. The images collected had been and screened for any abnormalities by two specialists in Oral maxillofacial pathology and Oral maxillofacial surgery due to the absence of maxillofacial radiologist. The data was collected by using the Planmeca Promax 3D and the Planmeca Romexis software version (2.1.1.R). The radiographs that has artifact, ghost image, distorted and blurred will be excluded from the study. After data has been collected and was visually checked for correction, the data was analyzed using SPSS Version 16.0.

Results

Out of the 1,405 OPG collected, 96 OPGs were excluded from the study due to technical defects and 1,309 were included. The number of radiographs with radiographic lesions were 77 and the rest, 1,232 were presented with normal radiographic findings. Out of the 77

images, 34 (44.2%) images showed bone lesion were from male patients and 43 (55.8%) were female. Regarding the race, the data showed that Malay patients were the highest 67 (87%), followed by 9 (11.7%) of Chinese and only 1 (1.3%) Indian respectively)

The prevalence of radiographic lesions of the jawbone in this study was 5.9%.

Table 1 shows the radiographic presentation radiolucent lesions were 30 (39%), (Figure 1) and radiopaque lesions were 41 (53.2%), (Figure 2). The remaining 6 (7.8%) were mixed lesions (Figure 3).

Sixty five radiographic lesions were found in the mandible, 37 on the right

body and seven on the left body. However, less lesion occurred in the maxilla as shown in Table 2.

Most of the radiographic bone lesions found in this study had a well-defined margin which was 63 (81.8%) and the rest 14 (18.2%) were not clearly defined. Most of bone lesions 71 (92.2%) appeared as unilocular, whereas six (7.8%) were presented as multilocular.

Size of lesions were divided into 4 groups, small (2.5-7.5mm), (62%), moderate (7.6-10mm), (21%) and large (above 10 mm). (17). Minimum size observed was 2.5 mm, and the maximum size was 19 mm, mean value was 7.344 and standard deviation was 3.19.



Figure 1. Panoramic radiograph showing a radiolucent lesion in the lower anterior region of the mandible.



Figure 2. Panoramic radiograph showing radiopaque lesion in the lower left and right body of the mandible.



Figure 3. Panoramic radiograph showing a mixed radiolucent- radiopaque lesion in the lower posterior right angle of the mandible.

Table 1. Distribution of different types of radiographic lesions according to gender, loculation and appearance.

Total	Gender		Loculation		Appearance		
	Male	Female	Uni-locular	Multi-locular	Radiolucent	Radiopaque	Mixed
25 (36.4%)	17 (22%)	11 (14.3%)	28	0	28 (93%)	0	0
5 (6.5%)	1 (1.3%)	4 (5.2%)	5	0	0	5 (12.2%)	0
25 (32.5%)	9 (11.7%)	16 (20.8%)	23	2	0	25 (61%)	0
3 (3.9%)	1 (1.3%)	2 (2.6%)	1	2	1 (16.7%)	2 (4.9%)	0
3 (3.9%)	2 (2.6%)	1 (1.3%)	3	0	0	3 (7.3%)	0
13 (16.9%)	4 (5.2%)	9 (11.7%)	11	2	2 (6.7%)	6 (14.6%)	5 (53.3%)
77 (100%)	34 (44.2%)	43 (58.2%)	71	6	30 (100%)	41 (100%)	6 (100%)

Table 2. Site of radiographic lesions seen in OPG

Site	Lesions	Percentage (%)
RT mandible ramus	1	1.3
RT mandible body	36	46.8
LT mandible ramus	3	3.9
LT mandible body	25	32.5
RT maxilla posterior	3	3.9
RT maxilla anterior	4	5.1
LT maxilla posterior	2	2.6
LT maxilla anterior	3	3.9
Total	77	100

Discussion

Based on the result, it showed that 77 (6.78 %) among 1232 patients attending the dental clinic of Kulliyah of Dentistry, IIUM, have jaw lesion. From all the patients having jaw lesion, 44.2 % were male and 55.8 % were female. The result may not show the exact number of the lesions affecting each gender since certain diseases are related to gender. For instance, in odontogenic kerato cyst, the

ratio of male to female is 2.1 to 1.2 (Al-Moraissi et al., 2017).

The characteristic appearance of each lesion varies, and it was categorized into three; radiolucent, radiopaque and mixed radiolucent and radiopaque. From the collected results, the radiopaque lesion was the highest among the collected sample with a percentage of 53.2 %, however, Eldaya in 2017 stated that more

than 80 % of the lesions should be of radiolucent (Eldaya et al., 2017).

In this study, the mandible was found to be affected by (84.5 %) of the jaw lesions. Similarly, Araki *et al.* (2011) found that radiopaque lesions were mostly found in molar and premolar region of mandible.

Conclusion

This study showed that 6.25 % of the radiographs exhibited pathological lesions. Moreover, most of the lesions were discovered accidentally in the radiograph. More studies are recommended with a larger sample size. In conclusion, most of the pathological lesion occur in the mandible, and less in the maxilla.

References

- Al-Moraissi, E. A., Dahan, A. A., Alwadeai, M. S., Oginni, F. O., Al-Jamali, J. M., Alkhutari, A. S., Al-Sanabani, J. S. (2017). What surgical treatment has the lowest recurrence rate following the management of keratocystic odontogenic tumor?: A large systematic review and meta-analysis. *Journal of Cranio-Maxillofacial Surgery*. 45(1) , 131-144.
- Deepthi, P., Beena, V., Padmakumar, S., Rajeev, R., & Sivakumar, R. (2016). A study of 1177 odontogenic lesions in a South Kerala population. *Journal of Oral and Maxillofacial Pathology*. 20(2), 202-207.
- Diz, P., Scully, C., & Sanz, M. (2013). Dental implants in the medically compromised patient. *Journal of Dentistry*. 41(3), 195-206.
- ED, O. (2017). Clinico-radiological Presentations of Maxillofacial Hard-Tissue Swellings in a Tertiary Health Facility in West African Suburb. *Biomedical Journal of Scientific & Technical Research*. 1(3), 2
- Eldaya, R., MD, M. B. A., Eissa, O., Herrmann, S., Pham, J., Calle, S., & Uribe, T. (2017). Mandibular Lesions: A Practical Approach for Diagnosis Based on Multimodality Imaging Findings. *Contemporary Diagnostic Radiology*. 40(6), 1-7.
- Hall, G. (2017). Fibro-osseous lesions of the head and neck. *Diagnostic Histopathology*. 23(5), 200-210.
- Kapila, S., Conley, R. S., & Harrell, W. E. (2011). The current status of cone beam computed tomography imaging in orthodontics. *Dentomaxillofacial Radiology*. 40(1), 24-34.
- Lee, C. S., Nagy, P. G., Weaver, S. J., & Newman-Toker, D. E. (2013). Cognitive and system factors contributing to diagnostic errors in radiology. *American Journal of Roentgenology*. 201, 611-617.
- Regezi, J., Sciubba, J., & Jordan, R. (2015). Oral Pathology Clinical Pathologic Correlations. The effects of brief mindfulness intervention on acute pain experience: An examination of individual difference (Vol.1).
- Sirisha, K., Srilalitha, A., Srinivas, M., & Chakrapani, S. (2013). Choice of diagnostic and therapeutic imaging in periodontics and implantology. *Journal of Indian Society of Periodontology*. 17(6), 711-718.
- Subbulakshmi, Ac., Mohan, N., Thiruneervannan, R., Naveen, S., & Gokulraj, S. (2016). Positioning errors in digital panoramic radiographs: a study. *Journal of Orofacial Sciences*. 8(1), 22-26.
- Zegalie, N., Speight, P. M., & Martin, L. (2015). Ossifying fibromas of the jaws and craniofacial bones. *Diagnostic Histopathology*. 21(9), 351-358.

Instruction to Authors

Submission

Complete manuscripts, written in English along with tables and illustrations should be submitted as softcopy to zainulrajion@iium.edu.my

Conditions

All manuscripts submitted for publication must be accompanied by a cover letter declaring originality of the work and certifies that the manuscript has not been published or submitted elsewhere, free from conflict of interest and conduct of research in accordance to ethical guidelines established for human subjects and animal welfare. In addition, a statement of agreement of all authors to transfer all the copyrights to the journal should also accompany the manuscript.

A copy of the certification form, which certifies that the manuscript has not been published or submitted before and all authors have contributed and are in agreement with the content of the manuscript, must be submitted to the journal's editorial office by uploading it as a PDF file.

Manuscripts submitted for publication are understood to be offered only to **IJOHS** and which have not been sent to other journals for consideration. There is no publication fee to submit or publish content in **IJOHS**.

Manuscript should be typed in **Arial, font size 11 with 1.5 spacing**.

The manuscript should be divided into the following sections with the sequence; Title Page, Abstract, Introduction, Materials and Methods, Results, Discussion, Acknowledgements, References, Tables, Illustrations and Figure Legends must be incorporated near to the text they are cited

Title Page:

Title page should be arranged in the following order: (1) a concise and informative title not exceeding 80 characters; (2) authors' full names (without degrees and titles) and affiliation including city and country. Superscript numbers may be used to affiliate authors to different departments or institutions; and (3) a complete mailing address, telephone, fax and e-mail address of the corresponding author.

Abstract:

Abstract should be unstructured and should not exceed 250 words. It should be a self-standing summary of the work that may be republished by information retrieval services. A maximum of 5 keywords should be listed at the end of the abstract.

Introduction:

Introduction should briefly and clearly describe the background and objective(s) of the study. Avoid exhaustive review of the literature and include only relevant recent studies.

Materials and Methods:

This section should be described in sufficient details so that it is reproducible. It should include technical information such as the study design, and specific procedures. Sub-headings may be used to enhance the clarity or to categorize the methods. Established methods or procedures should be named and cited. New methods or modifications of old methods should be described with complete details. Authors are advised to use generic names and terms rather than commercial names. Statistical methods applied, software used must also be stated clearly and concisely.

Results:

This section should be presented in a logical sequence in text, tables and illustrations. Subheadings may be used

to enhance clarity or to call attention to the most significant findings. Data appearing in tables or figures may be summarized but not duplicated in the text. Tables and figures should be numbered in the order in which they are described and cited in the text. Example, For tables, examples are Table 1, Table 2 and for figures, examples are Figure 1, Figure 2.

Discussion:

This section should summarize, explain and interpret the results with a scientifically critical view of previously published works in the field. Avoid repetition of the data already presented in the result section.

Acknowledgements:

This section is to be given to those who have provided assistance to the project and sponsors who have given financial support and funds.

References:

This section should list all sources cited in the paper. The citations should be arranged in alphabetical order by the surname of the first author without numbering in the text. Cite article as e.g. (Ahmad, 2016) for a single author, (Chu & Kim, 2019) for two authors and (Karim *et al.*, 2013) for more than three authors.

Standard journal article

References to journals should provide the name(s) of the author(s), year, title of the paper and journal, volume and issue number, page numbers. Please provide full name of journal. If there are more than six authors, the first six authors should be listed and followed by *et al.* in the list of References.

Sudbery, P. E. (2011). Growth of *Candida albicans* Hyphae. *Nature Reviews Microbiology*, 9(10), 737.

Book

References to books/monographs should give the name(s) of the author(s), year, title of book, edition number, place of publication, publisher and page numbers.

Conn, E.E., Stumpf, P.K., Brueing, G., Doi, R.H. (1987). *Outlines of Biochemistry*, 3rd edn. New York: John Wiley & Sons, pp. 45–52.

Chapter in book

References to chapter in books/monographs should give the name(s) of the author(s), year, chapter title, editors, title of book, edition number, place of publication, publisher and page numbers.

Fejerskov O, Nyvad B, Kidd EAM (2003). Clinical and histological manifestations of dental caries. In: Fejerskov O, Kidd EAM (eds.), *Dental Caries – The Disease and Its Clinical Management*. London: Blackwell Munksgaard, pp. 71–97.

Internet resources

Authors are required to include as much electronic retrieval information as needed for others to locate the sources they have cited.

López-Jornet P (2006). Labial mucocele: a study of eighteen cases. *The Internet Journal of Dental Science*, 3(2). Retrieved 7 February 2007, from <http://www.ispub.com/ostia/index.php?xmlFilePath=journals/ijds/vol3n2/labial.xml>

Tables

Each table should have a suitable title with footnotes wherever necessary. Do not use vertical lines in the table.

Illustrations

All figures and illustrations could be original photographs, artwork or high-quality digital images (submitted as CMYK - 8 bits per channel in TIFF format). Images must be at least 600 by 450 pixels (proportional height) in size when in landscape orientation with a resolution of at least 300 pixels per inch. Graphs should be approximately 500 pixels wide so that all labeling can be read with data points clearly visible. Figures should be numbered

consecutively in the order in which they appear in the manuscript, using Arabic numerals. A list of figure legends must be included on a separate page following the illustrations. The legend should explain each figure in detail. All figures will be printed as black and white. Colour figures will only appear in the PDF file.

Page proofs

On acceptance of the manuscripts for publication, page proofs should be reviewed meticulously by the contributors. Changes made in proof should be limited to the correction of typographical errors. Proofs must be returned for publication with corrections and responses to queries on the date specified by the Editor.



الجامعة الإسلامية العالمية ماليزيا
INTERNATIONAL ISLAMIC UNIVERSITY MALAYSIA
يُونِسُورِيَتِي اِسْلَامِيَّةَ اِنْدُونِيسِيَا
Garden of Knowledge and Virtue