



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE
Repository istituzionale dell'Università degli Studi di
Firenze

Existence and regularity for eddy current system with non-smooth conductivity

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Existence and regularity for eddy current system with non-smooth conductivity / Francini E., Franzina G., Vessella S.. -
In: SIAM JOURNAL ON MATHEMATICAL ANALYSIS. - ISSN 0036-1410. - STAMPA. - 52(2020), pp. 2134-2157.
[10.1137/19M1276984]

Availability:

This version is available at: 2158/1184190 since: 2020-12-15T08:14:56Z

Published version:

DOI: 10.1137/19M1276984

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

EXISTENCE AND REGULARITY FOR EDDY CURRENT SYSTEM WITH NON-SMOOTH CONDUCTIVITY

ELISA FRANCCINI, GIOVANNI FRANZINA, AND SERGIO VESSELLA

ABSTRACT. We discuss the well-posedness of the “transient eddy current” magneto-quasistatic approximation of Maxwell’s initial value problem with bounded and measurable conductivity, with sources, on a domain. We prove existence and uniqueness of weak solutions, and we provide global Hölder estimates for the magnetic part.

1. INTRODUCTION

Let Ω be a bounded $C^{1,1}$ domain in \mathbb{R}^3 (see Section 2.1 for definitions), and let n denote the outward unit normal to its boundary. We consider electromagnetic signals throughout a medium, filling the region Ω , with *magnetic permeability* being given by a Lipschitz continuous scalar function μ and *electric conductivity* being described by a bounded measurable function σ taking values in the real symmetric 3×3 matrices. We will assume the validity of the conditions

$$(1.1i) \quad \Lambda^{-1} \leq \mu \leq \max\{\mu, |\nabla\mu|\} \leq \Lambda, \quad \text{a.e. in } \Omega,$$

$$(1.1ii) \quad \Lambda^{-1}|\eta|^2 \leq \sigma\eta \cdot \eta \leq \Lambda|\eta|^2, \quad \text{for all } \eta \in \mathbb{R}^3, \text{ a.e. in } \Omega,$$

for an appropriate constant $\Lambda \geq 1$.

Given $T > 0$, $\mathbf{H}_0 \in L^2(\Omega; \mathbb{R}^3)$, $\mathbf{G} \in L^2(0, T; H^1(\Omega; \text{curl}))$, with $\partial_t \mathbf{G} \in L^2(0, T; L^2(\Omega; \mathbb{R}^3))$, and $\mathbf{J}^E, \mathbf{J}^M \in L^2(0, T; L^2(\Omega; \mathbb{R}^3))$, we consider weak solutions $(\mathbf{E}, \mathbf{H}) \in L^2(0, T; H^1(\Omega; \text{curl}) \times H_0^1(\Omega; \text{curl}))$, with $\partial_t \mathbf{H} \in L^2(0, T; L^2(\Omega; \mathbb{R}^3))$ (see Section 2 for definitions), of the initial value problem

$$(1.2) \quad \begin{cases} \nabla \times \mathbf{H} - \sigma \mathbf{E} = \mathbf{J}^E, & \text{in } \Omega \times (0, T), \\ \nabla \times \mathbf{E} + \mu \partial_t \mathbf{H} = \mathbf{J}^M, & \text{in } \Omega \times (0, T), \\ \mathbf{H} \times n = \mathbf{G} \times n, & \text{on } \partial\Omega \times (0, T), \\ \mathbf{H} = \mathbf{H}_0, & \text{in } \Omega \times \{0\}, \end{cases}$$

under the assumption that

$$(1.3) \quad \nabla \cdot \left(\mu \mathbf{G} - \mu \mathbf{H}_0 - \int_0^t \mathbf{J}^M ds \right) = 0, \quad \text{in } \Omega \times (0, T).$$

The meaning of (1.2) and of (1.3) will be understood in a suitable weak sense in Section 2.

Formally, the so-called eddy current system (1.2) is obtained from Maxwell’s equations when neglecting displacement currents and is equivalent to the parabolic system

$$(1.4) \quad \mu \partial_t \mathbf{H} + \nabla \times \left(\sigma^{-1} \nabla \times \mathbf{H} \right) = \nabla \times (\sigma^{-1} \mathbf{J}^E) + \mathbf{J}^M, \quad \text{in } \Omega \times (0, T),$$

2010 *Mathematics Subject Classification.* 35M33, 35M60, 35R05, 35Q61, 35B65.

Key words and phrases. eddy currents, non-smooth coefficients, initial-boundary value problem, global Hölder estimates.

with the conditions $\mathbf{H} \times n = \mathbf{G} \times n$ on $\partial\Omega \times (0, T)$ and $\mathbf{H} = \mathbf{H}_0$ in $\Omega \times \{0\}$, provided that

$$\mathbf{E} = \sigma^{-1} \left(\nabla \times \mathbf{H} - \mathbf{J}^E \right), \quad \text{in } \Omega \times (0, T).$$

To make an example, if σ is constant and $\mathbf{J}^E = \mathbf{J}^M = 0$, then (1.4) reads as

$$\mu\sigma\partial_t\mathbf{H} + \nabla \times \nabla \times \mathbf{H} = 0, \quad \text{in } \Omega \times (0, T),$$

and $\nabla \times \nabla \times \mathbf{H} = \nabla(\nabla \cdot \mathbf{H}) - \Delta\mathbf{H}$, where the Laplace operator is understood componentwise. Hence, in this case the problem is equivalent to the heat equation for the Hodge-Laplacian on vector fields, and the components of divergence-free solutions solve the classical heat equation (up to a weight).

Our interest in this *parabolic* magneto-quasistatic approximation of the laws of classical electromagnetism with possibly *discontinuous* electric conductivity tensor comes from *diffusive* models in applied seismo-electromagnetic studies [17, 19]. In geophysics, the importance of modelling *slowly varying* electromagnetic fields throughout the *stratified* lithosphere is due to the possibility that some of them may be generated by co-seismic subsurface electric currents, and hence have some rôle in the seismic precursor signal recognition. For a very general survey on eddy currents with discontinuous conductivity and related numerics, with applications to advanced medical diagnostics, the interested reader is referred instead to the nice treatise [3], where inverse problems are also considered. We refer to [4] for issues related to the source identification from boundary EM measurement.

The main results of this manuscript concern some qualitative properties of weak solutions of (1.2), i.e., their existence and uniqueness, as well as the Hölder continuity of their magnetic part. For expositional purposes, we limit ourselves to the case of homogeneous boundary conditions, which causes no restriction (see Section 2.4).

In Theorem 3.1 (see Section 3), we prove the well-posedness of (1.2); for, we make use of Galerkin's method and of the Hilbert basis that we manufacture in Section 3.1 by solving an auxiliary problem of spectral type. This special system of vector fields has the expedient feature of being independent of the conductivity stratification, at variance with the natural basis for the associated parabolic problem. Existence and uniqueness results are available in the literature for problems similar to (1.2); for example, in the time-harmonic regime the issue of well-posedness was addressed in [16], and in [6] (where it is also proved to be a good approximation of the complete set of Maxwell's equations), and the time-harmonic variant of (1.2) is also dealt with in the more recent paper [7], providing existence and uniqueness results and asymptotic expansions in terms of the size of the conductor in this context, whereas in [8] the well-posedness of the variant of this problem focused on the electric field is discussed using a different approach, in the time domain, with applications to the asymptotic behaviour of solutions in the non-conductive limit.

In Theorem 4.1 (see Section 4), inspired by the work [1] on Maxwell's system, we prove Hölder continuity estimates for the magnetic field, valid up to the boundary. In the literature, we could not find either global or local estimate of this kind; we refer to the paper [11] for some related result.

Plan of the paper. In Section 2 we make precise assumptions on the domain and on the structure of the problem, we introduce the reader to some useful functional-analytic tools, we state some Helmholtz-type decompositions (proved in Appendix), and we define the weak solutions of the eddy current system (1.2). In Section 3 we prove existence and uniqueness of weak solutions (\mathbf{E}, \mathbf{H}) , and in Section 4 we provide global *a-priori* Hölder estimates on the magnetic field \mathbf{H} .

Acknowledgments. This research is supported by the MIUR-FOE-INdAM 2014 grant "Strategic Initiatives for the Environment and Security - SIES".

2. TECHNICAL TOOLS

We recall that the tangential trace, defined by $\phi \times n$ for all $\phi \in C^1(\overline{\Omega}; \mathbb{R}^3)$, extends to a bounded operator from the Hilbert space $H^1(\Omega; \text{curl})$, consisting of all vector fields in $L^2(\Omega; \mathbb{R}^3)$ whose (distributional) curl is also in $L^2(\Omega; \mathbb{R}^3)$, endowed with the scalar product

$$(2.1) \quad (\varphi, \psi)_{H^1(\Omega; \text{curl})} = (\varphi, \psi)_{L^2(\Omega; \mathbb{R}^3)} + (\nabla \times \varphi, \nabla \times \psi)_{L^2(\Omega; \mathbb{R}^3)},$$

to the dual space $H^{-\frac{1}{2}}(\partial\Omega; \mathbb{R}^3)$ of $H^{\frac{1}{2}}(\partial\Omega; \mathbb{R}^3)$ (see, e.g., [12]). Indeed, the Green-type formula

$$(2.2) \quad \int_{\Omega} \varphi \cdot \nabla \times \psi \, dx - \int_{\Omega} \psi \cdot \nabla \times \varphi \, dx = - \int_{\partial\Omega} \varphi \cdot (\psi \times n) \, dS$$

holds for all $(\varphi, \psi) \in (C^1(\overline{\Omega}; \mathbb{R}^3))^2$. Moreover, given $\psi \in H^1(\Omega; \text{curl})$, by Sobolev extension and trace theorems, the left hand-side of (2.2) defines a bounded linear operator on $H^{\frac{1}{2}}(\partial\Omega; \mathbb{R}^3)$ and for every $\varphi \in H^1(\Omega; \text{curl})$ formula (2.2) holds valid provided that the right hand-side is understood in a suitable weak sense, replacing the boundary integral with a duality pairing.

The closed subspace $H_0^1(\Omega; \text{curl})$ of all $\psi \in H^1(\Omega; \text{curl})$ for which, in the previous weak sense, we have $\psi \times n = 0$ on $\partial\Omega$ is also a Hilbert space with respect to (2.1).

Throughout the paper, the spaces of L^2 scalar-valued, vector-valued, and tensor-valued functions will be denoted by $L^2(\Omega)$, $L^2(\Omega; \mathbb{R}^3)$, $L^2(\Omega; \mathbb{R}^{3 \times 3})$, respectively. For the sake of readability, we shall denote by $(\cdot, \cdot)_{L^2}$ and $\|\cdot\|_{L^2}$ the scalar product and the norm in all these spaces.

2.1. Regularity of the domain. An open set Ω is said to satisfy the uniform two-sided ball condition with radius r if for every $z \in \partial\Omega$ there exist a ball $B_r(x)$ contained in Ω and a ball $B_r(y)$ contained in its complement with z belonging to the closure of both $B_r(x)$ and of $B_r(y)$. If that is the case and we assume, in addition, that $\partial\Omega = \partial(\overline{\Omega})$, then Ω is a locally $C^{1,1}$ -domain, i.e., for every $z \in \partial\Omega$ there exist two positive constants $\rho_0, L_0 > 0$, and a rigid change of coordinates in \mathbb{R}^3 , under which $z = 0$ and

$$\Omega \cap B_{\rho_0}(0) = \{y \in B_{\rho_0}(0) : y_3 > \varphi(y_1, y_2)\},$$

for some $C^{1,1}$ function φ on $B'_{\rho_0} = \{(y_1, y_2) \in \mathbb{R}^2 : y_1^2 + y_2^2 < \rho_0^2\}$, with $\varphi(0) = |\nabla\varphi(0)| = 0$, such that

$$\|\varphi\|_{L^\infty(B'_{\rho_0})} + \rho_0 \|\nabla\varphi\|_{L^\infty(B'_{\rho_0})} + \rho_0^2 \text{Lip}(\nabla\varphi; B'_{\rho_0}) \leq L_0 \rho_0,$$

where

$$\text{Lip}(\nabla\varphi; B'_{\rho_0}) = \sup_{\substack{x, y \in B'_{\rho_0} \\ y \neq x}} \frac{|\nabla\varphi(x) - \nabla\varphi(y)|}{|x - y|}.$$

If Ω is bounded and the property described above holds with constants ρ_0, L_0 independent of z , then we say that Ω is of class $C^{1,1}$ with constants ρ_0, L_0 . In that case, it is easily seen that Ω satisfies the uniform two-sided ball condition with radius r , provided that $r < \min\{1, L_0^{-1}\}\rho_0$.

Throughout this paper we shall always assume the following condition to be in force:

$$(2.3) \quad \Omega \text{ is bounded, with uniform two-sided ball condition with radius } r, \text{ and } \partial\Omega = \partial(\overline{\Omega}).$$

We observe that (2.3) implies that Ω is of class $C^{1,1}$ with appropriate constants ρ_0, L_0 , satisfying $L_0 r < \rho_0$ (see [5, Corollary 3.14]), and we shall assume that $\rho_0 = 1$ with no loss of generality.

2.2. Gaffney inequality. The following result is proved in [14] in the case of domains with smooth boundaries but its validity is also well known on open sets satisfying assumption (2.3) (see, e.g., [10]).

Lemma 2.1 (Gaffney inequality). *Let $\psi \in L^2(\Omega; \mathbb{R}^3)$, with $\nabla \cdot \psi \in L^2(\Omega)$ and $\nabla \times \psi \in L^2(\Omega; \mathbb{R}^3)$. If either $\psi \times n = 0$ in $H^{-\frac{1}{2}}(\partial\Omega; \mathbb{R}^3)$ or $\psi \cdot n = 0$ in $H^{-\frac{1}{2}}(\partial\Omega)$, then $\psi \in H^1(\Omega; \mathbb{R}^3)$. Moreover,*

$$(2.4) \quad \int_{\Omega} (\nabla \cdot \psi)^2 dx + \int_{\Omega} |\nabla \times \psi|^2 dx + \int_{\Omega} |\psi|^2 dx \geq C \int_{\Omega} |\nabla \psi|^2 dx,$$

where the constant C depends on r , only.

For every $\mu \in L^\infty(\Omega)$, we set

$$(2.5) \quad X_\mu = \left\{ \psi \in L^2(\Omega; \mathbb{R}^3) : \int_{\Omega} \mu \psi \cdot \nabla u dx = 0, \text{ for all } u \in H_0^1(\Omega) \right\}, \quad Y_\mu = H_0^1(\Omega; \text{curl}) \cap X_\mu.$$

If $\mu = 1$ then, to shorten the notation, we write X, Y instead of X_μ, Y_μ .

Clearly if (1.1i) holds then X_μ is a Hilbert space with respect to the $L^2(\mu)$ -scalar product, i.e.

$$(2.6) \quad (\varphi, \psi)_{X_\mu} := \int_{\Omega} \mu \varphi \cdot \psi dx, \quad \text{for all } \varphi, \psi \in X_\mu.$$

The space Y_μ is closed in $H_0^1(\Omega; \text{curl})$ with respect to the topology induced by (2.6) which in fact is the standard topology of $L^2(\Omega; \mathbb{R}^3)$, as $\mu \in L^\infty(\Omega)$. It is straightforward to deduce the following result from Lemma 2.1.

Lemma 2.2. *Let Ω satisfy the uniform interior and exterior ball condition with radius r and let μ satisfy (1.1i). Then, every $\psi \in Y_\mu$ belongs to the Sobolev space $H^1(\Omega; \mathbb{R}^3)$ and we have*

$$\int_{\Omega} |\nabla \psi|^2 dx \leq C \left(\int_{\Omega} |\psi|^2 dx + \int_{\Omega} |\nabla \times \psi|^2 dx \right),$$

for a suitable constant C , depending only on Ω and r .

Remark 2.3. By Lemma 2.2, if (1.1i) holds then the norm

$$\|\psi\|_{Y_\mu} := \left(\int_{\Omega} \mu |\psi|^2 dx + \int_{\Omega} \mu |\nabla \times \psi|^2 dx \right)^{\frac{1}{2}},$$

is equivalent to that induced on Y_μ by $H^1(\Omega; \mathbb{R}^3)$.

Remark 2.4. By Remark 2.3, the compactness of the embedding of $H^1(\Omega; \mathbb{R}^3)$ into $L^2(\Omega; \mathbb{R}^3)$ implies that the embedding of Y_μ into X_μ is compact if condition (1.1i) holds.

2.3. Helmholtz decomposition. We shall make use of the following Helmholtz-type decompositions. The interested reader may find in the appendix their proofs, that are however standard.

Lemma 2.5. *Let $\mathbf{F} \in L^2(\Omega; \mathbb{R}^3)$. Then there exist $u \in H^1(\Omega)$ and $\eta \in L^2(\Omega; \mathbb{R}^3)$ such that*

$$(2.7a) \quad \mathbf{F} = \nabla u + \eta,$$

$$(2.7b) \quad \int_{\Omega} \eta \cdot \nabla v dx = 0, \quad \text{for all } v \in H^1(\Omega),$$

$$(2.7c) \quad \max \left\{ \|\nabla u\|_{L^2}, \|\eta\|_{L^2} \right\} \leq \|\mathbf{F}\|_{L^2}.$$

If in addition $\mathbf{F} \in H^1(\Omega; \text{curl})$, then $\eta \in H^1(\Omega; \mathbb{R}^3)$ and $\|\nabla \eta\|_{L^2} = \|\nabla \times \mathbf{F}\|_{L^2}$.

Lemma 2.6. *Let μ satisfy (1.1i). Given $\mathbf{F} \in L^2(\Omega; \mathbb{R}^3)$, let $q \in H_0^1(\Omega)$ be the solution of the problem*

$$(2.8) \quad \int_{\Omega} \mu \nabla q \cdot \nabla v \, dx = \int_{\Omega} \mu \mathbf{F} \cdot \nabla v \, dx, \quad \text{for all } v \in H_0^1(\Omega).$$

Then, writing

$$(2.9a) \quad \mathbf{F} = \nabla q + \zeta,$$

we have $\zeta \in X_{\mu}$ and

$$(2.9b) \quad \|\nabla q\|_{L^2} \leq \Lambda \|\mathbf{F}\|_{L^2}, \quad \|\zeta\|_{L^2} \leq \Lambda \|\mathbf{F}\|_{L^2}.$$

Moreover, if $\mathbf{F} \in H^1(\Omega; \mathbb{R}^3)$, with $\mathbf{F} \times n = 0$ in $H^{-\frac{1}{2}}(\partial\Omega; \mathbb{R}^3)$, then $q \in H^2(\Omega) \cap H_0^1(\Omega)$ and we may take $\zeta \in Y_{\mu}$.

Remark 2.7. Clearly Lemma 2.6 is valid also if μ is replaced by any other function for which property (1.1i) holds true; for example, it applies to constants. More precisely, we can decompose any L^2 vector field in the form $\mathbf{F} = \nabla q + \zeta$, where $q \in H_0^1(\Omega)$ is the weak solution of $\Delta q = \nabla \cdot \mathbf{F}$. In this case, ζ has null (distributional) divergence, and if \mathbf{F} belongs to $H^1(\Omega; \mathbb{R}^3)$ then so does ζ .

2.4. Weak formulation. We fix a Lipschitz continuous function μ satisfying (1.1i), we define the spaces X_{μ} , X , Y_{μ} , and Y , as in (2.5), and we denote by Y'_{μ} the dual space of Y_{μ} . For $p \in [1, +\infty]$ and for every Hilbert space Z we denote by $L^p(0, T; Z)$ the space of all measurable functions $\mathbf{F}: [0, T] \rightarrow Z$ such that

$$\|\mathbf{F}\|_{L^p(0, T; Z)} := \begin{cases} \left(\int_0^T \|\mathbf{F}(t)\|_Z^p \, dt \right)^{\frac{1}{p}} & \text{if } p < +\infty, \\ \text{ess sup}_{t \in [0, T]} \|\mathbf{F}(t)\|_Z & \text{if } p = +\infty, \end{cases}$$

is finite. We recall that $L^p(0, T; Z)$ is a Banach space (uniformly convex if $p < +\infty$). We shall need the following generalisation of a well known property of Sobolev space-valued mappings. For a proof, one can repeat verbatim the argument used in the proof of the analogous result in Sobolev spaces, see [13, Theorem 3, §5.9.2].

Proposition 2.8. *Suppose that $\mathbf{F} \in L^2(0, T; Y_{\mu})$, with $\partial_t \mathbf{F} \in L^2(0, T; Y'_{\mu})$. Then, by possibly redefining it on a negligible subset of $(0, T)$, the function \mathbf{F} belongs to $C([0, T]; X_{\mu})$. Moreover, the mapping $t \mapsto \|\mathbf{F}(t)\|_{X_{\mu}}^2$ is absolutely continuous and for a.e. $t \in (0, T)$ we have*

$$\frac{1}{2} \frac{d}{dt} \|\mathbf{F}(t)\|_{X_{\mu}}^2 = \langle \partial_t \mathbf{F}(t), \mathbf{F}(t) \rangle_{Y'_{\mu} \times Y_{\mu}}.$$

Eventually, there exists a constant C , depending only on T , such that

$$\sup_{t \in [0, T]} \|\mathbf{F}(t)\|_{X_{\mu}} \leq C \left(\|\mathbf{F}\|_{L^2(0, T; Y_{\mu})} + \|\partial_t \mathbf{F}\|_{L^2(0, T; Y'_{\mu})} \right).$$

If $\mathbf{F} \in L^2(0, T; H^1(\Omega; \text{curl}))$ and $\sigma \partial_t \mathbf{F} \in L^2(0, T; H^1(\Omega; \text{curl})')$ then $\mathbf{F} \in C([0, T]; L^2(\Omega; \mathbb{R}^3))$, for a.e. $t \in (0, T)$ we have

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} \sigma \mathbf{F}(t) \cdot \mathbf{F}(t) \, dx = \langle \sigma \partial_t \mathbf{F}(t), \mathbf{F}(t) \rangle,$$

where $\langle \cdot, \cdot \rangle$ denotes now the pairing between $H^1(\Omega; \text{curl})$ and its dual space $H^1(\Omega; \text{curl})'$, and

$$\sup_{t \in [0, T]} \|\mathbf{F}(t)\|_{L^2} \leq C \left(\|\mathbf{F}\|_{L^2(0, T; H^1(\Omega; \text{curl}))} + \|\partial_t \mathbf{F}\|_{L^2(0, T; H^1(\Omega; \text{curl})'} \right),$$

where the constant C depends on Λ and T , only.

Definition 2.9. Given

$$(2.10) \quad \mathbf{J}^E \in L^2(0, T; L^2(\Omega; \mathbb{R}^3)), \quad \mathbf{J}^M \in L^2(0, T; X),$$

and

$$(2.11) \quad \mathbf{H}_0 \in Y_\mu,$$

we say that $(\mathbf{E}, \mathbf{H}) \in L^2(0, T; H^1(\Omega; \text{curl}) \times H_0^1(\Omega; \text{curl}))$, with $\partial_t \mathbf{H} \in L^2(0, T; L^2(\Omega; \mathbb{R}^3))$, is a weak solution of the eddy current system

$$(2.12) \quad \begin{cases} \nabla \times \mathbf{H} - \sigma \mathbf{E} = \mathbf{J}^E, & \text{in } \Omega \times (0, T), \\ \nabla \times \mathbf{E} + \mu \partial_t \mathbf{H} = \mathbf{J}^M, & \text{in } \Omega \times (0, T), \\ \mathbf{H} \times n = 0, & \text{on } \partial\Omega \times (0, T), \\ \mathbf{H} = \mathbf{H}_0, & \text{in } \Omega \times \{0\}, \end{cases}$$

if for all $\varphi \in H^1(\Omega; \text{curl})$ and for all $\psi \in H_0^1(\Omega; \text{curl})$ we have

$$(2.13i) \quad \int_{\Omega} \mathbf{H} \cdot \nabla \times \varphi \, dx - \int_{\Omega} \sigma \mathbf{E} \cdot \varphi \, dx = \int_{\Omega} \mathbf{J}^E \cdot \varphi \, dx$$

$$(2.13ii) \quad \int_{\Omega} \mathbf{E} \cdot \nabla \times \psi \, dx + \int_{\Omega} \mu \partial_t \mathbf{H} \cdot \psi \, dx = \int_{\Omega} \mathbf{J}^M \cdot \psi \, dx$$

for a.e. $t \in [0, T]$, and in addition we have

$$(2.13iii) \quad \mathbf{H}(0) = \mathbf{H}_0.$$

Remark 2.10. We note that (2.10), (2.11), (2.13ii), and (2.13iii) imply that $\mathbf{H} \in L^2(0, T; Y_\mu)$ and $\partial_t \mathbf{H} \in L^2(0, T; X_\mu)$. Then, $\partial_t \mathbf{H} \in L^2(0, T; Y'_\mu)$, due to the isometric embedding of X_μ into the dual Y'_μ of Y_μ . Hence, in view of Proposition 2.8, we see that $\mathbf{H} \in C([0, T]; X_\mu)$ and thus equality (2.13iii) makes sense.

Remark 2.11. Let equation (2.13ii) hold for all $\psi \in Y_\mu$. Then, it holds for all $\psi \in H_0^1(\Omega; \text{curl})$. Indeed, by Lemma 2.6 we can write every $\psi \in C_0^1(\Omega; \mathbb{R}^3)$ in the form $\psi = \nabla q + \zeta$ where $\zeta \in Y_\mu$ and

$$(2.14) \quad \int_{\Omega} \mathbf{E} \cdot \nabla \times (\nabla q) \, dx = \int_{\Omega} \mu \partial_t \mathbf{H} \cdot \nabla q \, dx = \int_{\Omega} \mathbf{J}^M \cdot \nabla q \, dx = 0,$$

because $\nabla \times (\nabla q) = 0$, and $\mu \partial_t \mathbf{H}, \mathbf{J}^M \in X$ for a.e. $t \in (0, T)$. Then, (2.13ii) holds for all test fields in $C_0^1(\Omega; \mathbb{R}^3)$, which by [12, Remark 4.2] is dense in $H_0^1(\Omega; \text{curl})$.

Formally, in view of the integration by parts formula (2.2), a weak solution in the sense of Definition 2.9 is a solution to (1.2) with $\mathbf{G} = 0$, satisfying the additional condition $\nabla \cdot (\mu \mathbf{H}) = 0$. Weak solutions in case of non-homogeneous boundary conditions are defined in the following sense.

Definition 2.12. Given $\mathbf{J}^E, \mathbf{J}^M \in L^2(0, T; L^2(\Omega; \mathbb{R}^3))$, given $\mathbf{H}_0 \in H^1(\Omega; \text{curl})$, and given $\mathbf{G} \in L^2(0, T; H^1(\Omega; \text{curl}))$, with $\partial_t \mathbf{G} \in L^2(0, T; L^2(\Omega; \mathbb{R}^3))$, such that for a.e. $t \in (0, T)$ we have

$$(2.15) \quad \int_{\Omega} \left(\mu \mathbf{G}(x, t) - \mu \mathbf{H}_0(x) - \int_0^t \mathbf{J}^M(x, s) ds \right) \cdot \nabla u(x) dx = 0,$$

for all $u \in H_0^1(\Omega)$, we say that $(\mathbf{E}, \mathbf{H}) \in L^2(0, T; H^1(\Omega; \text{curl})^2)$, with $\partial_t \mathbf{H} \in L^2(0, T; L^2(\Omega; \mathbb{R}^3))$, is a *weak solution of the eddy current system (1.2)* if $\mathbf{F} := \mathbf{H} - \mathbf{G}$ belongs to $L^2(0, T; H_0^1(\Omega; \text{curl}))$ and (\mathbf{E}, \mathbf{F}) solves, in the sense of Definition 2.9, the system

$$(2.16) \quad \begin{cases} \nabla \times \mathbf{F} - \sigma \mathbf{E} = \mathbf{J}^E - \nabla \times \mathbf{G} & \text{in } \Omega \times (0, T), \\ \nabla \times \mathbf{E} + \mu \partial_t \mathbf{F} = \mathbf{J}^M - \mu \partial_t \mathbf{G} & \text{in } \Omega \times (0, T), \\ \mathbf{F} \times n = 0 & \text{on } \partial\Omega \times (0, T), \\ \mathbf{F} = \mathbf{H}_0 - \mathbf{G} & \text{in } \Omega \times \{0\}. \end{cases}$$

We observe that Definition 2.12 makes sense, because under the assumptions made in Definition 2.12 on $\mathbf{J}^E, \mathbf{J}^M, \mathbf{H}_0$, and \mathbf{G} , it makes sense to consider weak solutions of (2.12) in the sense of Definition 2.9, relative to the sources

$$\widetilde{\mathbf{J}}^E = \mathbf{J}^E - \nabla \times \mathbf{G}, \quad \widetilde{\mathbf{J}}^M = \mathbf{J}^M - \mu \partial_t \mathbf{G},$$

and to the initial datum

$$\widetilde{\mathbf{H}}_0 = \mathbf{H}_0 - \mathbf{G}(0).$$

Indeed, by (2.15), $\widetilde{\mathbf{J}}^E, \widetilde{\mathbf{J}}^M$ satisfy conditions (2.10). Moreover, since $\mathbf{G} \in L^2(0, T; H^1(\Omega; \text{curl}))$ and $\partial_t \mathbf{G} \in L^2(0, T; L^2(\Omega; \mathbb{R}^3))$, arguing as done in Remark 2.10 we see that \mathbf{G} belongs to $C([0, T]; L^2(\Omega; \mathbb{R}^3))$, hence $\widetilde{\mathbf{H}}_0$ is well-defined. Eventually, again by (2.15), $\widetilde{\mathbf{H}}_0$ satisfies (2.11).

3. EXISTENCE AND UNIQUENESS OF SOLUTIONS

The goal of the present section is to prove the following result.

Theorem 3.1. *Let $\mathbf{H}_0 \in Y_\mu$, let $\mathbf{J}^E \in L^2(0, T; L^2(\Omega; \mathbb{R}^3))$, with $\partial_t \mathbf{J}^E \in L^2(0, T; L^2(\Omega; \mathbb{R}^3))$, and let $\mathbf{J}^M \in L^2(0, T; X)$. Then, there exists a unique weak solution (\mathbf{E}, \mathbf{H}) of (2.12). Moreover,*

$$(3.1) \quad \begin{aligned} & \sup_{t \in [0, T]} \|\mathbf{E}(t)\|_{L^2}^2 + \sup_{t \in [0, T]} \|\mathbf{H}(t)\|_{L^2}^2 + \int_0^T \|\partial_t \mathbf{H}(t)\|_{L^2}^2 dt \\ & \leq C \left(\|\mathbf{H}_0\|_{H^1(\Omega; \text{curl})}^2 + \|\mathbf{J}^E(0)\|_{L^2}^2 + \int_0^T [\|\mathbf{J}^E(t)\|_{L^2}^2 + \|\mathbf{J}^M(t)\|_{L^2}^2 + \|\partial_t \mathbf{J}^E(t)\|_{L^2}^2] dt \right), \end{aligned}$$

where the constant C depends on Λ, T , only.

Remark 3.2. When considering initial data \mathbf{H}_0 that belong merely to X_μ , it is still possible to define solutions of (1.2) in a weaker sense than that of Definition 2.9, just requiring $\partial_t \mathbf{H}$ to take values in Y'_μ rather than in X_μ , and replacing the scalar product $(\partial_t \mathbf{H}, \psi)_{X_\mu}$ in the left hand-side of (2.13i) with the duality pairing $\langle \partial_t \mathbf{H}, \psi \rangle_{Y'_\mu \times Y_\mu}$. For a given $\mathbf{H}_0 \in X_\mu \setminus Y_\mu$, the existence of solutions (\mathbf{E}, \mathbf{H}) in this weaker sense could be proved arguing similarly as done below to prove Theorem 3.1,

except that the final apriori estimate would be the following one

$$(3.2) \quad \int_0^T \|\mathbf{E}(t)\|_{L^2}^2 dt + \sup_{t \in [0, T]} \|\mathbf{H}(t)\|_{L^2}^2 + \int_0^T \|\partial_t \mathbf{H}(t)\|_{Y'_\mu}^2 dt \leq C \left(\|\mathbf{H}_0\|_{L^2}^2 + \int_0^T [\|\mathbf{J}^E(t)\|_{L^2}^2 + \|\mathbf{J}^M(t)\|_{L^2}^2] dt \right),$$

for a suitable constant C , again depending on Λ and T , only.

3.1. Magnetic eigenbase. We fix $\mu \in W^{1, \infty}(\Omega)$ satisfying conditions (1.1i).

Lemma 3.3. *The space Y_μ is dense in X_μ , with respect to the weak convergence in X_μ .*

Proof. We fix $\phi \in X_\mu$. By standard density results, there exists a sequence $(\phi_i) \subset C_0^1(\Omega; \mathbb{R}^3)$ with

$$(3.3) \quad \lim_{i \rightarrow \infty} \int_\Omega (\phi - \phi_i) \cdot \eta dx = 0, \quad \text{for all } \eta \in L^2(\Omega; \mathbb{R}^3).$$

By Lemma 2.6, there exist $(q_i) \subset H^2(\Omega) \cap H_0^1(\Omega)$ and $(\zeta_i) \subset Y_\mu$ with $\phi_i = \nabla q_i + \zeta_i$, and we have

$$(3.4) \quad \int_\Omega \mu \nabla q_i \cdot \nabla v dx = \int_\Omega \mu \phi_i \cdot \nabla v dx, \quad \text{for all } v \in H_0^1(\Omega).$$

We prove that (ζ_i) converges to ϕ weakly in $L^2(\Omega; \mathbb{R}^3)$. To do so, by (3.3), it suffices to prove

$$(3.5) \quad \lim_{i \rightarrow \infty} \int_\Omega \mu \nabla q_i \cdot \eta dx = 0,$$

for all $\tilde{\eta} \in L^2(\Omega; \mathbb{R}^3)$. We fix a test field $\tilde{\eta}$ and, using again Lemma 2.6, we write $\tilde{\eta} = \nabla q + \zeta$ for suitable $q \in H_0^1(\Omega)$ and $\zeta \in X_\mu$. Inserting $v = q$ in (3.4) we obtain

$$\int_\Omega \mu \nabla q_i \cdot \nabla q dx = \int_\Omega \mu \phi_i \cdot \nabla q dx.$$

Passing to the limit in the latter, using (3.3), and recalling that $\phi \in X_\mu$, we get

$$(3.6) \quad \lim_{i \rightarrow \infty} \int_\Omega \mu \nabla q_i \cdot \nabla q dx = \int_\Omega \mu \phi \cdot \nabla q dx = 0.$$

Since $q_i \in H_0^1(\Omega)$ for all $i \in \mathbb{N}$ and $\zeta \in X_\mu$, we also have

$$(3.7) \quad \lim_{i \rightarrow \infty} \int_\Omega \mu \nabla q_i \cdot \zeta dx = 0.$$

Summing (3.6) and (3.7) and recalling that $\tilde{\eta} = \nabla q + \zeta$ we get (3.5). Since $\tilde{\eta}$ was arbitrary, we deduce that (ζ_i) converges to ϕ weakly in $L^2(\Omega; \mathbb{R}^3)$. By (1.1i), this implies that (ζ_i) converges to ϕ with respect to the weak topology in X_μ relative to the scalar product (2.6), too, as desired. \square

The proof of the following spectral decomposition is based on standard methods, but we present it for sake of completeness.

Lemma 3.4. *There exists a sequence $0 \leq \lambda_1 \leq \lambda_2 \leq \dots$, with $\lambda_i \rightarrow +\infty$ as $i \rightarrow \infty$, and a sequence $(\psi_i) \subset Y_\mu$, such that (ψ_i) is a complete orthonormal system in X_μ and for all $i \in \mathbb{N}$ we have*

$$(3.8) \quad \int_\Omega \mu \nabla \times \psi_i \cdot \nabla \times \phi dx = \lambda_i \int_\Omega \mu \psi_i \cdot \phi dx, \quad \text{for all } \phi \in H^1(\Omega; \mathbb{R}^3),$$

and $\psi_i \times n = 0$ in $H^{-\frac{1}{2}}(\partial\Omega)$. Moreover, for every $i, j \in \mathbb{N}$ we have

$$(3.9) \quad \int_{\Omega} \mu \nabla \times \psi_i \cdot \nabla \times \psi_j \, dx = \lambda_j \delta_{ij}$$

where $\delta_{ij} = 1$ if $i = j$ and $\delta_{ij} = 0$ otherwise.

Proof. By Remark 2.3 and Lax-Milgram Lemma, the linear operator \mathcal{R} from X_μ to X_μ that takes every $\mathbf{F} \in X_\mu$ to the corresponding solution $\psi \in Y_\mu$ of the following variational problem

$$(3.10) \quad \int_{\Omega} \mu \nabla \times \psi \cdot \nabla \times \phi \, dx + \int_{\Omega} \mu \psi \cdot \phi \, dx = \int_{\Omega} \mu \mathbf{F} \cdot \phi \, dx, \quad \text{for all } \phi \in Y_\mu,$$

is well defined. Moreover, for every $\mathbf{F} \in X_\mu$, plugging in $\psi = \mathcal{R}\mathbf{F}$ in (3.10) yields

$$(3.11) \quad \|\mathcal{R}\mathbf{F}\|_{Y_\mu} \leq \|\mathbf{F}\|_{X_\mu}.$$

Clearly $\|\mathcal{R}\mathbf{F}\|_{Y_\mu} \geq \|\mathcal{R}\mathbf{F}\|_{X_\mu}$. Then, by (3.11), \mathcal{R} has operator norm bounded by 1.

We observe that \mathcal{R} is injective. Indeed, by definition if \mathbf{F} belongs to the kernel of \mathcal{R} then $\psi = 0$ is the solution of (3.10). Thus, $(\mathbf{F}, \phi)_{X_\mu} = 0$ for all $\phi \in Y_\mu$. By Lemma 3.3, the latter holds in fact for all $\phi \in X_\mu$, hence $\mathbf{F} = 0$.

Also, $(\mathbf{F}, \mathcal{R}\mathbf{F})_{X_\mu} \geq 0$ and $(\mathbf{F}, \mathcal{R}\mathbf{G})_{X_\mu} = (\mathbf{G}, \mathcal{R}\mathbf{F})_{X_\mu}$, for every $\mathbf{F}, \mathbf{G} \in X_\mu$, i.e., \mathcal{R} is a positive and symmetric operator.

In addition, \mathcal{R} is compact. Indeed, given a bounded sequence $(\mathbf{F}_i) \subset X_\mu$, the sequence $(\mathcal{R}\mathbf{F}_i)$ is bounded in Y_μ by (3.11). By Remark 2.4, it follows that $(\mathcal{R}\mathbf{F}_i)$ is precompact in X_μ .

Therefore, \mathcal{R} is a positive, compact, self-adjoint operator with trivial kernel from X_μ to itself, having operator norm bounded by 1. By the Spectral Theorem, there exists a sequence $(\tau_i) \subset (0, 1]$ and a Hilbert basis (ψ_i) of X_μ with $\psi_i \in Y_\mu$ and $\mathcal{R}\psi_i = \tau_i \psi_i$ for all $i \in \mathbb{N}$, and the first statement follows just setting $\lambda_i = \tau_i^{-1} - 1$.

Eventually, we fix $i, j \in \mathbb{N}$, we test equation (3.8) with $\phi = \psi_j$, and we get

$$\int_{\Omega} \mu \nabla \times \psi_i \cdot \nabla \times \psi_j \, dx = \lambda_i \int_{\Omega} \mu \psi_i \cdot \psi_j \, dx.$$

Since (ψ_i) is orthonormal in X_μ with respect to (2.6), this gives (3.9) and concludes the proof. \square

Remark 3.5. Incidentally, Lemma 3.4, implies in particular that the vector space

$$(3.12) \quad H_\mu = \left\{ h \in L^2(\Omega; \mathbb{R}^3) : \nabla \cdot (\mu h) = 0, \nabla \times h = 0, h \times n = 0 \right\}$$

is finite-dimensional, because it consists of solutions of (3.8) corresponding to the null eigenvalue. In other words, the least eigenvalue either equals zero or is positive depending on whether or not Ω supports non-trivial vector fields within (3.12).

We note that (3.12) is trivial if Ω is *contractible*, i.e., if there exists $x_0 \in \Omega$ and a function $g \in C^\infty([0, 1] \times \Omega; \Omega)$ with $g(0, \cdot) = \text{id}_\Omega$ and $g(1, x) = x_0$ for all $x \in \Omega$. For example, Ω has this property if it is simply connected and $\partial\Omega$ is connected; in this case, every $h \in H_\mu$ is the gradient of a scalar potential w , and w is a weak solution of the elliptic equation $\nabla \cdot (\mu \nabla w) = 0$ with homogeneous Dirichlet boundary conditions, hence it is a constant.

To prove Theorem 3.1, we observe that $H^1(\Omega; \text{curl})$, with the scalar product induced by (2.1), is a separable Hilbert space. Thus it admits a complete orthonormal system; we pick one, and we denote it by (φ_i) . Then, let (ψ_i) be the complete orthonormal system of X_μ introduced in Section 3.1, with (λ_i) being the sequence of all corresponding eigenvalues, counted with multiplicity.

3.2. Approximate solutions. Given $\mathbf{J}^E \in L^2(0, T; L^2(\Omega; \mathbb{R}^3))$, $\mathbf{J}^M \in L^2(0, T; X)$, and $\mathbf{H}_0 \in X_\mu$, we set

$$(3.13) \quad \mathbf{H}_{0m} = \sum_{j=1}^m (\mathbf{H}_0, \psi_j)_{X_\mu} \psi_j,$$

and following Galerkin's scheme, we seek approximate solutions having the structure

$$(3.14) \quad \mathbf{E}_m(t) = \sum_{j=1}^m e_{jm}(t) \varphi_j, \quad \mathbf{H}_m(t) = \sum_{j=1}^m h_{jm}(t) \psi_j.$$

More precisely, we prescribe the validity of the following $2m$ equations

$$(3.15i) \quad \int_{\Omega} \nabla \times \mathbf{H}_m \cdot \varphi_i \, dx - \int_{\Omega} \sigma \mathbf{E}_m \cdot \varphi_i \, dx = \int_{\Omega} \mathbf{J}^E \cdot \varphi_i \, dx, \quad i = 1, 2, \dots, m$$

$$(3.15ii) \quad \int_{\Omega} \nabla \times \mathbf{E}_m \cdot \psi_i \, dx + \int_{\Omega} \mu \partial_t \mathbf{H}_m \cdot \psi_i \, dx = \int_{\Omega} \mathbf{J}^M \cdot \psi_i \, dx, \quad i = 1, 2, \dots, m$$

and of the initial conditions

$$(3.16) \quad \mathbf{H}_m(0) = \mathbf{H}_{0m}.$$

Lemma 3.6. *Let $\mathbf{H}_0 \in X_\mu$. Then, there exists a unique solution*

$$(3.17) \quad (\mathbf{E}_m, \mathbf{H}_m) \in C^1([0, T]; \text{Span}\{\varphi_1, \dots, \varphi_m\} \times \text{Span}\{\psi_1, \dots, \psi_m\})$$

of the system (3.15) satisfying (3.16). If in addition we have $\mathbf{H}_0 \in Y_\mu$, then

$$(3.18) \quad \|\mathbf{E}_m(0)\|_{L^2} \leq C(\|\nabla \times \mathbf{H}_0\|_{L^2} + \|\mathbf{J}^E(0)\|_{L^2}).$$

for a constant C depending only on Λ .

Proof. We write the system (3.15) in the form

$$(3.19) \quad \begin{aligned} (\nabla \times \mathbf{H}_m, \varphi_i)_{L^2} - (\sigma \mathbf{E}_m, \varphi_i)_{L^2} &= (\mathbf{J}^E, \varphi_i)_{L^2}, \\ (\mu^{-1} \nabla \times \mathbf{E}_m, \psi_i)_{X_\mu} + (\partial_t \mathbf{H}_m, \psi_i)_{X_\mu} &= (\mu^{-1} \mathbf{J}^M, \psi_i)_{X_\mu}. \end{aligned}$$

Seeking solution with the structure (3.14) we are led to the $2m$ equations

$$(3.20a) \quad \sum_{j=1}^m (\nabla \times \psi_j, \varphi_i)_{L^2} h_{jm}(t) - \sum_{j=1}^m (\sigma \varphi_j, \varphi_i)_{L^2} e_{jm}(t) = (\mathbf{J}^E(t), \varphi_i)_{L^2} \quad \text{for } i = 1, \dots, m$$

$$(3.20b) \quad \sum_{j=1}^m (\nabla \times \varphi_j, \psi_i)_{L^2} e_{jm}(t) + \sum_{j=1}^m (\mu \psi_j, \psi_i)_{L^2} \frac{d}{dt} h_{jm}(t) = (\mathbf{J}^M(t), \psi_i)_{L^2} \quad \text{for } i = 1, \dots, m.$$

By (1.1ii) and thanks to the fact that (φ_i) is a linearly independent system, the quadratic form defined on \mathbb{R}^m by

$$(3.21) \quad \mathcal{Q}(v) = \sum_{i,j=1}^m (\sigma \varphi_j, \varphi_i)_{L^2} v_i v_j, \quad \text{for all } v \in \mathbb{R}^m,$$

is positive definite and $\mathcal{Q}(v) \geq \Lambda^{-1}|v|^2$, for all $v \in \mathbb{R}^m$. The matrix $\{(\sigma \varphi_j, \varphi_i)_{L^2}\}_{i,j=1}^m$ is symmetric because so is σ . Moreover, it is invertible and, denoting by M^σ the inverse matrix (which is also symmetric), we have

$$(3.22) \quad |M^\sigma v| \leq \Lambda |v|^2, \quad \text{for all } v \in \mathbb{R}^m.$$

Then, (3.20a) becomes

$$(3.23) \quad e_{im}(t) = \sum_{j,k=1}^m M_{ik}^\sigma(\nabla \times \psi_j, \varphi_k)_{L^2} h_{jm}(t) - \sum_{j=1}^m M_{ij}^\sigma(\mathbf{J}^E(t), \varphi_j)_{L^2}, \quad i = 1, \dots, m.$$

Since (ψ_i) is an orthonormal system in X_μ with respect to the scalar product introduced in (2.6), $(\mu\psi_i, \psi_j)_{L^2} = \delta_{ij}$ for all $i, j = 1, \dots, m$. Then (3.20b) gives

$$(3.24) \quad \frac{d}{dt} h_{im} = - \sum_{j=1}^m (\nabla \times \varphi_j, \psi_i)_{L^2} e_{jm} + (\mathbf{J}^M, \psi_i)_{L^2}, \quad i = 1, \dots, m.$$

Using (3.20a) to get rid of e_{jm} in (3.24), we obtain

$$(3.25) \quad \begin{aligned} \frac{d}{dt} h_{im} = & - \sum_{j,k,\ell=1}^m (\nabla \times \varphi_j, \psi_i)_{L^2} M_{jk}^\sigma(\nabla \times \psi_\ell, \varphi_k)_{L^2} h_{\ell m} \\ & + \sum_{j,k=1}^m (\nabla \times \varphi_j, \psi_i)_{L^2} M_{jk}^\sigma(\mathbf{J}^E, \varphi_k)_{L^2} + (\mathbf{J}^M, \psi_i)_{L^2}, \quad i = 1, \dots, m. \end{aligned}$$

We set $\vec{e}_m = (e_{11}, \dots, e_{1m})$ and $\vec{h}_m = (h_{1m}, \dots, h_{mm})$. We observe that, by (2.2), for all $i, j = 1, \dots, m$ the scalar products $(\nabla \times \psi_j, \varphi_i)_{L^2}$ and $(\nabla \times \varphi_i, \psi_j)_{L^2}$ are equal and we denote by A_{ij} their common value. Then, the m equations appearing in (3.25) can be recast in the form

$$(3.26) \quad \frac{d}{dt} \vec{h}_m = -A^T M^\sigma A h_m + \vec{b}_m,$$

for a suitable $\vec{b}_m \in L^2([0, T]; \mathbb{R}^m)$. By the standard existence theory for linear systems, there exists $\vec{h}_m \in C^1([0, T]; \mathbb{R}^m)$ that solves (3.26) for a.e. $t \in (0, T)$, with the initial conditions

$$\vec{h}_m(0) = ((\mathbf{H}_0, \psi_1)_{X_\mu}, \dots, (\mathbf{H}_0, \psi_m)_{X_\mu}).$$

Then, we use (3.23) to define $\vec{e}_m \in C^1([0, T]; \mathbb{R}^m)$. Therefore, by construction the functions \mathbf{E}_m and \mathbf{H}_m introduced in (3.14) are such that (3.19) is valid, and the initial conditions (3.16) hold.

Now, we assume that $\mathbf{H}_0 \in Y_\mu$. By (3.14) and (3.21), we have

$$(3.27) \quad (\sigma \mathbf{E}_m, \mathbf{E}_m)_{L^2} = \mathcal{Q}(\vec{e}_m).$$

Then we observe that (3.23) implies

$$(3.28) \quad \mathcal{Q}(\vec{e}_m) = A \vec{h}_m \cdot \vec{e}_m - \sum_{i=1}^m (\mathbf{J}^E, \varphi_i)_{L^2} e_{im} = (\nabla \times \mathbf{H}_m, \mathbf{E}_m)_{L^2} - (\mathbf{J}^E, \mathbf{E}_m)_{L^2},$$

where in the second equality we simply used (3.14). Since (3.27) and (3.28) holds, in particular, for $t = 0$, we deduce that

$$(3.29) \quad (\sigma \mathbf{E}_m(0), \mathbf{E}_m(0))_{L^2} = (\nabla \times \mathbf{H}_{0m}, \mathbf{E}_m(0))_{L^2} - (\mathbf{J}^E(0), \mathbf{E}_m(0))_{L^2}.$$

By Cauchy-Schwartz inequality, we have

$$(\nabla \times \mathbf{H}_{0m}, \mathbf{E}_m(0))_{L^2} - (\mathbf{J}^E(0), \mathbf{E}_m(0))_{L^2} \leq \left[\|\nabla \times \mathbf{H}_{0m}\|_{L^2} + \|\mathbf{J}^E(0)\|_{L^2} \right] \|\mathbf{E}_m(0)\|_{L^2}.$$

Using this and (1.1i), from (3.29) we deduce

$$(3.30) \quad \Lambda^{-1} \|\mathbf{E}_m(0)\|_{L^2} \leq \|\nabla \times \mathbf{H}_{0m}\|_{L^2} + \|\mathbf{J}^E(0)\|_{L^2}.$$

By (1.1i), we have

$$(3.31) \quad \|\nabla \times \mathbf{H}_{0m}\|_{L^2}^2 \leq \Lambda(\nabla \times \mathbf{H}_{0m}, \nabla \times \mathbf{H}_{0m})_{X_\mu}.$$

Thanks to (3.13), (3.16), and recalling (3.9), we obtain that

$$(3.32) \quad (\nabla \times \mathbf{H}_{0m}, \nabla \times \mathbf{H}_{0m})_{X_\mu} = \sum_{i,j=1}^m (\mathbf{H}_0, \psi_i)_{X_\mu} (\mathbf{H}_0, \psi_j)_{X_\mu} (\nabla \times \psi_i, \nabla \times \psi_j)_{X_\mu} = \sum_{i=1}^m \lambda_i |(\mathbf{H}_0, \psi_i)_{X_\mu}|^2.$$

Since $\mathbf{H}_0 \in Y_\mu$, by (3.8) we also have $\lambda_i(\mathbf{H}_0, \psi_i)_{X_\mu} = (\nabla \times \mathbf{H}_0, \nabla \times \psi_i)_{X_\mu}$. Hence

$$(3.33) \quad \sum_{i=1}^{\infty} \lambda_i |(\mathbf{H}_0, \psi_i)_{X_\mu}|^2 = \sum_{\lambda_i > 0} \left| (\nabla \times \mathbf{H}_0, \lambda_i^{-\frac{1}{2}} \nabla \times \psi_i)_{X_\mu} \right|^2 \leq \Lambda \|\nabla \times \mathbf{H}_0\|_{L^2}^2,$$

where in the last passage we also used Bessel's inequality and the fact that $(\lambda_i^{-1/2} \nabla \times \psi_i)$ is an orthonormal system in X_μ , by (3.9). Clearly, (3.30), (3.31), (3.32), and (3.33) imply (3.18) and this concludes the proof. \square

3.3. Energy estimates. We provide ourselves with standard a priori bounds for the approximate solutions, so as to construct weak solutions by compactness.

Proposition 3.7. *Let $\mathbf{H}_0 \in X_\mu$ and let $(\mathbf{E}_m, \mathbf{H}_m)$ be as in Lemma 3.6. Then*

$$(3.34) \quad \int_0^T \|\mathbf{E}_m(t)\|_{L^2}^2 dt + \sup_{t \in [0, T]} \|\mathbf{H}_m(t)\|_{L^2}^2 \leq C \left(\|\mathbf{H}_0\|_{L^2}^2 + \int_0^T (\|\mathbf{J}^E(t)\|_{L^2}^2 + \|\mathbf{J}^M(t)\|_{L^2}^2) dt \right),$$

for a constant $C > 0$ depending on Λ , and T , only. If in addition $\mathbf{H}_0 \in Y_\mu$ then

$$(3.35) \quad \begin{aligned} & \sup_{t \in [0, T]} \|\mathbf{E}_m(t)\|_{L^2}^2 + \sup_{t \in [0, T]} \|\mathbf{H}_m(t)\|_{L^2}^2 + \int_0^T \|\partial_t \mathbf{H}_m(t)\|_{L^2}^2 dt \\ & \leq C \left(\|\mathbf{H}_0\|_{H^1(\Omega; \text{curl})}^2 + \|\mathbf{J}^E(0)\|_{L^2}^2 + \int_0^T (\|\mathbf{J}^E(t)\|_{L^2}^2 + \|\mathbf{J}^M(t)\|_{L^2}^2 + \|\partial_t \mathbf{J}^E(t)\|_{L^2}^2) dt \right), \end{aligned}$$

for a (possibly different) constant $C > 0$ depending on Λ , and T , only.

Proof. By (3.15), for all $(\varphi, \psi) \in \text{Span}\{\varphi_1, \dots, \varphi_m\} \times \text{Span}\{\psi_1, \dots, \psi_m\}$ we have

$$(3.36i) \quad (\nabla \times \mathbf{H}_m, \varphi)_{L^2} - (\sigma \mathbf{E}_m, \varphi)_{L^2} = (\mathbf{J}^E, \varphi)_{L^2}$$

$$(3.36ii) \quad (\nabla \times \mathbf{E}_m, \psi)_{L^2} + (\mu \partial_t \mathbf{H}_m, \psi)_{L^2} = (\mathbf{J}^M, \psi)_{L^2}.$$

We divide now the proof into two steps.

Step 1. Core Energy inequality

We observe that

$$(\mu \partial_t \mathbf{H}_m, \mathbf{H}_m)_{L^2} = \frac{1}{2} \frac{d}{dt} (\mu \mathbf{H}_m, \mathbf{H}_m)_{L^2}.$$

Then, choosing $\varphi = \mathbf{E}_m$ in (3.36i) and $\psi = \mathbf{H}_m$ in (3.36ii), integrating on $(0, t)$, and using (3.16), we obtain the energy identity

$$(3.37) \quad \frac{1}{2} \|\mathbf{H}_m(t)\|_{X_\mu}^2 + \int_0^t (\sigma \mathbf{E}_m, \mathbf{E}_m)_{L^2} = \frac{1}{2} \|\mathbf{H}_{0m}\|_{X_\mu}^2 + \int_0^t (\mathbf{J}^M, \mathbf{H}_m)_{L^2} - \int_0^t (\mathbf{J}^E, \mathbf{E}_m).$$

By Cauchy Schwartz and Young inequality we have

$$(\mathbf{J}^M, \mathbf{H}_m)_{L^2} \leq \frac{1}{2} \|\mathbf{H}_m\|_{X_\mu}^2 + \frac{1}{2} \|\mu^{-1} \mathbf{J}^M\|_{X_\mu}^2.$$

Using Cauchy-Schwartz inequality for the scalar product $(\phi, \psi) \mapsto (\sigma\phi, \psi)_{L^2}$ induced by the symmetric matrix σ and then using Young's inequality again, we also have

$$(\mathbf{J}^E, \mathbf{E}_m)_{L^2} \leq \frac{1}{2} (\sigma \mathbf{E}_m, \mathbf{E}_m)_{L^2} + \frac{1}{2} (\sigma^{-1} \mathbf{J}^E, \mathbf{J}^E)_{L^2}.$$

Also, $\|\mathbf{H}_{0m}\|_{X_\mu} \leq \|\mathbf{H}_0\|_{X_\mu}$ by (3.13). Using these inequalities in (3.37), together with (1.1i), we get

$$(3.38) \quad \|\mathbf{H}_m(t)\|_{X_\mu}^2 + \int_0^t (\sigma \mathbf{E}_m, \mathbf{E}_m)_{L^2} ds \leq \|\mathbf{H}_0\|_{X_\mu}^2 + \Lambda \int_0^t (\|\mathbf{J}^E\|_{L^2}^2 + \|\mathbf{J}^M\|_{L^2}^2) ds + \int_0^t \|\mathbf{H}_m(s)\|_{X_\mu}^2 ds.$$

By (3.17), $t \mapsto \|\mathbf{H}_m(t)\|^2$ is continuous. Thus, by Grönwall's Lemma, (3.38) implies the inequality

$$(3.39) \quad \|\mathbf{H}_m(t)\|_{X_\mu}^2 + \int_0^t (\sigma \mathbf{E}_m, \mathbf{E}_m)_{L^2} ds \leq C \left[\|\mathbf{H}_0\|_{X_\mu}^2 + \Lambda \int_0^t (\|\mathbf{J}^E\|_{L^2}^2 + \|\mathbf{J}^M\|_{L^2}^2) ds \right],$$

where C is a constant depending on T , only. Using (1.1), from (3.39) we deduce that

$$(3.40) \quad \|\mathbf{H}_m(t)\|_{L^2}^2 + \int_0^t \|\mathbf{E}_m\|_{L^2}^2 ds \leq C \left[\|\mathbf{H}_0\|_{L^2}^2 + \int_0^t (\|\mathbf{J}^E\|_{L^2}^2 + \|\mathbf{J}^M\|_{L^2}^2) ds \right], \quad \text{for all } t \in [0, T].$$

for an appropriate constant C , depending only on Λ and T . This implies (3.34).

Step 2. Estimate of $\partial_t \mathbf{H}_m$

Differentiating in (3.36i) with respect to t and taking $\varphi = \mathbf{E}_m$ in the resulting equation, we get

$$(3.41) \quad (\nabla \times \partial_t \mathbf{H}_m, \mathbf{E}_m)_{L^2} - (\sigma \mathbf{E}_m, \partial_t \mathbf{E}_m)_{L^2} = (\partial_t \mathbf{J}^E, \mathbf{E}_m)_{L^2}.$$

Choosing $\psi = \partial_t \mathbf{H}_m$ in (3.36ii), we obtain

$$(3.42) \quad (\nabla \times \mathbf{E}_m, \partial_t \mathbf{H}_m)_{L^2} + (\mu \partial_t \mathbf{H}_m, \partial_t \mathbf{H}_m)_{L^2} = (\mathbf{J}^M, \partial_t \mathbf{H}_m)_{L^2}.$$

Moreover, $\partial_t \mathbf{H}_m$ takes values in $H_0^1(\Omega; \text{curl})$. Hence, $(\nabla \times \mathbf{E}_m, \partial_t \mathbf{H}_m)_{L^2} = (\mathbf{E}_m, \nabla \times \partial_t \mathbf{H}_m)_{L^2}$. Then, subtracting (3.41) from (3.42) and integrating over $(0, t)$ we obtain

$$\int_0^t \|\partial_t \mathbf{H}_m\|_{X_\mu}^2 + \frac{1}{2} (\sigma \mathbf{E}_m(t), \mathbf{E}_m(t))_{L^2} = \frac{1}{2} (\sigma \mathbf{E}_m(0), \mathbf{E}_m(0))_{L^2} + \int_0^t [(\mathbf{J}^M, \partial_t \mathbf{H}_m)_{L^2} - (\partial_t \mathbf{J}^E, \mathbf{E}_m)_{L^2}].$$

By Cauchy-Schwartz and Young inequality,

$$(\mathbf{J}^M, \partial_t \mathbf{H}_m)_{L^2} \leq \frac{1}{2} \|\partial_t \mathbf{H}_m\|_{X_\mu}^2 + \frac{1}{2} (\mu^{-1} \mathbf{J}^M, \mathbf{J}^M)_{L^2}, \quad \text{and} \quad (\partial_t \mathbf{J}^E, \mathbf{E}_m)_{L^2} \leq \frac{1}{2} \|\partial_t \mathbf{J}^E\|_{L^2}^2 + \frac{1}{2} \|\mathbf{E}_m\|_{L^2}^2.$$

By these inequalities and (1.1ii), the previous identity implies that for a.e. $t \in (0, T)$ the inequality

$$\|\mathbf{E}_m(t)\|_{L^2}^2 + \int_0^t \|\partial_t \mathbf{H}_m\|_{L^2}^2 \leq C \left[\|\mathbf{E}_m(0)\|_{L^2}^2 + \int_0^T \|\mathbf{E}_m\|^2 + \int_0^T [\|\partial_t \mathbf{J}^E\|_{L^2}^2 + \|\mathbf{J}^M\|_{L^2}^2] \right],$$

holds, with a constant C depending only on Λ . Eventually, recalling (3.18), from the last inequality and (3.40) we deduce (3.35), as desired. \square

3.4. Proof of Theorem 3.1. We first assume that $\mathbf{J}^E = \mathbf{J}^M = 0$ for a.e. $t \in (0, T)$, and that $\mathbf{H}_0 = 0$. Then we test equation (2.13i) with $\varphi = \mathbf{E}$ and (2.13ii) with $\psi = \mathbf{H}$. By (2.2) and by an integration in time we arrive at

$$(\mu \mathbf{H}(t), \mathbf{H}(t))_{L^2} + \int_0^t (\sigma \mathbf{E}(s), \mathbf{E}(s))_{L^2} ds = 0, \quad \text{for all } t \in [0, T].$$

By (1.1), both the first summand and the integrand in the second one are positive quantities. Then, $\mathbf{H}(t) = \mathbf{E}(t) = 0$ for a.e. $t \in [0, T]$. By linearity this implies at once the uniqueness statement.

Now we prove the existence of solutions. For every $m \in \mathbb{N}$, let $(\mathbf{E}_m, \mathbf{H}_m)$ be as in Lemma 3.6. The energy estimate (3.35) of Proposition 3.7 implies that, by possibly passing to a subsequence,

$$(3.43) \quad \begin{aligned} \mathbf{E}_m &\rightharpoonup \mathbf{E} \quad \text{weakly-* in } L^\infty(0, T; X_\mu), \\ \mathbf{H}_m &\rightharpoonup \mathbf{H} \quad \text{weakly-* in } L^\infty(0, T; X_\mu), \\ \partial_t \mathbf{H}_m &\rightharpoonup \partial_t \mathbf{H} \quad \text{weakly in } L^2(0, T; X_\mu). \end{aligned}$$

Clearly (3.35) and (3.43) imply the estimate (3.1). We are left to prove that the limit (\mathbf{E}, \mathbf{H}) is a weak solution of (2.12).

For all functions $\varphi \in H^1(\Omega; \text{curl})$ and $\psi \in Y_\mu$ that take the form

$$(3.44) \quad \varphi(x, t) = \sum_{i=1}^N \alpha_i(t) \varphi_i(x), \quad \psi(x, t) = \sum_{i=1}^N \beta_i(t) \psi_i(x)$$

for some $\alpha_i, \beta_i \in C^\infty([0, T])$ and $N \in \mathbb{N}$, by (2.2) and (3.15) for all $m \geq N$ we have

$$(3.45a) \quad \int_0^T \int_\Omega \mathbf{H}_m \cdot \nabla \times \varphi \, dx \, dt - \int_0^T \int_\Omega \sigma \mathbf{E}_m \cdot \varphi \, dx \, dt = \int_0^T \int_\Omega \mathbf{J}^E \cdot \varphi \, dx \, dt$$

$$(3.45b) \quad \int_0^T \int_\Omega \mathbf{E}_m \cdot \nabla \times \psi \, dx \, dt + \int_0^T \int_\Omega \mu \partial_t \mathbf{H}_m \cdot \psi \, dx \, dt = \int_0^T \int_\Omega \mathbf{J}^M \cdot \psi \, dx \, dt.$$

Owing to (3.43), from (3.45) we infer that

$$(3.46) \quad \begin{aligned} \int_0^T \int_\Omega \mathbf{H} \cdot \nabla \times \varphi \, dx \, dt - \int_0^T \int_\Omega \sigma \mathbf{E} \cdot \varphi \, dx \, dt &= \int_0^T \int_\Omega \mathbf{J}^E \cdot \varphi \, dx \, dt, \\ \int_0^T \int_\Omega \mathbf{E} \cdot \nabla \times \psi \, dx \, dt + \int_0^T \int_\Omega \mu \partial_t \mathbf{H} \cdot \psi \, dx \, dt &= \int_0^T \int_\Omega \mathbf{J}^M \cdot \psi \, dx \, dt. \end{aligned}$$

The pairs (φ, ψ) of the form (3.44) form a dense set in $L^2(0, T; H^1(\Omega; \text{curl}) \times Y_\mu)$. Thus, from (3.46) we deduce that, for a.e. $t \in [0, T]$, (2.13i) holds for all $\varphi \in H^1(\Omega; \text{curl})$ and (2.13ii) holds for all $\psi \in Y_\mu$. In view of Remark 2.11, it follows that (2.13iii) holds for all $\psi \in H_0^1(\Omega; \text{curl})$.

For a.e. $t \in (0, T)$, (2.13) holds for all $\varphi \in H^1(\Omega; \text{curl})$ and for all $\psi \in H_0^1(\Omega; \text{curl})$ and this implies that $(\mathbf{E}, \mathbf{H}) \in L^2(0, T; H^1(\Omega; \text{curl}) \times H_0^1(\Omega; \text{curl}))$. By (3.43) we also have $\partial_t \mathbf{H} \in L^2(0, T; L^2(\Omega; \mathbb{R}^3))$.

Then, according to Definition 2.9 (see also Remark 2.10) we are left to prove that (2.13iii) holds. To do so, we fix $\psi \in C^1([0, T]; H_0^1(\Omega; \text{curl}))$, with $\psi(T) = 0$. By (2.13ii), we have

$$(3.47) \quad \int_0^T \int_\Omega \mathbf{E} \cdot \nabla \times \psi \, dx \, dt - \int_0^T \int_\Omega \mu \mathbf{H} \cdot \partial_t \psi \, dx \, dt = \int_0^T \int_\Omega \mathbf{J}^M \cdot \psi \, dx \, dt + \int_\Omega \mu \mathbf{H}(0) \cdot \psi(0) \, dx.$$

Also, by (3.45b) we have

$$(3.48) \quad \int_0^T \int_\Omega \mathbf{E}_m \cdot \nabla \times \psi \, dx \, dt - \int_0^T \int_\Omega \mu \mathbf{H}_m \cdot \partial_t \psi \, dx \, dt = \int_0^T \int_\Omega \mathbf{J}_m^M \cdot \psi \, dx \, dt + \int_\Omega \mu \mathbf{H}_{0m} \cdot \psi(0) \, dx.$$

By (3.43), passing to weak limits in (3.48) and comparing with (3.47) we get that

$$(\mathbf{H}(0), \psi(0))_{X_\mu} = (\mathbf{H}_0, \psi(0))_{X_\mu}.$$

Since $\psi(0)$ can be any element of Y_μ , by Lemma 3.3 we deduce (2.13iii) and this ends the proof. \square

4. GLOBAL HÖLDER ESTIMATES FOR THE MAGNETIC FIELD

Given $\alpha \in (0, 1]$, by $C^{0,\alpha}(\overline{\Omega})$ we denote the space of all continuous functions u that are α -Hölder continuous on $\overline{\Omega}$, meaning that

$$\|u\|_{C^{0,\alpha}(\overline{\Omega})} := \|u\|_{L^\infty(\Omega)} + \sup_{\substack{x,y \in \overline{\Omega} \\ x \neq y}} \frac{|u(x) - u(y)|}{|x - y|^\alpha} < +\infty.$$

We recall that $C^{0,\alpha}(\overline{\Omega})$ is a Banach space with this norm. The previous definition extends obviously to the case of vector-valued, and tensor-valued functions.

Theorem 4.1. *There exists $\alpha_0 \in (0, \frac{1}{2}]$, only depending on Λ , such that for every $\alpha \in (0, \alpha_0]$ the following holds: for every $\mathbf{H}_0 \in C^{0,\alpha}(\overline{\Omega}; \mathbb{R}^3)$ and for every $\mathbf{J}^E, \mathbf{J}^M \in L^2(0, T; C^{0,\alpha}(\overline{\Omega}; \mathbb{R}^3))$, if (\mathbf{E}, \mathbf{H}) is a weak solution of (2.12), then $\mathbf{H} \in L^2(0, T; C^{0,\alpha}(\overline{\Omega}; \mathbb{R}^3))$, and we have*

$$(4.1) \quad \begin{aligned} \|\mathbf{H}(t)\|_{C^{0,\alpha}(\overline{\Omega}; \mathbb{R}^3)} &\leq C \left[\|\mu \mathbf{H}_0\|_{C^{0,\alpha}(\overline{\Omega}; \mathbb{R}^3)} + \|\mathbf{E}(t)\|_{L^2} + \|\mathbf{H}\|_{L^2} + \|\mu \partial_t \mathbf{H}(t)\|_{L^2} \right. \\ &\quad \left. + \int_0^t \|\mathbf{J}^M(s)\|_{C^{0,\alpha}(\overline{\Omega}; \mathbb{R}^3)} ds + \|\mathbf{J}^M(t)\|_{L^2} + \|\mathbf{J}^E(t)\|_{C^{0,\alpha}(\overline{\Omega}; \mathbb{R}^3)} \right], \end{aligned}$$

for a.e. $t \in (0, T)$, where the constant C depends on Λ and on r .

4.1. Tools: Morrey and Campanato spaces. For every $\lambda > 0$, given $u \in L^2(\Omega)$ we say that u belongs to Morrey's space $L^{2,\lambda}(\Omega)$ if

$$[u]_{L^{2,\lambda}(\Omega)}^2 := \sup_{\substack{x_0 \in \Omega \\ \rho > 0}} \rho^{-\lambda} \int_{B_\rho(x_0) \cap \Omega} |u|^2 dx < +\infty.$$

In this case we also write $\|u\|_{L^{2,\lambda}(\Omega)} = \|u\|_{L^2(\Omega; \mathbb{R}^3)} + [u]_{L^{2,\lambda}(\Omega)}$. We say that $u \in \mathcal{L}^{2,\lambda}(\Omega)$ if

$$[u]_{\mathcal{L}^{2,\lambda}(\Omega)}^2 := \sup_{\substack{x_0 \in \Omega \\ \rho > 0}} \rho^{-\lambda} \int_{B_\rho(x_0) \cap \Omega} \left| u(x) - \frac{1}{|B_\rho(x_0) \cap \Omega|} \int_{B_\rho(x_0) \cap \Omega} u(y) dy \right|^2 dx < +\infty,$$

and in this case $\|u\|_{\mathcal{L}^{2,\lambda}(\Omega)} = \|u\|_{L^2(\Omega; \mathbb{R}^3)} + [u]_{\mathcal{L}^{2,\lambda}(\Omega)}$. For vector- and tensor-valued functions, Morrey's and Campanato's spaces are defined similarly.

The space $\mathcal{L}^{2,\lambda}(\Omega)$ was introduced by Campanato in [9]. If for all $x_0 \in \partial\Omega$ and for all $\rho > 0$ we have¹ $|\Omega \cap B_\rho(x_0)| \geq K\rho^3$, with a constant K depending only on Ω , then Campanato's space is isomorphic to $L^{2,\lambda}(\Omega)$ for every $\lambda \in (0, 3)$, to $C^{0, \frac{\lambda-3}{2}}(\overline{\Omega})$ for every $\lambda \in (3, 5]$. It can be seen that it only consists of constant functions for every $\lambda > 5$ and that it coincides with the space of BMO functions if $\lambda = 3$, but this will be of no use in the sequel.

¹For example, this measure density requirement is met by all open set satisfying an interior cone condition. In particular, clearly, it follows from assumption (2.3).

4.2. Energy estimates. In this section we provide some elementary a priori estimate for the eddy current system.

Lemma 4.2. *Let $\mathbf{H}_0 \in X_\mu$, and let (\mathbf{E}, \mathbf{H}) be a weak solution of (2.12) in the sense of Remark 3.2. Then estimate (3.2) holds with a constant C depending on μ , Λ , and T , only.*

Proof. Let $t \in (0, T)$ be such that (2.13) holds for all $(\varphi, \psi) \in H^1(\Omega; \text{curl}) \times H_0^1(\Omega; \text{curl})$. Inserting $\varphi = \mathbf{E}$ in (2.13i) and $\psi = \mathbf{H}$ in (2.13ii) and using (2.2) we obtain

$$\int_{\Omega} \mu \partial_t \mathbf{H} \cdot \mathbf{H} \, dx + \int_{\Omega} \sigma \mathbf{E} \cdot \mathbf{E} \, dx = \int_{\Omega} \mathbf{J}^M \cdot \mathbf{H} \, dx - \int_{\Omega} \mathbf{J}^E \cdot \mathbf{E} \, dx.$$

Using (1.1ii) to estimate from below the left hand-side, and Young inequality to estimate from above the right hand-side, we obtain, for all given $\delta \in (0, 1)$, that

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} \mu |\mathbf{H}|^2 \, dx + \frac{1}{\Lambda} \int_{\Omega} |\mathbf{E}|^2 \, dx \leq \frac{1}{2} \int_{\Omega} |\mathbf{J}^M|^2 \, dx + \frac{1}{2} \int_{\Omega} |\mathbf{H}|^2 \, dx + \frac{\delta}{\Lambda} \int_{\Omega} |\mathbf{E}|^2 \, dx + \frac{\Lambda}{4\delta} \int_{\Omega} |\mathbf{J}^E|^2 \, dx.$$

Choosing $\delta = 1/2$ we absorb a term in the left hand-side. Then an integration gives

$$(4.2) \quad \begin{aligned} & \int_{\Omega} |\mathbf{H}(t)|^2 \, dx - \int_{\Omega} |\mathbf{H}_0|^2 \, dx + \int_0^t \int_{\Omega} |\mathbf{E}|^2 \, dx \, ds \\ & \leq \Lambda^2 \left[\int_0^t \int_{\Omega} |\mathbf{H}|^2 \, dx \, ds + \int_0^t \int_{\Omega} |\mathbf{J}^M|^2 \, dx \, ds + \int_0^t \int_{\Omega} |\mathbf{J}^E|^2 \, dx \, ds \right]. \end{aligned}$$

By definition of weak solution (see Definition 2.9 and Remark 2.10), $\mathbf{H} \in L^2(0, T; Y_\mu)$ and $\partial_t \mathbf{H} \in L^2(0, T; Y'_\mu)$. In view of Proposition 2.8, we have $\mathbf{H} \in C([0, T]; L^2)$, and the function

$$t \longmapsto \int_{\Omega} |\mathbf{H}(t)|^2 \, dx,$$

appearing in (4.2), is absolutely continuous. Then, applying Grönwall's Lemma, we obtain that

$$\int_{\Omega} |\mathbf{H}(t)|^2 \, dx - \int_{\Omega} |\mathbf{H}_0|^2 \, dx + \int_0^t \int_{\Omega} |\mathbf{E}|^2 \, dx \, ds \leq C \left[\int_0^T \int_{\Omega} |\mathbf{J}^M|^2 \, dx \, ds + \int_0^T \int_{\Omega} |\mathbf{J}^E|^2 \, dx \, ds \right]$$

for a suitable constant $C > 0$, depending on μ , Λ , and T , only. Since this procedure can be repeated for a.e. $t \in (0, T)$, we deduce (3.2). \square

Theorem 4.3. *Let $\mathbf{H}_0 \in Y_\mu$, let $\mathbf{J}^E \in L^2(0, T; L^2(\Omega; \mathbb{R}^3))$, with $\partial_t \mathbf{J}^E \in L^2(0, T; L^2(\Omega; \mathbb{R}^3))$, let $\mathbf{J}^M \in L^2(0, T; X)$, and let (\mathbf{E}, \mathbf{H}) be a weak solution of (2.12) in the sense of Definition 2.9. Then*

$$\begin{aligned} & \sup_{t \in [0, T]} \|\mathbf{E}(t)\|_{L^2}^2 + \sup_{t \in [0, T]} \|\mathbf{H}(t)\|_{L^2}^2 + \int_0^T \|\partial_t \mathbf{H}(t)\|_{L^2}^2 \, dt \leq C \left[\|\mathbf{H}_0\|_{L^2}^2 \right. \\ & \quad \left. + \int_0^T (\|\mathbf{J}^E(t)\|_{L^2}^2 + \|\mathbf{J}^M(t)\|_{L^2}^2 + \|\partial_t \mathbf{J}^E(t)\|_{L^2}^2) \, dt \right] \end{aligned}$$

where the constant C depends on Λ and T , only.

Proof. Let $\varphi \in H^1(\Omega; \text{curl})$. Differentiating with respect to t in (2.13i) we obtain

$$(4.3) \quad \int_{\Omega} \partial_t \mathbf{H} \cdot \nabla \times \varphi \, dx - \langle \sigma \partial_t \mathbf{E}, \varphi \rangle = \int_{\Omega} \partial_t \mathbf{J}^E \cdot \varphi \, dx,$$

where $\langle \cdot, \cdot \rangle$ stands for the pairing between $H^1(\Omega; \text{curl})$ and its dual space. Since, in (4.3), φ is arbitrary, by (1.1ii) and by a density argument we deduce that that

$$(4.4) \quad \int_0^T \langle \sigma \partial_t \mathbf{E}, v \rangle \leq \left[\int_0^T \|\partial_t \mathbf{H}\|_{(H^1(\Omega; \text{curl}))'}^2 + \int_0^T \|\partial_t \mathbf{J}^E\|_{(H^1(\Omega; \text{curl}))'}^2 \right]^{\frac{1}{2}} \|v\|_{L^2(0, T; H^1(\Omega; \text{curl}))},$$

for all $v \in L^2(0, T; H^1(\Omega; \text{curl}))$. Then, as a function taking values in the dual space of $H^1(\Omega; \text{curl})$, $\partial_t \mathbf{E}$ is L^2 on the interval $(0, T)$. In view of Proposition 2.8, this gives $\mathbf{E} \in C([0, T]; L^2(\Omega; \mathbf{R}^3))$ and

$$(4.5) \quad \frac{d}{dt} \int_{\Omega} \sigma \mathbf{E}(t) \cdot \mathbf{E}(t) dt = 2 \langle \sigma \partial_t \mathbf{E}, \mathbf{E} \rangle, \quad \text{for a.e. } t \in (0, T),$$

where $\langle \cdot, \cdot \rangle$ denotes the duality pairing between $H^1(\Omega; \text{curl})$ and its dual space. Now we take $\varphi = \mathbf{E}$ in (4.3), which we can do for a.e. $t \in (0, T)$. As a result, by (4.5) we get

$$\int_{\Omega} \partial_t \mathbf{H} \cdot \nabla \times \mathbf{E} dx - \frac{1}{2} \frac{d}{dt} \int_{\Omega} \sigma \mathbf{E} \cdot \mathbf{E} dx = \int_{\Omega} \partial_t \mathbf{J}^E \cdot \mathbf{E} dx.$$

Also, for a.e. $t \in (0, T)$ we can test (2.13ii) with $\psi = \partial_t \mathbf{H}$, and doing so we get

$$\int_{\Omega} \mathbf{E} \cdot \nabla \times \partial_t \mathbf{H} dx + \int_{\Omega} \mu \partial_t \mathbf{H} \cdot \partial_t \mathbf{H} dx = \int_{\Omega} \mathbf{J}^M \cdot \partial_t \mathbf{H} dx.$$

We observe that (2.2) implies

$$\int_{\Omega} \mathbf{E} \cdot \nabla \times \partial_t \mathbf{H} dx = \int_{\Omega} \partial_t \mathbf{H} \cdot \nabla \times \mathbf{E} dx.$$

Combining the last three identities we get

$$\int_{\Omega} \mu \partial_t \mathbf{H} \cdot \partial_t \mathbf{H} dx + \frac{1}{2} \frac{d}{dt} \int_{\Omega} \sigma \mathbf{E} \cdot \mathbf{E} dx = - \int_{\Omega} \partial_t \mathbf{J}^E \cdot \mathbf{E} dx + \int_{\Omega} \mathbf{J}^M \cdot \partial_t \mathbf{H} dx.$$

Integrating this energy identity over the interval $[0, t]$, using (1.1) and Young's inequality we obtain

$$\int_0^t \|\partial_t \mathbf{H}\|_{L^2}^2 + \|\mathbf{E}(t)\|_{L^2}^2 \leq C \left[\|\mathbf{E}(0)\|_{L^2}^2 + \int_0^t \|\mathbf{E}\|^2 + \int_0^t (\|\partial_t \mathbf{J}^E\|^2 + \|\mathbf{J}^M\|^2) \right]$$

for a suitable C depending only on Λ . By Grönwall's Lemma, we deduce that

$$(4.6) \quad \int_0^t \|\partial_t \mathbf{H}\|_{L^2}^2 + \|\mathbf{E}(t)\|_{L^2}^2 \leq C \left[\|\mathbf{E}(0)\|_{L^2}^2 + \int_0^t (\|\partial_t \mathbf{J}^E\|^2 + \|\mathbf{J}^M\|^2) \right],$$

where the constant depends now on Λ and T , only.

In order to get rid of the term depending on $\mathbf{E}(0)$ in the right hand-side of (4.6), we note that by Proposition 2.8 we also have

$$\sup_{t \in [0, T]} \|\mathbf{E}(t)\|_{L^2}^2 \leq C \left[\int_0^T \|\mathbf{E}\|_{H^1(\Omega; \text{curl})}^2 + \int_0^T \|\partial_t \mathbf{E}\|_{(H^1(\Omega; \text{curl}))'}^2 \right],$$

with a constant depending only on Λ , and T . We also recall that by (2.13ii) we have

$$\|\mathbf{E}\|_{H^1(\Omega; \text{curl})}^2 = \|\mathbf{E}\|_{L^2}^2 + \|\nabla \times \mathbf{E}\|_{L^2}^2 \leq \|\mathbf{E}\|_{L^2}^2 + \|\mu \partial_t \mathbf{H}\|_{L^2}^2 + \|\mathbf{J}^M\|_{L^2}^2,$$

whereas (4.4) implies

$$\int_0^T \|\partial_t \mathbf{E}\|_{(H^1(\Omega; \text{curl}))'}^2 \leq \int_0^T \|\partial_t \mathbf{H}\|^2 + \|\partial_t \mathbf{J}^E\|^2.$$

Then, by Grönwall Lemma it follows that

$$(4.7) \quad \sup_{t \in [0, T]} \|\mathbf{E}(t)\|_{L^2}^2 \leq C \left[\int_0^T \|\partial_t \mathbf{H}\|_{L^2}^2 + \|\partial_t \mathbf{J}^E\|_{L^2}^2 + \|\mathbf{J}^M\|_{L^2}^2 \right],$$

where C depends on Λ and T , only.

Inserting (4.7) in (4.6) we arrive at

$$\int_0^t \|\partial_t \mathbf{H}\|_{L^2}^2 + \|\mathbf{E}(t)\|_{L^2}^2 \leq C \int_0^t (\|\partial_t \mathbf{H}\|_{L^2}^2 + \|\partial_t \mathbf{J}^E\|_{L^2}^2 + \|\mathbf{J}^M\|_{L^2}^2),$$

for a.e. $0 \leq t \leq T$. □

4.3. Proof of Theorem 4.1. We set

$$\mathcal{I} = \{t \in [0, T] : \mathbf{E}(t) \in H^1(\Omega; \text{curl}), \mathbf{H}(t) \in Y_\mu, \mathbf{J}^E(t), \mathbf{J}^M(t) \in C^{0, \alpha}(\bar{\Omega}; \mathbb{R}^3)\}$$

and we recall that $[0, T] \setminus \mathcal{I}$ is a negligible set (see Remark 2.10). We drop the dependance on t of the vector fields, so as to abbreviate the notations.

By Lemma 2.5, there exist $u \in H^1(\Omega)$ and $\eta \in H^1(\Omega; \mathbb{R}^3)$ with

$$(4.8a) \quad \mathbf{E} = \nabla u + \eta$$

$$(4.8b) \quad \|\nabla \eta\|_{L^2} = \|\nabla \times \mathbf{E}\|_{L^2}$$

$$(4.8c) \quad \max\{\|\nabla u\|_{L^2}, \|\eta\|_{L^2}\} \leq \|\mathbf{E}\|_{L^2}.$$

Recalling equation (2.13ii), from (4.8b) and (4.8c) we deduce

$$(4.9) \quad \|\eta\|_{H^1(\Omega; \mathbb{R}^3)} \leq \|\mathbf{E}\|_{L^2} + \|\mu \partial_t \mathbf{H}\|_{L^2} + \|\mathbf{J}^M\|_{L^2}.$$

By Sobolev embedding Theorem, the inclusion of $H^1(\Omega; \mathbb{R}^3)$ into $L^6(\Omega; \mathbb{R}^3)$ is continuous, and so is the embedding of $L^6(\Omega; \mathbb{R}^3)$ into Morrey's space $L^{2,2}(\Omega; \mathbb{R}^3)$, thanks to Hölder inequality. Thus, $\|\eta\|_{L^{2,2}(\Omega; \mathbb{R}^3)} \leq C \|\eta\|_{H^1(\Omega; \mathbb{R}^3)}$ for a constant $C > 0$ that depends on r , only. Hence, by (4.9) we get

$$(4.10) \quad \|\eta\|_{L^{2,2}(\Omega; \mathbb{R}^3)} \leq C \left[\|\mathbf{E}\|_{L^2} + \|\mu \partial_t \mathbf{H}\|_{L^2} + \|\mathbf{J}^M\|_{L^2} \right].$$

Next, we pick $w \in H^1(\Omega)$ and we test equation (2.13i) with $\varphi = \nabla w$. By (4.8a), we obtain

$$\int_{\Omega} \sigma \nabla u \cdot \nabla w \, dx = - \int_{\Omega} (\sigma \eta + \mathbf{J}^E) \cdot \nabla w \, dx.$$

By [18, Theorem 2.19] with $\Gamma = \partial\Omega$ (see also Lemma 2.18 therein), there exists $\bar{\lambda} \in (1, 2]$, depending only on Λ , such that for all $\lambda \in (1, \bar{\lambda}]$ we have

$$\|\nabla u\|_{L^{2, \lambda}(\Omega; \mathbb{R}^3)} \leq C \left[\|\nabla u\|_{L^2} + \|\sigma \eta + \mathbf{J}^E\|_{L^{2, \lambda}(\Omega; \mathbb{R}^3)} \right],$$

for a suitable $C > 0$, depending on Λ and on r , only. By (1.1ii) and (4.8c), the latter implies

$$(4.11) \quad \|\nabla u\|_{L^{2, \lambda}(\Omega; \mathbb{R}^3)} \leq C \left[\|\mathbf{E}\|_{L^2} + \|\eta\|_{L^{2, \lambda}(\Omega; \mathbb{R}^3)} + \|\mathbf{J}^E\|_{L^{2, \lambda}(\Omega; \mathbb{R}^3)} \right].$$

Fix $\lambda \in (1, \bar{\lambda}]$. By (4.8a), (4.10), and (4.11), there exists $C > 0$, depending only on Λ and r , with

$$(4.12) \quad \|\mathbf{E}\|_{L^{2, \lambda}(\Omega)} \leq C \left[\|\mathbf{E}\|_{L^2} + \|\mu \partial_t \mathbf{H}\|_{L^2} + \|\mathbf{J}^M\|_{L^2} + \|\mathbf{J}^E\|_{L^{2, \lambda}(\Omega; \mathbb{R}^3)} \right].$$

We recall that $\mathbf{H} \in Y_\mu$. By Lemma 2.2, this gives $\mathbf{H} \in H^1(\Omega; \mathbb{R}^3)$. In view of Remark 2.7, there exist $q \in H_0^1(\Omega)$, and $\zeta \in H_0^1(\Omega; \text{curl})$, with

$$(4.13) \quad \int_{\Omega} \zeta \cdot \nabla v \, dx = 0, \quad \text{for all } v \in H_0^1(\Omega),$$

such that $\mathbf{H} = \nabla q + \zeta$ and

$$(4.14) \quad \max \left\{ \|\nabla q\|_{L^2}, \|\zeta\|_{L^2} \right\} \leq \|\mathbf{H}\|_{L^2}.$$

Then, by [1, Lemma 6], for a constant C depending only on Λ, r we have

$$\|\nabla \zeta\|_{L^{2,\lambda}(\Omega; \mathbb{R}^{3 \times 3})} \leq C \|\nabla \times \zeta\|_{L^{2,\lambda}(\Omega; \mathbb{R}^3)}.$$

Thus, recalling that $\nabla \times \zeta = \nabla \times \mathbf{H}$ and using equation (2.13i), we arrive at

$$(4.15) \quad \|\nabla \zeta\|_{L^{2,\lambda}(\Omega; \mathbb{R}^{3 \times 3})} \leq C \left[\|\mathbf{E}\|_{L^{2,\lambda}(\Omega; \mathbb{R}^3)} + \|\mathbf{J}^E\|_{L^{2,\lambda}(\Omega; \mathbb{R}^3)} \right]$$

where C depends on Λ, r , only.

We note that, by (2.3), there exists $\bar{\rho}_0 > 0$, depending only on r , such that if $0 < \rho < \bar{\rho}_0$ then

- (i) the boundary of $\Omega \cap B(x_0, \rho)$, in the sense of [2, Definition 3.2], is of Lipschitz class with constants $c\rho, L$, with c and L depending on r , only;
- (ii) $\Omega \cap B(x_0, \rho)$ satisfies the *scale-invariant fatness condition*, in the sense of [2, equation (2.3)].

Thus, by [2, Proposition 3.2], for every $0 < \rho < \bar{\rho}_0$ the following Poincaré inequality

$$\int_{B_\rho(x_0) \cap \Omega} \left| \zeta(x) - \fint_{B_\rho(x_0) \cap \Omega} \zeta(y) \, dy \right|^2 dx \leq C\rho^2 \int_{B_\rho(x_0) \cap \Omega} |\nabla \zeta|^2 dx,$$

holds for all $x_0 \in \Omega$, for a constant C depending only on r . Hence,

$$(4.16) \quad \|\zeta\|_{\mathcal{L}^{2,\lambda+2}(\Omega; \mathbb{R}^3)} \leq C \|\nabla \zeta\|_{L^{2,\lambda}(\Omega; \mathbb{R}^{3 \times 3})}.$$

By (4.14), (4.15), (4.16), there exists a constant $C > 0$ depending on Λ and r such that

$$(4.17) \quad \|\zeta\|_{\mathcal{L}^{2,\lambda+2}(\Omega; \mathbb{R}^3)} \leq C \left[\|\mathbf{H}\|_{L^2} + \|\mathbf{E}\|_{L^{2,\lambda}(\Omega; \mathbb{R}^3)} + \|\mathbf{J}^E\|_{L^{2,\lambda}(\Omega; \mathbb{R}^3)} \right].$$

We recall that Campanato's space $\mathcal{L}^{2,\lambda+2}(\Omega)$, as a Banach space, is isomorphic to $C^{0,\alpha}(\bar{\Omega})$, where $\alpha \in (0, \frac{1}{2})$ is given by $\alpha = (\lambda - 1)/2$. Incidentally, we set $\alpha_0 = (\bar{\lambda} - 1)/2$, we observe that $\alpha \in (0, \alpha_0)$ and $\alpha_0 \in (0, \frac{1}{2}]$, because $\bar{\lambda} \in (1, 2]$. Then, (4.17) implies

$$(4.18) \quad \|\zeta\|_{C^{0,\alpha}(\bar{\Omega}; \mathbb{R}^3)} \leq C \left[\|\mathbf{H}\|_{L^2} + \|\mathbf{E}\|_{L^{2,\lambda}(\Omega; \mathbb{R}^3)} + \|\mathbf{J}^E\|_{L^{2,\lambda}(\Omega; \mathbb{R}^3)} \right].$$

We take $w \in H_0^1(\Omega)$ and we test equation (2.13ii) with $\psi = \nabla w$. By Fubini's Theorem and integrations by parts, we get

$$\int_{\Omega} \mu \mathbf{H} \cdot \nabla w \, dx - \int_{\Omega} \mu \mathbf{H}_0 \cdot \nabla w \, dx = \int_{\Omega} \int_0^t \mathbf{J}^M \cdot \nabla w \, ds \, dx.$$

Since $\mathbf{H} = \nabla q + \zeta$ and w can be any element of $H_0^1(\Omega)$, it follows that $q \in H_0^1(\Omega)$ is a weak solution of the elliptic equation

$$\nabla \cdot (\mu \nabla q) = \nabla \cdot \left(\int_0^t \mathbf{J}^M \, ds + \mu \mathbf{H}_0 - \mu \zeta \right).$$

Then, classical global Schauder estimates (see, e.g., [18, Theorem 2.19] with $\Gamma = \emptyset$, and Lemma 2.18 therein) imply

$$(4.19) \quad \|\nabla q\|_{C^{0,\alpha}(\overline{\Omega};\mathbb{R}^3)} \leq C \left[\int_0^t \|\mathbf{J}^M\|_{C^{0,\alpha}(\overline{\Omega};\mathbb{R}^3)} ds + \|\mu \mathbf{H}_0\|_{C^{0,\alpha}(\overline{\Omega};\mathbb{R}^3)} + \|\zeta\|_{C^{0,\alpha}(\overline{\Omega};\mathbb{R}^3)} \right],$$

where the constant depends on Λ , on r .

Since $\mathbf{H} = \nabla q + \zeta$, from (4.18) and (4.19) we deduce that the estimate (4.1) is valid for all t that belong to the set \mathcal{I} defined at the beginning of the proof. Since \mathcal{I} has full measure in $(0, T)$, clearly it follows that (4.1) holds for a.e. $t \in (0, T)$. \square

APPENDIX A. HELMOLTZ DECOMPOSITIONS

A.1. Proof of Lemma 2.5. We define

$$V = \left\{ u \in H^1(\Omega) : \int_{\Omega} u \, dx = 0 \right\}$$

and we observe that V is a closed subspace of the Hilbert space $H^1(\Omega)$. By Poincaré's inequality and Lax-Milgram Lemma, there exists a (unique) solution $u \in V$ to the variational problem

$$(A.1) \quad \int_{\Omega} \nabla u \cdot \nabla v \, dx = \int_{\Omega} \mathbf{F} \cdot \nabla v \, dx, \quad \text{for all } v \in V.$$

Since every $v \in H^1(\Omega)$ differs from some element of V by a constant, from (A.1) we can infer

$$(A.2) \quad \int_{\Omega} \nabla u \cdot \nabla v \, dx = \int_{\Omega} \mathbf{F} \cdot \nabla v \, dx, \quad \text{for all } v \in H^1(\Omega).$$

Setting $\eta = \mathbf{F} - \nabla u$, we have (2.7a) trivially, and (A.2) implies (2.7b). To conclude the proof, we test (A.2) with $v = u$ and get

$$(A.3) \quad \int_{\Omega} |\nabla u|^2 \, dx = \int_{\Omega} \mathbf{F} \cdot \nabla u \, dx.$$

Therefore, Cauchy-Schwartz inequality implies $\|\nabla u\|_{L^2} \leq \|\mathbf{F}\|_{L^2}$. Then, we note that

$$\int_{\Omega} |\eta|^2 \, dx = \int_{\Omega} |\mathbf{F}|^2 \, dx + \int_{\Omega} |\nabla u|^2 \, dx - 2 \int_{\Omega} \mathbf{F} \cdot \nabla u \, dx.$$

Hence, recalling (A.3), we have $\|\eta\|_{L^2}^2 \leq \|\mathbf{F}\|_{L^2}^2 - \|\nabla u\|_{L^2}^2 \leq \|\mathbf{F}\|_{L^2}^2$ and we deduce (2.7c).

Now, we also assume that $\mathbf{F} \in H^1(\Omega; \text{curl})$. Since $\nabla \times \eta = \nabla \times (\mathbf{F} - \nabla u) = \nabla \times \mathbf{F}$, the (distributional) curl of η belongs to L^2 . Since (2.7b) holds, in particular, for all $v \in H_0^1(\Omega)$, the (distributional) divergence $\nabla \cdot \eta$ of η equals 0. Moreover, again by (2.7b), for every $v \in H^1(\Omega)$

$$\langle \gamma_{\partial\Omega}(v), \eta \cdot n \rangle = \int_{\Omega} \nabla v \cdot \eta \, dx,$$

where γ is the trace operator from $H^1(\Omega)$ to $H^{\frac{1}{2}}(\partial\Omega)$ and $\langle \cdot, \cdot \rangle$ is the duality pairing between $H^{-\frac{1}{2}}(\partial\Omega)$ and $H^{\frac{1}{2}}(\partial\Omega)$. Hence $\eta \cdot n = 0$ in $H^{-\frac{1}{2}}(\partial\Omega)$. Then, by an integration by parts, we deduce that $\eta \in H^1(\Omega; \mathbb{R}^3)$ and $\|\nabla \eta\|_{L^2} = \|\nabla \times \eta\|_{L^2}$. Since $\nabla \times \eta = \nabla \times \mathbf{F}$, we conclude that $\|\nabla \eta\|_{L^2} = \|\nabla \times \mathbf{F}\|_{L^2}$ as desired.

A.2. **Proof of Lemma 2.6.** Equation (2.8) with $v = q$ reads as

$$(A.4) \quad \int_{\Omega} \mu |\nabla q|^2 dx = \int_{\Omega} \mu \mathbf{F} \cdot \nabla q dx.$$

Using Cauchy-Schwartz inequality and (1.1i), from (A.4) we obtain $\|\nabla q\|_{L^2} \leq \Lambda \|\nabla \mathbf{F}\|_{L^2}$, which gives the first inequality in (2.9b); setting $\zeta = \mathbf{F} - \nabla q$ and using (A.4) again we also get

$$\int_{\Omega} \mu |\zeta|^2 dx = \int_{\Omega} \mu |\mathbf{F}|^2 dx + \int_{\Omega} \mu |\nabla q|^2 dx - 2 \int_{\Omega} \mu \mathbf{F} \cdot \nabla q dx = \int_{\Omega} \mu |\mathbf{F}|^2 dx - \int_{\Omega} \mu |\nabla q|^2 dx \leq \int_{\Omega} \mu |\mathbf{F}|^2 dx,$$

which gives the second inequality, too. Since $\zeta = \mathbf{F} - \nabla q$, clearly (2.9a) holds, $\zeta \in L^2(\Omega; \mathbb{R}^3)$, and by (2.8) we also have $\zeta \in X_{\mu}$.

If, in addition, $\mathbf{F} \in H^1(\Omega; \mathbb{R}^3)$, then $\nabla \cdot \mathbf{F} \in L^2(\Omega; \mathbb{R}^3)$. Hence, by (2.8) and Elliptic Regularity we have $q \in H^2(\Omega)$ (see, e.g., [15, §8.3]). By difference, $\zeta \in H^1(\Omega; \mathbb{R}^3)$. Moreover,

$$(A.5) \quad \begin{aligned} \int_{\Omega} \varphi \cdot \nabla \times \zeta dx - \int_{\Omega} \zeta \cdot \nabla \times \varphi dx &= \int_{\Omega} \varphi \cdot \nabla \times (\mathbf{F} - \nabla q) dx - \int_{\Omega} (\mathbf{F} - \nabla q) \cdot \nabla \times \varphi dx \\ &= \int_{\Omega} \varphi \cdot \nabla \times \mathbf{F} dx - \int_{\Omega} \mathbf{F} \cdot \nabla \times \varphi dx + \int_{\Omega} \nabla q \cdot \nabla \times \varphi dx, \end{aligned}$$

for all given $\varphi \in C^1(\overline{\Omega}; \mathbb{R}^3)$. Now we also assume that $\mathbf{F} \times n = 0$ in $H^{-\frac{1}{2}}(\partial\Omega; \mathbb{R}^3)$. Then, by (2.2),

$$\int_{\Omega} \varphi \cdot \nabla \times \mathbf{F} dx - \int_{\Omega} \mathbf{F} \cdot \nabla \times \varphi dx = 0.$$

Since $q \in H_0^1(\Omega)$, by divergence theorem we also have

$$\int_{\Omega} \nabla q \cdot \nabla \times \varphi dx = 0.$$

Inserting the last two identities in (A.5) we obtain

$$\int_{\Omega} \varphi \cdot \nabla \times \zeta dx - \int_{\Omega} \zeta \cdot \nabla \times \varphi dx = 0.$$

Since φ was arbitrary, by (2.2) we deduce that $\zeta \times n = 0$ in $H^{-\frac{1}{2}}(\partial\Omega; \mathbb{R}^3)$. Thus, $\zeta \in H_0^1(\Omega; \text{curl})$. Recalling that $\zeta \in X_{\mu}$ and that by definition $Y_{\mu} = H_0^1(\Omega; \text{curl}) \cap X_{\mu}$, this concludes the proof.

REFERENCES

- [1] G. S. ALBERTI, Hölder regularity for Maxwell's equations under minimal assumptions on the coefficients, *Calc. Var. and Part. Diff. Equations* **57** (2018), pp. 1–11.
- [2] G. ALESSANDRINI, A. MORASSI, AND E. ROSSET, Detecting an inclusion in an elastic body by boundary measurements, *SIAM J. Math. Anal.* **33** no. 6 (2002), pp. 1247–1268.
- [3] A. ALONSO RODRÍGUEZ, A. VALLI, *Eddy Current Approximation of Maxwell Equations*, Springer-Verlag Italia, Milan, 2010.
- [4] A. ALONSO RODRÍGUEZ, J. CAMAÑO, A. VALLI, Inverse source problems for eddy current equations, *Inverse Problems* **28** (2012), no. 1, 015006, 15 pp.
- [5] R. ALVARADO, D. BRIGHAM, V. MAZ'YA, M. MITREA, E. ZIADÉ, On the regularity of domains satisfying a uniform hour-glass condition and a sharp version of the Hopf-Oleinik boundary point principle, *Journ. of Math. Sci.* **176**, No. 3 (2011), pp. 281–360.
- [6] H. AMMARI, A. BUFFA, J.C. NÉDÉLEC, A justification of eddy currents model for the Maxwell equations, *SIAM Journ. on Appl. Math.*, **60** No. 5 (2000), pp. 1805–1823.
- [7] H. AMMARI, J. JUNQING CHEN, Z. CHEN, J. GARNIER, AND D. VOLKOV Target detection and characterization from electromagnetic induction data, *J. Math. Pures Appl.* **101** (2014) 54–75.

- [8] L. ARNOLD, B. HARRACH, A unified variational formulation for the parabolic-elliptic eddy current equations, *SIAM J. Appl. Math.* **72** 2 (2012), pp. 558–576.
- [9] S. CAMPANATO, Equazioni ellittiche del II° ordine e spazi $\mathcal{L}^{2,\lambda}$, *Annal. Mat. Pura e Appl.* **69** No. 1 (1965), pp. 321–381.
- [10] M. COSTABEL, A coercive bilinear form for Maxwell’s equations, *J. Math. Anal. Appl.*, **157** (1991), 527–541.
- [11] M. COSTABEL, M. DAUGE, AND S. NICAISE, Singularities of eddy current problems, *M2AN Math. Model. Numer. Anal.* **37** (2003), pp. 807–831.
- [12] G. DUVAUT, J.L. LIONS, *Inequalities in Mathematical Physics*, Springer-Verlag Berlin, Heidelberg, 1976.
- [13] L.C.EVANS, *Partial Differential Equations*, American Mathematical Society, Providence, R.I., 2010.
- [14] K. O. FRIEDRICHS, Differential Forms on Riemannian Manifolds, *Comm. Pure and Appl. Math.* **8** (1955), pp. 551–590.
- [15] D. GILBARG AND N.S. TRUDINGER, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag Berlin, Heidelberg, 1977.
- [16] R. HIPTMAIR, Symmetric coupling for eddy current problems, *SIAM J. Numer. Anal.* **40** (2002) 41–65.
- [17] P. PALANGIO, C. DI LORENZO, F. MASCI, AND M. DI PERSIO, The study of the electromagnetic anomalies linked with the Earth’s crustal activity in the frequency band [0.001 Hz–100 kHz], *Nat. Hazards Earth Syst. Sci.* **7** (2007), pp. 507–511.
- [18] G.M. TROIANELLO, *Elliptic Differential Equations and Obstacle Problems*, Plenum Press, New York, 1987.
- [19] K. YAMAZAKI, An analytical expression for early electromagnetic signals generated by impulsive line-currents in conductive Earth crust, with numerical examples, *Annals of Geophysics [S.l.]* **59** No. 2 (2016), pG0212 1–9.

(G. Franzina) UNITÀ DI RICERCA INdAM DI FIRENZE %o DiMaI “ULISSE DINI”
 UNIVERSITÀ DEGLI STUDI DI FIRENZE
 VIALE MORGAGNI 67/A, 50134 FIRENZE, ITALY
 E-mail address: franzina@math.unifi.it

(E. Francini, S. Vessella) DiMaI “ULISSE DINI”
 UNIVERSITÀ DEGLI STUDI DI FIRENZE
 VIALE MORGAGNI 67/A, 50134 FIRENZE, ITALY
 E-mail address: francini@math.unifi.it, vessella@unifi.it