

1 **Rapid cooling and increased storminess triggered by**
2 **freshwater in the North Atlantic**

3 M. Oltmanns¹, J. Karstensen², G. W. K. Moore³, S. A. Josey¹

4 ¹National Oceanography Centre, Southampton, UK

5 ²GEOMAR - Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

6 ³University of Toronto, Toronto, Canada

7 **Key Points:**

- 8 • Large freshwater events result in distinct cold anomalies with sharp temperature
9 gradients in the subpolar North Atlantic in winter.
- 10 • A strong, freshwater-induced cold anomaly promotes an enhanced storminess, which
11 reinforces the anomaly by modulating the surface flow.
- 12 • Consistent with this mechanism, large freshwater events in the past have preceded
13 positive North Atlantic Oscillation periods in winter.

Corresponding author: Marilena Oltmanns, marilena.oltmanns@noc.ac.uk

Abstract

Recent winters have been unique due to the rapid and extreme cooling of the subpolar North Atlantic. Here, we present a novel view on its causes and consequences. Combining in-situ observations with remote sensing and atmospheric reanalysis data, we show that increased freshening of the subpolar region gives rise to a faster surface cooling in fall and winter. Large freshwater events, in particular, result in pronounced cold anomalies with sharp temperature gradients that promote an enhanced storminess. The storms reinforce the cooling by driving stronger heat losses and modulating the surface flow. Consistent with this mechanism, past freshwater events have been followed by cold anomalies in winter of ~ -2 °C and increases in the North Atlantic Oscillation index of up to ~ 0.6 within 3 years. We expect that future freshwater discharges into the North Atlantic will amplify the cold anomaly and trigger an enhanced wintertime storminess with far-reaching climatic implications.

Plain Language Summary

Recent winters have been unique due to a rapid and extreme cooling of the subpolar North Atlantic. Combining ocean and atmospheric data, we show that increased freshwater in this region leads to shallower surface layers that adjust faster to the lower air temperature in fall and winter. The faster surface cooling increases the south-north temperature gradient which promotes the development of storms. The storms, in turn, reinforce the cooling by modulating the surface flow. Accordingly, past freshwater events have been followed by an extremely cold ocean surface in the subpolar North Atlantic in winter and major changes in large-scale weather patterns. We expect that future freshwater discharges from Greenland and the Arctic will amplify the cooling and trigger an enhanced wintertime storminess with far-reaching implications for the climate.

1 Introduction

Recent winters have been characterised by a rapid and extreme cooling of the subpolar North Atlantic, which has been unprecedented in over 30 years and stands in marked contrast to the warming observed over most of the Earth's surface (Josey et al., 2018). Given the importance of the North Atlantic sea surface temperature (SST) for the large-scale weather and climate (Czaja & Frankignoul, 2002; Sutton & Dong, 2012), it is critical to understand the causes and effects of this anomaly.

Previous studies have attributed the cooling to a slowdown of the Atlantic overturning circulation due to increased freshwater fluxes from Greenland (Rahmstorf et al., 2015; Caesar et al., 2018). This idea is motivated by paleoclimate records, suggesting that past cooling events were caused by a freshwater-forced shutdown of deep ocean convection in the subpolar North Atlantic and a subsequent collapse of the overturning (Barber et al., 1999; Clark et al., 2001, 2002). However, observations show that this buoyancy-driven mechanism cannot easily explain the recent heat transport into the subpolar region (Lozier et al., 2019).

The influences of the cold anomaly on the climate are likewise uncertain. Yet, earlier studies have found that *increased* SSTs in the subpolar region initially trigger a transient baroclinic response in the atmosphere, forcing enhanced ocean heat losses (Kushnir et al., 2002; Deser et al., 2007). After a few weeks, a barotropic equilibrium response emerges that is associated with reduced ocean heat losses and thus represents a positive feedback to the SST anomaly (Kushnir et al., 2002; Deser et al., 2007).

However, we hypothesise that freshwater modulates this atmospheric response by strengthening the stratification, allowing for a faster adjustment of the surface to the lower air temperatures in fall and winter. By eroding the SST anomaly during the transient

Table 1. Main data products and indices involved in this study.

Data/index	Period
Hydrographic observations from Argo floats	2002–2018
Remote sensing-based SST data	1982–2018
Absolute dynamic topography from altimetry	1993–2018
Atmospheric data from the reanalysis ERA-Interim	1979–2018
Freshwater index (F_{NA}) ^a	1979–2018
Cold anomaly index (CAI) ^b	1982–2019

^a F_{NA} : Mean NAO in July and August, multiplied by -1 .

^b CAI: Mean SST in the cold anomaly region, multiplied by -1 .

62 response, the faster surface cooling interferes with the setup of the equilibrium response
 63 and instead, increases the meridional SST gradient, key source of baroclinic instability
 64 (Hoskins et al., 1985). To test this hypothesis, we investigate the chain of mechanisms
 65 initiated by freshwater. The main data products involved in this study are listed in Ta-
 66 ble 1, while a detailed data description is provided in the Supporting Information.

67 2 Results

68 2.1 Detection of surface freshwater

69 Motivated by earlier studies that discovered a significant anti-correlation of the North
 70 Atlantic Oscillation (NAO) in summer with the melting over Greenland (Hanna et al.,
 71 2013) and the Arctic sea ice export (Haine et al., 2015), both potential sources of fresh-
 72 water, we use the mean NAO in July and August, multiplied by -1 , as an index (F_{NA})
 73 to describe the freshwater variability in the subpolar North Atlantic (Fig. 1a).

74 Based on a scaling analysis of the surface mass balance, we find that the surface
 75 freshening associated with F_{NA} can be estimated from:

$$\beta\Delta SSS \approx \alpha\Delta SST, \quad (1)$$

76 where both sides have been regressed on F_{NA} , α and β are the thermal and haline ex-
 77 pansion coefficients, SSS is the surface salinity and Δ refers to the change from sum-
 78 mer to winter (see Supporting Information for details).

79 The inferred surface freshening is most pronounced in the western subpolar region,
 80 off the coast of Newfoundland, from where it expands eastward into the central gyre, fol-
 81 lowing the mean geostrophic flow (Fig. 1b). Both F_{NA} and the inferred freshening are
 82 characterised by a significantly positive trend over the investigated period and their cor-
 83 relation is largest on time scales above 5 years (Fig. 1c). Thus, F_{NA} identifies periods
 84 of increased freshwater rather than individual years.

85 2.2 Influence of surface freshwater on the SST

86 The surface mass balance states that an enhanced surface freshening strengthens
 87 the stratification, which allows for an increased surface cooling in fall and winter before
 88 the freshwater is mixed down. This freshwater-induced surface cooling is one order of
 89 magnitude larger than the cooling resulting from the anomalous surface heat flux asso-
 90 ciated with F_{NA} .

91 Hydrographic observations from the Labrador Sea over the period 2002–2018 show
 92 that an increased surface freshening not only leads to an anomalous cooling relative to




Figure 1. (a) Variability of F_{NA} , which is the mean NAO index in July and August, multiplied by -1 , and the inferred surface freshening ($-\Delta SSS$) from summer (August) to winter (January through March) in the region enclosed by the blue 95% confidence line in b. Also shown are the trends. (b) Regression of the inferred SSS change on F_{NA} in the region where an anomalous surface cooling is observed. Thick contours mark the 95% confidence level and arrows indicate the direction of the mean geostrophic flow. (c) Multi-tapered coherence and phase shift between F_{NA} and the seasonal SSS change in the region enclosed by the blue 95% confidence line in b, computed using 8 tapers. Significance analyses are provided in the Supporting Information.




Figure 2. Regressions of (a) the SST change from summer (August) to winter (January through March) and (b,c) the absolute salinity and potential temperature in the Labrador Sea on F_{NA} , based on 17-year long mooring and Argo float observations. The black contour in a delineates the Argo float sampling region and the circle marks the mooring location. Coloured contours indicate the 95% confidence levels.

93 the summer (Fig. 2a) but also to a cold surface anomaly relative to the climatological
 94 mean (Fig. 2b and c). While the positive subsurface temperature and salinity anom-
 95 alies are expected from the increased advection of heat and salt in negative NAO peri-
 96 ods (Sarafanov, 2009; Reverdin, 2010), the cold surface anomaly can only be explained
 97 by the freshwater-induced surface cooling.

98 However, the hydrographic observations were acquired over a time with an over-
 99 all elevated freshening (Fig. 1a). When considering the full period of satellite observa-
 100 tions, the anti-correlation between the freshening in summer and the SST in the subse-
 101 quent winter only holds for positive freshwater anomalies (Fig. 3a-c), reflecting the tran-
 102 sition from the regime where the surface heat fluxes control the SST anomaly to the regime
 103 where the freshwater dominates.

104 The freshwater-induced cold anomaly is most pronounced in the eastern subpolar
 105 region (Fig. 3a and b), where the correlation of the SST, multiplied by -1 (the cold anomaly
 106 index ‘CAI’), with the positive F_{NA} values amounts to ~ 0.57 , which increases for higher
 107 values of F_{NA} (Fig. 3c). The associated heat flux anomaly is directed upwards and thus
 108 cannot account for the cold anomaly. On the contrary, the cold anomaly weakens the
 109 heat loss to the atmosphere by reducing the air-sea temperature contrast (Fig. 3d).

110 Southeast of the cold anomaly over the Gulf Stream, the SST is increased (Fig. 3a),
 111 giving rise to a sharper meridional SST gradient, key source of baroclinic instability in
 112 the atmosphere (Hoskins et al., 1985). Thus, we next investigate the interaction of the
 113 cold anomaly with the atmosphere.

114 **2.3 Atmospheric feedbacks**

115 Regressing the sea level pressure (SLP) onto the cold anomaly, we find that the cold
 116 anomaly is accompanied by an anomalous low over and southeast of Greenland (Fig. 4a),
 117 representative of a positive NAO. In addition, the standard deviation of the 2-6 day band-
 118 pass filtered SLP is increased (Fig. 4b), implying an enhanced storminess (Blackmon,
 119 1976; Blackmon et al., 1977; Ulbrich et al., 2008). Consistent with the increased cyclonic
 120 activity, the heat loss over the Labrador Sea, south of the sea ice edge, is amplified (Fig.
 121 4c), and the wind stress curl under the low is higher (Fig. 4d).




Figure 3. (a,b,d) Regression of the SST, the inferred SSS in the cold anomaly region, and the surface heat fluxes in winter (January through March) on F_{NA} from the preceding summer, with only positive F_{NA} included. A positive heat flux anomaly means that the ocean loses less heat. Thick contours indicate the 95% confidence levels. (c) Relationship between F_{NA} and the SST in the region delineated by the thick blue contour in a. When multiplied by -1 , this SST anomaly corresponds to the cold anomaly index (CAI). Also shown are the correlation coefficients and regression lines for $F_{NA} > 0$ and $F_{NA} > 0.5$.




Figure 4. (a–d) Regressions of (a) the SLP, (b) the 2 to 6-day band-pass filtered standard deviation of the SLP, (c) the surface heat flux (positive downward) and (d) the wind stress curl on CAI, multiplied by the regression of CAI on positive F_{NA} . (e) Regressions of the absolute dynamic topography on the wind stress curl within the red contour in d (‘WSC’), multiplied by the regressions of WSC on CAI and CAI on positive F_{NA} . (f) Correlations of CAI with F_{NA} and the WSC, with the envelopes indicating the 95% confidence levels. A positive lag implies that the cold anomaly lags F_{NA} and the WSC. All variables are in winter (January through March).

122 A high wind stress curl has been identified as an important driver of the subpo-
 123 lar gyre circulation, which retains cold, fresh polar water in the interior gyre (Häkkinen
 124 et al., 2011; Chafik et al., 2019). Altimetry data confirm that the obtained wind stress
 125 curl is associated with a stronger cyclonic circulation, implying an increased advection
 126 of polar water off the coast of Newfoundland towards the anomaly, hence reinforcing it
 127 (Fig. 4e). The correlation between the surface flow pattern and the cold anomaly is ~ 0.80 .

128 2.4 Summary of the feedback mechanism

129 Closer inspection of the timing of the correlations reveals that the cold anomaly
 130 emerges in the winter immediately *following* a high F_{NA} summer and occurs in *the same*
 131 winter with the enhanced wind stress curl (Fig. 4f). It then continues to intensify and
 132 peaks one year *after* the wind stress curl reaches its maximum (Fig. 4f), as expected from
 133 the subpolar gyre response (Lohmann et al., 2009). We sum up:

- 134 1. Increased freshening of the subpolar region gives rise to a faster surface cooling.
- 135 2. After large freshwater anomalies, a distinct cold anomaly appears.
- 136 3. The cold anomaly promotes an enhanced storminess.
- 137 4. The resulting higher wind stress curl strengthens the subpolar gyre circulation.
- 138 5. The stronger subpolar gyre circulation reinforces the cold anomaly.

139 This chain of events is supported by the change in the sign of the heat flux anomaly,
 140 which is initially downward due to the reduced air-sea contrast (Fig. 3d) and then up-
 141 ward and driven by the atmospheric forcing (Fig. 4c). However, the discrepancy between
 142 the two heat flux anomalies also reveals a nonlinearity in the direct relation between fresh-
 143 water and the atmosphere: There exists a threshold for the freshwater, after which the
 144 sign of the heat flux anomaly reverses, reflecting the transition from the ocean-driven
 145 to the atmospherically-driven heat flux anomaly.

146 A composite of the largest freshwater events over the investigated period, included
 147 in the Supporting Information, shows the transition of the heat flux anomaly from be-
 148 ing positive in January and February to negative in March. It shows that the most *neg-*
 149 *ative* NAO summers were followed by a *positive* NAO in winter. A regression on these
 150 large freshwater events reveals, in addition, a high sensitivity of the atmospheric response
 151 to small variations in the freshwater when the freshwater concentration is already high,
 152 consistent with the nonlinear relation between F_{NA} and the SST (Figs. 3c and S7–S10).

153 2.5 Implications of freshwater events for the North Atlantic climate

154 Considering that the observational record is still short with regard to the occur-
 155 rence of large freshwater events, we inspect the evolution of these events individually.
 156 Two large freshwater events are the Great Salinity Anomaly in 1969–1972 (Lazier, 1980)
 157 and the recent freshening (Holliday et al., 2020) peaking in 2008–2012 and 2015/2016
 158 (Supporting Information). The Great Salinity Anomaly preceded a cold anomaly of \sim -
 159 1.5 °C while the recent freshening culminated in an anomalous \sim -2 °C (Supporting In-
 160 formation). Both events were followed by an enhanced storminess, reflected in increases
 161 of the winter NAO of ~ 0.61 after 1970 and ~ 0.72 after 2010 (Fig. 5a).

162 Additional salinity anomalies have been reported for the 1980’s and 1990’s (Belkin
 163 et al., 1998). Although the sparse sampling prevents the determination of the exact start-
 164 ing dates, compilations of historical salinity data suggest that they appeared in the Labrador
 165 Sea in 1980 (Reverdin et al., 1997; Yashayaev & Loder, 2016) and 1993 (Yashayaev &
 166 Clarke, 2006; Yashayaev & Loder, 2016), consistent with increased values of F_{NA} (Fig.
 167 1a). Both events were followed by a reduced SLP and a negative SST anomaly in the
 168 subpolar region within three years (Fig. 5).




Figure 5. (a) Variability of the 3-winter low-pass filtered NAO index. The solid lines and shadings indicate the times of large documented freshwater anomalies and the dashed line marks the time of an inferred event. (b-g) SLP and SST anomalies following the events in 1980, 1986 and 1993. While the anomalies typically last for several winters, we show them at the time when they reach their maximum amplitude: (b) SLP 1982, (c) SST 1983, (d) SLP 1989, (e) SST 1990, (f) SLP 1994, (g) SST 1994. All anomalies are with respect to the climatological winter mean.

169 The detailed evolution of the cold anomaly differed between the investigated fresh-
 170 water events, which we attribute to variations in the freshwater volumes, their pathways
 171 (Belkin et al., 1998) and the strength of the surface fluxes, mixing freshwater down. Also,
 172 the event in 1980 preceded an increase in the winter NAO of ~ 0.46 , similar to the other
 173 events, whereas the event in 1993 occurred in a phase, in which the winter NAO was al-
 174 ready high (Fig. 5a). Thus, the subpolar gyre circulation was already intensified (Belkin,
 175 2004) and the maximum SST anomaly was reached sooner (Fig. 5g).

176 Examining the winter NAO more closely, we detect another period of increase start-
 177 ing in 1986 (Fig. 5a), concurrent with an elevated value of F_{NA} (Fig. 1a). The follow-
 178 ing winters were characterised by a strongly reduced SLP and a distinct cold anomaly,
 179 exceeding ~ -2 °C in the Labrador Sea (Fig. 5d and e). We conclude that, despite dif-
 180 ferences in their amplitude and detailed evolution, all freshwater events show a link with
 181 an enhanced wintertime storminess and the emergence of a cold anomaly. Conversely,
 182 all major NAO increases in the last 50 years show a link with a freshwater event.

183 3 Conclusion

184 Combining in-situ observations with remote sensing and atmospheric reanalysis data,
 185 we have shown that enhanced freshening of the subpolar North Atlantic gives rise to an
 186 increased surface cooling in fall and winter. Over the last four decades, the freshwater-
 187 induced surface cooling has been characterised by a significantly positive trend, reflect-
 188 ing a growing discrepancy between the summer and winter SSTs.

189 Superimposed on the trend, individual strong freshwater events have triggered pro-
 190 nounced cold anomalies in winter. By increasing the meridional SST gradient, cold anoma-
 191 lies in the subpolar region promote the development of cyclones, which reinforce the cool-
 192 ing by modulating the subolar gyre circulation. In agreement with this chain of events,
 193 past freshwater events were followed by cold anomalies of ~ -2 °C and an enhanced win-
 194 tertime storminess, reflected in increases of the NAO of up to ~ 0.6 within three years.

195 Our findings suggest that the cold anomaly in winter can be explained by an en-
 196 hanced surface freshening and the resulting regional atmosphere-ocean interactions, with-
 197 out the need for a buoyancy-driven slowdown of the large-scale overturning circulation.
 198 Thus, this study reconciles the proposed inconsistencies between the recent strong con-
 199 vection in the Labrador Sea and the reduced northward heat transport (Lozier et al., 2019).
 200 However, the results do not exclude the possibility of a buoyancy-driven slowdown of the
 201 overturning on longer time scales.

202 The identified, freshwater-induced SST pattern resembles the negative phase of the
 203 Atlantic Multidecadal Oscillation, which has been found to drive a positive NAO response
 204 in winter (Peings & Magnusdottir, 2014; Gastineau & Frankignoul, 2015). Our findings
 205 are therefore consistent with earlier studies and suggest that freshwater plays a key role
 206 in modulating the low-frequency climate variability over the North Atlantic. In partic-
 207 ular, they show that large freshwater events, arising from the most negative NAO sum-
 208 mers, were followed by rapid transitions to the most positive NAO phases in winter.

209 With regard to the influences of the winter NAO on large-scale weather patterns
 210 (Hurrell & Deser, 2010), freshwater-induced NAO increases have far-reaching implica-
 211 tions. For instance, they can trigger cold spells over Canada (Shabbar & Bonsal, 2004),
 212 precipitation events over the UK (Lavers et al., 2011) and storms over northwest Europe
 213 (Feser et al., 2015). Considering the growing freshwater discharges from Greenland (Bamber
 214 et al., 2012) and the Arctic (Haine et al., 2015), and the high sensitivity to small vari-
 215 ations in the freshwater, when the freshwater concentration is already high, our results
 216 raise caution that weather extremes associated with deep cyclones in the subpolar re-
 217 gion in winter will increase.

Acknowledgments

The Argo data were collected and made freely available by the international Argo project and the national programs that contribute to it (<http://doi.org/10.17882/42182>). We further thank the National Oceanic and Atmospheric Administration and the Hadley Centre for providing the SST data (<https://psl.noaa.gov/data/gridded/data.noaa.oisst.v2.html> and <https://www.metoffice.gov.uk/hadobs/hadisst/>), the Copernicus Marine Service for distributing the altimeter products (<https://marine.copernicus.eu>) and the European Centre for Medium-Range Weather Forecasts for their reanalysis products (<https://www.ecmwf.int/en/forecasts/datasets>).

This study was supported by the EU Horizon 2020 programme under grant agreements 727852 and 633211, the German BMBF project “RACE” and the UK NERC programmes NE/N018044/1 and NE/R015953/1.

References

- Bamber, J., van den Broeke, M., Ettema, J., Lenaerts, J., & Rignot, E. (2012). Recent large increases in freshwater fluxes from Greenland into the North Atlantic. *Geophysical Research Letters*, *39*(19).
- Barber, D. C., Dyke, A., Hillaire-Marcel, C., Jennings, A. E., Andrews, J. T., Kerwin, M. W., ... others (1999). Forcing of the cold event of 8,200 years ago by catastrophic drainage of Laurentide lakes. *Nature*, *400*(6742), 344.
- Barnston, A. G., & Livezey, R. E. (1987). Classification, seasonality and persistence of low-frequency atmospheric circulation patterns. *Monthly weather review*, *115*(6), 1083–1126.
- Belkin, I. M. (2004). Propagation of the “great salinity anomaly” of the 1990s around the northern north atlantic. *Geophysical Research Letters*, *31*(8).
- Belkin, I. M., Levitus, S., Antonov, J., & Malmberg, S.-A. (1998). “great salinity anomalies” in the North Atlantic. *Progress in Oceanography*, *41*(1), 1–68.
- Blackmon, M. L. (1976). A climatological spectral study of the 500 mb geopotential height of the northern hemisphere. *Journal of the Atmospheric Sciences*, *33*(8), 1607–1623.
- Blackmon, M. L., Wallace, J. M., Lau, N.-C., & Mullen, S. L. (1977). An observational study of the Northern Hemisphere wintertime circulation. *Journal of the Atmospheric Sciences*, *34*(7), 1040–1053.
- Caesar, L., Rahmstorf, S., Robinson, A., Feulner, G., & Saba, V. (2018). Observed fingerprint of a weakening atlantic ocean overturning circulation. *Nature*, *556*(7700), 191.
- Chafik, L., Nilsen, J. E. Ø., Dangendorf, S., Reverdin, G., & Frederikse, T. (2019). North atlantic ocean circulation and decadal sea level change during the altimetry era. *Scientific reports*, *9*(1), 1–9.
- Clark, P. U., Marshall, S. J., Clarke, G. K., Hostetler, S. W., Licciardi, J. M., & Teller, J. T. (2001). Freshwater forcing of abrupt climate change during the last glaciation. *Science*, *293*(5528), 283–287.
- Clark, P. U., Pisias, N. G., Stocker, T. F., & Weaver, A. J. (2002). The role of the thermohaline circulation in abrupt climate change. *Nature*, *415*(6874), 863.
- Cronin, M., & Sprintall, J. (2009). Wind-and buoyancy-forced upper ocean. *Elements of Physical Oceanography: A derivative of the Encyclopedia of Ocean Sciences*, 237–245.
- Czaja, A., & Frankignoul, C. (2002). Observed impact of atlantic sst anomalies on the North Atlantic Oscillation. *Journal of Climate*, *15*(6), 606–623.
- Dee, D. P., Uppala, S., Simmons, A., Berrisford, P., Poli, P., Kobayashi, S., ... others (2011). The era-interim reanalysis: Configuration and performance of the data assimilation system. *Quarterly Journal of the royal meteorological society*, *137*(656), 553–597.

- 270 Deser, C., Tomas, R. A., & Peng, S. (2007). The transient atmospheric circula-
 271 tion response to north atlantic sst and sea ice anomalies. *Journal of Climate*,
 272 *20*(18), 4751–4767.
- 273 Feser, F., Barcikowska, M., Krueger, O., Schenk, F., Weisse, R., & Xia, L. (2015).
 274 Storminess over the north atlantic and northwestern europe—a review. *Quar-*
 275 *terly Journal of the Royal Meteorological Society*, *141*(687), 350–382.
- 276 Foukal, N. P., & Lozier, M. S. (2018). Examining the origins of ocean heat content
 277 variability in the eastern north atlantic subpolar gyre. *Geophysical Research*
 278 *Letters*, *45*(20), 11–275.
- 279 Gastineau, G., & Frankignoul, C. (2015). Influence of the north atlantic sst variabil-
 280 ity on the atmospheric circulation during the twentieth century. *Journal of Cli-*
 281 *mate*, *28*(4), 1396–1416.
- 282 Gill, A. E. (2016). *Atmosphere—ocean dynamics*. Elsevier.
- 283 Griffies, S. M., & Greatbatch, R. J. (2012). Physical processes that impact the evo-
 284 lution of global mean sea level in ocean climate models. *Ocean Modelling*, *51*,
 285 37–72.
- 286 Haine, T. W., Curry, B., Gerdes, R., Hansen, E., Karcher, M., Lee, C., . . . others
 287 (2015). Arctic freshwater export: Status, mechanisms, and prospects. *Global*
 288 *and Planetary Change*, *125*, 13–35.
- 289 Häkkinen, S., Rhines, P. B., & Worthen, D. L. (2011). Warm and saline events em-
 290 bedded in the meridional circulation of the northern north atlantic. *Journal of*
 291 *Geophysical Research: Oceans*, *116*(C3).
- 292 Hanna, E., Jones, J. M., Cappelen, J., Mernild, S. H., Wood, L., Steffen, K., & Huy-
 293 brechts, P. (2013). The influence of north atlantic atmospheric and oceanic
 294 forcing effects on 1900–2010 greenland summer climate and ice melt/runoff.
 295 *International Journal of Climatology*, *33*(4), 862–880.
- 296 Holliday, N. P., Bersch, M., Berx, B., Chafik, L., Cunningham, S., Florindo-López,
 297 C., . . . others (2020). Ocean circulation causes the largest freshening event for
 298 120 years in eastern subpolar north atlantic. *Nature Communications*, *11*(1),
 299 1–15.
- 300 Holte, J., Talley, L. D., Gilson, J., & Roemmich, D. (2017). An argo mixed layer cli-
 301 matology and database. *Geophysical Research Letters*, *44*(11), 5618–5626.
- 302 Hoskins, B. J., McIntyre, M. E., & Robertson, A. W. (1985). On the use and sig-
 303 nificance of isentropic potential vorticity maps. *Quarterly Journal of the Royal*
 304 *Meteorological Society*, *111*(470), 877–946.
- 305 Hurrell, J. W., & Deser, C. (2010). North atlantic climate variability: the role of the
 306 north atlantic oscillation. *Journal of Marine Systems*, *79*(3-4), 231–244.
- 307 Josey, S. A., Hirschi, J. J.-M., Sinha, B., Duchez, A., Grist, J. P., & Marsh, R.
 308 (2018). The recent atlantic cold anomaly: Causes, consequences, and related
 309 phenomena. *Annual review of marine science*, *10*, 475–501.
- 310 Kennedy, J., Rayner, N., Smith, R., Parker, D., & Saunby, M. (2011). Reassessing
 311 biases and other uncertainties in sea surface temperature observations mea-
 312 sured in situ since 1850: 1. measurement and sampling uncertainties. *Journal*
 313 *of Geophysical Research: Atmospheres*, *116*(D14).
- 314 Kushnir, Y., Robinson, W., Bladé, I., Hall, N., Peng, S., & Sutton, R. (2002). Atmo-
 315 spheric gcm response to extratropical sst anomalies: Synthesis and evaluation.
 316 *Journal of Climate*, *15*(16), 2233–2256.
- 317 Lavers, D. A., Allan, R. P., Wood, E. F., Villarini, G., Brayshaw, D. J., & Wade,
 318 A. J. (2011). Winter floods in britain are connected to atmospheric rivers.
 319 *Geophysical Research Letters*, *38*(23).
- 320 Lazier, J. R. (1980). Oceanographic conditions at ocean weather ship bravo, 1964–
 321 1974. *Atmosphere-ocean*, *18*(3), 227–238.
- 322 Le Traon, P., Nadal, F., & Ducet, N. (1998). An improved mapping method of
 323 multisatellite altimeter data. *Journal of atmospheric and oceanic technology*,
 324 *15*(2), 522–534.

- 325 Lohmann, K., Drange, H., & Bentsen, M. (2009). Response of the north atlantic
326 subpolar gyre to persistent north atlantic oscillation like forcing. *Climate dy-*
327 *namics*, *32*(2-3), 273–285.
- 328 Lozier, M., Li, F., Bacon, S., Bahr, F., Bower, A., Cunningham, S., . . . others
329 (2019). A sea change in our view of overturning in the subpolar north atlantic.
330 *Science*, *363*(6426), 516–521.
- 331 Peings, Y., & Magnusdottir, G. (2014). Forcing of the wintertime atmospheric cir-
332 culation by the multidecadal fluctuations of the north atlantic ocean. *Environ-*
333 *mental Research Letters*, *9*(3), 034018.
- 334 Percival, D. B., Walden, A. T., et al. (1993). *Spectral analysis for physical applica-*
335 *tions*. cambridge university press.
- 336 Rahmstorf, S., Box, J. E., Feulner, G., Mann, M. E., Robinson, A., Rutherford, S.,
337 & Schaffernicht, E. J. (2015). Exceptional twentieth-century slowdown in
338 atlantic ocean overturning circulation. *Nature climate change*, *5*(5), 475.
- 339 Reverdin, G. (2010). North atlantic subpolar gyre surface variability (1895–2009).
340 *Journal of climate*, *23*(17), 4571–4584.
- 341 Reverdin, G., Cayan, D., & Kushnir, Y. . (1997). Decadal variability of hydrography
342 in the upper northern north atlantic in 1948–1990. *Journal of Geophysical Re-*
343 *search: Oceans*, *102*(C4), 8505–8531.
- 344 Reynolds, R. W., Rayner, N. A., Smith, T. M., Stokes, D. C., & Wang, W. (2002).
345 An improved in situ and satellite sst analysis for climate. *Journal of climate*,
346 *15*(13), 1609–1625.
- 347 Sarafanov, A. (2009). On the effect of the north atlantic oscillation on temperature
348 and salinity of the subpolar north atlantic intermediate and deep waters. *ICES*
349 *Journal of Marine Science*, *66*(7), 1448–1454.
- 350 Shabbar, A., & Bonsal, B. (2004). Associations between low frequency variability
351 modes and winter temperature extremes in canada. *Atmosphere-Ocean*, *42*(2),
352 127–140.
- 353 Spall, M. A., & Pickart, R. S. (2003). Wind-driven recirculations and exchange
354 in the labrador and irvinger seas. *Journal of Physical Oceanography*, *33*(8),
355 1829–1845.
- 356 Sutton, R. T., & Dong, B. (2012). Atlantic ocean influence on a shift in european
357 climate in the 1990s. *Nature Geoscience*, *5*(11), 788.
- 358 Ulbrich, U., Pinto, J. G., Kupfer, H., Leckebusch, G., Spangehl, T., & Reyers, M.
359 (2008). Changing northern hemisphere storm tracks in an ensemble of ipcc
360 climate change simulations. *Journal of Climate*, *21*(8), 1669–1679.
- 361 Yashayaev, I., & Clarke, A. (2006). Recent warming of the labrador sea. *AZMP Bul-*
362 *letin PMZA*, *5*, 12–20.
- 363 Yashayaev, I., & Loder, J. W. (2016). Recurrent replenishment of labrador sea wa-
364 ter and associated decadal-scale variability. *Journal of Geophysical Research:*
365 *Oceans*, *121*(11), 8095–8114.