



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Enhanced fluorescence from X-ray line coincidence pumping

J. Nilsen, D. Burrridge, L. M. R. Hobbs, D. Hoarty, P.
Beiersdorfer, G. V. Brown, N. Hell, D. Panchenko, M. F. Gu, A.
M. Saunders, H. A. Scott, P. Hatfield, M. P. Hill, L. Wilson, R.
Charles, C. R. D. Brown, S. Rose

November 26, 2018

16th International Conference on X-ray Lasers
Prague, Czech Republic
October 7, 2018 through October 12, 2018

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

Enhanced fluorescence from X-ray line coincidence pumping

Joseph Nilsen¹, D. Burrige², L. M. R. Hobbs², D. Hoarty², P. Beiersdorfer¹, G. V. Brown¹, N. Hell¹, D. Panchenko¹, M. F. Gu³, A. M. Saunders¹, H. A. Scott¹, P. Hatfield⁴, M. P. Hill², L. Wilson², R. Charles², C. R. D. Brown², and S. Rose⁴

¹Lawrence Livermore National Laboratory, Livermore, CA 94551

²AWE Plc, UK

³University of California, Berkeley, CA

⁴Imperial College, UK

Abstract. Many resonant photo-pumped X-ray laser schemes that use a strong pump line such as Ly- α or He- α to populate the upper laser state of a separate lasing material have been proposed over the last four decades but none have been demonstrated. As a first step to creating a photo-pumped X-ray laser we have decided to reinvestigate some of these schemes at the Orion laser facility with the goal to show enhanced fluorescence. In particular we look at using the Ly- α or He- α K lines to pump the 1s – 3p and 4p transitions in H-like Cl and see fluorescence on the 4f – 3d line at 65 Å and the 3d – 2p line at 23 Å. Preliminary experiments are presented that show a modest enhancement. As an alternative we also look at enhancing the 2p – 2s line in Ne-like Ge at 65Å using the Ly- α Mg line to photo-pump the 2s – 3p line of Ne-like Ge. Calculations are presented that suggest modest enhancements of 2.5.

1 Introduction

One early approach for creating X-ray lasers was the idea of a resonantly photo-pumped laser where a strong emission line in one material could be used to photo-excite a transition in another material and create lasing. The Na-pumped Ne X-ray laser scheme proposed 40 years ago by Vinogradov and colleagues [1,2] is a classic example that used the strong Na He- α line at 1127 eV to resonantly photo-pump the Ne He- γ line and lase on the 4f – 3d transition at 231 Å in He-like Ne. This scheme was studied extensively and weak gain [3] was inferred in several experiments. There are a series of similar resonances [2] that use Ly- α or He- α lines to photo-pump Ly- γ or He- γ lines and lase on the 4f – 3d lines of H or He-like ions. In this paper we discuss the next higher Z pair of possible resonances that use the strong Ly- α line at 3699 eV from H-like K

to resonantly photo-pump the $1s - 4p$ transition in Cl and enhance the emission of the $4f - 3d$ line at 65 \AA in H-like Cl.

One can also pump the Ly- β or He- β lines and lase or enhance the fluorescence of the $3d - 2p$ lines. The K-Cl combination is interesting because it also has a resonance between the strong K He- α line to resonantly photo-pump the Cl He- β transition and enhance the emission of the $3d - 2p$ line at 23 \AA in H-like Cl.

Experiments were done at the Orion laser facility in the UK to look at enhanced emission on both the 23 and 65 \AA Cl lines.

An alternative experiment is being designed to look at the emission of the $2p - 2s$ line in Ne-like Ge at 65 \AA that could be photo-pumped by the Ly- α line of H-like Mg at 1472 eV . In this case the Ly- α line photo-pumps the inner shell $2s - 3p$ ground state transition. This resonance was measured in previous experiments at the EBIT facility at LLNL, the NPO laser facility at VNIIFTRI, and the Princeton Large Torus (PLT) tokamak. Experiments are being planned at the Orion laser facility to investigate this scheme.




Fig. 1. Energy level diagram for the Ly- α and He- α lines of K photo-pumping the $1s - 3p$ and $4p$ lines of H-like Cl to enhance the fluorescence of the $4f - 3d$ line at 64.8 \AA and the $3d - 2p$ line at 22.7 \AA .

2 K-pumped Cl experiments

Figure 1 shows the energy level diagram for the Potassium (K)-pumped Chlorine (Cl) schemes. The 4f – 3d line at 64.8 Å in H-like Cl is enhanced by having the strong Ly- α line of K photo-pump the 1s – 4p (Ly- γ) transition of H-like Cl. The $n = 4$ states of Cl are all very close in energy and become statistically populated by collisional processes at the temperatures and densities expected in the plasma. The 4f level has the largest statistical weight ($2J + 1$) which results in strong emission and potential gain on the 4f – 3d line. Potassium is rather unique because the He- α line, which is usually the strongest line one can produce in a hot plasma, is nearly resonant with the 1s – 3p (Ly- β) line of Cl. This photo-pumps population into the 3p level which then equilibrates through collisional processes to statistically populate the $n=3$ level resulting in a large population in the 3d level. One then looks for strong emission on the 3d – 2p line at 22.7 Å. This line is also part of the cascade from the 4f-3d emission which also enhances the line. This scheme was originally proposed and modelled [3,4] many years ago.

To look for potential enhancements, experiments were done at the Orion laser facility [5] at the atomic weapons establishment (AWE) in the UK. The laser systems consist of ten beams, optimized around 1 ns pulse duration, which each provide a nominal 500 J at a wavelength of 351 nm. There are also two short pulse beams, which each provide 500 J in 0.5 ps at 1054 nm, that were not used in these experiments.

The targets consisted a 0.5 μm thick microdot of KCl with a diameter of 300 μm . This was coated on each side with 0.5 μm thick layer of KF and then an




Fig. 2. Emission from the NaCl target and KCl target showing the strong Ly- α and He- α emission lines from K and Cl.

outer protective coating of $0.3 \mu\text{m}$ thick layer of parylene-N, which is a CH plastic polymer. The goal of the KF layer was to have a thick layer of K surrounding the Cl plasma that would enhance the strength of the Ly- α and He- α K lines pumping the Cl. As a null experiment we used targets that replaced the KCl with NaCl and the KF with NaF. In the null experiment there are no K lines to photo-pump the Cl plasma.

The targets were hit from both side by 250-1150 J of energy in a $300 \mu\text{m}$ diameter beam with a 1-1.5 ns FWHM pulse duration. Hydrodynamic calculations done with the 1D NYM code predict that the plasma would expand and blow down to densities of about 1-2 mg/cc or electron density of $3-6 \times 10^{20}/\text{cc}$ and reach temperatures up to several keV. Under these conditions we expect to ionize the plasmas to H and He-like states. We observe strong emission, as seen in Fig. 2, on the K and Cl Ly- α and He- α lines for the nominal target and strong emission on the Cl Ly- α and He- α lines for the null target indicating that we have reached the correct ionization conditions. This data was recorded on the Mark 2 time-integrated spectrometer with a resolution of 7 eV using CsAP crystals.

To look at the spectral lines near 23 and 65 Å we used a grating instrument spectrometer that covered the range of 100 – 1000 eV (120-12 Å) with a resolution of 1000. Figure 3 shows the spectrum from 21-27 Å that looks at the enhancement of the Cl 3d-2p line at 22.7 Å. When we normalize to the other Cl lines we believe we observe a modest enhancement of about 20% on this line due to the photo-pumping. The spectrum near the 65 Å line is more




Fig. 3. Emission from the NaCl and KCl targets shows modest enhancement of the H-like Cl 3d-2p line at 22.7 Å in the KCl target.

complicated and still being analysed as there are nearby lines in Li-like Cl in the same spectral region.

3 Mg-pumped Ge experiments

The second scheme that we are actively pursuing, shown in Fig. 4, uses the Magnesium (Mg) Ly- α lines to photo-pump the Neon (Ne)-like Germanium (Ge) $2s - 3p$ line (B) and enhance the emission of the $2p - 2s$ line in Ne-like Ge at 65.1 \AA . The resonance between the Mg and Ge lines were measured in previous experiments at the EBIT facility at LLNL, the NPO laser facility at VNIIFTRI, and the Princeton Large Torus (PLT) tokamak. [6] Experiments are being planned at the Orion laser facility to investigate this scheme.

These experiments would be very similar to the KCL experiments where we use microdot targets with a $0.2 \mu\text{m}$ thick Ge target surrounded by $0.3 \mu\text{m}$ of Mg for the pumped experiment and $0.3 \mu\text{m}$ of Al for the null experiment. Both sets of targets are overcoated with $0.3 \mu\text{m}$ of Parylene-N on each side as a protective layer from the atmosphere. To investigate the viability of this scheme we have done a series of kinetics and radiation transport calculations with the Cretin




Fig. 4. Energy level diagram for the Mg Ly- α photo-pumping the Ne-like Ge $2s - 3p$ line to enhance the fluorescence of the $2p-2s$ line at 65.1 \AA




Fig. 5. Line strength for the Mg Ly- α line vs photon energy centered around the Ly- α 2 line.




Fig. 6. Predicted emission from the Ge target comparing the case where it is pumped by the Mg Ly- α line to the case without any pumping.

code [7] to estimate enhancements that we might see in the upcoming

experiments at the Orion facility.

If we assume the targets have blown down in density to an electron density of $5 \times 10^{20}/\text{cc}$ (1.8 mg/cc) at a temperature of 500 eV we can first estimate the strength and width of the Mg Ly- α alpha line. Doing a 1D line transport calculation assuming a Mg target with a radius of $350 \mu\text{m}$, Fig 5 shows the line strength for the Mg Ly- α in the center of the target. One sees that the line has a width of 2.1 eV with the Ge line located at an energy of 0.6 eV relative to the Ly- α 2 line. This makes for essentially a perfect resonance between the Mg and Ge lines. The Mg line has a peak strength of 1.2×10^{-3} photons per mode in the center of the target and falls off by a factor of 6 at the outer surface of the target. When pumping a line, the line strength in photons per mode is very similar to the fractional population of the upper state being pumped.

The next step is to do a kinetics calculation for the spectra of Ge near 65 \AA with and without the presence of the Mg line using the pump strength calculated above. To best estimate the spectra, we created an atomic model for Ge that uses many body perturbation theory calculations for the energy levels of Ne-like Ge $n=2$ and $n=3$ levels. These calculations have been compared with detail spectra taken at the EBIT facility [8] at LLNL.

Taking the same plasma conditions described above for Ge we use Cretin to calculate the Ge spectra. Without the Mg pump present we observe a series of lines that are dominated by the very strong F-like Ge $2p - 2s$ line at 65.9 \AA . When we do a similar calculation including the Mg pump line the spectra are very similar except for a significant enhancement of about 2.5 on the $2p - 2s$ Ne-like Ge line at 65.06 \AA , as shown in Fig. 6. This is the type of enhancement we are trying to observe in the future experiments. One advantage of looking for enhanced fluorescence instead of gain is that it can take a more substantial enhancement of the line before it has enough gain to lase.

4 Conclusions

In this paper we look at using the K Ly- α or He- α lines to photo-pump the $1s - 3p$ and $4p$ transitions in H-like Cl and see fluorescence on the $4f - 3d$ line at 64.8 \AA and the $3d - 2p$ line at 22.7 \AA . Preliminary experiments are presented that show a modest enhancement of about 20% on the 22.7 \AA line. As an alternative we also look at enhancing the $2p - 2s$ line in Ne-like Ge at 65.1 \AA using the Mg Ly- α line to photo-pump the $2s - 3p$ line of Ne-like Ge. Calculations are presented that suggest modest enhancements of 2.5. Experiments are planned in the near future to test the Mg-pumped Ge scheme.

Acknowledgements. This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344. Add AWE or other acknowledgments.

References

1. Vinogradov, A. V., Sobelman, I. I., Yukov, E. A., *Sov. J. Quantum Electron.* **5**, 59–63, 1975.
2. Nilsen, J., Scofield, J. H., Chandler, E. A., *Appl. Opt.* **31**, 4950 – 4956, 1992.
3. Nilsen J., Chandler, E. A., *Phys. Rev. A* **44**, 4591-4598, 1991.
4. Rose, S. J., *J. Phys. B* **31**, 2129-2144, 1998.
5. Hopps, N. et al., *Plasma Phys. Control Fusion* **57**, 064002, 2015.
6. Nilsen, J. et al., *Phys. Rev A* **50**, 2143, 1994.
7. Scott, H. A., *JQSRT* **71**, 689–701, 2001.
8. Beiersdorfer, P., *Nuclear Instruments and Methods in Physics Research B* **205**, 173, 2003.