

## Interventions for attentional disruption in pain

1

1 Interventions for attentional disruption in pain:  
2 cognition-general, mechanism-specific or exercise-based?

3

4 Nina Attridge and Nils Niederstrasser

5 Loughborough University

6

7 Keywords: cognition; attention; disruption; interventions; cognitive training

8

9 In the current issue of PAIN, Baker et al. [6] report that a computerised cognitive training  
10 programme was able to improve chronic pain patients' performance on a neurocognitive  
11 assessment composite. Decades' worth of research has supported the hypothesis that pain  
12 disrupts attention [3-5; 9-14; 17; 20; 21; 28], with the ultimate goal being to reverse this  
13 effect in people with pain. As far as we are aware, this is the first study that attempts to do  
14 so.

15

16 Improving attention in people with chronic pain is a very worthwhile goal. Patients often  
17 report that pain makes it difficult to think clearly [1; 2; 7; 14; 16], and disruption to attention  
18 has the potential to impact on many, if not all, areas of life, such as work, study, socialising  
19 and mood, to name a few. We are not always able to reduce patients' pain, so many people  
20 must learn to live with their pain long-term. By improving their cognitive function, we may  
21 be able to improve quality of life for people living with chronic pain.

22

23 How should we go about developing interventions to improve attention in people with  
24 pain? One way is to use commercially available computerised cognitive training

25 programmes, as done by Baker et al [6]. Such intervention programmes are usually targeted  
26 at older adults to prevent cognitive decline, and operate under the premise that practice on  
27 tasks that load heavily on core cognitive domains leads to improvements on other everyday  
28 tasks that rely on these domains. This concept is often compared to how exercise improves  
29 physical fitness.

30

31 Definitive statements pertaining to the efficacy of computerised cognitive training are  
32 contentious and beyond the scope of this commentary. Generally, computerised cognitive  
33 training is believed to be effective in improving function on the practised cognitive domains,  
34 but efficacy varies by domain and improvements seldom transfer far beyond the practiced  
35 domain or task [18; 23; 24]. The efficacy for improving attention in particular has been  
36 contested [18] and it remains to be determined whether the effects that have been  
37 reported represent improvements in attentional capacity, the acquisition of a new strategy,  
38 or increased motivation. This has implications for applying computerised cognitive training  
39 to improving chronic pain patients' attention: acquired strategies are highly context- or task-  
40 dependent and unlikely to transfer to novel contexts.

41

42 Aside from disputes pertaining to its efficacy, the approach of using computerised cognitive  
43 training in chronic pain patients has the drawback that interventions tend to be very broad,  
44 i.e. training a wide range of cognitive domains at the same time. This may produce small  
45 positive effects, but the training will be time-consuming for users and difficult for them to  
46 sustain over time. This may be an appropriate approach for programmes that aim to  
47 improve or maintain cognitive function in general, as is the case in computerised cognitive

48 training. But when the aim is to reverse an effect caused by a specific trigger, such as pain,  
49 many aspects of a catch-all intervention could be unnecessary.

50

51 Another approach to developing interventions for people with pain, and the one we wish to  
52 advocate, would be to target the specific mechanisms of the effect of pain on attention. For  
53 example, if pain and ongoing tasks are processed simultaneously (i.e. if pain acts as a  
54 working memory load), then perhaps interventions should specifically target working  
55 memory capacity. Alternatively, if attention is disrupted because our focus shifts frequently  
56 from our ongoing tasks to our pain, then perhaps interventions should target attentional  
57 control and inhibitory control. These are just two examples of possible mechanisms in the  
58 cognitive domain: the mechanisms may instead be metacognitive, motivational [27], or  
59 otherwise.

60

61 Unfortunately, we do not yet know enough about the mechanisms of the disruptive effect  
62 of pain on attention, and what the individual differences may be, to develop such targeted  
63 interventions. But if we could develop targeted interventions, the demand on patients' time  
64 could be substantially reduced or spent more effectively, which could reduce the risk of  
65 attrition over the course of programmes. If time demands are minimised and patients are  
66 provided with strong theoretical reasons why an intervention will work, alongside evidence  
67 that it does, they may be more motivated to stick with it for long enough to see results.

68

69 If the improvement or maintenance of cognitive function in general is indeed the goal,  
70 rather than targeting a specific mechanism, a third approach would be to increase exercise

71 within the constraints of the patient's condition. Not only does exercise improve cognitive  
72 function, it can also help prevent chronic diseases and improve mood [8; 15; 19; 25].

73

74 The specific nature of interventions aside, we can only develop effective interventions if the  
75 research they are based on and the studies testing their effectiveness are conducted in a  
76 rigorous manner. Pre-registration of study protocols, methods, hypotheses, measures and  
77 analyses is vital to ensure transparency and reproducibility of research. Furthermore, it aids  
78 in the fight against publication bias and p-hacking. Sample size calculations, active control  
79 groups, delayed post-tests and replication studies will also allow us to place trust in research  
80 findings. [22; 26]

81

82 Baker et al. [6] have shown that there is potential for improving cognition in people with  
83 chronic pain, which gives hope that as we develop and refine interventions, we will be able  
84 to substantially improve the everyday lives of patients. However, in our view, we need to  
85 fully understand the mechanisms of the effect of pain on attention before we can develop  
86 interventions that will be as effective, time-effective, and cost-effective as we would like.

87

88

#### 89 Conflict of Interest Statement

90 The authors have no conflicts of interest.

## 91 References

92

- 93 [1] Arnold LM, Crofford LJ, Mease PJ, Burgess SM, Palmer SC, Abetz L, Martin SA. Patient  
94 perspectives on the impact of fibromyalgia. *Patient Education and Counseling*  
95 2008;73(1):114-120.
- 96 [2] Attridge N, Crombez G, Van Ryckeghem D, Keogh E, Eccleston C. The Experience of  
97 Cognitive Intrusion of Pain: scale development and validation. *PAIN*  
98 2015;156(10):1978-1990.
- 99 [3] Attridge N, Eccleston C, Noonan D, Wainwright E, Keogh E. Headache impairs attentional  
100 performance: a conceptual replication and extension. *The Journal of Pain*  
101 2017;18:29-41.
- 102 [4] Attridge N, Keogh E, Eccleston C. The effect of pain on task switching: pain reduces  
103 accuracy and increases reaction times across multiple switching paradigms. *PAIN*  
104 2016.
- 105 [5] Attridge N, Noonan D, Eccleston C, Keogh E. The disruptive effects of pain on n-back task  
106 performance in a large general population sample. *PAIN* 2015;156(10):1885-1891.
- 107 [6] Baker KS, Georgiou-Karistianis N, Lampit A, Valenzuela M, Gibson SJ, Giummarra MJ.  
108 Computerised Training Improves Cognitive Performance in Chronic Pain: A  
109 Participant-Blinded Randomised Active-Controlled Trial with Remote Supervision.  
110 *PAIN* 2018.
- 111 [7] Bennett RM, Jones J, Turk DC, Russell IJ, Matallana L. An internet survey of 2,596 people  
112 with fibromyalgia. *BMC Musculoskeletal Disorders* 2007;8(1):27.
- 113 [8] Bherer L, Erickson KI, Liu-Ambrose T. A review of the effects of physical activity and  
114 exercise on cognitive and brain functions in older adults. *Journal of Aging Research*  
115 2013;2013.
- 116 [9] Buhle J, Wager TD. Performance-dependent inhibition of pain by an executive working  
117 memory task. *PAIN* 2010;149(1):19-26.
- 118 [10] Crombez G, Eccleston C, Baeyens F, Eelen P. The disruptive nature of pain: an  
119 experimental investigation. *Behaviour Research and Therapy* 1996;34(11):911-918.
- 120 [11] Crombez G, Eccleston C, Baeyens F, Eelen P. Attentional disruption is enhanced by the  
121 threat of pain. *Behaviour Research Therapy* 1998;36(2):195-204.
- 122 [12] Crombez G, Eccleston C, Van den Broeck A, Van Houdenhove B, Goubert L. The effects  
123 of catastrophic thinking about pain on attentional interference by pain: no mediation  
124 of negative affectivity in healthy volunteers and in patients with low back pain. *Pain*  
125 *Research and Management* 2002;7(1):31-39.
- 126 [13] Eccleston C. Chronic pain and distraction: an experimental investigation into the role of  
127 sustained and shifting attention in the processing of chronic persistent pain.  
128 *Behaviour Research and Therapy* 1995;33(4):391-405.
- 129 [14] Glass JM, Park DC. Cognitive dysfunction in fibromyalgia. *Current Rheumatology*  
130 *Reports* 2001;3(2):123-127.
- 131 [15] Gomez-Pinilla F, Hillman C. The influence of exercise on cognitive abilities.  
132 *Comprehensive Physiology* 2013.
- 133 [16] Katz RS, Heard AR, Mills M, Leavitt F. The prevalence and clinical impact of reported  
134 cognitive difficulties (fibrofog) in patients with rheumatic disease with and without  
135 fibromyalgia. *JCR: Journal of Clinical Rheumatology* 2004;10(2):53-58.

- 136 [17] Keogh E, Cavill R, Moore DJ, Eccleston C. The effects of menstrual-related pain on  
137 attentional interference. *PAIN* 2014;155(4):821-827.
- 138 [18] Lampit A, Hallock H, Valenzuela M. Computerized cognitive training in cognitively  
139 healthy older adults: a systematic review and meta-analysis of effect modifiers. *PLoS*  
140 *Medicine* 2014;11(11):e1001756.
- 141 [19] Liao Y, Shonkoff ET, Dunton GF. The acute relationships between affect, physical feeling  
142 states, and physical activity in daily life: a review of current evidence. *Frontiers in*  
143 *Psychology* 2015;6.
- 144 [20] Moore DJ, Keogh E, Eccleston C. The interruptive effect of pain on attention. *The*  
145 *Quarterly Journal of Experimental Psychology* 2012;65(3):565-586.
- 146 [21] Moore DJ, Keogh E, Eccleston C. Headache impairs attentional performance. *PAIN*  
147 2013;154(9):1840-1845.
- 148 [22] Munafò MR, Nosek BA, Bishop DV, Button KS, Chambers CD, du Sert NP, Simonsohn U,  
149 Wagenmakers E-J, Ware JJ, Ioannidis JP. A manifesto for reproducible science.  
150 *Nature Human Behaviour* 2017;1:0021.
- 151 [23] Owen AM, Hampshire A, Grahn JA, Stenton R, Dajani S, Burns AS, Howard RJ, Ballard  
152 CG. Putting brain training to the test. *Nature* 2010;465(7299):775-778.
- 153 [24] Simons DJ, Boot WR, Charness N, Gathercole SE, Chabris CF, Hambrick DZ, Stine-  
154 Morrow EA. Do “brain-training” programs work? *Psychological Science in the Public*  
155 *Interest* 2016;17(3):103-186.
- 156 [25] Smith PJ, Blumenthal JA, Hoffman BM, Cooper H, Strauman TA, Welsh-Bohmer K,  
157 Browndyke JN, Sherwood A. Aerobic exercise and neurocognitive performance: a  
158 meta-analytic review of randomized controlled trials. *Psychosomatic medicine*  
159 2010;72(3):239.
- 160 [26] Tackett JL, Lilienfeld SO, Patrick CJ, Johnson SL, Krueger RF, Miller JD, Oltmanns TF,  
161 Shrout PE. It’s time to broaden the replicability conversation: Thoughts for and from  
162 clinical psychological science. *Perspectives on Psychological Science* 2017;12(5):742-  
163 756.
- 164 [27] Van Damme S, Legrain V, Vogt J, Crombez G. Keeping pain in mind: A motivational  
165 account of attention to pain. *Neuroscience & Biobehavioral Reviews* 2010;34(2):204-  
166 213.
- 167 [28] Van Ryckeghem DM, Crombez G, Eccleston C, Liefoghe B, Van Damme S. The  
168 interruptive effect of pain in a multitask environment: an experimental investigation.  
169 *The Journal of Pain* 2012;13(2):131-138.  
170