

Oscheius myriophila (Nematoda: Rhabditida) isolated in sugar cane soils in Mexico with potential to be used as entomopathogenic nematode

Iveth del Rocio Castro-Ortega¹,
Juan Manuel Caspeta-Mandujano²,
Ramón Suárez-Rodríguez³,
Guadalupe Peña-Chora⁴,
José Augusto Ramírez-Trujillo³,
Karina Cruz-Pérez, Iván Arenas
Sosa⁵ and Víctor Manuel
Hernández-Velázquez^{1,*}

¹Universidad Autónoma del Estado de Morelos, Laboratorio de Control Biológico, Centro de Investigación en Biotecnología, Cuernavaca, México.

²Laboratorio de Parasitología Animal, Facultad de Ciencias Biológicas, Av. Universidad No. 1001, Col. Chamilpa, Cuernavaca C.P. 6220, Morelos, México.

³Laboratorio de Fisiología Molecular de Plantas, Centro de Investigación en Biotecnología, Cuernavaca, México.

⁴Laboratorio de Parasitología Vegetal, Centro de Investigaciones Biológicas, Cuernavaca, México.

⁵Universidad Nacional Autónoma de México Departamento de Medicina Molecular y Bioprocesos del IBT-UNAM. Av. Universidad 2001, Chamilpa, 62210 Cuernavaca, Morelos, México.

*E-mail: vmanuelh@uaem.mx

This article was edited by Eyuaalem Abebe.

Received for publication February 25, 2019.

Abstract

A survey of entomopathogenic nematodes was conducted in sugar cane crops in a total of 14 soils, and positive results were obtained for strain MC5-2014 in the municipality of Tepalcingo, Morelos, in soil with a sandy loam texture and a pH of 6.4. Species determination was performed via morphological and morphometric techniques by searching for a tubular stoma with a swollen cylindrical pharyngeal body and a metacarpus in the basal part. The range of body length (*L*) was 750 to 1200 μm in females and 720 to 910 μm in males, while the corresponding maximum widths (*W*) of the body were 30 to 60 μm and 20 to 30 μm, respectively. Males exhibited bursa with a 1+1+3+3 distribution of papillae, and females exhibited a vulva located at the mid-body. For molecular identification, the ITS region of ribosomal DNA was used. Virulence tests (LC_{50}) were conducted with *Galleria mellonella*, and a value of 4.732 was obtained for infective juveniles (IJs). Taking taxonomic and molecular characteristics into account, the isolate was determined to be *Oscheius myriophila*. The isolation of this strain represents the first geographic report of *O. myriophila* in Mexico, and it should be noted that the cultivation of sugar cane occurs with constant application of insecticides, herbicides, fungicides, and fertilizers as well as harvesting activities such as burning of the crop for harvest. The *O. myriophila* isolate has the potential to be used in the future as a method of biological control in our country.

Keywords

First report, México, *Oscheius myriophila*, sugar cane, virulence.

Entomopathogenic nematodes (EPNs) exhibit a cosmopolitan distribution and have been isolated on five continents in different habitats around the world (Griffin et al., 1990; Hominik et al., 1996). In Mexico, EPNs, particularly species of the genera *Steinernema* and *Heterorhabditis*, have been isolated in the desert, forests, and arable areas (Lezama et al., 2001; Stock et al., 2009). Worldwide, the main studies and reported isolates have come from these genera. Recently, two new species of EPNs of the genus *Oscheius* have been reported: *Oscheius chongmingensis* and *O. carolinensis*, which are associated with the pathogenic bacteria of insects of the genus *Serratia* (Nguyen

and Hunt, 2007; Abebe et al., 2010). The genus *Oscheius* was described as a sister taxon of *Rhabditis* (Dujardin, 1845) Sudhaus (1976). The genus *Oscheius* belongs to the Rhabditidae family and is divided into two groups: insectivora and dolichura. The species *O. carolinensis*, *O. amsactae*, *O. niazii*, and *O. siddiqii* of the insectivora group and one species of the dolichura group, *O. onirici*, have been recognized as EPNs (Tabassum and Shahina, 2010; Ye et al., 2010; Torrini et al., 2015). In 2017, it was reported that *O. myriophila* was associated with the European mole cricket, *Gryllotalpa gryllotalpa* (Orthoptera: Gryllotalpidae) (Zeynep et al., 2017). The objective of the present study was to isolate EPNs in sugar cane crops, and strain MC5-2014 was obtained, after which the *Serratia* bacterial strain was also isolated. This work provides the first geographic report of *Oscheius myriophila* in Mexico and the first global report of its entomopathogenic nature.

Materials and methods

Collection of soil samples

Soil samples were collected from 14 plots of sugar cane located in the state of Tepalcingo, Morelos in Mexico using the technique proposed by Stock et al. (1999), and a modified method described by Orozco et al. (2014). For soil sample collection, within a plot, samples were taken at five points, with three subsamples per site; for sampling, a grid of 3 m² was established around the sugar cane plant, and 1 kg of soil was collected at a depth of approximately 15 cm; the soil samples were placed in plastic bags. The sites were sampled in September 2014.

Soil analysis was performed to determine soil physicochemical characteristics such as texture, pH, electrical conductivity, and % organic matter (MO) in the bioremediation laboratory of CEIB/UAEM. EPNs were extracted with the insect baiting technique as described by Orozco et al. (2014). Each soil sample was baited by placing 12 fifth-instar *Galleria mellonella* Linnaeus (Lepidoptera: Pyralidae) wax worms in speech boxes containing sampled soil, and they were stored at room temperature in the dark for a week. Subsequently, the boxes were checked daily until the wax worms began to show signs of infection, at which time the insects were removed from the boxes, rinsed with water and stored in modified white traps (Orozco et al., 2014) until the EPNs emerged. After emergence, the EPNs were collected and stored in 50 mL Corning® cell culture flasks at 12 °C.

Nematode identification

Morphological and morphometric characterization

Morphological and morphometric identification was performed using an optical microscope equipped with a micrometric eyepiece. The nematodes were extracted through the dissection of wax worm larvae, and the IJs were collected from the white trap according to the method proposed by Nguyen and Smart (1996).

Molecular characterization of nematodes

For identification with molecular techniques, three hundred IJs were frozen in liquid nitrogen and macerated with a plastic pestle. Their DNA was purified with a Puregene® DNA Purification Kit following the manufacturer's instructions. An 850 bp fragment corresponding to the D2D3 region of the 23S rDNA gene was PCR amplified with the primers pair D2F (5'CCTTAGTAACGGCGAGTGAAA-3') and 536 (5'-CAGCTATCCTGAGGGAAAC-3') (Nguyen et al., 2006). The reaction was conducted in a final volume of 50 µl containing each primer at 0.5 µM, 0.16 mM dNTPs, 1.5 mM MgCl₂, 2 units of Taq DNA polymerase (Thermo Fisher Scientific Inc.) and 1X reaction buffer. The PCR product was verified by electrophoresis in a 1% agarose gel, and the bands were excised and purified using a QIAquick® Gel Extraction kit (QIAGEN, Germany). The PCR conditions were 94 °C for 2 min initial denaturation; 37 cycles of 30 s of denaturation at 94 °C, 45 s of hybridization at 48 °C and extension at 72 °C for 90 s; with a final extension of 72 °C for 5 min. The PCR product was sent to the Biotechnology Institute of the National Autonomous University of Mexico for sequencing. The D2D3 region was analyzed using the BLAST program of the GenBank database of the National Institutes of Health.

Isolation and identification of the bacterial symbiont

For the extraction of symbiotic bacteria from MC5-2014, two methods were used. In the first method, 5000 freshly emerged IJs were obtained from the white trap, and were placed in Eppendorf tubes. The tubes were centrifuged at 1,000 revolutions per minute (rpm) for 7 min, and the supernatant was discarded. Thereafter, washes were conducted with 5% sodium hypochlorite, and three washes were performed with sterile distilled water (Lee and Stock, 2010). The IJ nematodes were macerated in a mortar and pestle for a period of

30min and plated in solid Luria-Bertani (LB) and HCT (Bacto Tryptone (Difco) 5; Casamino acids (Difco) 2; pH adjusted to 7.5) media. After sterilization, culture medium consisting of KH_2PO_4 , 3.4g/L; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.012g/L; $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$, 0.003g/L; $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 0.0028g/L; $\text{Fe}(\text{SO}_4) \cdot 3.7\text{H}_2\text{O}$, 0.02g/L; $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 0.147g/L; and glucose, 3g/L was used. For the second method of extraction from hemolymph, 10 larvae of *G. mellonella* at the 5th instar were infected with 10 IJs from a stock solution of MC5-2014 nematodes. After 96hr, larvae with characteristics corresponding to death caused by EPNs were selected, and the outer parts of the larvae were disinfected with 70% alcohol for 10min, followed by drying for 2min inside a laminar flow hood (Woodring and Kaya, 1988).

Subsequently, a 0,3ml insulin syringe with a 31 × 6mm needle were used to extract the hemolymph by inserting the needle between the 6th and 7th inter-segments of the larvae. An aliquot of the hemolymph was plated on solid LB and HCT culture media, and the plates were incubated for 24hr at 27 °C. All colonies were sampled, and the cross-streak technique was used to obtain single colonies. The bacteria were identified by biochemical methods using Gram staining, motility tests, and assays of protease activity and lecithinase and antibiotic production as suggested by Boemare et al. (1997) and Akhurst (1980). For molecular identification of the isolated bacteria, the cells were incubated in LB liquid medium for 12h at 30 °C, and DNA extraction was then carried out with the Easy-DNA™ kit following the instructions recommended by the manufacturer. Then, amplification of a partial 16S rRNA region (600bp) was performed by PCR using the primer 63f (5'-CAG GCC TAA CAC ATG CAA GTC-3') (Nübel et al., 1996). The PCR conditions were 95 °C for 3min initial denaturation; 37 cycles of 35s of denaturation at 95 °C, 42s of hybridization at 59 °C and extension at 72 °C for 1min 30s; with a final extension of 72 °C for 5min. The PCR products were identified at the Biotechnology Institute of the National Autonomous University of Mexico with the primer 63f. The sequences were analyzed using the BLAST program of the GenBank database, National Institutes of Health.

Pathogenicity of the symbiont bacterium in *Galleria mellonella*

Pathogenicity bioassays of bacterial strain MC5-2014 were carried out in 10 larvae of the 5th stage of *Galleria mellonella*, and the treatments were carried out in triplicate. The bacterial isolates were cultured in solid LB medium for 18hr at 27 ± 2 °C. The surfaces of the larvae were previously disinfected with 70% ethanol and inoculated by injection using

10µl of the bacterial suspension at a concentration of 1×10^5 cells. The bacterial concentration was obtained via volumetric quantification using a Neubauer chamber.

Strain Bar 86 of *S. marcescens* was used as a positive control, while the *Escherichia coli* ATCC strain and distilled water were used as negative controls. The injected larvae were placed in 60 × 15mm Petri dishes with a meridian diet (bee honey 97.5ml, glycerol 120ml, wheat bran (sterile) 37.5ml, rice cereal 300g, yeast 75g). The pathogenicity and the changes in the coloration of the cuticle were evaluated at 24 and 48hr. The results were subjected to analysis of variance (ANOVA) and the Tukey test at the 0.05 probability. The SAS statistical package was applied.

Virulence of the bacteria

The virulence bioassays of the bacteria isolated from the MC5-2014 strain were performed on 30 larvae of *G. mellonella* in the 5th instar, and the treatment was carried out in triplicate. Bacterial strains were cultured in solid LB medium for 18hr at 28 ± 2 °C prior to quantification with the help of a Neubauer camera; volumetric calculations were performed and verified, and four concentrations were used to determine virulence for *G. mellonella* (10^5 , 10^3 , 50, and 5 cells). The surfaces of the larvae were previously disinfected with 70% ethanol. The same positive and negative controls mentioned above were used. The larvae were placed in 60 × 15mm Petri dishes with sterile filter paper with a meridian diet for *G. mellonella*. The percentage of mortality and the change in the coloration of the cuticle were evaluated at 24 and 48hr. Virulence (LC_{50}) was determined by PROBIT analysis using the POLO PLUS version 1.0 program (LeOra Software LLC).

Virulence of EPNs

Virulence was determined using 6, 10, 14, 18, or 20 infected juveniles (IJs) plus the control, which contained only sterile distilled water. Virulence was determined by obtaining the average lethal concentration (LC_{50}) via PROBIT analysis using the POLO PLUS version 1.0 program. (LeOra Software LLC).

Results and discussion

Isolation of EPN in sugar cane soils

Among the 14 soil samples taken from different locations in the state of Morelos, Mexico, positive

results were found in the municipality of Teplacingo, which has an altitude of 1100mm, a warm subhumid climate, an average annual temperature of 24 °C and a minimum temperature of 14 °C, from a site with sandy loam soil with a pH of 6.4 and an organic material (OM) percentage of 2.64%. We obtained the MC5-2014 isolate from sugar cane crops, and there have been reports of positive isolation in sugar cane cultures of *O. maqbooli* and *O. sacchari*. in Pakistan (Tabassum and Shahina, 2002; Tabassum et al., 2016). Our MC5-2014 isolate provides an additional isolate of nematodes belonging to the genus *Oscheius* from sugar cane, but it is the first member of this genus to be identified in Mexico.

Nematode identification through morphological and morphometric characterization

Description

Female: Small-sized nematodes. Slender body 760 to 1160µm long slightly curved ventrally after fixation. Cuticles are finely annulated with fine longitudinal striations. The head is continuous with the body contour, with six separate well-developed lips. Pore-like amphids on the lateral lips. Cylindrical pharyngeal body with cheilorhabdions that are not notable. The median bulb was absent. The metacorpus is swollen and pharyngeal. The isthmus is distinct, cylindrical, and slightly narrow. Nerve rings usually surround the mid portion of the isthmus. Excretory pores are conspicuously located at the level of the basal bulb. The excretion duct is circularized and curved anteriorly, then posteriorly. Vulva near the mid body and lips protruding.

Male: Similar to the female in general morphology except for a smaller size of 720 to 910µm in length. The body is straight after fixation. Spicules are paired, separate, symmetrical, and slightly curved ventrally with hooked tips. Head of the spicules with a rounded anterior end, lamina expanded in their proximal part and velum prominent. Bursa leptoderan and eight genital papillae (rays) of different lengths are with a 1 + 1 + 3 + 3 arrangement. There were 3 precloacal papillae and 6 postcloacal papillae. Papillae 8 and 7 form 6 groups, and pairs 5, 4, and 3 form the second postcloacal group. Papillae two and one are separated near the base or back of the spicules. The spicules are paired and separate. They are slightly curved and have a triangular head and a rounded tip. The lengths of the spicules are between 30 and 80µm. The gubernaculum is ventrally flattened, and it follows the contour of the spicules. The proximal tip

is curved upward. The length of the gubernaculum is 20 to 30µm.

Habitat and locality: soil around the roots of sugar cane (*Saccharum officinarum*) in Morelos, Mexico.

The male, female and IJ specimens were deposited in the Nacional Collection of Helminthes of the Biology Institute of the National Autonomous University of Mexico (CNHE/UNAM) with registration number 11085.

Diagnosis: *O. myriophila* belongs to the insectivora group and exhibits a cylindrical pharynx, a body with a leptoderan bursa and eight genital papillae of different lengths arranged in a 1 + 1 + 3 + 3 formation.

O. sacchari was isolated from sugar cane soil and belongs to the insectivora group. It presents similarity in the morphology of the anterior part that is a characteristic of the genus. *O. sacchari* is similar to *O. myriophila* in the body length of males (average: 843 vs 862µm), but exhibits a different body width (average 61.6 vs 28µm) and longer spicules (average: 41 vs 50µm). Posterior part arrangement with the papillae in a 1 + 1 + 1 + 3 + 3. In females, shorter body lengths (average: 1607 vs 964.5µm) and narrower body widths (average: 37 vs 88µm) were observed.

O. necromenus is similar to *O. myriophila* in the body length of males (average: 843 vs 790µm), but those of females differ (average: 964.6 vs 1179µm) as does their body width (average 28-46µm), and they exhibit longer spicules and a longer gubernaculum (average: 41 vs 40; 22 vs 15µm, respectively). Detailed species comparisons for the genus of *Oscheius* are given in Table 1.

Based on the morphometric characteristics of males and females that are commonly used for the identification of nematodes, 10 females and 10 males were observed under a microscope (Nikon eclipse e200), and morphometric measurements were performed (Table 1). The maximum and minimum ranges of the body length were obtained: L, body width: W, spicules, distance from the part before the vulva, and length of eggs. The average values were calculated.

Molecular identification

For molecular identification, an 832 bp fragment of the D2D3 region of the 28s ribosomal gene was amplified by PCR. The sequence of the MC5 2014 isolate was deposited in GenBank under accession number MK418537, showing 99% identity with DF5020 of *O. myriophila* isolated in the USA.

In Turkey (Zeynep et al., 2017), a nematode associated with *G. gryllotalpa* was isolated, and similar to our work, these authors used IJs for DNA extraction.

Table 1. Comparison of morphometrics with the MC5-2014 isolate.

Characteristic	<i>Rhabditis myriophila</i> Poinar, (1986)	<i>Oscheius myriophila</i> Tabassum et al. (2016)	<i>Oscheius myriophila</i> MC5-2014	<i>Oscheius sacchari</i> Tabassum et al. (2016)	<i>Oscheius rugaroensis</i> Zhang et al. (2012)	<i>Oscheius necromenus</i> Sudhaus and Schulte (1989)
Locality	California	Turkey	Mexico	Pakistan	China	Iran
Type host/habitat	<i>Oxidis gracilis</i>		Soil,	Soil,	Soil	
1. Female						
L	- (830-1500)	- (972-1530)	964.5 (760-1160)	1607 (1362-2015)	1042 (920-1179)	1179
W	-	- (52-100)	37 (30-60)	88 (72-125)	49.5 (39.8-58.2)	69 (54-90)
V	-	-	554 (410-760)	-	-	-
Eggs	-	-	52 (50-56)	-	-	-
1. Male						
L	- (830-1470)	- (841-1175)	843 (720-910)	862 (760-1390)	1396 (1195-1692)	790 (671-950)
W	- (380-800)	- (52-72)	28 (20-30)	61.6 (53-75)	62.4 (46.3-66.2)	46 (36-50)
Spicules	-	(27-39)	41 (30-40)	50 (47-55)	49.2 (35.2-60.9)	40 (34-44)
G	-	-	22	20.3	19.7	15
-	-	-	(20-30)	(18-22)	(19.9-26.5)	(12-23)

Note: L: total body length, W: maximum body width, V: vulva, and G: gubernaculum.

They obtained 99% identity with *O. myriophila* Gg1 and an isolate from the USA (AY602176).

Notably, Torrini et al. (2015) reported a new species of EPN belonging to the genus *Oscheius* using the D2D3 region of 28s rDNA, as in our investigation.

Bacterial symbiont isolation and identification

For the macerated extraction and the hemolymph extraction methods, the bacterial colonies were isolated at 48 and 72h, grown on solid LB and HCT agar plates, and initially assigned as MC5-R based on red colonies. The biochemical tests showed that the bacterial strain MC5-R was Gram negative and was positive for motility tests, protease activity and lecithin activity. From the genomic DNA extraction product, a fragment of 600bp was obtained. Once obtained, the PCR products were sent to the Institute of Biotechnology of UNAM (National Autonomous University of Mexico). The sequence of the isolated MC5-R was deposited in GenBank with the accession number MK463930. A BLAST search indicated 100% similarity with an MK463930 sequence and *S. marcescens* with accession number KM099142.1.

Pathogenicity of the bacterium

The percentage of mortality in *G. mellonella* larvae of the 5th instar resulting from treatment with a concentration of 1×10^5 was 100% for bacterial isolate MC5-R, 96.66% for *S. marcescens* Bar 86 and 0% for the *E. coli* ATCC and sterile distilled water negative controls.

Virulence of the bacterium

The percentage of mortality for *G. mellonella* larvae of the 5th instar resulting from treatment at a

concentration of five cells per larvae was 76.66% for the bacterial isolate MC5-R and 0% and 1.11% for the *E. coli* ATCC and sterile distilled water negative controls, respectively (Figure 1). The result obtained with the bacterial strain MC5-R was an $LD_{50} < 10$ cells injected in *G. mellonella* larvae at the 5th instar and the positive control *S. marcescens* Bar 86 showed an $LD_{50} < 10$ cells, while that for the negative controls *E. coli* ATCC and distilled water was 0%.

Virulence of the nematode MC5-2014 isolate

The mean lethal concentration was 4.732 IJ in *G. mellonella*. In EPN virulence evaluations in sugar cane borers, Grifaldo (Grifaldo et al., 2010) evaluated steinernematids in *D. saccharalis* in 2010 and obtained an LC_{50} of 3.7 IJ. The values obtained in this investigation were greater, and it should be taken into account that the behavior of the entomopathogens was different; this difference in virulence is influenced by different factors, such as the search strategy (Grewal et al., 1994) as well as their penetration and infectivity. In this study, *O. myriophila* was isolated, identified and evaluated, and solid evidence that it has potential to be used as an EPN is provided; it is worth mentioning that it is necessary to carry out an evaluation of the transmission of the bacteria to subsequent generations of nematodes (Dillman et al., 2012). The MC5-2014 isolate was virulent to *G. mellonella* larvae as quickly and efficiently as the isolated bacteria. An LC_{50} of 4.73 IJs/ml and an LD_{50} of < 10 cells were obtained for the bacterium. The existing reports mention that species of the family Rhabditidae are pathogenic against pests with the aid of bacterial strains of the genus *Serratia*, which is the same bacterial species as the nematode MC5-2014 (Abebe et al., 2010; Ye et al., 2010; Torres-Barragan et al., 2011; Zhang et al., 2012).




Figure 1: Mortality of *Galleria mellonella* larvae exposed to the MC5-R bacterium after 48 hours. 1: MC5-R, 2 *S. marcescens*, 3: *E. coli*, 4: water, means \pm SD (range).

Currently, species belonging to the genus *Oscheius* have been reported as EPNs, and the results obtained in our research highlight the potential of *O. myriophila* to be used in the future in biological control. However, to consider *O. myriophila* as an EPN, it is necessary to perform more evaluations on the type of association that exists with the bacteria isolated from this nematode.

Acknowledgments

The authors acknowledge Dr Veronica Obregón Barbosa and M in C. Laura Patricia Lina García for providing the biological material. The State Committee of Plant Health of the State of Morelos A. C provides information on and the locations of crops infected with *D. magnificella*. Conacyt/México provided a scholarship for doctoral studies to I.R.C.O.

References

- Abebe, E., Jumba, M., Bonner, K., Gray, V., Morris, K. and Thomas, W. K. 2010. An entomopathogenic *Caenorhabditis briggsae*. *Journal Experimental Biology* 213:3223–9.
- Akhurst, R. J. 1980. Morphological and functional dimorphisms in *Xenorhabdus* spp., bacteria symbiotically associated with the insect pathogenic nematodes *Neoplectana* and *Heterorhabditis*. *Journal of General Microbiology* 121:303–9.
- Boemare, N., Givaudan, A., Brehelin, M. and Laumond, C. 1997. Symbiosis and pathogenicity of nematode bacterium complexes. *Symbiosis* 22:2145.
- Dillman, A. R., Chaston, J. M., Adams, B. J., Ciche, T. A., Goodrich-Blair, H., Stock, S. P. and Sternberg, P. W. 2012. An entomopathogenic nematode by any other name. *PLOS Pathogens* 8:e1002527.
- Dujardin, F. 1845. *Histoire naturelle des helminthes ou vers intestinaux* Paris, 654 pp.
- Grewal, P. S., Selvan, S. and Gaugler, R. 1994. Thermal adaptation of entomopathogenic nematodes: Niche breadth for infection, establishment, and reproduction. *Journal Thermal Biol* 19:245–53.
- Grifaldo, A. F., Alatorre, R. R., Segura, L. O. and Hernandez, R. F. 2010. *Steinernema ralatorei* n. sp. isolated from sugarcane areas at Veracruz, Mexico. *Source Southwestern Entomologist* 42:171–90.
- Griffin, C. T., Downes, M. J. and Block, W. 1990. Test of Antarctic soils for insect parasitic nematodes. *Antarctic Science* 2:221–2.
- Hominik, W. M., Reid, A. P., Bohan, D. A. and Briscoe, B. R. 1996. Entomopathogenic nematodes: biodiversity, geographical distribution and the convention on biological diversity. *Biocontrol Science and Technology* 6:317–31.
- Lee, M. M. and Stock, S. P. 2010. A multigene approach for assessing evolutionary relationships of *Xenorhabdus* spp. (γ -Proteobacteria), the bacterial symbionts of entomopathogenic *Steinernema* nematodes. *Journal of Invertebrate Pathology* 104:67–74.
- Lezama, G. R., Hamm, J. J., Molina, O. J., Lopez, E. M., Pescador, R. A., Gonzalez, R. M. and Styler, E. L. 2001. Occurrence of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in the Mexican states of Michoacan, Colima, Jalisco and Tamaulipas. *Florida Entomologist* 84:23–30.
- Nguyen, K. B. and Hunt, D. J. 2007. “Entomopathogenic nematodes: systematics, phylogeny and bacterial symbionts”, In Hunt, D. J. and Perry, R. N. (Eds), Leiden-Boston: Brill.
- Nguyen, K. B. and Smart, G. C. 1996. Identification of entomopathogenic nematodes in the Steinernematidae and Heterorhabditidae (Nemata: Rhabditidae). *Nematology* 28:286–300.
- Nguyen, K. B., Gozel, U., Koppenhofer, H. S. and Adams, B. J. 2006. *Heterorhabditis floridensis* n. sp. (Rhabditida: Heterorhabditidae) from Florida. *Zootaxa* 1177:1–19.
- Nübel, U., Engelen, B., Felske, A., Snaidr, J., Wieshuber, A., Amann, R., Ludwig, W. and Backhaus, A. 1996. Sequence heterogeneities of genes encoding 16S rRNAs in *Paenibacillus polymyxa* detected by temperature gradient gel electrophoresis. *Journal of Bacteriology* 178:5636–43.
- Orozco, R. A., Lee, M. M. and Stock, S. P. 2014. Soil sampling and isolation of entomopathogenic nematodes (Steinernematidae, Heterorhabditidae). *Journal of Visualized Experiments* 11:89, doi: 10.3791/52083.
- Poinar, G. O. 1986. *Rhabditis myriophila* sp. (Rhabditidae:Rhabditida), associated with the millipede, *Oxidis gracilis* (Polydesmida: Diplopoda). *Proceedings of the Helminthological Society of Washington* 53:232–6.
- Stock, S. P., Pryor, B. M. and Kaya, H. K. 1999. Distribution of entomopathogenic nematodes (Steinernematidae and Heterorhabditidae) in natural habitats in California, USA. *Biodiversity and Conservation* 8:535–49.
- Stock, S. P., Rivera, O. P. and Flores, L. Y. 2009. *Heterorhabditis sonorensis* n. sp. (Nematoda: Heterorhabditidae), a natural pathogen of the seasonal cicada *Diceroprocta ornea* (Walker) (Homoptera: Cicadidae) in the Sonorandesert. *Journal of Invertebrate Pathology* 100:175–84.
- Sudhaus, W. 1976. Vergleichende untersuchungen zur phylogenie, Systematik, Ökologie, Biologie and Ethologie der Rhabditidae (Nematoda). *Zoologica* 43:1–229.
- Sudhaus, W. and Schulte, F. 1989. *Rhabditis (Rhabditis) necromena* sp. n. (Nematoda: Rhabditidae) from South Australian Diplopoda with notes on its siblings *R. myriophila* Poinar, 1986 and *R. caulleryi* Maupas, 1919. *Nematologica* 35:15–24.
- Tabassum, K. A. and Shahina, F. 2010. *Oscheius siddiqii* and *O. niazii*, two new entomopathogenic

nematode species from Pakistan, with observation on *O. shamimi*. International Journal of Nematology 20:75–84.

Tabassum, K. A. and Shahina, F. 2002. *Oscheius maqbooli* n. sp. and observation on three known Rhabditis species (Nematoda: Rhabditida) from sugarcane fields of Balochistan, Pakistan. Pakistan Journal of Nematology 20:1–21.

Tabassum, K. A., Shahina, F. N. and Erum, Y. I. 2016. Description of six new species of *Oscheius* Andrassy, 1976 (Nematoda: Rhabditida) from Pakistan with a key and diagnostic compendium to species of the genus. Pakistan Journal of Nematology 34:109–61.

Torres-Barragan, A., Suazo, A., Buhler, W. G. and Cardoza, Y. J. 2011. Studies on the entomopathogenicity and bacterial associates of the nematode *Oscheius carolinensis*. Journal Biological Control 59:123–9.

Torrini, G., Mazza, G., Carletti, B., Benvenuti, C., Roversi, P. F., Fanelli, E., Luca, F. D., Troccoli, A. and Tarasco, E. 2015. *Oscheius* Description of six new species of *Oscheius* Andrassy, 1976 (Nematoda: Rhabditida) from Pakistan 161 *O. onirici* sp. n. (Nematoda: Rhabditidae);

a new entomopathogenic nematode from an Italian cave. Journal Zootaxa 26:533–48.

Woodring, J. L. and Kaya, H. K. 1988. Steinernematid and Heterorhabditid nematodes: A handbook of biology and techniques. Southern Cooperative Series Bulletin 331, Fayetteville, AR: Arkansas Agricultural.

Ye, W., Torres-Barragan, A. and Cardoza, Y. J. 2010. *Oscheius carolinensis* n. sp. (Nematoda: Rhabditidae), a potential entomopathogenic nematode from vermicompost. Nematology 12:121–35.

Zeynep, E., İsmail, D. and Zihni, D. 2017. Isolation and characterization of a parasitic nematode, *Oscheius myriophila* (Nematoda: Rhabditida), associated with European mole cricket, *Gryllotalpa gryllotalpa* (Orthoptera: Gryllotalpidae). Journal Biology and Chemistry 45:197–2.

Zhang, K. Y., Liu, X. H., Tan, J., Wang, Y., Qiao, L., Yedid, G., Lai, R. and Gao, G. F. 2012. *Heterorhabditoides rugaroensis* n. sp. (Rhabditida:Rhabditidae), a novel highly pathogenic entomopathogenic nematode member of Rhabditidae. Journal of Nematology 44:348–60.