

เทคโนโลยีจำลองตัวละครในการสอนและการให้ความรู้แก่ผู้ป่วย: การทบทวนวรรณกรรมแบบบูรณาการอย่างเป็นระบบ Avatar-based Technology Interventions in Patient Education: A Systematic Integrative Review

นิพนธ์ปริทัศน์

Review Article

จันทนา ทองเพชร^{1*} และ ณัฐรภัฏดา เขาวนารักษ์²¹ วิทยาลัยพยาบาลพระจอมเกล้า จังหวัดเพชรบุรี อ. เมือง จ. เพชรบุรี 76000² สำนักงานสาธารณสุขจังหวัดเพชรบุรี อ.เมือง จ. เพชรบุรี 76000

* Corresponding author: jintana@pckpb.ac.th

วารสารไทยเภสัชศาสตร์และวิทยาการสุขภาพ 2563;15(2):144-153.

Jintana Tongpeth^{1*} and Natkritta Chaowararak²¹ Prachomkiao College of Nursing, Muang district, Phetchaburi, 76000 Thailand² Phetchaburi Provincial Public Health Office, Muang district, Phetchaburi, 76000 Thailand

* Corresponding author: jintana@pckpb.ac.th

Thai Pharmaceutical and Health Science Journal 2020;15(2):144-153.

บทคัดย่อ

วัตถุประสงค์: เพื่อทบทวนวรรณกรรมและหลักฐานงานวิจัยเกี่ยวกับการใช้ นวัตกรรมเทคโนโลยีจำลองตัวละครในการสื่อสาร ส่งต่อข้อมูลและความรู้ทาง สุขภาพให้กับผู้ป่วย **วิธีการศึกษา:** การทบทวนวรรณกรรมแบบบูรณาการอย่าง เป็นระบบนี้คัดเลือกเฉพาะงานวิจัยเชิงทดลองแบบมีกลุ่มควบคุม โดยสืบค้นจาก ฐานข้อมูลอิเล็กทรอนิกส์ ได้แก่ CINAHL, MEDLINE, PubMed, Scopus, Cochrane และ EMBASE รวมทั้งมีการประเมินคุณภาพของระเบียบวิธีวิจัยของ งานวิจัยที่คัดเลือก **ผลการศึกษา:** มีงานวิจัยจำนวน 6 เรื่องที่ตรงตามเกณฑ์การ คัดเข้า ผลการวิเคราะห์และสังเคราะห์พบว่าการใช้เทคโนโลยีจำลองตัวละครใน การเพิ่มความรู้ด้านสุขภาพที่เกี่ยวข้องกับการปรับเปลี่ยนพฤติกรรมสุขภาพของ ผู้ป่วย แต่พบว่ามีหลักฐานงานวิจัยที่ยังไม่ชัดเจนและเพียงพอที่จะประเมิน ประสิทธิภาพของรูปแบบเทคโนโลยีจำลองตัวละครต่อการรับรู้ความสามารถของ ตนเองของผู้ป่วย **สรุป:** การทบทวนวรรณกรรมนี้พบงานวิจัยเพื่อพิจารณาพัฒนา รูปแบบการให้ความรู้แก่ผู้ป่วยโดยใช้เทคโนโลยีจำลองตัวละคร และเทคโนโลยี จำลองตัวละครเป็นเครื่องมือให้ความรู้แก่ผู้ป่วยในบริบททางสุขภาพต่าง ๆ ได้ มากกว่าการให้ความรู้ในรูปแบบสื่อสิ่งพิมพ์ พบองค์ประกอบของการใช้รูปแบบ เทคโนโลยีการจำลองตัวละครที่สามารถประยุกต์ใช้พัฒนารูปแบบการให้ความรู้ ทางสุขภาพแก่ผู้ป่วยในการปฏิบัติการพยาบาลได้อย่างมีประสิทธิภาพต่อไป

คำสำคัญ: การทบทวนวรรณกรรมแบบบูรณาการอย่างเป็นระบบ, เทคโนโลยี จำลองตัวละคร, การปฏิบัติการพยาบาลในคลินิก, การให้ความรู้แก่ ผู้ป่วย

Editorial note

Manuscript received in original form on November 24, 2019;
revised February 15, 2020;
and accepted in final form on March 25, 2020

Abstract

Objective: To review the evidence for the avatar-based technology as an innovative and emerging intervention for delivery of health education in clinical practice. **Method:** This review followed a systematic review methodology. Randomised controlled trials in English language were searched in CINAHL, MEDLINE, PubMed, Scopus, Cochrane and EMBASE. Articles were assessed for quality of methodology. **Results:** Six randomised controlled trials met the eligibility criteria and were included in the review. Avatar-based technology interventions improved health knowledge, and health-related behaviour. There was insufficient evidence to suggest the effectiveness of Avatar-based interventions on, self-efficacy, and of the interventions among the participants were unclear. **Conclusion:** This review established a scientific basis for avatar-based technology as an effective intervention for the transfer of knowledge and skill development for patients. Avatar-based education tools have a clear advantage over paper-based education materials. This review provides nurses with the state of science in relation for the avatar-based technology interventions in patient education. A summary of intervention components are provided to assist nurses to develop and use the avatar-based interventions in clinical practice.

Keywords: systematic integrative review, avatar-based technology, nursing practice, patient education

Journal website: <http://ejournals.swu.ac.th/index.php/pharm/index>

Introduction

Self-care behaviour is the process of maintaining health through health-promoting practices in states of both health and illness.¹ Individuals are able to limit symptoms when they understand the disease management and self-management and learn to undertake simple interventions.² The ability of a patient to control their health and illness through an effective care plan is significantly influenced by cognitive behavioural, social or situation factors.^{3,4} A major problem of unhealthy behaviour is the resistance of patients to lifestyle advice, such as eating a healthy diet, doing more exercise and quitting smoking.⁵⁻⁷ Numerous studies have shown that a large

percentage of patients do not adhere to healthy lifestyle change, self-care behaviour or disease management.^{8,9} The process of behaviour change is complicated and requires more than simply the acquisition of health knowledge.¹⁰⁻¹² To improve the effectiveness and efficacy of care, patient education plays a significant role in the decision-making processes of patients, seeking of medical care and behaviour change to improve health-related quality of life.¹²⁻¹⁴

Patient education is an important component of many disease management and health promotion programs.^{1,8,15-18} Health care providers are under increasing pressure to provide

health information and resources to patients to encourage them to follow disease management programs and make informed decisions about their health behaviour.^{1,3,19} Patient education has often been delivered face-to-face and supported by paper-based flyers, brochures and sometimes videos or DVDs.²⁰ However, as information technology has rapidly advanced, computer-based information and resources for supporting patient teaching, decision-making and information retrieval have become increasingly available.²¹⁻²⁵ Several authors have claimed that these new types of systems have many advantages compared to the more traditional means used for patient education.²⁶⁻²⁹ The main advantage is that health information can be delivered in a more interactive and engaging way and has the ability to provide individually tailored information at a lower cost.^{6,30-32}

An avatar is a computer-generated character representing people and interactive behaviours.³¹ It is designed to carry out simulated 'human-like character' conversations with player or user, including appropriate use of mimicking human behaviours interaction such as talking and hand gestures.³³

Moreover, avatars are computational human-like behaviour technology designed to build and maintain long-term, social-emotional relationships with their users.³⁴⁻³⁶ They are particularly effective as relational agents, as many of the interaction strategies used to build and support interpersonal relationships involve non-verbal communication.^{19,34,37} The use of avatar-based in patient education is growing, creating a new simulated avatar characters for patients during bedside education session and discharge plan from the comfort of their own home. As health and patients education changes to meet the needs of patients or caregivers, options such as avatar-based technologies need to be considered.^{38,39} However, there is only limited research on the effectiveness and applicability of avatar-based technology interventions for patient education. This article used an integrative review approach that allowed the inclusion of randomised controlled trials (RCTs) to provide readers with a better understanding on the topic. It could allow readers a greater capacity to explore the relevance and applicability of avatar-based technologies as interventions for improving outcomes for patient education in clinical practice.

To date, the literature has demonstrated that the integration of avatar-based technology interventions to deliver health information has improved the health behaviour change process in a range of different health conditions. The aim of

this review was to explore the effectiveness of avatar-based technology interventions in patient education. Specifically, there were two objectives. The first objective was to report and summarise the evidence on avatar-based technology interventions in patient education. The second one was to report how avatar-based technology interventions have improved knowledge, changed behaviours and enhanced self-efficacy and health related quality of life (HRQoL), as well as to report the participating patients' acceptability of and satisfaction with the application. The findings could be applied for further development of the practical intervention.

Methods

Database and search terms

This systematic integrative review used a systematic approach, including the methodological quality assessment outlined by Whitemore and Knaf⁴⁰ and a checklist for writing an systematic integrative review recommended by Torraco.⁴¹ A systematic approach has been used to identify the included studies, extract and analyse the information, appraise the study quality, and present the findings in a usable format.^{40,42} The synthesis of the data was conducted through a narrative analysis.⁴³ Finally, the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flowchart was used to present the process and result of this review.⁴⁴

A systematic search of the literature was conducted for peer-reviewed articles written in Thai and English and published between 2009 and 2019. Informed by the review steps,⁴⁰ the literature was searched for published articles on avatar-based education interventions in clinical practice. A search of the CINAHL, MEDLINE, EMBASE, PubMed, PsycINFO, Web of Knowledge, Cochrane Library, ProQuest (health sub-set), Scopus and Web of Science databases was conducted using the following keywords: "avatar-based," "avatar-based education," "agent-based," "conversation agent," "intelligent agent," "computer agent," "conversational agent," "interactive computer agent," "embodies conversational agent," "patient education," "health education," "health knowledge," "self-care," "health related quality of life," "self-efficacy" and "consumer health information."

Criteria for study selection

In terms of types of studies, this review considered studies that evaluated the effectiveness of avatar-based technologies

in patient education. The avatar-based technology interventions could be either used alone or in combination with the “usual care.” For types of participants, studies with participants with any medical conditions that required patient education were selected, regardless of age, ethnicity, gender or geographical location (i.e., metropolitan, rural or remote area). For intervention types, the selected studies used avatar-based technologies that were delivered either by nurse, physician or allied health educator via a computer, tablet-based device or mobile phone in a hospital (i.e., inpatient or outpatient) or in the community. Finally, in terms of types of outcomes, studies with knowledge as the primary outcome that was measured by valid and reliable instruments were selected. We also selected studies that measured secondary outcomes including self-care or behaviour change, self-efficacy, HRQoL and acceptability of or satisfaction with avatar-based technology interventions.

Methodological quality assessment

Two independent reviewers extracted data from and critically appraised the six selected studies. The data items extracted included the country in which the study was conducted, participant characteristics, objectives, intervention and control conditions and the length of follow-up. The extracted data from the reviewed studies were displayed and compared in tables based on the outcome measures. Two independent reviewers assessed all the included full-text papers through the standardised critical appraisal instrument from Critical Appraisal Skills Programme (CASP).⁴⁵ The CASP Randomised Controlled Trials Checklist was used to check the methodological validity of the included studies.

Results

Search outcomes

The initial search identified a total of 2,165 articles, while a manual search of other sources (i.e., Google search and the grey literature) identified a further 58 articles. In total, 2,223 articles were identified through all methods of searching. After removing duplicates, 1,946 articles remained for screening and another 1,925 articles were excluded. Full-text manuscripts were obtained from the remaining 21 articles. Based on a full-text review, 15 articles were excluded. Six RCT studies were identified. After assessment of the studies against the inclusion criteria, these six articles were included

in the review (Table 1). The PRISMA flow diagram presents the overall process in Figure 1.




Figure 1 PRISMA flowchart showing the identification of studies.

Study characteristics

The six articles in this systematic integrative review were published between 2005 and 2017. All of the studies were RCTs that were conducted in the USA, with each being designed to evaluate the effectiveness of avatar-based technologies for the delivery of health information for patient education in the clinical practice area. Two studies were implemented in a hospital setting, while the other four were implemented in a community setting. The characteristics of the included studies are presented in Table 1.

Participant characteristics

This review included a total of 2,204 participants from the six selected studies. All the participants were adults ranging from 18 to 91 years of age. Five studies included both genders while one consisted of only women.³⁰ In total, there were 1,565 women and 648 men. The six studies in this review presented research on health education relating to general lifestyle health behaviours⁴⁶, self-management, including fruit and vegetable consumption and physical activity³³, breakfast eating habits, alcohol consumption and smoking⁴⁶, overactive bladder (OAB) education³⁰, depressive symptoms⁴⁷, and explaining medical documents to patients.⁴⁸ The outcome measures of all the included studies were defined differently through differing measurement methods and timeframes. The

outcomes were divided into primary and secondary outcomes (Table 2). The primary outcome of this review was that of the acquisition of knowledge. The secondary outcomes included improved self-care or behaviour change, increased HRQoL scores, improved self-efficacy and satisfaction with and acceptability of the intervention.

Table 1 Summary of included studies.

Study	Participants	Objectives	Intervention/control condition	Length of follow-up
Bickmore et al. ⁴⁸ USA	Inpatients n = 29 adults Age: 28-91 years 19 females 10 males	Explaining consent form document	Three-armed RCT I: Document explanation by avatar I: Self-study C: Usual care (human explanation)	Until signed
An et al. ⁴⁶ USA	Previous 30 days smokers n = 1,698 Age: 18-30 years 1,230 females 468 males	Health behaviour change: eating breakfast, exercise, alcohol consumption and cigarette smoking	3 armed RCT I: Online video-based with peer avatar coaching I: Personally tailored plus online video-based peer avatar coaching C: Usual care (personally tailored)	30 days and 12 weeks
Bickmore et al. ³³ USA	Sedentary older adults n = 263 Mean age: 71.3±5.4 years 161 females 102 males	Daily walking step count Walking goal setting	Two-armed RCT I: Avatar exercise coach with take-home tablet computer and pedometer record C: Usual care with pedometer record	30 days and 12 weeks
Andrade et al. ³⁰ USA	Women with at least three months of OAB symptoms n = 41 Age: 55-75 years	OAB self-management behaviours	Two-armed RCT I: Avatar-based C: Voice only online program	1, 6, and 12 weeks
Bickmore et al. ⁴⁹ USA	Walk unaided n = 122 Age: 18 or older 48 males 74 females	Health behaviour change: physical activity, step walk with pedometer record, fruit and vegetable consumption	Four-armed RCT I: Online physical activity with avatar I: Online fruit and vegetable consumption with avatar I: Online physical activity plus fruit and vegetable consumption with avatar C: Usual care	2 months
Pinto et al. ⁴⁷ USA	Young adults with two weeks depressive symptoms n = 60 Age: 18-25 years 40 females 20 males	Reduce depressive symptoms among young adults	Two-armed RCT I: Avatar-based C: Usual care	12 weeks

Note: I = intervention group, C = control group; OAB = overactive bladder; RCT = randomised controlled trial.

Study outcomes

1) Primary outcomes: Knowledge

The acquisition of knowledge was explicitly referred to in two of the six selected studies.^{30,48} Both studies reported that avatar-based technology interventions were effective in delivering information and that the participants were generally able to use them without difficulty.^{30,48} Each of the two studies indicated an improvement in health knowledge scores when the information was communicated through an avatar compared to face-to-face communication and self-study instruction. Bickmore et al found that participants who received information through a conversational avatar to explain health documents and a consent form were just as likely to improve their knowledge scores as those who received explanations from health care providers or through self-study.⁴⁸ In addition, the study reported that avatar-based delivery was useful in identifying those patients who needed additional assistance on health information.⁴⁸ Andrade et al demonstrated that at 12-

week follow-up, the intervention resulted in an increase in OAB symptom knowledge scores when compared with usual care (P -value < 0.001).³⁰

Table 2 Summary of primary and secondary outcomes.

Study	Outcomes	Intervention	Control / Usual care	Follow-up	P-value*
1) Primary outcome: Knowledge					
Bickmore et al ⁴⁸	Medical document knowledge scores	42.20 (20.33)	25.91 (11.36)	1 day	0.006*
Andrade et al ³⁰	OAB symptoms knowledge scores (mean average)	5.77	2.61	12 wks	< 0.001*
2) Secondary outcomes					
2.1 Behaviour change					
Bickmore et al ³³	Daily step count (adjusted mean)	4,041 steps/day 3,861 steps/day	3,499 steps/day 3,383 steps/day	2 mos 12 mos	0.01* 0.09
An et al ⁴⁶	Smoking abstinence rate Number of days of change in drinking (mean change)	31% -1.87	11% -1.00	12 wks 12 wks	< 0.0001* < 0.0001*
	Number of days of eating breakfast (mean change) Number of days of exercise (mean change)	7.58 5.08	2.20 1.74	12 wks 12 wks	< 0.0001* 0.0058*
Bickmore et al ³³	Daily step count Fruit and vegetable consumption (mean change)	7,398 steps/day 3.4	8,003 steps/day 0.6	2 mos 2 mos	0.367 0.0005*
2.2 HRQoL					
Andrade et al ³⁰	OAB HRQoL scores (mean average)	90.9	79.52	12 wks	0.002*
2.3 Self-efficacy					
Andrade et al ³⁰	Self-efficacy for urinary incontinence scores (mean average)	8.27	7.41	12 wks	0.005*
2.4 Satisfaction/acceptability					
Bickmore et al ³³	Satisfaction (score range 1-7, mean average)	6.56	N/A	-	-
Pinto et al ⁴⁷	Acceptability (score range 1-5, mean average)	4.45	N/A	-	-

Note: *Statistical significance when P -value \leq .05; F/U = follow-up, N/A = not applicable; OAB = overactive bladder; HRQoL = health-related quality of life; mos = months; wks = weeks.

2) Secondary outcomes

2.1 Behaviour change

Three of the six studies reported on the effectiveness of an avatar-based technology intervention for changing behaviour.^{33,46} One study demonstrated the effectiveness of an avatar-based technology intervention for the daily step count of older adults³³, while another study detailed an intervention for daily step count and fruit and vegetable consumption.³³ One study reported multiple behaviour changes in young adult smokers, consisting of number of days of exercise, smoking, alcohol consumption and breakfast consumption.⁴⁶ Two studies measured daily step count^{33,49}, but only one study demonstrated a significant difference in the daily step count of older adults with an avatar coach (two-month follow-up, P -value = 0.01).³³ In this study, the authors compared the efficacy of an avatar-based exercise coach interacting on a daily basis with an intervention group in the

usual care using a pedometer record to increase walking among sedentary older adults.³³

These studies demonstrated that those coached with an avatar walked significantly more steps on a daily basis than control-group participants at two-month follow-up (P -value = 0.01). The qualitative data from the interviews with the same participants also indicated high levels of satisfaction with the intervention in 90% of the participants. An et al conducted a three-group RCT measuring the delivery of health behaviour information to young adult smokers.⁴⁶ The participants were assigned to the following groups: 1) online health messages, 2) online health messages plus avatar peer support, and 3) usual care. The results showed that the second group had increased abstinence rates from smoking (P -value < 0.001), increased number of days of breakfast consumption (P -value < 0.001) and days of exercise (P -value = 0.0058) and reduced number of days of alcohol consumption (P -value < 0.001) at 12-week follow-up.⁴⁶

2.2 Health-Related Quality of Life

Only one study assessed the effectiveness of an avatar-based technology intervention on HRQoL compared to a “voice only” online program.³⁰ In this study, it was found that avatar-based technologies used with women with OABs resulted in increased HRQoL scores compared to a control group at 12-week follow-up (P -value = 0.002).³⁰

2.3 Self-Efficacy

One of the six studies examined the effectiveness of an avatar-based technology intervention on self-efficacy.³⁰ The results showed that the avatar-based intervention was statistically significant as measured by self-efficacy for urinary incontinence in OAB patients.³⁰

2.4 Acceptability and Satisfaction

Two studies reported participants' satisfaction³³ with and acceptability⁴⁷ of an avatar-based technology. One study reported a higher mean average for acceptability in the intervention group in younger people with depressive symptoms (mean average = 4.45 out of 5).⁴⁷ Bickmore et al showed that an avatar-based exercise coach contributed to high levels of satisfaction for 90% of older sedentary adult participants (mean average = 6.56 out of 7).³³

Discussions and Conclusion

This integrative review aimed to synthesise existing evidence on the effectiveness of avatar-based technology interventions for patient education. It has demonstrated that these types of interventions can improve knowledge acquisition in patient education in clinical practice.^{30,48} Avatar-based technology interventions have also been shown to be effective for behaviour change.^{33,46} However, there was insufficient evidence to recommend these interventions for improving HRQoL, self-efficacy³⁰ and acceptability of or satisfaction with the intervention.

In terms of **study primary outcome** which was **knowledge**, the available research on the use of avatar-based technology interventions demonstrates that positive change in knowledge may be beneficial for managing disease and changing behaviours. Two studies revealed that the intervention yielded significant results in terms of improving knowledge.^{30,48} These studies reported that interactive avatar-based technology interventions could be potential tools to educate patients about medical documents, the consent process⁴⁸ and overactive bladder conditions.³⁰ This approach could increase knowledge in patients with specific health conditions because avatar-based technology interventions are able to improve health literacy in individuals using visual and auditory signals. Thus, avatar-based technology interventions may promote health knowledge and have a positive effect on patient outcomes.

Two systematic reviews reported that interactive media information technologies could be a potential tool for educating patients on their health conditions.^{50,51} Interactive media technologies⁵⁰ and serious game interactive design⁵¹ are able to account for health literacy through the use of visual and auditory signals that provide messages about various diseases. Thus, these technologies may have a positive effect on health outcomes. A number of authors have evaluated the effectiveness of interactive avatar-based interventions on health knowledge.⁵²⁻⁵⁴ Three other studies, including two pilot studies and one pre-test post-test design evaluated avatar-based technology interventions for improving heart failure knowledge⁵³, insulin treatment knowledge⁵⁴ and ileostomy care knowledge.⁵² Clark et al developed an avatar-based educational resource that was delivered by tablet computers and designed as an adjunct to traditional heart failure education.⁵³ The educational resource was provided to five

Aboriginal participants, resulting in a 20.8% increase in heart failure knowledge.⁵³ Another pilot study demonstrated improved insulin treatment knowledge via an online interactive video game education tool on type 1 diabetes mellitus patients at six-month follow-up.⁵⁴ One pre-test post-test study⁵² explored the use of an interactive avatar-based intervention to educate new stoma patients about ileostomy care. After using the intervention, the knowledge scores of participants showed significant improvement when compared with their baseline knowledge scores.⁵² In addition, interactive health teaching tools have been shown to be able to adapt content based on personal interests and specific diseases. One study reported, in accordance with one systematic review and meta-analysis, improved cancer care knowledge through the use of interactive technologies and videotapes for cancer patients.⁵⁵

The above evidence endorsed the findings of other studies that patients often forget the information that health care providers and professionals tell them during education sessions, bedside teaching or consultations. Some patients have difficulty understanding health information in booklets and brochures.⁵⁶ Interactive media technology interventions have been shown to have a variable effect in enhancing patient involvement in decision-making processes on self-care, self-management and disease management.^{57,58} Avatar-based technologies can provide a flexible learning environment in which patients can learn about their health condition in a personalised health setting through decision-making tools that provide tailored health information.^{59,60} Additionally, a systematic review by Friedman et al reported that the use of multiple teaching techniques with specific information for patients should be provided rather than general health information.⁶¹

Overall, this present systematic review has revealed that positive and effective teaching strategies and delivery methods for patient education would include integrating technologies via visual messages to enhance learning and understanding in patients with low health literacy to increase their knowledge.⁶¹ This review of RCTs supports the findings of previous research into the potential effectiveness of avatar-based technology interventions for patient education.

Regarding **behaviour change** and **self-care behavior**, which were two of the **secondary outcomes**, avatar-based technology interventions for patient education have also been shown to improve health behaviour change or self-care behaviour in patients with multiple health risks.^{34,62} These

interventions enhance activities that change behaviours such as daily step counting⁴⁹ and activities that lower alcohol and tobacco consumption.⁴⁶ As the results of this review, technological methods for maintaining self-care behaviour need to be included in patient education.⁶² A study by Lo et al demonstrated that their chosen multimedia education program improved self-care behaviour in post-operative stoma patients.⁶³ Another study reported that the use of interactive technology interventions via serious games improved the self-care behaviour of insulin administration in patients with type 1 diabetes mellitus.⁵⁴ The reason for these findings could well be that the most effective way to change behaviour is through delivery of health information by empathic avatar characters via multimedia or interactive technologies to increase user engagement, persuasion and motivation similar to coaching by nurses or health care professional educators.^{60,64} Health coaching is a behaviour change intervention that helps individuals to achieve individual health goals to change lifestyle behaviours, with the goal of decreasing the risk of health conditions and improving self-management and HRQoL.^{65,66} In addition, interactive technologies have the capacity to facilitate involvement and learning in an active way because the information can be matched to the patients' preferences, needs, diseases and health risk behaviours.^{67,68}

In relation to self-care behaviour, one pre-test post-test study demonstrated an avatar-based nursing intervention via a tablet-based educational resource about heart failure.⁵³ The results showed an increase in self-care behaviour in three domains, specifically, a 95% increase in self-care confidence, 26.1% increase in self-maintenance and 1.9% increase in self-management.⁵³ Another single group study demonstrated improved diabetes self-care behaviour via a virtual world with an avatar on a website about diabetes self-care education.⁶⁹ The results showed improved diabetes self-care behaviour at six-month follow-up.⁶⁹ These findings could be due to the participants having long-term chronic conditions or health risks that may have contributed to their lack of knowledge and understanding of the importance of health risk factors and the negative effects of not changing their behaviours.⁷⁰ This review has revealed that avatar-based technology interventions have been used to track daily health behaviours and make positive lifestyle changes such as quitting smoking, increasing exercise and decreasing alcohol consumption.

In terms of self-efficacy, one of the secondary outcomes, one study in this review showed that the use of avatar-based

technology interventions in patient education significantly improved patients' self-efficacy.³⁰ Other studies about integrating interactive technologies for patient education revealed significant improvements in self-efficacy.^{54,71,72} One pilot study reported that using serious game design in type 1 diabetes mellitus patients significantly improved insulin therapy self-efficacy scores at six-month follow-up.⁵⁴ A pre-test post-test intervention by Johnson et al demonstrated the effectiveness of virtual technologies in various diabetes management situations including selecting dishes from the menu in restaurants and making decisions in pharmacies, fitness centres, bookshops and convenience stores significantly improved self-efficacy in adults with type 2 diabetes mellitus.⁷¹ Another RCT demonstrated improved self-efficacy in a group of young adults (aged 13 - 29 years old) with cancer via a three-dimensional interactive video game at three-month follow-up.⁷² The reason for these positive findings could be that avatar-based technology interventions provide significant opportunities in health care settings through providing decision-making support rather than substituting for health care professionals.³⁰

For the last two secondary outcomes, **satisfaction** and **acceptability**, two studies reported on participants' satisfaction³³ with and acceptability⁴⁷ of an avatar-based technology. One study reported a higher mean average of satisfaction for the avatar-based technology intervention group,³³ while another study reported higher levels of acceptability for avatar-based technology interventions in the intervention group.⁴⁷ One pre-test post-test study showed an 83% level of satisfaction when using a heart failure avatar peer mentor through a tablet- and computer-based educational resource.⁵³ Another study about technology interventions showed high levels of satisfaction when using an online video game about insulin therapy in younger patients with type 1 diabetes mellitus.⁵⁴

In relation to the acceptability of these technologies, one pilot study demonstrated that 76% of participants accepted using an avatar mentor via a virtual world for diabetes education.⁷³ Therefore, interactive technologies for patient education were well obtained and patients were satisfied with the health information presented to assist their decision-making processes.⁷⁴ Through the use of interactive technologies, health care providers are able to tailor health information to patients' needs, and patients can control the

flow of information and select what they want to view and when to log out of the system.⁷⁵

In conclusion, this review is the first step in investigating the effectiveness of using avatar-based technologies. The results of this review suggest that avatar-based technologies can extend personal health information communication by nurses or health care professionals. Avatar-based technology interventions have been shown to increase knowledge in adults, elderly individuals and women with overactive bladder and have determined high levels of acceptability and satisfaction. The available research on the use of the avatar-based technologies shows that it can help patients and other individuals to achieve a greater understanding and recognition of disease management, healthy lifestyles and decision-making processes for changing behaviours to increase HRQoL. To capitalise on the advantages of these new patient educational learning technologies approach, health care professionals need to broaden their understandings of how people most appropriately learn through the avatar-based technologies and determine the effect of this knowledge on outcomes in health care context over time. The findings of this review may have implications for both nursing practice and patient education in relation to the delivery of health information interventions. This can only be accomplished through further research.

In terms of practice implications, avatar-based technology interventions have been shown to be an effective method of delivery for health information and an alternative tool for improving health knowledge, changes in behaviour and patient outcomes. Health care providers should actively select and adapt avatar-based technology methods to extend the reach of health behaviour and self-management principles to the clinical environment, community and home to provide individualised care and just-in-time information.^{30,33,46} In this review, the avatar-based educational technologies have been shown to be effective among people with different conditions, with most outcomes being superior to the use of traditional methods. It remains a challenge to find the most suitable avatar-based technology design for particular patient groups. Although all avatar-based technologies were found to be effective, their design and content varied due to the intrinsic differences among different conditions and characteristics of the patient population. This may need to be considered when implementing or testing an avatar-based intervention in a specific population.^{30,46,52}

The increased use of avatar-based technology interventions indicates that people living with health risks are interested in using methods to improve their knowledge, self-care, decision-making processes and clinical outcomes, in addition to changing their behaviours.^{30,50} Although it has been found that the avatar-based technologies in patient education are an effective and powerful way to provide information and teach skills to patients and, it is not clear whether it is as beneficial when the health information needs to be regularly updated or when the skills gained require ongoing rectification.³³ Finally, if knowledge acquisition depreciates over time, the overall usefulness of the avatar-based technologies for patient education would be greatly reduced. This adds support to the need for customised health information that is adaptable enough to be modified to the dynamic nature of patients' ongoing health information.^{30,33}

References

- Mohammadpour A, Rahmati Sharghi N, Khosravan S, Alami A, Akhond M. The effect of a supportive educational intervention developed based on the Orem's self-care theory on the self-care ability of patients with myocardial infarction: a randomised controlled trial. *J Clin Nurs* 2015; 24(2):1686-1692.
- Barnason S, Zimmerman L, Young L. An integrative review of interventions promoting self-care of patients with heart failure. *J Clin Nurs* 2012;21(3-4):448-475.
- Glanz K, Rimer BK, Viswanath K. Health behavior and health education: theory, research, and practice. San Francisco, CA. John Wiley & Sons, 2008.
- Bagheri-Saweh MI, Lotfi A, Salawati Ghasemi S. Self-care behaviors and related factors in chronic heart failure patients. *Int J Biomed Pub Health* 2018;1(1):42-47.
- Fjeldsoe B, Neuhaus M, Winkler E, Eakin E. Systematic review of maintenance of behavior change following physical activity and dietary interventions. *Health Psychol* 2011;30(1):99-109.
- Webb TL, Joseph J, Yardley L, Michie S. Using the internet to promote health behavior change: a systematic review and meta-analysis of the impact of theoretical basis, use of behavior change techniques, and mode of delivery on efficacy. *J Med Internet Res* 2010;12(1):e4-12. (doi:10.2196/jmir.1376)
- Pearson ES. Goal setting as a health behavior change strategy in overweight and obese adults: a systematic literature review examining intervention components. *Patient Educ Couns* 2012;87(1):32-42.
- Ahmed MM, El Degwy HM, Ali MI, Hegazy NH. The effect of educational intervention on knowledge, attitude and glycemic control in patients with type 2 diabetes mellitus. *Int J Commu Med Pub Health* 2015;2(3):302-307.
- Nielsen D, Ryg J, Nielsen W, Knold B, Nissen N, Brixen K. Patient education in groups increases knowledge of osteoporosis and adherence to treatment: a two-year randomized controlled trial. *Patient Educ Couns* 2010;81(2):155-160.
- Bartholomew LK, Mullen PD. Five roles for using theory and evidence in the design and testing of behavior change interventions. *J Pub Health Dent* 2011;71(S1):S20-S33.
- Ciechanowski PS, Katon WJ, Russo JE, Walker EA. The patient-provider relationship: attachment theory and adherence to treatment in diabetes. *Am J Psychiatry* 2001;158(1):29-35.
- Gruman J, Rovner MH, French ME, et al. From patient education to patient engagement: implications for the field of patient education. *Patient Educ Couns* 2010;78(3):350-356.
- Jamshidi N, Abbaszadeh A, Kalyani MN, Sharif F. Effectiveness of video information on coronary angiography patients' outcomes. *Collegian* 2013;20(3):153-159.
- Kav S, Yilmaz AA, Bulut Y, Dogan. Self-efficacy, depression and self-care activities of people with type 2 diabetes in Turkey. *Collegian* 2015;24(1):27-35.
- Comodore-Mensah Y, Himmelfarb CR. Patient education strategies for hospitalized cardiovascular patients: a systematic review. *J Cardiovasc Nurs* 2012;27(2):154-174.
- Stenberg U, Haaland-Ørverby M, Fredriksen K, Westermann KF, Kvisvik T. A scoping review of the literature on benefits and challenges of participating in patient education programs aimed at promoting self-management for people living with chronic illness. *Patient Educ Couns* 2016;99(11):1759-1771.
- Warsi A, Wang PS, LaValley MP, Avorn J, Solomon DH. Self-management education programs in chronic disease: a systematic review and methodological critique of the literature. *Arch Intern Med* 2004;164(15):1641-1649.
- Barnason S, White-Williams C, Rossi LP, et al. Evidence for therapeutic patient education interventions to promote cardiovascular patient self-management: A scientific statement for healthcare professionals from the American Heart Association. *Circ Cardiovasc Qual Outcomes* 2017;10(6):e000025. (doi:10.1161/HCQ.0000000000000025)
- Neville RG, Greene AC, Lewis S. Patient and health care professional views and experiences of computer agent-supported health care. *Inform Prim Care* 2006;14(1):11-15.
- Veneri DA, Zdanis K. Are technology-informed methods better than traditional approaches in educating patients? A meta-analysis. *Health Educ J* 2018;77(3):261-276.
- Caligtan CA, Carroll DL, Hurley AC, Gersh-Zaremski R, Dykes PC. Bedside information technology to support patient-centered care. *Int J Med Inform* 2012;81(7):442-451.
- Heapy AA, Higgins DM, Cervone D, Wandner L, Fenton BT, Kerns RD. A systematic review of technology-assisted self-management interventions for chronic pain. *Clin J Pain* 2015;31(6):470-492.

23. Isern D, Moreno A. A systematic literature review of agents applied in healthcare. *J Med Syst* 2016;40(2):43-59. (doi:10.1007/s10916-015-0376-2)
24. Or CKL, Karsh BT. A systematic review of patient acceptance of consumer health information technology. *J Am Med Inform Assoc* 2009; 16(4):550-560.
25. Or CKL, Tao D, Wang H. The effectiveness of the use of consumer health information technology in patients with heart failure: A meta-analysis and narrative review of randomized controlled trials. *J Telemed Telecare* 2017;23(1):155-166.
26. Bernhardt JM, Chaney JD, Chaney BH, Hall AK. New media for health education a revolution in progress. *Health Educ Behav* 2013;40(2):129-132.
27. Kooij L, Groen WG, van Harten WH. The effectiveness of information technology-supported shared care for patients with chronic disease: A systematic review. *J Med Internet Res* 2017;19(6):e221. (doi:10.2196/jmir.7405)
28. Suhling H, Rademacher J, Zinowsky I, et al. Conventional vs. tablet computer-based patient education following lung transplantation—a randomized controlled trial. *PLoS One* 2014;9(6):e90828-39. (doi:10.1371/journal.pone.0090828)
29. Gandapur Y, Kianoush S, Kelli HM, et al. The role of mHealth for improving medication adherence in patients with cardiovascular disease: a systematic review. *Eur J Qual Care Clin Outcomes* 2016;2(4):237-244.
30. Andrade AD, Anam R, Karanam C, Downey P, Ruiz JG. An overactive bladder online self-management program with embedded avatars: A randomized controlled trial of efficacy. *Urology* 2015;85(3):561-567.
31. Provoost S, Lau HM, Ruwaard J, Riper H. Embodied conversational agents in clinical psychology: A scoping review. *J Med Internet Res* 2017;19(5):e151. (doi: 10.2196/jmir.6553)
32. Xinyu D, Pin J. Development and application of virtual avatar interactive technology. 2008 International Conference on Advanced Computer Theory and Engineering. IEEE. (doi: 10.1109/ICACTE.2008.19)
33. Bickmore TW, Schulman D, Sidner C. Automated interventions for multiple health behaviors using conversational agents. *Patient Educ Couns* 2013;92(2):142-148.
34. Anam R, Andrade AD, Ruiz JG. Promoting lifestyle change through medical avatars (Chapter 26). In: Encyclopedia of E-Health and Telemedicine. IGI Global, 2016. (doi: 10.4018/978-1-4666-9978-6.ch026)
35. Abbott MB, Shaw P. Virtual nursing avatars: Nurse roles and evolving concepts of care. *Online J Issues Nurs* 2016;21(3):7. (doi:10.3912/OJIN.Vol21No03PPT39,05)
36. Lyles AA, Amresh A, Huberty J, Todd M, Lee RE. A mobile, avatar-based app for improving body perceptions among adolescents: A pilot test. *JMIR Serious Games* 2017;5(1):e4. (doi:10.2196/games.6354)
37. Song H, Kim J, Kwon RJ, Jung Y. Anti-smoking educational game using avatars as visualized possible selves. *Comput Human Behav* 2013;29(5):2029-2036.
38. Schulman D, Bickmore T. Persuading users through counseling dialogue with a conversational agent. Persuasive '09: Proceedings of the 4th International Conference on Persuasive Technology. ACM, 2009. (doi:10.1145/1541948.1541983)
39. Kothgassner OD, Hlavacs H, Beutl L, Glenk LM, Palme R, Felnhofer A. Two experimental virtual paradigms for stress research: Developing avatar-based approaches for interpersonal and evaluative stressors. International Conference on Entertainment Computing (ICEC) 2016. Springer, 2016: pp.51-62. (doi:10.1007/978-3-319-46100-7_5)
40. Whittemore R, Knaf K. The integrative review: updated methodology. *J Adv Nurs* 2005;52(5):546-553.
41. Torraco RJ. Writing integrative literature reviews: Guidelines and examples. *Hum . Resource Dev Rev* 2005;4(3):356-367. (doi:10.1177/1534484305278283)
42. Souza MT, Silva MD, Carvalho RD. Integrative review: what is it? How to do it? *Einstein (São Paulo)* 2010;8(1):102-106.
43. Popay J, Roberts H, Sowden A, et al. Guidance on the conduct of narrative synthesis in systematic reviews. A product from the ESRC Methods Programme. Version 1. 2006. (Accessed on Jan. 15, 2019, at <https://www.semanticscholar.org/paper/Guidance-on-the-conduct-of-narrative-synthesis-in-A-Popay-Roberts/ed8b23836338f6fdea0cc55e161b0fc5805f9e27>)
44. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6(7):e1000097. (doi: 10.1371/journal.pmed.1000097)
45. CASP. Critical Appraisal Skills Programme (CASP) (Randomised Controlled Trial) Checklist. (Accessed on Jan. 15, 2018, at <https://casp-uk.net/wp-content/uploads/2018/01/CASP-Randomised-Controlled-Trial-Checklist.pdf>)
46. An LC, Demers MR, Kirch MA, et al. A randomized trial of an avatar-hosted multiple behavior change intervention for young adult smokers. *J Natl Cancer Inst Monogr* 2013;2013(47):209-215.
47. Pinto MD, Hickman RL Jr, Clochesy J, Buchner M. Avatar-based depression self-management technology: promising approach to improve depressive symptoms among young adults. *Appl Nurs Res* 2013;26(1):45-48.
48. Bickmore TW, Pfeifer LM, Paasche-Orlow MK. Using computer agents to explain medical documents to patients with low health literacy. *Patient Educ Couns* 2009;75(3):315-320.
49. Bickmore TW, Silliman RA, Nelson K, et al. A randomized controlled trial of an automated exercise coach for older adults. *J Am Geriatr Soc* 2013;61(4):1676-1683.
50. Annaim A, Lassiter M, Viera AJ, Ferris M. Interactive media for parental education on managing children chronic condition: a systematic review

- of the literature. *BMC Pediatr* 2015;15:201. (doi:10. 1186/s12887-015-0517-2)
51. Charlier N, Zupancic N, Fieuws S, Denhaerynck K, Zaman B, Moons P. Serious games for improving knowledge and self-management in young people with chronic conditions: a systematic review and meta-analysis. *J Am Med Inform Assoc* 2015;23(3):230-239.
 52. Bedra M, Wick E, Brotman D, Finkelstein J. Avatar-based interactive ileostomy education in hospitalized patients. *Stud Health Technol Inform* 2013;190:83-85.
 53. Clark RA, Fredericks B, Buitendyk NJ, et al. Development and feasibility testing of an education program to improve knowledge and self-care among Aboriginal and Torres Strait Islander patients with heart failure. *Rural Remote Health* 2015;15(3):3231. (<https://www.rhh.org.au/journal/article/3231>)
 54. Joubert M, Armand C, Morera J, Tokayeva L, Guillaume A, Reznik Y. Impact of a serious videogame designed for flexible insulin therapy on the knowledge and behaviors of children with type 1 diabetes: the LUDIDIAB pilot study. *Diabetes Technol Ther* 2016;18(2):52-58.
 55. Gysels M, Higginson IJ. Interactive technologies and videotapes for patient education in cancer care: systematic review and meta-analysis of randomised trials. *Support Care Cancer* 2007;15(1):7-20.
 56. Livne Y, Peterfreund I, Sheps J. Barriers to patient education and their relationship to nurses' perceptions of patient education climate. *Clin Nurs Stud* 2017;5(4):65-72. (doi:10.5430/cns.v5n4p65)
 57. Azevedo R, Bernardes M, Fonseca J, Lima A. Smartphone application for rheumatoid arthritis self-management: cross-sectional study revealed the usefulness, willingness to use and patients' needs. *Rheumatol Int* 2015;35(10):1675-1685.
 58. Chiou CP, Chung YC. Effectiveness of multimedia interactive patient education on knowledge, uncertainty and decision-making in patients with end-stage renal disease. *Clin Nurs Stud* 2012;21(9-10):1223-1231.
 59. Blok AC, May CN, Sadasivam RS, Houston TK. Virtual patient technology: Engaging primary care in quality improvement innovations. *JMIR Med Educ* 2017;3(1):e3. (doi:10.2196/mededu.7042)
 60. Lisetti CL. 10 advantages of using avatars in patient-centered computer-based interventions for behavior change. *SIGHIT Record* 2012;2(1). (Accessed on Oct. 15, 2018, at <http://ascl.cis.fiu.edu/uploads/1/3/4/2/13423859/lisetti-10advantages-ih12.pdf>)
 61. Friedman AJ, Cosby R, Boyko S, Hatton-Bauer J, Turnbull G. Effective teaching strategies and methods of delivery for patient education: A systematic review and practice guideline recommendations. *J Cancer Educ* 2011;26(1):12-21.
 62. Napolitano MA, Hayes S, Russo G, Muresu D, Giordano A, Foster GD. Using avatars to model weight loss behaviors: participant attitudes and technology development. *J Diabetes Sci Technol* 2013;7(4):1057-1065.
 63. Lo SF, Wang YT, Wu LY, Hsu MY, Chang SC, Hayter M. Multimedia education programme for patients with a stoma: effectiveness evaluation. *J Adv Nurs* 2011;67(1):68-76.
 64. Lisetti C, Amini R, Yasavur U, Rishe N. I can help you change! an empathic virtual agent delivers behavior change health interventions. *ACM Transactions on Management Information Systems* 2013;4(4). (doi: <https://doi.org/10.1145/2544103>)
 65. Butterworth SW, Linden A, McClay W. Health coaching as an intervention in health management programs. *Dis Manag Health Outcomes* 2007;15(5):299-307.
 66. Olsen JM, Nesbitt BJ. Health coaching to improve healthy lifestyle behaviors: an integrative review. *Am J Health Promot* 2010;25(1):e1-e12.
 67. Boren SA, Gunlock TL, Peebles MM, Krishna S. Computerized learning technologies for diabetes: a systematic review. *J Diabetes Sci Technol* 2008;2(1):139-146.
 68. List TP. How technology is shaping patient education. 2013. (Accessed on Jan. 16, 2019, at http://www.pmlive.com/pmhub/medical_communications/fti_consulting/white_papers_and_resources/how_technology_is_shaping_patient_education)
 69. Rosal MC, Heyden R, Mejilla R, et al. A virtual world versus face-to-face intervention format to promote diabetes self-management among african american women: a pilot randomized clinical trial. *JMIR Res Protoc* 2014; 3(4):e54. (doi:10.2196/resprot.3412)
 70. Plow M, Golding M. Using mHealth technology in a self-management intervention to promote physical activity among adults with chronic disabling conditions: Randomized controlled trial. *JMIR Mhealth Uhealth* 2017;5(12):e185. (doi: 10.2196/mhealth.6394)
 71. Johnson C, Feinglos M, Pereira K, et al. Feasibility and preliminary effects of a virtual environment for adults with type 2 diabetes: pilot study. *JMIR Res Protoc* 2014;3(2):e23. (doi: 10.2196/resprot.3045)
 72. Kato PM, Cole SW, Bradlyn AS, Pollock BH. A video game improves behavioral outcomes in adolescents and young adults with cancer: a randomized trial. *Pediatrics* 2008;122(2):e305-e317.
 73. Ruggiero L, Moadsiri A, Quinn LT, et al. Diabetes island: preliminary impact of a virtual world self-care educational intervention for african americans with type 2 diabetes. *JMIR Serious Games* 2014;2(2):e10. (doi:10.2196/games.3260)
 74. Välimäki M, Hätönen H, Lahti M, Kuosmanen L, Adams CE. Information and communication technology in patient education and support for people with schizophrenia. *Cochrane Database Syst Rev* 2012;10: D007198. (doi:10.1002/14651858.CD007198.pub2)
 75. Edwards RA, Bickmore T, Jenkins L, Foley M, Manjourides J. Use of an interactive computer agent to support breastfeeding. *Matern Child Health J* 2013;17(10):1961-1968.