

*Int. J. Aquat. Biol.* (2020) 8(2): 143-147  
ISSN: 2322-5270; P-ISSN: 2383-0956  
Journal homepage: [www.ij-aquaticbiology.com](http://www.ij-aquaticbiology.com)  
© 2020 Iranian Society of Ichthyology

## Review Article

# Alkaline phosphatase activity as a biochemical biomarker in aqua-toxicological studies

Mahdi Banaee\*

Aquaculture Department, Faculty of Natural Resources and the Environment, Behbahan Khatam Alanbia University of Technology, Behbahan, Iran.

**Abstract:** Alkaline phosphatase (ALP) is a glycoprotein with a metallophosphatase structure that catalyzes the hydrolysis of monophosphate esters of biomolecule esters at alkaline pH. ALP activity is a useful bioindicator to assess the physiological health of cellular membranes, cell growth, apoptosis and cell migration, cellular metabolic status, hepatocyte function, and detoxification activity in hepatocytes. ALP activity is detected in a colorimetric method using the para-nitrophenyl phosphate substrate (p-NPP) at a wavelength of 405 nm in biological samples. Cell hemolysis, especially erythrocytes; increased levels of sex hormones and corticosteroids, biological infections, and poor nutrition can adversely affect ALP activity.

*Article history:*

Received 16 January 2020

Accepted 20 April 2020

Available online 25 April 2020

*Keywords:* Detoxification, Alkaline phosphatase, Interfering factors, Aquatic organisms.

## Introduction

Alkaline phosphatase (ALP) is a metalloenzyme that is anchored to the cell membrane by a glycosylphosphatidylinositol (GPI) (Atkins et al., 2011). ALP is a glycoprotein with an external domain and a catalytic domain with a serine base. The activity of ALP depends on the binding of two  $Zn^{+2}$  ions and one  $Mg^{+2}$  ion to the active site of the enzyme. The serine base can inhibit the activity of the ALP enzyme (Banaee and Taheri, 2019). Therefore, ALP activity depends on the concentration of these ions in cells and intercellular fluids (Banaee et al., 2011; Nafisi Bahabadi et al., 2014; Soleimany et al., 2016). Therefore, gender, health status, ontogeny (Kim et al., 2001), oral supplements administration (Taheri et al., 2017; Gholizadeh Zare Tavana et al., 2018), or exposure to pollutants, and environmental factors can have a significant effect on ALP activity (Woo et al., 2018).

Alkaline phosphatase plays a vital role in phosphate hydrolysis and membrane transfer (Derikvandy et al., 2020), and is also involved in transphosphorylation of many biomolecules, such as nucleotides, proteins, and alkaloids in alkaline pH

(Nematdoost Haghi and Banaee, 2017). The activity of this enzyme is essential for cell growth, planned cell death (apoptosis), and cell migration (Sharma et al., 2014; Banaee et al., 2019c). ALP activity often increases during differentiation of B lymphocytes to plasmocytes (Akcakaya et al., 2007).

Alkaline phosphatase is often synthesized in the liver, bones, and to a lesser extent, in the intestine and kidneys (Telega, 2018). Additionally, the ALP gene is expressed in the endothelial blood vessels and the brain tissue of fish (Goishi and Klagsbrun, 2004). ALP activity is essential for bone mineralization (Atkins et al., 2011). So far, 12 isoenzymes from ALP have been identified that may have different origins and synthesis sites.

Alkaline phosphatase plays a vital role in glycogen metabolism (Kim et al., 2001). The activity of this enzyme in hepatocytes can inactivate phosphorylase and increase the rate of glycogen breakdown in the liver (Raisi et al., 2018; Rezaei Shadegan and Banaee, 2018). Increased levels of synthesis and ALP activity can play an essential role in providing energy for stressed cells (Agrahari et al., 2007; Saha and Kaviraj, 2009).

\*Correspondence: Mahdi Banaee  
E-mail: [mahdibanaee@yahoo.com](mailto:mahdibanaee@yahoo.com)

**Purpose of measuring ALP activity:** Alkaline phosphatase is a non-specific indicator for the diagnosis of liver and bone damage. Furthermore, ALP is an excellent biological indicator for assessing cellular stress (Banday et al., 2019). It plays a vital role in the xenobiotic detoxification process in hepatocytes (Vaziryan et al., 2017; Hatami et al., 2019).

**ALP activity measurement method:** Alkaline phosphatase activity is measured by a quantitative colorimetric method using the para-nitrophenyl phosphate substrate (p-NPP) at a wavelength of 405 nm (Moss and Henderson, 1999).

**Reasons for increased ALP activity:** Tumor formation in the liver and bone tissue, hyperthyroidism, bile duct obstruction, liver nodules, liver cysts, liver failure, and increased ALP biosynthesis rates in hepatocytes, and intestinal dysfunction can lead to increased ALP activity in the serum or plasma. For example, ALP activity in rainbow trout, *Oncorhynchus mykiss*, and *Cyprinus carpio* fish plasma increased significantly after exposure to diazinon (Banaee et al. 2011, 2012). Increased ALP activity in the body tissues of freshwater fish of *Alburnus sellal* exposed to fenpropathrin indicates damage to cell membranes (Banaee et al., 2014). Additionally, damage to the cell membrane led to the release of ALP into the fish plasma exposed to titanium dioxide nanoparticles (Banaee et al., 2019b). Increased ALP activity in fish hepatocytes exposed to dimethoate and bacillar fertilizer may also be a physiological response to increased glycogen breakdown in the liver (Rezaei Shadegan and Banaee, 2018).

Nematdoost Haghi and Banaee (2017) and Wan Do et al. (2019) stated that damage to the hepatocyte membrane, gallbladder ducts, intestinal epithelial cells, and renal ducts increased ALP in fish plasma exposed to microplastics and paraquat. Banaee et al. (2019b) reported an increase in ALP activity in plasma of fish exposed to paraquat and nano-TiO<sub>2</sub>. Research shows that sometimes pollutants may increase ALP activity in cells by affecting the process of transphosphorylation in cells (Banaee et al., 2019a). Increased ALP activity in cells of *C. carpio* following oral exposure to zinc oxide nanoparticles may be

related to the dysfunction of cells (Taheri et al., 2017). Furthermore, an increase in the production rate of reactive oxygen species (ROS) may elevate ALP activity (Banday et al., 2020; Banaee et al., 2017). Woo et al. (2018) observed elevated activity of ALP in plasma of *C. carpio* exposed to trichlorfon (Woo et al., 2018). An increase in ALP activity was observed after exposure of *Clarias gariepinus* to deltamethrin (Amin and Hashem, 2012; Eni et al., 2019), cypermethrin (Eni et al., 2019), heavy metals (Banday et al., 2019). Similar results have been reported after exposing *Oreochromis niloticus* to deltamethrin (Abdel-Daim et al., 2015; Abdelkhalek et al., 2015), diazinon (Abdelkhalek et al., 2017). An increase in ALP activity was observed in the blood of *Channa punctatus* exposed to monocrotophos (Agrahari et al., 2007).

**Reasons for decreased ALP activity:** ALP activity may be reduced by malnutrition, especially protein, zinc, magnesium and vitamin C deficiency, and increased vitamin D levels in the diet. Anemia and hypothyroidism can lead to decreased ALP activity. Severe cellular necrosis can be another reason for decreased ALP activity in fish plasma (Banaee et al., 2016). Moreover, disturbance in the transport of zinc, magnesium, and phosphorus ions from cell membranes can lead to decreased ALP activity (Banaee et al., 2019c; Banaee et al., 2020). Decreased absorption of essential ions for ALP biosynthesis can lead to decreased enzyme activity (Veerappan et al., 2012). Besides, the interaction of environmental pollutants with ALP can reduce its activity (Banaee et al., 2019a; b). For example, decreased ALP activity in gill cells of the rainbow trout exposed to cadmium chloride may be due to the effect of cadmium on cell membranes (Evaz-Zadeh Samani et al., 2017).

Exposure of *C. carpio* to Cd reduced ALP activity in muscle tissue (Banaee et al., 2015). Decreased plasma ALP activity may be due to damage to erythrocyte membranes and their hemolysis (Farah et al., 2012). Vaziryan et al. (2017) suggested that reduced ALP activity may be associated with impaired ALP biosynthesis, hepatic cell necrosis, and damage to the intestinal epithelial cells of fish exposed to oral

aflatoxins. Decreased ALP activity may affect bone marrow cell activity (Prins et al., 2014). Decrease in ALP activity was reported in serum of *Rhamdia quelen* fed with a feed artificially contaminated with aflatoxin B1 (Anater et al., 2020).

**Interventional factors in measuring ALP activity:** Sex hormones (Ahmad et al., 2002), viral and bacterial infections (Banaee et al., 2017; Xia et al., 2017), parasites infection, administration of antibiotics (Gora et al., 2018), dietary supplements, hemolysis of blood cells (Farah et al., 2012) and feeding fish with oxidized and fatty foods can interfere with ALP activity in blood or tissue samples.

### Conclusion

Measurement of the ALP activity in the intercellular and extracellular fluids can be indirectly useful for evaluating the membrane function of cells exposed to xenobiotics. Interaction of environmental pollutants with the biochemical structure of ALP or lipid peroxidation of cell membranes can cause fluctuations in ALP activity in cells or extracellular fluids. Therefore, each change in ALP activity outside the normal range can indicate stress on the cells.

### References

- Balon E.K. (1984). Reflections on some decisive events in the early life of fishes. Transactions of the American Fisheries Society, 113: 178-185.
- Berg L.S. (1965). Freshwater fishes of the U.S.S.R. and adjacent countries. Volume 3, 4th edition. Israel Program for Scientific Translations Ltd, Jerusalem. (Russian version published 1949).
- Abdel-Daim M.M., Abdelkhalek N.K., Hassan A.M. (2015). Antagonistic activity of dietary allicin against deltamethrin-induced oxidative damage in freshwater Nile tilapia; *Oreochromis niloticus*. Ecotoxicology and Environmental Safety, 111: 146-52.
- Abdelkhalek N.K.M., Eissa I.A.M., Ahmed E., Kilany O.E., El-Adl M., Dawood M.A.O., Hassan A.M., Abdel-Daim M.M. (2017). Protective role of dietary *Spirulina platensis* against diazinon-induced Oxidative damage in Nile tilapia; *Oreochromis niloticus*. Environmental Toxicology and Pharmacology, 54: 99-104.
- Abdelkhalek N.K., Ghazy E.W., Abdel-Daim M.M. (2015). Pharmacodynamic interaction of *Spirulina platensis* and deltamethrin in freshwater fish Nile tilapia, *Oreochromis niloticus*: impact on lipid peroxidation and oxidative stress. Environmental Science and Pollution Research International, 24(4): 3023-3031.
- Agrahari S., Pandey K.C., Gopal K. (2007). Biochemical alteration induced by monocrotophos in the blood plasma of fish, *Channa punctatus* (Bloch). Pesticide Biochemistry and Physiology, 88(3): 268-272.
- Ahmad M., Abdel-Tawwab M., Shalaby A., Khattab Y. (2002). Effects of 17 a-methyltestosterone on growth performance and some physiological changes of Nile tilapia (*Oreochromis niloticus* L.) fingerlings. Egyptian Journal of Aquatic Biology and Fisheries, 6(2): 1-23.
- Akcakaya H., Aroymak A., Gokce S. (2007). A quantitative colorimetric method of measuring alkaline phosphatase activity in eukaryotic cell membranes. Cell Biology International, 31(2): 186-190.
- Amin K.A., Hashem K.S. (2012). Deltamethrin-induced oxidative stress and biochemical changes in tissues and blood of catfish (*Clarias gariepinus*): antioxidant defense and role of alpha-tocopherol. BMC Veterinary Research, 8: 45.
- Anater A., Domingues Araújo C.M.T., Rocha D.C.C., Ostrensky A., Filho J.R.E., Ribeiro D.R., Pimpão C.T. (2020). Evaluation of growth performance, hematological, biochemical and histopathological parameters of *Rhamdia quelen* fed with a feed artificially contaminated with aflatoxin B1. Aquaculture Reports, 17: 100326.
- Atkins G.J., Findlay D.F., Anderson P.H., Morris H.A. (2011). Target genes: bone proteins. In: D. Feldman, J.W. Pike, J.S. Adams (Eds). Vitamin D, 3<sup>rd</sup> edition, Academic Press. pp: 411-424.
- Banaee M., Akhlaghi M., Soltanian S., Gholamhosseini A., Heidarieh H., Fereidouni M.S. (2019a). Acute exposure to chlorpyrifos and glyphosate induces changes in hemolymph biochemical parameters in the crayfish, *Astacus leptodactylus* (Eschscholtz, 1823). Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology, 222: 145-155.
- Banaee M., Akhlaghi M., Soltanian S., Sureda A., Gholamhosseini A., Rakhshaninejad M. (2020). Combined effects of exposure to sub-lethal concentration of the insecticide chlorpyrifos and the herbicide glyphosate on the biochemical changes in the freshwater crayfish *Pontastacus leptodactylus*.

Ecotoxicology, ( In press)

- Banaee M., Mehrpak M., Nematdoost Haghi B., Noori A. (2015). Amelioration of cadmium-induced changes in biochemical parameters of the muscle of Common Carp (*Cyprinus carpio*) by Vitamin C and Chitosan. *International Journal of Aquatic Biology*, 3(6); 362-371.
- Banaee M., Mirvaghefi A.R., Mojazi Amiri B., Rafiee G.R. (2012). Biochemical characteristics of blood and histopathological study of experimental diazinon poisoning in common carp (*Cyprinus carpio*). *Journal of Fisheries (Iranian Journal of Natural Resource)*, 64 (1): 1-12.
- Banaee M., Shahafve S., Tahery S., Nematdoost Haghi B., Vaziriyani M. (2016). Sublethal toxicity of TiO<sub>2</sub> nanoparticles to common carp (*Cyprinus carpio*, Linnaeus, 1758) under visible light and dark conditions. *International Journal of Aquatic Biology*, 4(6): 370-377.
- Banaee M., Soleimany V., Nematdoost Haghi B. (2017). Therapeutic effects of marshmallow (*Althaea officinalis* L.) extract on plasma biochemical parameters of common carp infected with *Aeromonas hydrophila*. *Veterinary Research Forum*, 8(2): 145-153.
- Banaee M., Soltanian S., Sureda A., Gholamhosseini B N.H., Akhlaghi M., Derikvandy A. (2019c). Evaluation of single and combined effects of cadmium and microplastic particles on biochemical and immunological parameters of common carp (*Cyprinus carpio*). *Chemosphere*, 236: 124335.
- Banaee M., Sureda A., Mirvaghefi A.R., Ahmadi K. (2011). Effects of diazinon on biochemical parameters of blood in rainbow trout (*Oncorhynchus mykiss*). *Pesticide Biochemistry and Physiology*, 99(1): 1-6.
- Banaee M., Sureda A., Taheri S., Hedayatzadeh F. (2019). Sub-lethal effects of dimethoate alone and in combination with cadmium on biochemical parameters in freshwater snail, *Galba truncatula*. *Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology*, 220: 62-70.
- Banaee M., Sureda A., Zohiery F., Nematdoust Hagi B., Garanzini D.S. (2014). Alterations in biochemical parameters of the freshwater fish, *Alburnus mossulensis*, exposed to sub-lethal concentrations of Fenpropathrin. *International Journal of Aquatic Biology*, 2(2): 58-68.
- Banaee M., Taheri S. (2019). Metal bioaccumulation, oxidative stress, and biochemical alterations in the freshwater snail (*Galba truncatula*) exposed to municipal sewage. *Journal of Advances in Environmental Health Research*, 7(1): 8-17.
- Banaee M., Tahery S., Nematdoost Haghi B., Shahafve S., Vaziriyani M. (2019b). Blood biochemical changes in common carp (*Cyprinus carpio*) upon co-exposure to titanium dioxide nanoparticles and paraquat. *Iranian Journal of Fisheries Sciences*, 8(2): 242-255.
- Banday U.Z., Swaleh S.B., Usmani N. (2019). Insights into the heavy metal-induced immunotoxic and genotoxic alterations as health indicators of *Clarias gariepinus* inhabiting a rivulet. *Ecotoxicology and Environmental Safety*, 183: 109584.
- Banday U.Z., Swaleh S.B., Usmani N. (2020). Heavy metal toxicity has an immunomodulatory effect on metallothionein and glutathione peroxidase gene expression in *Cyprinus carpio* inhabiting a wetland lake and a culture pond. *Chemosphere*, 251: 126311.
- Derikvandy A., Pourkhabbaz H.R., Banaee M., Sureda A., Nematdoost Haghi B., Pourkhabbaz A.R. (2020). Genotoxicity and oxidative damage in zebrafish (*Danio rerio*) after exposure to effluent from ethyl alcohol industry. *Chemosphere*, 251: 126609.
- Eni G., Ibor O.R., Andem A.B., Oku E.E., Chukwuka A.V., Adeogun A.O., Arukwe A. (2019). Biochemical and endocrine-disrupting effects in *Clarias gariepinus* exposed to the synthetic pyrethroids, cypermethrin and deltamethrin. *Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology*, 225: 108584.
- Evaz-Zadeh Samani H., Banaee M., Shoukat P., Noori A., Mousavi Dehmoredi L. (2017). Protective effects of dietary *Spirulina platensis* against cadmium-induced oxidative stress in gills of rainbow trout. *Iranian Journal of Toxicology*, 11(4): 5-12.
- Farah H.S., Al-Atoom A.A., Shehab G.M. (2012). Explanation of the decrease in alkaline phosphatase (ALP) activity in hemolysed blood samples from the clinical point of view : In vitro study. *Jordan Journal of Biological Sciences*, 5(2): 125-128.
- Gholizadeh Zare Tavana B., Banaee M., Yousefi Jourdehi A., Nematdoost Haghi B., Seyed Hassani M. (2018). Effects of selenium (Sel-Plex) supplement on blood biochemical parameters of juvenile Siberian sturgeon (*Acipenser baeri*). *Iranian Journal of Fisheries Sciences*, 17(2): 300-312.
- Goishi K., Klagsbrun M. (2004). Vascular endothelial growth factor and its receptors in embryonic zebrafish blood vessel development. *Current Topics in Developmental Biology*, 62: 127-152.

- Gora A.H., Sahu N.P., Sahoo S., Rehman S., Ahmad I., Agarwal D., Ahmad Dar S., Rasool S.I. (2018). Metabolic and haematological responses of *Labeo rohita* to dietary fucoidan. *Journal of Applied Animal Research*, 46(1): 1042-1050.
- Hatami M., Banaee M., Nematdoost Haghi B. (2019). Sub-lethal toxicity of chlorpyrifos alone and in combination with polyethylene glycol to common carp (*Cyprinus carpio*). *Chemosphere*, 219: 981-988.
- Kim B.G., Divakaran S., Brown C.L., Ostrowski A.C. (2001). Comparative digestive enzyme ontogeny in two marine larval fishes: Pacific threadfin (*Polydactylus sexfilis*) and bluefin trevally (*Caranx melampygus*). *Fish Physiology and Biochemistry*, 24: 225-241.
- Moss D.V., Henderson A.R. (1999). Clinical enzymology. In: C.A. Burtis, E.R. Ashwood (Eds.). *Tietz textbook of clinical chemistry*. 3rd ed., Philadelphia: W.B. Saunders Company. pp: 617-721.
- Nafisi Bahabadi M., Banaee M., Taghiyan M., Nematdoost Haghi B. (2014). Effects of dietary administration of yarrow extract on growth performance and blood biochemical parameters of rainbow trout (*Oncorhynchus mykiss*). *International Journal of Aquatic Biology*, 2(5): 275-285.
- Nematdoost Haghi B., Banaee M. (2017). Effects of microplastic particles on paraquat toxicity to common carp (*Cyprinus carpio*): biochemical changes. *International Journal of Environmental Science and Technology*, 14(3): 521-530.
- Prins H.J., Braat A.K., Gawlitta D., Dhert W.J., Egan D.A., Tijssen-Slump E., Yuan H., Coffey P.J., Rozemuller H., Martens A.C. (2014). In vitro induction of alkaline phosphatase levels predicts in vivo bone forming capacity of human bone marrow stromal cells. *Stem Cell Research*, 12(2): 428-40.
- Raisi M., Pourkhabaz H.R., Banaee M., Pourkhabaz A., Javanmardi S. (2018). Effects of Pirimicarb carbamate insecticide alone and in combination with lead (Pb) on biochemical parameters of soft tissues in freshwater snail, *Galba truncatula*. *International Journal of Aquatic Biology*, 6(3): 126-137.
- Rezaei Shadegan M., Banaee M. (2018). Effects of dimethoate alone and in combination with Bacilar fertilizer on oxidative stress in common carp, *Cyprinus carpio*. *Chemosphere*, 208: 101-107.
- Saha S., Kaviraj A. (2009). Effects of cypermethrin on some biochemical parameters and its amelioration through dietary supplementation of ascorbic acid in freshwater catfish *Heteropneustes fossilis*. *Chemosphere*, 74(9): 1254-1259.
- Sharma U., Pal D., Prasad R. (2014). Alkaline phosphatase: an overview. *Indian Journal of Clinical Biochemistry*, 29(3): 269-278.
- Soleimany V., Banaee M., Mohiseni M., Nematdoost Haghi B., Mousavi Dehbourdi L. (2016). Evaluation of pre-clinical safety and toxicology of *Althaea officinalis* extracts as naturopathic medicine for common carp (*Cyprinus carpio*). *Iranian Journal of Fisheries Sciences*, 15(2): 613-629.
- Taheri S., Banaee M., Nematdoost Haghi B., Mohiseni M. (2017). Effects of Dietary Supplementation of Zinc Oxide Nanoparticles on Some Biochemical Biomarkers in Common Carp (*Cyprinus carpio*). *International Journal of Aquatic Biology*, 5(5): 286-294.
- Telega G.W. (2018). Jaundice. In: R.M. Kliegma, P.S. Lye, B. Bordini, H. Toth, D. Basel (Eds.). *Nelson Pediatric Symptom-Based Diagnosis*, Elsevier. pp: 255-274.
- Vaziryan M., Banaee M., Nematdoost Haghi B., Mohiseni M. (2017). Effects of dietary exposure to aflatoxins on some plasma biochemical parameters of common carp (*Cyprinus carpio*). *Iranian Journal of Fisheries Sciences*, 17(3): 487-502.
- Veerappan M., Hwang I., Pandurangan M. (2012). Effect of cypermethrin, carbendazim and their combination on male albino rat serum. *International Journal of Experimental Pathology*, 93(5): 361-369.
- Wan Do J., Saravanan M., Nam S.E., Lim H.J., Rhee J.S. (2019). Waterborne manganese modulates immunity, biochemical, and antioxidant parameters in the blood of red seabream and black rockfish. *Fish and Shellfish Immunology*, 88: 546-555.
- Woo S.J., Kim N.Y., Kim S.H., Ahn S.J., Seo J.S., Jung S.H., Cho M.Y., Chung J.K. (2018). Toxicological effects of trichlorfon on hematological and biochemical parameters in *Cyprinus carpio* L. following thermal stress. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 209: 18-27.
- Xia H., Tang Y., Lu F., Luo Y., Yang P., Wang W., Jiang J., Li N., Han Q., Liu F., Liu L. (2017). The effect of *Aeromonas hydrophila* infection on the non-specific immunity of blunt snout bream (*Megalobrama amblycephala*). *Central European Journal of Immunology*, 42(3): 239-243.