Scientia Agropecuaria 11(2): 281 (2020)



# Scientia Agropecuaria

Website: http://revistas.unitru.edu.pe/index.php/scientiaagrop

Facultad de Ciencias Agropecuarias

provided by Universidad Nacional de Truiillo: Publicacione

Universidad Nacional de Trujillo

# Letter

## Will SARS-CoV-2 cause diseases in poultry?

Sebastián Iglesias-Osores<sup>\*</sup>; Johnny Leandro Saavedra-Camacho

Facultad de Ciencias Biológicas, Universidad Nacional Pedro Ruiz Gallo, Calle Juan XXIII, Lambayeque, Peru.

Received April 15, 2020. Accepted May 18, 2020.

### Dear Editor:

Pandemics are frequently caused by a group of viruses who causes infections in birds and mammals called coronavirus (Sahin et al., 2020). Its high capacity to infect humans, such as the case of Severe Acute Respiratory Syndrome (SARS) or Middle East Respiratory Syndrome (MERS), has taken great relevance in the outbreak of a new coronavirus disease (COVID-19) in a market in the city of Wuhan, China, spreading throughout the Asian continent and later to more than 180 countries in the world, causing a pandemic. The coronaviruses also cause diseases in mammals such as dogs, mice, horses, whales, cats, and with animals with economic importance and global consumption such as birds (including poultry such as broilers and turkeys) (Biswas et al., 2020). Birds are possible reservoirs of SARS-CoV-2 and can transmit it to humans or vice versa? COVID-19 is caused in humans by Betacoronavirus SARS-CoV-2, in poultry coronaviruses cause Avian Infectious Bronchitis by Gammacoronavirus which produces a highly contagious disease in chickens (Gorbalenya et al., 2020). Coronaviruses, in general, are spread throughout the planet and are highly infectious, in addition to being extremely difficult to control because they have high genetic diversity over large areas, short multiplication periods and a high rate of mutation (Sahin et al., 2020). SARS-CoV-2 uses a host cell receptor angiotensinconverting enzyme II (ACE2) and IBV (Infectious Bronchitis Virus) enters the body primarily by clathrinmediated endocytosis and requires a classical endosomal/lysosomal system (Wang et al., 2019). IBV produces respiratory tract infection, and affects the reproductive tract, and some strains can cause nephritis, SARS-CoV-2 cause severe acute respiratory syndrome (Ennaji et al., 2019; Sahin et al., 2020). IBV genotypes and serotypes are related to the vaccines strains (S1) and SARS-CoV-2 does not haver variant yet (Bande et al., 2017). The IBV incubation period is very short compared to SARS-CoV-2, which is 18 to 36 hours and depends on the dose of virus infection, and clinical signs appear within 24 and 48 hours of exposure to the virus (Cavanagh and Nagi, 2008). Information about zoonotic reservoirs and its transmission among them can help to understand the COVID-19 outbreaks and zoonotic transmission of IBV, we should have a clear knowledge of its reservoir host, distribution pattern and spreading routes of IBV. SARS-CoV-2 has not a probability of infecting chickens or any other poultry, the main reason for non-infection in birds it is both viruses have different receptors on the hosts and belong to phylogenetically different groups.

Keywords: IBV; COVID-19; SARS-CoV-2; poultry; zoonotic.

## ORCID

S. Iglesias-Osores Dhttps://orcid.org/0000-0002-4984-4656 J.L. Saavedra-Camacho D https://orcid.org/0000-0002-3842-4314

#### References

- Bande, F.; Arshad, S.S.; Rahman Omar, A.; et al. 2017. Global distributions and strain diversity of avian infectious bronchitis virus: a review. Animal Health Research Reviews 18(1): 70-83. Biswas, A.; Bhattacharjee, U.; Chakrabarti, A.K.; *et al.* 2020.
- Emergence of Novel Coronavirus and COVID-19: whether to stay or die out? Critical Reviews in Microbiology 46(2): 182-193. Cavanagh, D.; Naqi, S.A. 2003. Infectious bronchitis. En: Y.M. Saif
- (Editor). Diseases of Poultry 11ava Edición. Ames: Iowa State University Press. USA. Pp. 101-119.
- Ennaji, Y.; Khataby, K.; Ennaji, M.M. 2019. Infectious bronchitis virus in poultry: Molecular epidemiology and factors leading to the emergence and reemergence of novel strains of infectious emergence and reemergence of novel strains of infectious bronchitis virus. En: Moulay Mustapha Ennaji (Editor). Emerging and Reemerging Viral Pathogens: Volume 2: Applied Virology Approaches Related to Human, Animal and Environmental Pathogens. 1era Edición. Academic Press. USA. Pp. 31-44.
  Gorbalenya, A.E.; Baker, S.C.; Baric, R.S.; *et al.* 2020. The species Severe acute respiratory syndrome-related coronavirus: classificing. 2019. PCoV and parting it SAPS CoV/2. Network
- classifying 2019-nCoV and naming it SARS-CoV-2. Nature Microbiology 5(4): 536-544. Sahin, A.; Erdogan, A.; Mutlu Agaoglu, P.; et al. 2020. 2019 Novel
- Coronavirus (COVID-19) Outbreak: A Review of the Current Literature. Eurasian Journal of Medicine and Oncology 4(1): 1-7.
- Wang, H.; Yuan, X.; Sun, Y.; et al. 2019. Infectious bronchitis virus entry mainly depends on clathrin mediated endocytosis and requires classical endosomal/lysosomal system. Virology 528: 118-136.

#### **Cite this article:**

Iglesias-Osores, S.; Saavedra-Camacho, J.L. 2020. Will SARS-CoV-2 cause diseases in poultry?. Scientia Agropecuaria 11(2): 281.

\* Corresponding author:

siglesias@unprg.edu.pe (S. Iglesias-Osores).

© 2020 All rights reserved DOI: 10.17268/sci.agropecu.2020.02.17