

**CLOSED FORMULAS AND IDENTITIES FOR THE BELL
POLYNOMIALS AND FALLING FACTORIALS**

FENG QI, DA-WEI NIU, DONGKYU LIM, AND BAI-NI GUO

ABSTRACT. The authors establish a pair of closed-form expressions for special values of the Bell polynomials of the second kind for the falling factorials, derive two pairs of identities involving the falling factorials, find an equivalent expression between two special values for the Bell polynomials of the second kind, and present five closed-form expressions for the (modified) spherical Bessel functions.

1. MOTIVATIONS

The Bell polynomials of the second kind, also known as partial Bell polynomials, denoted by $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ for $n \geq k \geq 0$, are defined in [1, p. 134, Theorem A] by

$$(1.1) \quad B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = \sum_{\substack{1 \leq i \leq n-k+1 \\ \ell_i \in \{0\} \cup \mathbb{N} \\ \sum_{i=1}^{n-k+1} i \ell_i = n \\ \sum_{i=1}^{n-k+1} \ell_i = k}} \frac{n!}{\prod_{i=1}^{n-k+1} \ell_i!} \prod_{i=1}^{n-k+1} \left(\frac{x_i}{i!}\right)^{\ell_i}.$$

The Faà di Bruno formula [1, p. 139, Theorem C] can be described in terms of the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ by

$$(1.2) \quad \frac{d^n}{dx^n} f \circ h(x) = \sum_{k=0}^n f^{(k)}(h(x)) B_{n,k}(h'(x), h''(x), \dots, h^{(n-k+1)}(x)).$$

Received by the editors April 17, 2019, and in revised form August 28, 2019.

2010 *Mathematics Subject Classification.* Primary 11B83; Secondary 05A10, 11B65, 12E10, 26C05, 33C10, 43A90.

Key words and phrases. closed-form expression; Bell polynomial of the second kind; falling factorial; special value; identity; equivalent expression; spherical Bessel function; modified spherical Bessel function; Faà di Bruno formula.

The third author was supported by the National Research Foundation of Korea (Grant No. 2018R1D1A1B07041846)

Corresponding author: Bai-Ni Guo.

By (1.1), we can easily deduce that, for $n \geq k \geq 0$,

$$(1.3) \quad \begin{aligned} B_{n,k}(1, 0, \dots, 0) &= B_{n,k} \left(\frac{d}{dx} x, \frac{d^2}{dx^2} x, \dots, \frac{d^{n-k+1}}{dx^{n-k+1}} x \right) \\ &= \binom{0}{n-k} = \begin{cases} 1, & n = k; \\ 0, & n \neq k. \end{cases} \end{aligned}$$

In [10, Theorem 5.1] and [16, Section 3], it was established that the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ satisfy

$$(1.4) \quad B_{n,k}(x, 1, 0, \dots, 0) = \frac{1}{2^{n-k}} \frac{n!}{k!} \binom{k}{n-k} x^{2k-n},$$

where $n \geq k \geq 0$, $\binom{0}{0} = 1$, and $\binom{p}{q} = 0$ for $q > p \geq 0$. Since

$$(1.5) \quad B_{n,k}(abx_1, ab^2x_2, \dots, ab^{n-k+1}x_{n-k+1}) = a^k b^n B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$$

for $n \geq k \geq 0$, see [1, p. 135], we can rearrange (1.4) as

$$\begin{aligned} B_{n,k} \left(\frac{d}{dx} x^2, \frac{d^2}{dx^2} x^2, \dots, \frac{d^{n-k+1}}{dx^{n-k+1}} x^2 \right) &= B_{n,k}(2x, 2, 0, \dots, 0) \\ &= 2^k B_{n,k}(x, 1, 0, \dots, 0) = \frac{n!}{k!} \binom{k}{n-k} (2x)^{2k-n} \end{aligned}$$

for $n \geq k \geq 0$. This means that one can combine the Faà di Bruno formula (1.2) with the formula (1.4) to compute the n th derivative for functions of the type $f(ax^2 + bx + c)$, such as

$$(1.6) \quad \begin{aligned} e^{\pm x^2}, \quad \sin(x^2), \quad \cos(x^2), \quad \ln(1 \pm x^2), \\ (1 \pm x^2)^\alpha, \quad \arcsin x, \quad \arccos x, \quad \arctan x, \end{aligned}$$

and to investigate the generating functions

$$\begin{aligned} \frac{1}{\sqrt{1-2xt+t^2}}, \quad \frac{1}{1-2xt+t^2}, \quad \frac{1}{\sqrt{1-6x+x^2}}, \\ \frac{1-x-\sqrt{x^2-6x+1}}{2x}, \quad \frac{1+x-\sqrt{x^2-6x+1}}{4}, \\ \frac{1-x-\sqrt{1-2x-3x^2}}{2x^2}, \quad e^{2xt-t^2}, \quad \frac{2e^z}{e^{2z}+1} \end{aligned}$$

of the Legendre polynomials, the Chebyshev polynomials of the second kind, central Delannoy numbers, the large and little Schröder numbers, the Motzkin numbers, the Hermite polynomials, the Euler numbers, and the Rodrigues formulas for the Chebyshev polynomials of the first and second kinds. This idea has been carried out and applied in [7, 8, 10, 14, 16] and closely related references therein.

Now it is natural and significant to ask the following question: how to compute the n th derivative of functions of the type $f(x^\alpha)$ for $\alpha \in \mathbb{R}$ and to

apply the results? To answer such a question, by virtue of the Faà di Bruno formula (1.2), we need to calculate

$$\begin{aligned} & B_{n,k} \left(\frac{d}{dx} x^\alpha, \frac{d^2}{dx^2} x^\alpha, \dots, \frac{d^{n-k+1}}{dx^{n-k+1}} x^\alpha \right) \\ &= B_{n,k} (\langle \alpha \rangle_1 x^{\alpha-1}, \langle \alpha \rangle_2 x^{\alpha-2}, \dots, \langle \alpha \rangle_{n-k+1} x^{\alpha-(n-k+1)}) \\ &= x^{k\alpha-n} B_{n,k} (\langle \alpha \rangle_1, \langle \alpha \rangle_2, \dots, \langle \alpha \rangle_{n-k+1}), \end{aligned}$$

where we used identity (1.5) and

$$(1.7) \quad \langle \alpha \rangle_n = \prod_{k=0}^{n-1} (\alpha - k) = \begin{cases} \alpha(\alpha - 1) \cdots (\alpha - n + 1), & n \geq 1; \\ 1, & n = 0 \end{cases}$$

is called the falling factorial.

In recent years, the first author and his coauthors discovered and applied many closed-form expressions of special values for the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ in the papers [6, 7, 8, 9, 15] and closely related references therein.

In this paper, we will establish a pair of closed-form expressions for the Bell polynomials of the second kind $B_{n,k}(\langle \alpha \rangle_1, \langle \alpha \rangle_2, \dots, \langle \alpha \rangle_{n-k+1})$, derive two pairs of identities involving the falling factorials $\langle \alpha \rangle_\ell$, find an equivalent expression of $B_{n,k}(\langle \alpha \rangle_1, \langle \alpha \rangle_2, \dots, \langle \alpha \rangle_{n-k+1})$, and present five closed-form expressions for the (modified) spherical Bessel functions $j_n(z)$, $y_n(z)$, $i_n^{(1)}(z)$, $i_n^{(2)}(z)$, and $k_n(z)$.

2. A PAIR OF CLOSED-FORM EXPRESSIONS FOR BELL POLYNOMIALS

Now we state a pair of closed-form expressions for the Bell polynomials of the second kind $B_{n,k}(\langle \alpha \rangle_1, \langle \alpha \rangle_2, \dots, \langle \alpha \rangle_{n-k+1})$ and its proof.

Theorem 2.1. *For $n \geq k \geq 0$ and $\alpha \in \mathbb{R}$, the Bell polynomials of the second kind $B_{n,k}(\langle \alpha \rangle_1, \langle \alpha \rangle_2, \dots, \langle \alpha \rangle_{n-k+1})$ can be computed by*

$$(2.1) \quad B_{n,k}(\langle \alpha \rangle_1, \langle \alpha \rangle_2, \dots, \langle \alpha \rangle_{n-k+1}) = \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \langle \alpha \ell \rangle_n$$

and

$$(2.2) \quad \sum_{\ell=0}^k \frac{B_{n,\ell}(\langle \alpha \rangle_1, \langle \alpha \rangle_2, \dots, \langle \alpha \rangle_{n-\ell+1})}{(k-\ell)!} = \frac{\langle \alpha k \rangle_n}{k!}.$$

Proof. In [1, p. 133], it is listed that

$$(2.3) \quad \frac{1}{k!} \left(\sum_{m=1}^{\infty} x_m \frac{t^m}{m!} \right)^k = \sum_{n=k}^{\infty} B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) \frac{t^n}{n!}$$

for $k \geq 0$. Taking $x_m = \langle \alpha \rangle_m x^{\alpha-m}$ for $\alpha \in \mathbb{R}$ in (2.3) leads to

$$\begin{aligned} & \sum_{n=k}^{\infty} B_{n,k}(\langle \alpha \rangle_1 x^{\alpha-1}, \langle \alpha \rangle_2 x^{\alpha-2}, \dots, \langle \alpha \rangle_{n-k+1} x^{\alpha-(n-k+1)}) \frac{t^n}{n!} \\ &= \frac{x^{\alpha k}}{k!} \left[\sum_{m=1}^{\infty} \frac{\langle \alpha \rangle_m}{m!} \left(\frac{t}{x} \right)^m \right]^k = \frac{x^{\alpha k}}{k!} \left[\left(1 + \frac{t}{x} \right)^\alpha - 1 \right]^k. \end{aligned}$$

Taking $a = x^\alpha$, $b = \frac{1}{x}$, and $x_i = \langle \alpha \rangle_i$ for $1 \leq i \leq n-k+1$ in the identity (1.5), we obtain

$$\begin{aligned} & B_{n,k}(\langle \alpha \rangle_1 x^{\alpha-1}, \langle \alpha \rangle_2 x^{\alpha-2}, \dots, \langle \alpha \rangle_{n-k+1} x^{\alpha-(n-k+1)}) \\ &= x^{k\alpha-n} B_{n,k}(\langle \alpha \rangle_1, \langle \alpha \rangle_2, \dots, \langle \alpha \rangle_{n-k+1}). \end{aligned}$$

Consequently, it follows that

$$\sum_{n=k}^{\infty} x^{k\alpha-n} B_{n,k}(\langle \alpha \rangle_1, \langle \alpha \rangle_2, \dots, \langle \alpha \rangle_{n-k+1}) \frac{t^n}{n!} = \frac{x^{\alpha k}}{k!} \left[\left(1 + \frac{t}{x} \right)^\alpha - 1 \right]^k$$

which can be rearranged as

$$\begin{aligned} & \sum_{n=k}^{\infty} B_{n,k}(\langle \alpha \rangle_1, \langle \alpha \rangle_2, \dots, \langle \alpha \rangle_{n-k+1}) \frac{s^n}{n!} = \frac{[(1+s)^\alpha - 1]^k}{k!} \\ &= \frac{1}{k!} \sum_{\ell=0}^k (-1)^{k-\ell} \binom{k}{\ell} (1+s)^{\alpha\ell} = \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} (1+s)^{\alpha\ell}, \end{aligned}$$

where $s = \frac{t}{x}$. Differentiating $p \geq k \geq 0$ times with respect to s at both ends of the above equality gives

$$\begin{aligned} & \sum_{n=k}^{\infty} B_{n,k}(\langle \alpha \rangle_1, \langle \alpha \rangle_2, \dots, \langle \alpha \rangle_{n-k+1}) \frac{s^{n-p}}{(n-p)!} \\ &= \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \langle \alpha \ell \rangle_p (1+s)^{\alpha\ell-p}. \end{aligned}$$

Further letting $s \rightarrow 0$ on both sides of the above equality results in

$$B_{p,k}(\langle \alpha \rangle_1, \langle \alpha \rangle_2, \dots, \langle \alpha \rangle_{n-k+1}) = \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \langle \alpha \ell \rangle_p.$$

The formula (2.1) thus follows.

The binomial inversion theorem [1, pp. 143–144] reads

$$(2.4) \quad s_n = \sum_{k=0}^n \binom{n}{k} S_k \quad \text{if and only if} \quad S_n = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} s_k$$

for $n \geq 0$, where $\{s_n, n \geq 0\}$ and $\{S_n, n \geq 0\}$ are sequences of complex numbers. Applying this theorem to (2.1) readily produces (2.2). The proof of Theorem 2.1 is complete. \square

Let

$$(x)_n = \prod_{\ell=0}^{n-1} (x + \ell) = \begin{cases} x(x+1) \cdots (x+n-1), & n \geq 1; \\ 1, & n = 0 \end{cases}$$

denote the rising factorial of $x \in \mathbb{R}$. Since

$$(-x)_n = (-1)^n \langle x \rangle_n \quad \text{and} \quad \langle -x \rangle_n = (-1)^n (x)_n,$$

we can now rewrite Theorem 2.1 in terms of the rising factorial $(x)_n$ as follows.

Corollary 2.1. *For $n \geq k \geq 0$ and $\alpha \in \mathbb{R}$, the Bell polynomials of the second kind $B_{n,k}((\alpha)_1, (\alpha)_2, \dots, (\alpha)_{n-k+1})$ can be computed by*

$$B_{n,k}((\alpha)_1, (\alpha)_2, \dots, (\alpha)_{n-k+1}) = \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} (\alpha \ell)_n$$

and

$$\sum_{\ell=0}^k \frac{B_{n,\ell}((\alpha)_1, (\alpha)_2, \dots, (\alpha)_{n-\ell+1})}{(k-\ell)!} = \frac{(\alpha k)_n}{k!}.$$

3. TWO PAIRS OF IDENTITIES INVOLVING FALLING FACTORIALS

We now recover formula (1.3) from Theorem 2.1 as follows. Replacing α by 1 in (2.1) leads to

$$B_{n,k}(\langle 1 \rangle_1, \langle 1 \rangle_2, \dots, \langle 1 \rangle_{n-k+1}) = \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \langle \ell \rangle_n$$

which is equivalent to

$$\begin{aligned} B_{n,k}(1, 0, \dots, 0) &= \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \sum_{m=0}^n s(n, m) \ell^m \\ &= \sum_{m=0}^n s(n, m) \left[\frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \ell^m \right] \\ &= \sum_{m=0}^n s(n, m) S(m, k) = \binom{0}{n-k}, \end{aligned}$$

where $s(n, k)$ and $S(n, k)$ denote the Stirling numbers of the first and second kinds and we used the formulas

$$\langle x \rangle_n = \sum_{k=0}^n s(n, k) x^k, \quad S(n, k) = \frac{1}{k!} \sum_{\ell=1}^k (-1)^{k-\ell} \binom{k}{\ell} \ell^n,$$

and

$$\sum_{\alpha=0}^n s(n, \alpha) S(\alpha, k) = \binom{0}{n-k}$$

in [1, p. 206] and [17, p. 171, Eq. (12.19)]. Formula (1.3) is thus recovered.

Now we state and prove two pairs of identities involving falling factorials.

Theorem 3.1. *For $n \geq k \geq 0$, we have*

$$(3.1) \quad \sum_{\ell=0}^k (-1)^{k-\ell} \binom{k}{\ell} \langle 2\ell \rangle_n = \frac{n!}{2^{n-2k}} \binom{k}{n-k}$$

and

$$(3.2) \quad \sum_{\ell=0}^k 4^\ell \binom{k}{\ell} \binom{\ell}{n-\ell} = \frac{2^n}{n!} \langle 2k \rangle_n.$$

Proof. Replacing α by 2 in (2.1) and making use of (1.7) and (1.5) in sequence, we derive

$$\begin{aligned} B_{n,k}(\langle 2 \rangle_1, \langle 2 \rangle_2, \dots, \langle 2 \rangle_{n-k+1}) &= B_{n,k}(2, 2, 0, \dots, 0) \\ &= 2^k B_{n,k}(1, 1, 0, \dots, 0) = \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \langle 2\ell \rangle_n. \end{aligned}$$

Comparing this with (1.4) for $x = 1$ yields (3.1).

Applying the binomial inversion theorem recited in (2.4) to (3.1) leads to (3.2). The proof of Theorem 3.1 is complete. \square

Theorem 3.2. *For $n \geq k \geq 0$, we have*

$$(3.3) \quad \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \left\langle \frac{\ell}{2} \right\rangle_n = (-1)^n \frac{k! [2(n-k) - 1]!!}{2^n} \binom{2n-k-1}{2(n-k)}$$

and

$$(3.4) \quad \sum_{\ell=0}^k (-1)^\ell \frac{[2(n-\ell) - 1]!!}{(k-\ell)!} \binom{2n-\ell-1}{2(n-\ell)} = (-1)^n \frac{2^n}{k!} \left\langle \frac{k}{2} \right\rangle_n,$$

where the double factorial of negative odd integers $-(2n+1)$ is defined by

$$(-2n-1)!! = \frac{(-1)^n}{(2n-1)!!} = (-1)^n \frac{2^n n!}{(2n)!}, \quad n \geq 0.$$

Proof. The first paragraph in [14, Theorem 4] reads that the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ satisfy

$$(3.5) \quad \begin{aligned} B_{n,k}(g'(x), g''(x), \dots, g^{(n-k+1)}(x)) \\ = (-1)^{n+k} [2(n-k) - 1]!! \left(\frac{b}{2}\right)^n \binom{2n-k-1}{2(n-k)} \frac{1}{(a+bx)^{n-k/2}}, \end{aligned}$$

where $n \in \mathbb{N}$ and $g(x) = \sqrt{a+bx}$ for $a, b \in \mathbb{R}$ and $b \neq 0$. Taking $a = 0$, $b = 1$, and $x \rightarrow 1$ in (3.5) results in

$$(3.6) \quad B_{n,k} \left(\left\langle \frac{1}{2} \right\rangle_1, \left\langle \frac{1}{2} \right\rangle_2, \dots, \left\langle \frac{1}{2} \right\rangle_{n-k+1} \right) \\ = (-1)^{n+k} [2(n-k) - 1]!! \left(\frac{1}{2} \right)^n \binom{2n-k-1}{2(n-k)}$$

for $n \in \mathbb{N}$. On the other hand, taking $\alpha = \frac{1}{2}$ in (2.1) reduces to

$$B_{n,k} \left(\left\langle \frac{1}{2} \right\rangle_1, \left\langle \frac{1}{2} \right\rangle_2, \dots, \left\langle \frac{1}{2} \right\rangle_{n-k+1} \right) = \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \left\langle \frac{\ell}{2} \right\rangle_n.$$

Identity (3.3) thus follows.

Applying the above mentioned binomial inversion theorem [1, pp. 143–144] again to identity (3.3) gives

$$\sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} (-1)^n \frac{\ell! [2(n-\ell) - 1]!!}{2^n} \binom{2n-\ell-1}{2(n-\ell)} = \left\langle \frac{k}{2} \right\rangle_n$$

which can be reformulated as identity (3.4). The proof of Theorem 3.2 is complete. \square

4. AN EQUIVALENT EXPRESSION

Formula (2.1) in Theorem 2.1 has an equivalent expression.

Theorem 4.1. For $n \geq k \geq 0$ and $\lambda \in \mathbb{R}$, formula (2.1) and

$$(4.1) \quad B_{n,k} \left(1, 1 - \lambda, (1 - \lambda)(1 - 2\lambda), \dots, \prod_{\ell=0}^{n-k} (1 - \ell\lambda) \right) \\ = \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \prod_{q=0}^{n-1} (\ell - q\lambda)$$

are equivalent to each other.

Proof. The closed-form expression (4.1) has been applied in [8, Lemma 2.2], [10, Remark 6.1], [11, Lemma 3], and [12, Lemma 2.6].

By identity (1.5), it is easy to see that

$$B_{n,k} \left(1, 1 - \lambda, (1 - \lambda)(1 - 2\lambda), \dots, \prod_{\ell=0}^{n-k} (1 - \ell\lambda) \right) \\ = B_{n,k} \left(\frac{1}{\lambda}\lambda, \frac{1}{\lambda}\lambda^2 \left(\frac{1}{\lambda} - 1 \right), \frac{1}{\lambda}\lambda^3 \left(\frac{1}{\lambda} - 1 \right) \left(\frac{1}{\lambda} - 2 \right), \right. \\ \left. \dots, \frac{1}{\lambda}\lambda^{n-k+1} \prod_{\ell=1}^{n-k} \left(\frac{1}{\lambda} - \ell \right) \right)$$

$$\begin{aligned}
&= \lambda^n B_{n,k} \left(\frac{1}{\lambda}, \frac{1}{\lambda} \left(\frac{1}{\lambda} - 1 \right), \frac{1}{\lambda} \left(\frac{1}{\lambda} - 1 \right) \left(\frac{1}{\lambda} - 2 \right), \dots, \prod_{\ell=0}^{n-k} \left(\frac{1}{\lambda} - \ell \right) \right) \\
&= \lambda^n B_{n,k} \left(\left\langle \frac{1}{\lambda} \right\rangle_1, \left\langle \frac{1}{\lambda} \right\rangle_2, \left\langle \frac{1}{\lambda} \right\rangle_3, \dots, \left\langle \frac{1}{\lambda} \right\rangle_{n-k+1} \right).
\end{aligned}$$

Substituting this into the closed-form expression (4.1) and simplifying yield

$$\begin{aligned}
B_{n,k} \left(\left\langle \frac{1}{\lambda} \right\rangle_1, \left\langle \frac{1}{\lambda} \right\rangle_2, \left\langle \frac{1}{\lambda} \right\rangle_3, \dots, \left\langle \frac{1}{\lambda} \right\rangle_{n-k+1} \right) \\
= \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \prod_{q=0}^{n-1} \left(\frac{\ell}{\lambda} - q \right).
\end{aligned}$$

Replacing $\frac{1}{\lambda}$ by α in the above equation leads to formula (2.1). Conversely, each equality and every step above are invertible. The proof of Theorem 4.1 is complete. \square

5. CLOSED-FORM EXPRESSIONS FOR SPHERICAL BESSEL FUNCTIONS

In [2, 3, 5, 18, 19] and closely related references therein, the n th derivatives of the functions

$$\begin{aligned}
&\frac{1}{1 - \lambda e^{\mu x}}, \quad \cot x, \quad \tan x, \quad \sec x, \quad \csc x, \\
&\tanh x, \quad \coth x, \quad \operatorname{sech} x, \quad \operatorname{csch} x
\end{aligned}$$

were computed by various approaches. If replacing x by x^α for $\alpha \in \mathbb{R}$ in these functions and those listed in (1.6), we obtain the functions

$$\begin{aligned}
&e^{\pm x^\alpha}, \quad \sin(x^\alpha), \quad \cos(x^\alpha), \quad \ln(1 \pm x^\alpha), \quad (1 \pm x^\alpha)^\beta, \quad \arcsin(x^\alpha), \\
&\arccos(x^\alpha), \quad \arctan(x^\alpha), \quad \frac{1}{1 - \lambda e^{\mu x^\alpha}}, \quad \cot(x^\alpha), \quad \tan(x^\alpha), \\
&\sec(x^\alpha), \quad \csc(x^\alpha), \quad \tanh(x^\alpha), \quad \coth(x^\alpha), \quad \operatorname{sech}(x^\alpha), \quad \operatorname{csch}(x^\alpha).
\end{aligned}$$

The n th derivatives for these functions and others of the type $f(x^\alpha)$ can be alternatively and explicitly calculated by combining the Faà di Bruno formula (1.2), the identity (2.1), and other techniques appeared in the above mentioned references.

In [4, p. 266, Section 10.56], it is listed that

$$\begin{aligned}
\frac{\cos \sqrt{z^2 - 2zt}}{z} &= \frac{\cos z}{z} + \sum_{n=1}^{\infty} \frac{t^n}{n!} j_{n-1}(z), \\
\frac{\sin \sqrt{z^2 - 2zt}}{z} &= \frac{\sin z}{z} + \sum_{n=1}^{\infty} \frac{t^n}{n!} y_{n-1}(z), \\
\frac{\cosh \sqrt{z^2 + 2izt}}{z} &= \frac{\cosh z}{z} + \sum_{n=1}^{\infty} \frac{(it)^n}{n!} i_{n-1}^{(1)}(z),
\end{aligned}$$

$$\frac{\sinh \sqrt{z^2 + 2izt}}{z} = \frac{\sinh z}{z} + \sum_{n=1}^{\infty} \frac{(it)^n}{n!} i_{n-1}^{(2)}(z),$$

$$\frac{\exp(-\sqrt{z^2 + 2izt})}{z} = \frac{e^{-z}}{z} + \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{(-it)^n}{n!} k_{n-1}(z)$$

for $2|t| < |z|$, where $j_n(z)$ and $y_n(z)$ denote the spherical Bessel functions of the first and second kinds and $i_n^{(1)}(z)$, $i_n^{(2)}(z)$, and $k_n(z)$ denote the modified spherical Bessel functions.

Theorem 5.1. For $n \in \mathbb{N}$ and $z \neq 0$, we have

(5.1)

$$j_{n-1}(z) = (-1)^n \sum_{\ell=0}^n (-1)^\ell (2\ell - 1)!! \binom{n + \ell - 1}{2\ell} \cos\left(z + \frac{(n - \ell)\pi}{2}\right) \frac{1}{z^{\ell+1}},$$

(5.2)

$$y_{n-1}(z) = (-1)^n \sum_{\ell=0}^n (-1)^\ell (2\ell - 1)!! \binom{n + \ell - 1}{2\ell} \sin\left(z + \frac{(n - \ell)\pi}{2}\right) \frac{1}{z^{\ell+1}},$$

(5.3)

$$i_{n-1}^{(1)}(z) = \sum_{\ell=0}^n (-1)^\ell (2\ell - 1)!! \binom{n + \ell - 1}{2\ell} \times \left[\frac{1 - (-1)^{n-\ell}}{2} \sinh z + \frac{1 + (-1)^{n-\ell}}{2} \cosh z \right] \frac{1}{z^{\ell+1}},$$

(5.4)

$$i_{n-1}^{(2)}(z) = \sum_{\ell=0}^n (-1)^\ell (2\ell - 1)!! \binom{n + \ell - 1}{2\ell} \times \left[\frac{1 - (-1)^{n-\ell}}{2} \cosh z + \frac{1 + (-1)^{n-\ell}}{2} \sinh z \right] \frac{1}{z^{\ell+1}},$$

(5.5)

$$k_{n-1}(z) = (-1)^n \frac{\pi}{2} e^{-z} \sum_{\ell=0}^n (2\ell - 1)!! \binom{n + \ell - 1}{2\ell} \frac{1}{z^{\ell+1}}.$$

Proof. For $n \in \mathbb{N}$, by virtue of the formula (1.2), the identity (1.5), and the formula (3.6) in sequence, we have

$$j_{n-1}(z) = \frac{1}{z} \lim_{t \rightarrow 0} \frac{d^n \cos \sqrt{z^2 - 2zt}}{dt^n} = \frac{1}{z} \lim_{t \rightarrow 0} \sum_{k=0}^n (\cos u)^{(k)} \times B_{n,k} \left((-2z) \left\langle \frac{1}{2} \right\rangle_1 (z^2 - 2zt)^{-1/2}, (-2z)^2 \left\langle \frac{1}{2} \right\rangle_2 (z^2 - 2zt)^{-3/2}, \dots, \right. \\ \left. (-2z)^{n-k+1} \left\langle \frac{1}{2} \right\rangle_{n-k+1} (z^2 - 2zt)^{1/2-(n-k+1)} \right) \\ = \frac{1}{z} \lim_{t \rightarrow 0} \sum_{k=0}^n \cos \left(u + \frac{k\pi}{2} \right) (-2z)^n (z^2 - 2zt)^{k/2-n}$$

$$\begin{aligned}
& \times B_{n,k} \left(\left\langle \frac{1}{2} \right\rangle_1, \left\langle \frac{1}{2} \right\rangle_2, \dots, \left\langle \frac{1}{2} \right\rangle_{n-k+1} \right) \\
&= \frac{1}{z} \sum_{k=0}^n \cos \left(z + \frac{k\pi}{2} \right) (-2z)^n z^{k-2n} B_{n,k} \left(\left\langle \frac{1}{2} \right\rangle_1, \left\langle \frac{1}{2} \right\rangle_2, \dots, \left\langle \frac{1}{2} \right\rangle_{n-k+1} \right) \\
&= \frac{(-2)^n}{z^{n+1}} \sum_{k=0}^n (-1)^{n+k} [2(n-k)-1]!! \left(\frac{1}{2} \right)^n \binom{2n-k-1}{2(n-k)} \cos \left(z + \frac{k\pi}{2} \right) z^k \\
&= \frac{1}{z^{n+1}} \sum_{k=0}^n (-1)^k [2(n-k)-1]!! \binom{2n-k-1}{2(n-k)} \cos \left(z + \frac{k\pi}{2} \right) z^k,
\end{aligned}$$

where $u = u_z(t) = \sqrt{z^2 - 2zt}$. Formula (5.1) is thus proved.

By the same arguments as above, we can derive the formulas (5.2), (5.3), (5.4), and (5.5) immediately. The proof Theorem 5.1 is complete. \square

Remark 5.1. This paper is a revised version of the electronic preprint [13].

ACKNOWLEDGEMENTS

The authors are thankful to anonymous referees for their careful corrections and valuable comments on the original version of this paper.

REFERENCES

1. L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, Revised and Enlarged Edition, D. Reidel Publishing Co., 1974; available online at <https://doi.org/10.1007/978-94-010-2196-8>.
2. B.-N. Guo and F. Qi, *Explicit formulae for computing Euler polynomials in terms of Stirling numbers of the second kind*, J. Comput. Appl. Math. **272** (2014), 251–257; available online at <https://doi.org/10.1016/j.cam.2014.05.018>.
3. B.-N. Guo and F. Qi, *Some identities and an explicit formula for Bernoulli and Stirling numbers*, J. Comput. Appl. Math. **255** (2014), 568–579; available online at <https://doi.org/10.1016/j.cam.2013.06.020>.
4. F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark (eds.), *NIST Handbook of Mathematical Functions*, Cambridge University Press, New York, 2010; available online at <http://dlmf.nist.gov/>.
5. F. Qi, *Derivatives of tangent function and tangent numbers*, Appl. Math. Comput. **268** (2015), 844–858; available online at <http://dx.doi.org/10.1016/j.amc.2015.06.123>.
6. ———, *Diagonal recurrence relations for the Stirling numbers of the first kind*, Contrib. Discrete Math. **11** (2016), 22–30; available online at <https://doi.org/10.11575/cdm.v11i1.62389>.
7. ———, *Integral representations for multivariate logarithmic polynomials*, J. Comput. Appl. Math. **336** (2018), 54–62; available online at <https://doi.org/10.1016/j.cam.2017.11.047>.
8. F. Qi, V. Čerňanová, X.-T. Shi, and B.-N. Guo, *Some properties of central Delannoy numbers*, J. Comput. Appl. Math. **328** (2018), 101–115; available online at <https://doi.org/10.1016/j.cam.2017.07.013>.
9. F. Qi and B.-N. Guo, *A diagonal recurrence relation for the Stirling numbers of the first kind*, Appl. Anal. Discrete Math. **12** (2018), no. 1, 153–165; available online at <https://doi.org/10.2298/AADM170405004Q>.

10. ———, *Explicit formulas for special values of the Bell polynomials of the second kind and for the Euler numbers and polynomials*, *Mediterr. J. Math.* **14** (2017), no. 3, Article 140, 14 pages; available online at <https://doi.org/10.1007/s00009-017-0939-1>.
11. F. Qi, A.-Q. Liu, and D. Lim, *Explicit expressions related to degenerate Cauchy numbers and their generating function*, In: J. Singh, D. Kumar, H. Dutta, D. Baleanu, and S. Purohit (eds), *International workshop of Mathematical Modelling, Applied Analysis and Computation ICMMAAC 2018: Mathematical Modelling, Applied Analysis and Computation* (Jaipur, India, July 6–8, 2018), *Springer Proceedings in Mathematics & Statistics*, vol. 272, Chapter 2, pp. 41–52, Springer, Singapore, 2019; available online at https://doi.org/10.1007/978-981-13-9608-3_2.
12. F. Qi, D.-W. Niu, and B.-N. Guo, *Some identities for a sequence of unnamed polynomials connected with the Bell polynomials*, *Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Math. RACSAM* **113** (2019), no. 2, 557–567; available online at <https://doi.org/10.1007/s13398-018-0494-z>.
13. F. Qi, D.-W. Niu, D. Lim, and B.-N. Guo, *Explicit formulas and identities on Bell polynomials and falling factorials*, *ResearchGate Preprint* (2018), available online at <https://doi.org/10.13140/RG.2.2.34679.52640>.
14. F. Qi, X.-T. Shi, F.-F. Liu, and D. V. Kruchinin, *Several formulas for special values of the Bell polynomials of the second kind and applications*, *J. Appl. Anal. Comput.* **7** (2017), no. 3, 857–871; available online at <https://doi.org/10.11948/2017054>.
15. F. Qi and J.-L. Zhao, *Some properties of the Bernoulli numbers of the second kind and their generating function*, *Bull. Korean Math. Soc.* **55** (2018), no. 6, 1909–1920; available online at <https://doi.org/10.4134/BKMS.b180039>.
16. F. Qi and M.-M. Zheng, *Explicit expressions for a family of the Bell polynomials and applications*, *Appl. Math. Comput.* **258** (2015), 597–607; available online at <http://dx.doi.org/10.1016/j.amc.2015.02.027>.
17. J. Quaintance and H. W. Gould, *Combinatorial Identities for Stirling Numbers*. The unpublished notes of H. W. Gould. With a foreword by George E. Andrews. World Scientific Publishing Co. Pte. Ltd., Singapore, 2016.
18. A.-M. Xu and G.-D. Cen, *Closed formulas for computing higher-order derivatives of functions involving exponential functions*, *Appl. Math. Comput.* **270** (2015), 136–141; available online at <http://dx.doi.org/10.1016/j.amc.2015.08.051>.
19. A.-M. Xu and Z.-D. Cen, *Some identities involving exponential functions and Stirling numbers and applications*, *J. Comput. Appl. Math.* **260** (2014), 201–207; available online at <http://dx.doi.org/10.1016/j.cam.2013.09.077>.

COLLEGE OF MATHEMATICS, INNER MONGOLIA UNIVERSITY FOR NATIONALITIES,
 TONGLIAO 028043, INNER MONGOLIA, CHINA;
 SCHOOL OF MATHEMATICAL SCIENCES, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN
 300387, CHINA
E-mail address: qifeng618@gmail.com, qifeng618@hotmail.com
URL: <https://qifeng618.wordpress.com>

DEPARTMENT OF SCIENCE, HENAN UNIVERSITY OF ANIMAL HUSBANDRY AND
 ECONOMY, ZHENGZHOU 450046, HENAN, CHINA
E-mail address: nnddww@gmail.com
URL: <https://orcid.org/0000-0003-4033-7911>

DEPARTMENT OF MATHEMATICS EDUCATION, ANDONG NATIONAL UNIVERSITY, ANDONG
 36729, REPUBLIC OF KOREA
E-mail address: dgrim84@gmail.com, dklim@anu.ac.kr
URL: <https://orcid.org/0000-0002-0928-8480>

SCHOOL OF MATHEMATICS AND INFORMATICS, HENAN POLYTECHNIC UNIVERSITY,
JIAOZUO 454010, HENAN, CHINA

E-mail address: bai.ni.guo@gmail.com, bai.ni.guo@hotmail.com

URL: <https://orcid.org/0000-0001-6156-2590>