



## FURTHER GENERALIZATIONS OF THE PARALLELOGRAM LAW

ANTONIO M. OLLER-MARCÉN

**ABSTRACT.** In a recent work of Alessandro Fonda, a generalization of the parallelogram law in any dimension  $N \geq 2$  was given by considering the ratio of the quadratic mean of the measures of the  $(N - 1)$ -dimensional diagonals to the quadratic mean of the measures of the faces of a parallelotope. In this paper, we provide a further generalization considering not only  $(N - 1)$ -dimensional diagonals and faces, but the  $k$ -dimensional ones for every  $1 \leq k \leq N - 1$ .

### 1. INTRODUCTION

If we consider the usual Euclidean space  $(\mathbb{R}^n, \|\cdot\|)$ , the well-known identity

$$(1.1) \quad \|a + b\|^2 + \|a - b\|^2 = 2(\|a\|^2 + \|b\|^2)$$

is called the *parallelogram law*.

This identity can be extended to higher dimensions in several ways. For example, it is straightforward to see that

$$(1.2) \quad \|a+b+c\|^2 + \|a+b-c\|^2 + \|a-b+c\|^2 + \|a-b-c\|^2 = 4(\|a\|^2 + \|b\|^2 + \|c\|^2)$$

with the subsequent analogous identities arising inductively. There are many works devoted to provide generalizations of (1.1) in many different contexts [1, 3, 4].

Note that if we rewrite (1.1) as

$$(1.3) \quad \frac{\|a + b\|^2 + \|a - b\|^2}{2} = 2 \frac{(\|a\|^2 + \|b\|^2 + \|a\|^2 + \|b\|^2)}{4}$$

this means that in any parallelogram, the ratio of the quadratic mean of the lengths of its diagonals to the quadratic mean of the lengths of its sides equals  $\sqrt{2}$ . With this interpretation in mind, Alessandro Fonda [2] recently proved the following interesting generalization.

---

Received by the editors November 4, 2019, and in revised form December 4, 2019.

2000 *Mathematics Subject Classification.* 51M04, 52B05.

*Key words and phrases.* Parallelogram law, parallelotope.

**Theorem 1.1.** *Given linearly independent vectors  $a_1, \dots, a_N \in \mathbb{R}^n$ , it holds that*

$$\begin{aligned} & \sum_{i < j} \left( \left\| (a_i + a_j) \wedge \bigwedge_{k \neq i, j} a_k \right\|^2 + \left\| (a_i - a_j) \wedge \bigwedge_{k \neq i, j} a_k \right\|^2 \right) = \\ & = (N-1) \sum_{k=1}^N 2 \|a_1 \wedge \dots \wedge \widehat{a}_k \wedge \dots \wedge a_N\|^2. \end{aligned}$$

*In other words, for any  $N$ -dimensional parallelotope, the ratio of the quadratic mean of the  $(N-1)$ -dimensional measures of its diagonals to the quadratic mean of the  $(N-1)$ -dimensional measures of its faces is equal to  $\sqrt{2}$ .*

In this work we extend this result to faces of dimension  $k$  for every  $1 \leq k \leq N-1$  and to a suitable definition of the  $k$ -dimensional diagonal of a parallelotope. Then Theorem 1.1 will be a particular case of our result for  $k = N-1$ . Indeed, our result can be stated as follows.

**Theorem 1.2.** *Let us consider an  $N$ -dimensional parallelotope and let  $1 \leq k \leq N-1$ . The ratio of the quadratic mean of the  $k$ -dimensional measures of its  $k$ -dimensional diagonals to the quadratic mean of the  $k$ -dimensional measures of its  $k$ -dimensional faces is equal to  $\sqrt{N-k+1}$ .*

In fact, our generalization follows in line with the work [3] but instead considers the definition of a diagonal face given in [2].

## 2. NOTATION AND PRELIMINARIES

In this section, we introduce some notation and present some basic facts that will be useful in the sequel. Let us consider a parallelotope  $\mathcal{P}$  generated by a family of linearly independent vectors  $\mathcal{B} = \{a_1, a_2, \dots, a_N\} \subseteq \mathbb{R}^n$ . This means that

$$\mathcal{P} = \left\{ \sum_{i=1}^N \alpha_i a_i : \alpha_i \in [0, 1] \right\}.$$

Let us fix  $1 \leq k \leq N-1$ . Given  $k$  different vectors  $\mathcal{S} = \{a_{i_1}, \dots, a_{i_k}\} \subseteq \mathcal{B}$ , we can consider the face generated by them:

$$\mathcal{F}(\mathcal{S}) = \left\{ \sum_{v \in \mathcal{S}} \alpha_v v : \alpha_v \in [0, 1] \right\}.$$

This face can now be translated by one or more of the remaining vectors thus obtaining a face

$$\mathcal{F}^I(\mathcal{S}) = \left\{ \sum_{v \in \mathcal{S}} \alpha_v a_v + \sum_{w \in \mathcal{B} \setminus \mathcal{S}} \alpha_w w \in \mathcal{P} : \alpha_w \in \{0, 1\} \right\},$$

where  $I = (\alpha_v)_{v \notin \mathcal{S}} \in \{0, 1\}^{N-k}$ . Since each choice of a set  $\mathcal{S} \subseteq \mathcal{B}$  and a vector  $I \in \{0, 1\}^{N-k}$  leads to a different face and every face can be obtained in this way, it follows that  $\mathcal{P}$  has exactly  $2^{N-k} \binom{N}{k}$   $k$ -dimensional faces. Moreover, it is clear that all the  $2^{N-k}$  different faces  $\mathcal{F}^I(\mathcal{S})$  are congruent to the set generated by  $\mathcal{S}$ ,  $\mathcal{F}(\mathcal{S})$ .

Now, we focus on the  $k$ -dimensional diagonals which will be defined following the ideas in [2]. Let us consider  $N - k + 1$  different vectors  $\mathcal{T} = \{a_{i_1}, \dots, a_{i_{N-k+1}}\} \subseteq \mathcal{B}$  and let  $\mathcal{T} = \mathcal{T}_1 \cup \mathcal{T}_2$  be a decomposition of  $\mathcal{T}$  into two disjoint sets (either  $\mathcal{T}_1$  or  $\mathcal{T}_2$  could be empty). Then, the following set

$$\mathcal{D}_{\mathcal{T}}(\mathcal{T}_1, \mathcal{T}_2) = \left\{ \alpha \sum_{v \in \mathcal{T}_1} v + (1 - \alpha) \sum_{v \in \mathcal{T}_2} v + \sum_{w \in \mathcal{B} \setminus \mathcal{T}} \alpha_w w : \alpha, \alpha_w \in [0, 1] \right\}$$

is called the  $k$ -dimensional diagonal associated to  $(\mathcal{T}, \mathcal{T}_1, \mathcal{T}_2)$ . Clearly each choice of a set  $\mathcal{T} \subseteq \mathcal{B}$  and a decomposition  $\mathcal{T} = \mathcal{T}_1 \cup \mathcal{T}_2$  allows us to define a diagonal. Since it is clear that  $\mathcal{D}_{\mathcal{T}}(\mathcal{T}_1, \mathcal{T}_2) = \mathcal{D}_{\mathcal{T}}(\mathcal{T}_2, \mathcal{T}_1)$ , it readily follows that  $\mathcal{P}$  has exactly  $2^{N-k} \binom{N}{N-k+1}$  different  $k$ -dimensional diagonals. Moreover, if we define the vector

$$V_{\mathcal{T}}(\mathcal{T}_1, \mathcal{T}_2) = \sum_{v \in \mathcal{T}_1} v - \sum_{v \in \mathcal{T}_2} v,$$

we have that

$$\mathcal{D}_{\mathcal{T}}(\mathcal{T}_1, \mathcal{T}_2) = \left\{ \alpha V_{\mathcal{T}}(\mathcal{T}_1, \mathcal{T}_2) + \sum_{v \in \mathcal{T}_2} v + \sum_{w \in \mathcal{B} \setminus \mathcal{T}} \alpha_w w : \alpha, \alpha_w \in [0, 1] \right\}$$



and consequently, it is clear that the diagonal  $\mathcal{D}_{\mathcal{T}}(\mathcal{T}_1, \mathcal{T}_2)$  is a translation of the set generated by  $\{V_{\mathcal{T}}(\mathcal{T}_1, \mathcal{T}_2), w : w \in \mathcal{B} \setminus \mathcal{T}\}$  and hence it is congruent to it.

**Example.** Let us see how the definition of  $\mathcal{D}(\mathcal{T}_1, \mathcal{T}_2)$  applies in the case of lower dimensions; i.e, if  $N = 2, 3$ .

In the case  $N = 2$ , we only consider  $k = 1$ . If we consider the parallelogram  $\mathcal{P}$  generated by  $\mathcal{B} = \{a_1, a_2\} \subseteq \mathbb{R}^N$ , then clearly  $\mathcal{T} = \mathcal{B}$  (because  $k = 1$ ) and  $\mathcal{P}$  has two different diagonals which are defined by the two possible decompositions  $\mathcal{T} = \{a_1\} \cup \{a_2\}$  and  $\mathcal{T} = \mathcal{T} \cup \emptyset$ . In fact,

$$\begin{aligned} \mathcal{D}_{\mathcal{B}}(\{a_1\}, \{a_2\}) &= \{\alpha a_1 + (1 - \alpha)a_2 : \alpha \in [0, 1]\} \\ &= a_2 + \{\alpha(a_1 - a_2) : \alpha \in [0, 1]\}, \\ \mathcal{D}_{\mathcal{B}}(\mathcal{B}, \emptyset) &= \{\alpha(a_1 + a_2) : \alpha \in [0, 1]\}. \end{aligned}$$

Figure 1 shows how we obtain the two diagonals of the parallelogram. Note that, in this case,  $V_{\mathcal{B}}(\mathcal{B}, \emptyset) = a_1 + a_2$  and  $V_{\mathcal{B}}(\{a_1\}, \{a_2\}) = a_1 - a_2$ .

FIGURE 1. The case  $N = 2$ ,  $k = 1$ .FIGURE 2. The case  $N = 3$ ,  $k = 1$  (left) and  $k = 2$  (right).

Now, if  $N = 3$  and  $k = 1$ , let us consider the parallelepiped  $\mathcal{P}$  generated by  $\mathcal{B} = \{a_1, a_2, a_3\} \subseteq \mathbb{R}^N$ . Again,  $\mathcal{T} = \mathcal{B}$  but in this case there are four different 1-dimensional diagonals which are defined by the decompositions  $\mathcal{T} = \{a_1, a_2\} \cup \{a_3\}$ ,  $\mathcal{T} = \{a_1, a_3\} \cup \{a_2\}$ ,  $\mathcal{T} = \{a_2, a_3\} \cup \{a_1\}$ , and  $\mathcal{T} = \mathcal{T} \cup \emptyset$ . In fact,

$$\begin{aligned} \mathcal{D}_{\mathcal{B}}(\{a_1, a_2\}, \{a_3\}) &= \{\alpha(a_1 + a_2) + (1 - \alpha)a_3 : \alpha \in [0, 1]\} \\ &= a_3 + \{\alpha(a_1 + a_2 - a_3) : \alpha \in [0, 1]\}, \\ \mathcal{D}_{\mathcal{B}}(\{a_1, a_3\}, \{a_2\}) &= \{\alpha(a_1 + a_3) + (1 - \alpha)a_2 : \alpha \in [0, 1]\} \\ &= a_2 + \{\alpha(a_1 - a_2 + a_3) : \alpha \in [0, 1]\}, \\ \mathcal{D}_{\mathcal{B}}(\{a_2, a_3\}, \{a_1\}) &= \{\alpha(a_2 + a_3) + (1 - \alpha)a_1 : \alpha \in [0, 1]\} \\ &= a_1 + \{\alpha(-a_1 + a_2 + a_3) : \alpha \in [0, 1]\}, \\ \mathcal{D}_{\mathcal{B}}(\mathcal{B}, \emptyset) &= \{\alpha(a_1 + a_2 + a_3) : \alpha \in [0, 1]\}. \end{aligned}$$

On the left hand side of Figure 2, we can see the above four 1-dimensional diagonals of  $\mathcal{P}$  (in red, purple, green, and blue, respectively). Note that, in this case,  $V_{\mathcal{B}}(\mathcal{B}, \emptyset) = a_1 + a_2 + a_3$ ,  $V_{\mathcal{B}}(\{a_1, a_2\}, \{a_3\}) = a_1 + a_2 - a_3$ ,  $V_{\mathcal{B}}(\{a_1, a_3\}, \{a_2\}) = a_1 - a_2 + a_3$ , and  $V_{\mathcal{B}}(\{a_2, a_3\}, \{a_1\}) = -a_1 + a_2 + a_3$ .

In the same way, if  $N = 3$  and  $k = 2$ , we could define the six 2-dimensional diagonals of  $\mathcal{P}$ . On the right hand side of Figure 2 we see, for instance,  $\mathcal{D}_{\{a_1, a_3\}}(\{a_1\}, \{a_3\})$  in red and  $\mathcal{D}_{\{a_2, a_3\}}(\{a_2\}, \{a_2\})$  in blue.

### 3. PROOF OF THEOREM 1.2

After introducing the notation and the basic objects involved in this work, we are now ready to prove the main result of the paper.

Let  $\mathcal{P}$  be a parallelotope generated by  $\mathcal{B} = \{a_1, a_2, \dots, a_N\} \subseteq \mathbb{R}^n$ . We first compute the quadratic mean of the  $k$ -dimensional measures of its  $k$ -dimensional faces. We first note that for every  $\mathcal{S} = \{a_{i_1}, \dots, a_{i_k}\} \subseteq \mathcal{B}$ , the  $k$ -dimensional measure of the face  $\mathcal{F}(\mathcal{S})$  is  $\|a_{i_1} \wedge \dots \wedge a_{i_k}\|$ . In the previous section we have seen that  $\mathcal{P}$  has exactly  $2^{N-k} \binom{N}{k}$   $k$ -dimensional faces and moreover, there are exactly  $2^{N-k}$  copies of each face  $\mathcal{F}(\mathcal{S})$ . Consequently, the quadratic mean of the  $k$ -dimensional measures of the  $k$ -dimensional faces of  $\mathcal{P}$  is:

$$(3.1) \quad \sqrt{\frac{2^{N-k} \sum \|a_{i_1} \wedge \dots \wedge a_{i_k}\|^2}{2^{N-k} \binom{N}{k}}}.$$

Now we have to compute the quadratic mean of the  $k$ -dimensional measures of the  $k$ -dimensional diagonals of  $\mathcal{P}$ . First of all, recall that  $\mathcal{P}$  has exactly  $2^{N-k} \binom{N}{N-k+1}$  different  $k$ -dimensional diagonals. Each of them is a translation of the set generated by  $\{V_{\mathcal{T}}(\mathcal{T}_1, \mathcal{T}_2), w : w \in \mathcal{B} \setminus \mathcal{T}\}$  for exactly one choice of  $(\mathcal{T}, \mathcal{T}_1, \mathcal{T}_2)$ . The  $k$ -dimensional measure of this latter set is  $\|V_{\mathcal{T}}(\mathcal{T}_1, \mathcal{T}_2) \wedge \bigwedge_{w \in \mathcal{B} \setminus \mathcal{T}} w\|$ . Consequently, the quadratic mean of the  $k$ -dimensional measures of the  $k$ -dimensional diagonals of  $\mathcal{P}$  is:

$$(3.2) \quad \sqrt{\frac{\sum_{\mathcal{T}, \mathcal{T}_1, \mathcal{T}_2} \left\| V_{\mathcal{T}}(\mathcal{T}_1, \mathcal{T}_2) \wedge \bigwedge_{w \in \mathcal{B} \setminus \mathcal{T}} w \right\|^2}{2^{N-k} \binom{N}{N-k+1}}}.$$

Using the bilinearity of the scalar product and taking into account the definition of  $V_{\mathcal{T}}(\mathcal{T}_1, \mathcal{T}_2)$ , it can be easily seen that when we vary  $(\mathcal{T}, \mathcal{T}_1, \mathcal{T}_2)$ , we get the term  $\|a_{i_1} \wedge \dots \wedge a_{i_k}\|^2$  exactly  $2^{N-k}k$  times for every possible choice of  $\{a_{i_1}, \dots, a_{i_k}\} \subseteq \mathcal{B}$ . This implies that the quadratic mean of the  $k$ -dimensional measures of the  $k$ -dimensional diagonals of  $\mathcal{P}$  (3.2) can be written as:

$$(3.3) \quad \sqrt{\frac{2^{N-k}k \sum \|a_{i_1} \wedge \dots \wedge a_{i_k}\|^2}{2^{N-k} \binom{N}{N-k+1}}}.$$

Finally to obtain Theorem 1.2, it is enough to divide (3.3) by (3.1) to get

$$\sqrt{\frac{k \binom{N}{k}}{\binom{N}{N-k+1}}} = \sqrt{N - k + 1}.$$

#### REFERENCES

1. Ö. Egeciöglü, *Parallelogram-law-type identities*, Linear Algebra Appl. **225** (1995), 1–12.
2. A. Fonda, *A generalization of the parallelogram law to higher dimensions*, Ars Math. Contemp. **16** (2019), no. 2, 411–417.
3. A. Nash, *A generalized parallelogram law*, Amer. Math. Monthly **110** (2003), 52–57.
4. A. J. Penico, and Č. V. Stanojević, *An integral analogue to parallelogram law*, Proc. Amer. Math. Soc. **79** (1980), no. 3, 427–430.

CENTRO UNIVERSITARIO DE LA DEFENSA DE ZARAGOZA  
CTRA. HUESCA S/N, 50090 ZARAGOZA, SPAIN  
*E-mail address:* oller@unizar.es