

**The Effect of Separate-Sex Rearing of Slaughter Quail on  
Serum Corticosterone Level**Usama T. Mahmoud<sup>1\*</sup>, Mootaz A.M. Abdel-Rahman<sup>1</sup>, Madeha H.A. Darwish<sup>1</sup>, Gamal M.M. Mosaad<sup>2</sup><sup>1</sup>Department of Animal Hygiene, Faculty of Veterinary Medicine, Assiut University 71526, Assiut, Egypt<sup>2</sup>Department of Animal and Clinical Nutrition, Faculty of Veterinary Medicine, Assiut University 71526, Assiut, Egypt

(Received 4 June 2012/ Accepted 10 July 2012)

Japanese quails are used widely in biomedical and agricultural research as well as for the production of meat and eggs. Separate-sex rearing is commonly employed in the poultry industry because of differences in growth rates and nutritional requirements of male and female chickens (Mine *et al.*, 2002). The separate-sex rearing of growing birds allows achieving a better homogeneity with respect to live weight and high production traits (Mincheva *et al.*, 2012).

Gender of quails play a role on fattening performance and body weight of quail. Shrivastava *et al.* (1995) reported that the birds grown as separate had a higher live body weight at 5 weeks of age compared to those grown as mixed. Kul *et al.* (2006) suggested that rearing male and female quails separately can be recommended to the producers for a higher live body weight, body weight gain, and carcass weight.

Corticosteroid concentrations in blood have been used as a measure of environmental stress in birds (Edens and Siegel, 1976; Beuving *et al.*, 1989). Corticosterone is of particular interest because it is the primary hormone facilitating stress responses in birds, it is functionally important during maturation and ovulation of ovarian follicles (Etches and Cunningham, 1976). Mature males exhibited higher corticosteroid levels in response to ACTH than mature females (Sheila and Cheryl, 1987)

This study was undertaken to determine the effect of growing males and female Japanese quails separate or mixed together on serum corticosterone level. In order to assess whether this system had a

stressful effect on the bird health or not.

This study was carried out from October 2008 to December 2008 at the Faculty of Veterinary Medicine, Assiut University, Egypt. Forty five quail chicks at age of twenty one days were divided into 3 groups according to sex: Females only (15 birds), males only (15 birds) and mixed group (7 males and 8 females).

Twenty blood sample (5 from each group except in the mixed one where 10 blood samples were taken, 5 from females and 5 from males). Blood samples were collected in test tubes without anticoagulant to determine the corticosterone levels. The tubes were kept at the room temperature for 30 minutes then stored at a refrigerator for 60-90 minutes and then centrifuged at 3000 r.p.m for 10 minutes and the separated serum was transferred to another Epindorf's tube using micropipette. The serum samples were kept at -20 °C until analyses for its corticosterone levels using AssayMax corticosterone ELISA kits (obtained from ASSAYPRO, catalog number EC3001-1) following the procedure outlined by manufacturer. The results were expressed as the mean  $\pm$  SE. All data were analyzed using one way analysis of variances (ANOVA) followed by LSD TEST using SPSS 11.0 statistical software (spss, Inc, Chicago, IL,2001).

The previously mentioned results revealed that, males group had a significantly ( $p < 0.05$ ) higher level of serum corticosterone than females group (19.48 and 13.16 ng/ml, respectively) or mixed one (12.09 and 12.99 ng/ml for females and males, respectively). The results disagreed with Remage *et al.* (2003) who reported that, group housed birds which were separated from their pair mates and housed in a same sex group exhibited a non-significant increase in plasma Corticosterone in response

---

\*Corresponding author: Usama T. Mahmoud

E-mail address: [usamakomsan5@gmail.com](mailto:usamakomsan5@gmail.com)

to separation.

Table 1. Effect of sex separation on serum corticosterone level of quail chicks

Group	Serum corticosterone (ng/ml)
females	13.16 <sup>b</sup> ±1.26
males	19.48 <sup>a</sup> ±1.31
Mixed	
female	12.09 <sup>b</sup> ±2.11
male	12.99 <sup>b</sup> ±1.54

<sup>ab</sup> Means with different superscripts in the same column differ significantly (p<0.05).




Fig. 1. Effect of sex separation on serum corticosterone level of quail chicks .

The results of the present experiments could be explained as a social frustration that has been shown to elicit hypothalamic-pituitary axis (HPA) activation in mammals (Romero *et al.*, 1995; Lyons *et al.*, 2000). It has been established that subordinate male's rats have a reduced serum testosterone and higher corticosterone relative to dominant (Hardy *et al.*, 2002). It Could be concluded that, when quail used for studying the stress the examined group of birds must include both genders. The lowest serum corticosterone level was for the group which contains both sexes (males and females). Therefore, it is advisable to raise both sexes in the same group. The effect of sex separation as stress factor needs further work

## References

Beuving, G., Jones, R.B., Blokhuis, H.J., 1989. Adrenocortical and heterophil / lymphocyte responses to challenge in hens showing short or long tonic immobility reactions. *British Poultry Science* 30, 175-184.

Edens, F.W., Siegel, H.S., 1976. Modification of corticosterone and glucose responses by sympatholytic agents in young chickens during acute heat exposure. *Poultry Science* 55, 1704-1712.

Etches, R.J., Cunningham, F.J., 1976. Effect of pregnenolone, progesterone, deoxycorticosterone or corticosterone on time of ovulation and oviposition in hen. *British Poultry Science* 17, 637-642.

Hardy, M.P., Sottas, C.M., Ge, R.S., McKittrick, C.R., Tamashiro, K.L., McEwen, B.S., Haider, S.G., Markham, C.M., Blanchard, R.J., Blanchard, D.C., Sakai, R.R., 2002. Trends of reproductive hormones in male rats during psychosocial stress: role of glucocorticoid metabolism in behavioral dominance. *Biology of Reproduction* 67, 1750-1755.

Kul, S., Ibrahim, S., Ozge, Y., 2006. Effect of separate and mixed rearing according to sex on fattening performance and carcass characteristics in Japanese quails (*Coturnix coturnix Japonica*). *Arch. Tierz., Dummerstorf* 49, 607-614.

Romero, L.M., Levine, S., Sapolsky, R.M., 1995. Adrenocorticotropin secretagog release-Stimulation by frustration and paradoxically by reward presentation. *Brain Research* 676, 151-156.

Lyons, D.M., Fong, K.D., Schrieken, N., Levine, S., 2000. Frustrative nonreward and pituitary-adrenal activity in squirrel monkeys. *Physiology and Behavior* 71, 559-563.

Mincheva, N., Lalev, M., Oblakova, M., Hristakieva, P., Ivanova, I., 2012. Investigation on the frequency of alleles at the k locus and their effect on the growth of two lines of Plymouth Rock chickens. *Archiva Zootechnica* 15, 69-75.

Mine, O.M., Mochakana, M.E., Mpapho, T., Motlhanka, D.T.M., Kgwatalala, P., 2002. Application of a sex identification technique in juvenile ostriches and its potential application in Botswana. *South African Journal of Animal Science* 32, 160- 163.

Remage, L.H., Regan, E.A., Romeroc, L.M., 2003. Behavioral and adrenocortical responses to mate separation and reunion in the zebra finch. *Hormones and Behavior* 43, 108-114.

Sheila, K.S., Cheryl, F.N., 1978. Effects of Age, Sex, and Ascorbic Acid Ingestion on Chicken Plasma Corticosterone Levels. *Poultry Science* 57, 527-533.

Shrivastava, S.K., Ahuja, S.D., Bandyopadhyay, U.K., Singh, R.P., 1995. Influence of rearing mixed and separate sexes on growth performance and carcass yield of Japanese quail. *Indian Journal of Poultry Science*. 30, 158-160.

SPSS, 2001. Statistical software package for the social sciences. SPSS Inc. United States of America. Cited by <http://www.spss.com>.