




Journal of Advanced Veterinary Research

<https://advetresearch.com/index.php/avr/index>

Evaluation of Some Virulence Factors of Methicillin and Vancomycin Resistant *Staphylococcus epidermidis* Isolated from Cheese and Human Samples

Asmaa G. Mubarak¹, Mona A. El-Zamkan^{2*}¹Department of Zoonoses, Faculty of Veterinary Medicine, South Valley University, Qena 83523, Egypt.²Department of Food Hygiene and Control, Faculty of Veterinary Medicine, South Valley University, Qena 83523, Egypt.**ARTICLE INFO****ABSTRACT****Original Research****Received:**

15 September 2019

Accepted:

25 October 2019

Keywords:Biofilm, Dairy food, Enterotoxin, Food handlers, *S. epidermidis*

Biofilm formation and enterotoxin production represent major virulence factors of *S. epidermidis*. Also, biofilm formation is greatly associated with multidrug resistance. So the objective of the present study was to expand the current knowledge regarding the importance and virulence of methicillin and vancomycin resistant *S. epidermidis* originated from dairy food, food handlers and patients in a hospital, and highlight the possible transmission through foods and food handlers. Biofilm formation was evaluated phenotypically by the tube method and microtiter plate method and genotypically through detection of *icaA* and *icaD* genes, while *S. epidermidis* isolates were investigated for potential enterotoxin production through detection of enterotoxin encoding genes (*sea*, *seb* and *sec*). Among the investigated isolates, phenotypic and genotypic biofilm formation was confirmed in 78.4 and 66.7 % of the isolates, respectively. Regarding enterotoxin encoding genes, it was found that *seb* gene was the only prevailing gene in the three categories of samples with an incidence of 27.5 %. The findings of this study illustrated the prominent role that played by food handlers in transmission of virulent *S. epidermidis* to food and subsequently patients.

*J. Adv. Vet. Res. (2019), 9 (4), 187-196***Introduction**

Coagulase-negative *Staphylococci* (CoNS) constitute a major component of the normal human microfloras and have been evaluated as saprophytes. However, over the last 3 decades, the increased number of human infection cases due to CoNS has been documented representing one of the major nosocomial pathogens, with *S. epidermidis* as the serious significant species (Kloos and Bannerman, 1994; De Silva *et al.*, 2001; Vuong and Otto, 2002; Jean-Baptiste *et al.*, 2011; Becker *et al.*, 2014).

An important virulence factor associated with *S. epidermidis* is its ability to form biofilms and colonize biomaterials (Fey and Olson, 2010). Bacterial ability to form biofilms is of great importance and represents a big challenge not only for the food industry but also for the health facilities, as some strains in their sessile state may endure antimicrobial agents, making the bacterium immensely difficult to be erased (Basanisi *et al.*, 2017). In dairy industry the removal of permanently adhered cells is difficult and imposes the application of strong mechanical force or chemical interruption of the mi-

crobial adhesion through using surfactants, sanitizers or heat; even though, the probability that the irreversibly adhered cells will survive even after pasteurization is high (Schlegelova *et al.*, 2008). On the other hand, from the public health hazard point of view, biofilms are contributed to more than 80% of all infections in humans (Kot *et al.*, 2017); *S. epidermidis* is a major cause of nosocomial and implant-associated infections (Otto, 2009) due to its capacity to adhere to catheters, indwelling medical devices or to colonize different surfaces (Freireira *et al.*, 2012; Prakash *et al.*, 2016). Biofilm formation in *S. epidermidis*, is associated with the *icaABDC* operon presence, which encodes the synthesis of the polysaccharide intercellular adhesion (PIA), which considered an important component in the process of biofilm formation and consists of a b-1,6-linked homoglycan composed of N-acetylglucosamine (Gerke *et al.*, 1998; Mack *et al.*, 2004; Liberto *et al.*, 2009; Oliveira and de Cunha, 2010). The chromosomal intercellular adhesion (*ica*) locus, consisting of the *icaADBC* structural; and the *icaA* and *icaD* genes have been reported to play a fundamental role in biofilm consistence (Darwish and Asfour, 2013).

In the late '50s and early '70s of the twentieth century, a few reports suggested that CoNS might also produce enterotoxins in food poisoning cases (Omori and Kato, 1959; Breckinridge and Bergdoll, 1971). Enterotoxigenic CoNS including *S. epidermidis* were isolated from food and dairy products (Valle

*Corresponding author: Mona A. El-Zamkan
E-mail address: m_zam@vet.svu.edu.eg

et al., 1990; Marín et al., 1992; Rodríguez et al., 1996; Zell et al., 2008; Even et al., 2010; Rall et al., 2010a) and also from cases of human clinical infections (Da Cunha et al., 2007; Atee et al., 2011; Vasconcelos et al., 2011). The presence of sequences homologous to *S. aureus* enterotoxins has been affirmed in the genomes of CoNS strains used in food processing, associated with human infection and from other environments (Weir et al., 2007; Madhusoodanan et al., 2011). Several studies have shown that some CoNS species including *S. epidermidis* possess the genes for staphylococcal enterotoxins (SE) and can produce a functional toxin (Becker et al., 2001; Blaiotta et al., 2004; Oliveira et al., 2011). Staphylococcal enterotoxins can be classified into five main groups (*sea*, *seb*, *sec*, *sed*, and *see*) (Rojas et al., 2012; De Freitas et al., 2013; Pinheiro et al., 2015).

Antibiotic resistance in CoNS, including the clinically-significant species *S. epidermidis*, represents an important health problem worldwide (Cosgrove, 2006). The prevalence of methicillin-resistant *S. epidermidis* has been increased widely and glycopeptides particularly vancomycin, which was the recommended treatment has also exhibited a failure in the treatment of some cases as a result of its resistance (Chakraborty et al., 2011; Morgenstern et al., 2016).

Biofilms formed by *S. epidermidis* decrease the metabolic activity of the bacteria and protect it from phagocytosis by effector cells (Schommer et al., 2011; Spiliopoulou et al., 2012; Hanke et al., 2013), also reduced the diffusion of antibiotics that makes it difficult for antibiotics to affect this type of infection (Arciola et al., 2010). Since, there is an association between biofilm production and the antimicrobial resistance in *S. epidermidis* (Argudin et al., 2015), and many other authors stated that the pathogenicity of *S. epidermidis* has ascribed to the biofilm formation beside enterotoxins production (Irlinger, 2008; Podkowik et al., 2013; Chaves et al., 2018); so, the intention of our study was to expand the knowledge regarding the virulence potential of methicillin and vancomycin resistant *S. epidermidis* isolates from dairy foods, food handlers and patients in the hospital through: a) Detection of the ability of these strains to phenotypic formation of biofilm; b) Molecular detection of biofilm encoding genes (*icaA* and *icaD*). c) Molecular detection of staphylococcal enterotoxins genes *sea*, *seb*, and *sec*.

Materials and methods

S. epidermidis isolates

A total of 51 *S. epidermidis* isolates showed resistance to both methicillin and vancomycin antibiotics (MR/VR- *S. epidermidis*) were obtained in a study conducted by El-Zamkan et al. (2019) at a hospital in Qena City, Egypt, on 210 samples that examined for detection of methicillin and/or vancomycin resistant CoNS including 90 dairy food samples which offered to the patients in the hospital (soft cheese, processed cheese, and yoghurt; 30 samples for each), 60 nasal and hand swabs from food handlers working in the hospital (30 for each) and 60 nasal and diarrheal swabs from patients admitted to the hospital and suffered from diarrhea (30 for each). In this study, the isolates were evaluated for their enterotoxigenicity and ability to form biofilm representing the most important *S. epidermidis* hazardous virulence factors.

Ethical approval is not required in case of food samples while oral consent was obtained from each participant patient.

Isolation and detection of methicillin and vancomycin resistant *S. epidermidis*

Human samples were collected in sterile plastic containers

with sterile saline solution (0.9% NaCl) by using sterile swabs and transported to the laboratory, then one ml of each human sample was inoculated in 9 ml sterile buffered peptone water (Oxoid CM0509, Basingstoke, Hampshire, England), while 10 gm of each food sample was homogenized in 90 ml sterile buffered peptone water (Oxoid, CM0509). Buffered peptone water was incubated overnight at 37 °C. Then, a loopful of each sample was streaked on Mannitol Salt Agar (Merck, Germany) and incubated aerobically at 37 °C for 24 h for growth; pink colonies were considered as CoNS. Identification of CoNS species was done according to Kloos and Schleifer (1975) and Bannerman (2003). Resistance to methicillin and vancomycin was detected phenotypically according to the guidelines of the National Reference Centre for Antimicrobial Susceptibility and internationally recognized standards of the Clinical and Laboratory Standards Institute (CLSI, 2014) on Muller Hinton agar (Merck, Germany) using the diffusion disk method. Later, the isolates that showed phenotypic antibiotic resistance were submitted to PCR to confirm presence of *mecA* gene F (5'- GTA GAA ATG ACT GAA CGT CCG ATA A-3') and R (5'-CCA ATT CCA CAT TGT TTC GGT CTA A-3') (McClure et al., 2006), and *vanA* gene F (5'- GGGAAAACG ACAATTGC -3') and R (5'- GTACAAT-GCGGCCGTTA -3') (Depardieu et al., 2004).

Biofilm Formation tests

Tube method (TM)

Biofilm production was examined by the tube adherence test proposed by Christensen et al. (1982) in which 10 mL of Trypticase soy broth with 1% glucose was inoculated with a loopfull of the overnight culture on nutrient agar individually. Broths were incubated at 37 °C for 24 h. The cultures were decanted and tubes were washed with phosphate buffer saline (pH 7.3). The tubes were dried and stained with 0.1% crystal violet. Excess stain was washed with deionized water. Tubes were dried in an inverted position and observed for biofilm formation. Biofilm Production was considered positive when a visible film lined the wall and bottom of the tube. Ring formation at the liquid interface was not indicative of biofilm formation.

Microtiter plate (MTP) method

The technique was done as described by Stepanović et al. (2000). Each MR/VR- *S. epidermidis* isolate was grown on Tryptic-Soy Agar with 1% glucose at 37 °C overnight, as the growth medium. Then a dilution 1:100 was prepared in Tryptic soy broth (TSB) medium supplemented with glucose (1%) followed by filling three wells of a sterile 96- well flat-bottomed polystyrene tissue culture plate with a lid with 200 µL of aliquots of the diluted culture; negative control wells contained uninoculated sterile broth. The plates were covered and incubated for 24 h at 37 °C. The contents of each well were gently aspirated, and the wells were washed three times with 250 µL of phosphate-buffered saline (PBS). The remaining adhered bacterial cells were fixed with 200 µL of 99% methanol for 15 min., then plates were air-dried after methanol removal. After that wells were stained with 0.2 mL of 0.2% crystal violet (CV) for five min. Then, wells were rinsed twice with distilled water and dried thoroughly. For the quantification of biofilm growth, the adherent cells were resuspended with glacial acetic acid (33%). The OD of the resulting solutions in each well was measured at 570 nm with a microplate reader. Based on the OD of the bacterial film, all strains were classified into the following categories: "OD ≤ ODc: non-adherent, ODc < OD ≤ 2 × ODc: weakly adherent, 2 × ODc < OD ≤ 4 × ODc: moderately adherent, 4 × ODc < OD: strongly adherent". All

tests were carried out in triplicates and the results were averaged. Isolates were considered biofilm-positive when they have an OD570 nm > 0.2. Each isolate was tested in triplicate and the results were averaged.

Molecular detection of enterotoxins and biofilm associated genes

Biofilm encoding genes and enterotoxin genes detected in this study and their amplification conditions are listed in Table 1. Primers were supplied from Metabion (Germany).

DNA extraction

DNA extraction was performed using the QIAamp DNA Mini kit (Qiagen, Germany, GmbH) with modifications from the manufacturer's recommendations. Briefly, 200 µL of the sample suspension was incubated with 10 µL of proteinase K and 200 µL of lysis buffer at 56 °C for 10 min. After incubation, 200 µL of 100% ethanol was added to the lysate. The sample was then washed and centrifuged following the manufacturer's recommendations. Nucleic acid was eluted with 100 µL of elution buffer provided in the kit.

Biofilm production and analysis of icaA/D genes

Primers were utilized in a 25 µL reaction containing 12.5 µL of Emerald Amp Max PCR Master Mix (Takara, Japan), one µL of each primer of 20 pmol concentrations, 4.5 µL of water, and 6 µL of DNA template. The reaction was performed in an Applied Biosystem 2720 thermal cycler.

Enterotoxins detection

Primers of *sea*, *seb*, and *sec* genes were utilized in a 50 µL reaction containing 25 µL of EmeraldAmp Max PCR Master Mix (Takara, Japan), one µL of each primer of 20 pmol concentration, 8 µL of water, and 7 µL of DNA template. The reaction was performed in an Applied Biosystem 2720 thermal cycler.

Analysis of the PCR Products

The products of PCR were separated by electrophoresis on 1.5% agarose gel (Applichem, Germany, GmbH) in 1x TBE buffer at room temperature using gradients of 5V/cm. For gel analysis, 20 µL of the uniplex PCR products and 40 µL of the multiplex PCR products were loaded in each gel slot. Gelpilot 100 bp and 100 bp plus DNA ladders (Qiagen, Germany, GmbH) were used to determine the fragment sizes. The gel was photographed by a gel documentation system (Alpha Innotech, Biometra) and the data was analyzed through computer software.

Statistical analysis

Bacterial count and OD variables, significant factors (P≤0.05) and agglomerative hierarchical clustering (ACH) performed using XLSTAT 2016 software for Microsoft Excel® (Microsoft®, WA, USA).

Results

Of the all 51 methicillin and vancomycin resistant *S. epidermidis* isolates obtained from cheese, food handlers and pa-

Table 1. Primers sequences, target genes, amplicon sizes and cycling conditions.

	Primers sequences	Amplified segment (bp)	Primary denaturation	Amplification (35 cycles)			Final extension	References
				Secondary denaturation	Annealing	Extension		
<i>sea</i>	GGTTATCAATGTGCGGGTGG CGGCACCTTTTTCTCTTCGG	102						
<i>seb</i>	GTATGGTGGTGTAACGTGAGC CCAAATAGTGACGAGTTAGG	164	94°C 5 min.	94°C 30 sec.	57 °C 40 sec.	72°C 40 sec	72°C 10 min.	Mehrotra et al. (2000)
<i>sec</i>	AGATGAAGTAGTTGATGTGTATGG CACACTTTTAGAATCAACCG	451						
<i>icaA</i>	CCT AAC TAA CGA AAG GTA G AAG ATA TAG CGATAA GTG C	1315	94°C 5 min	94°C 30 sec	49°C 1 min.	72°C 1 min.	72°C 10 min	Ciftci et al. (2009)
<i>icaD</i>	AAA CGTAAG AGA GGT GG GGC AAT ATG ATC AAGATA	381	94°C 5 min	94°C 30 sec	49°C 30 sec.	72°C 30 sec.	72°C 10 min	

Table 2. Prevalence of *icaA* and *icaD* genes in *S. epidermidis* isolates and phenotypic biofilm production evaluated by microtitre plate method (MPT) and tube method.

Source of isolates (No.)		Biofilm genes						Biofilm Production									
		<i>icaA</i>		<i>icaD</i>		<i>icaA</i> and <i>icaD</i>		MTP			Tube Method						
		No.	%	No.	%	No.	%	SA ^a		MA ^b		WA ^c		+ve		-ve	
Dairy Food	Cheese (n.=13)	3	23.1	10	76.9	3	23.1	5	38.5	2	15.4	3	23.1	7	53.8	6	46.2
Food Handlers	Nasal Swab (n.= 6)	1	16.7	4	66.7	1	16.7	2	33.3	2	33.3	0	0.0	4	66.7	2	33.3
	Hand Swab (n.= 7)	1	14.3	5	71.4	1	14.3	2	33.3	2	33.3	1	14.3	4	57.1	3	42.9
Total	13	2	15.4	9	69.2	2	15.4	4	30.8	4	30.8	1	7.7	8	61.5	5	38.5
Hospitalized Patients	Nasal Swab (n.= 16)	5	31.3	7	43.8	3	18.8	4	25.0	2	12.5	8	50.0	8	50.0	8	50.0
	Diarrheal Swab (n.= 9)	2	22.2	6	66.7	2	22.2	3	33.3	2	22.2	2	22.2	4	44.4	5	55.6
Total	25	7	28.0	13	52.0	5	20.0	7	28.0	4	16.0	10	40.0	12	48.0	13	52.0
Overall Total	51	12	23.5	32	62.7	10	19.6	16	31.4	10	19.6	14	27.5	27	52.9	24	47.1

a=strong adherent; b= moderate adherent; c= weakly adherent

tients in a hospital in Qena Governorate, 23.5, 62.7 and 19.6% were positive for the presence of *icaA*, *icaD* and both *icaA* & *icaD* genes, respectively (Table 2 and Figs. 1 and 2). Out of the 51 isolates, 27 versus 40 isolates showed biofilm-forming phenotypes using tube and microtiter plate (MTP) methods, respectively; and the later ranged from weak to strong adherent (Table 4). The frequency distribution of genes and biofilm phenotype detected is shown in Fig. 3. There were no statistically significant differences between the presence of *icaA* and/or *icaD* gene and biofilm formation using MTP method ($P > 0.05$), while the accompanying presence of both *icaA* and *icaD* was significantly associated with biofilm phenotype ($P < 0.05$).




Fig. 1. Detection of *icaA* gene in methicillin and vancomycin resistant *S. epidermidis* isolates obtained from cheese and human samples. L: Ladder, +ve: positive control, -ve: negative control.

isolates 27.5% (14/51) were harboring *seb* gene, including 30.8% (4/13) of each dairy food and food handlers' isolates and 24% (6/25) of hospitalized patients, while *sea* gene was found in only one *S. epidermidis* isolate that originated from cheese sample (Fig. 4).




Fig. 2. Detection of *icaD* gene in methicillin and vancomycin resistant *S. epidermidis* isolates obtained from cheese and human samples. L: Ladder, +ve: positive control, -ve: negative control.

Discussion

There is a limited data regarding phenotypic and genotypic biofilm production patterns in *S. epidermidis* isolated from dairy food. Additionally, the incidences of strains that possessed *icaA* and/or *icaD* genes and showed a phenotypic biofilm formation in the current work should be interpreted

According to data showed in Table 3, of all the investigated




Fig. 3. Frequency distribution of biofilm genes in relation to phenotypic biofilm formation (*SA=strong adherent; MA= moderate adherent; WA= weakly adherent; *icaD**=*icaD* gene only; *icaA*= *icaA* gene only).




Fig. 4. Detection of *sea*, *seb* and *sec* genes encoding staphylococcal enterotoxin production in methicillin and vancomycin resistant *S. epidermidis* isolates obtained from cheese and human samples. L: Ladder, +ve: positive control, -ve: negative control

Table 3. Prevalence of enterotoxin genes in *S. epidermidis* isolates

Source of samples (No.)		<i>sea</i>		<i>seb</i>		<i>sec</i>	
		No.	%	No.	%	No.	%
Dairy foods	Cheese (n.=13)	1	7.7	4	30.8	0	0.0
	Nasal swab (n.=6)	0	0.0	2	33.3	0	0.0
Food handlers	Hand swab (n.=7)	0	0.0	2	28.6	0	0.0
	Total	13	0	4	30.8	0	0.0
Hospitalized patients	Nasal swab (n.=16)	0	0.0	2	12.5	0	0.0
	Diarrheal swab (n.=9)	0	0.0	4	44.4	0	0.0
Total	25	0	0.0	6	24.0	0	0.0
Over all total	51	1	1.9	14	27.5	0	0.0

with caution because the study was confined to a certain group of small *S. epidermidis* isolates number, which all shared resistance to both methicillin and vancomycin, which may be the reason for reporting such higher results in cheese than Schlegelová *et al.* (2008); Chajęcka-Wierzchowska *et al.* (2019), and in food handlers than Udo *et al.* (2009), while in patients it was higher than Ninin *et al.* (2006); Gad *et al.* (2009); Saising *et al.* (2012); Ebrahimi *et al.* (2014) and Kord *et al.* (2018). However, in human samples, many other authors like Smith *et al.* (2008) and Piechota *et al.* (2018) recorded higher results as *S. epidermidis* is more abundant in human including patients and healthy persons (Schlegelova *et al.*, 2008; Jaglic *et al.* 2010).

From the presented results, it is clear that *icaD* gene was the most detected biofilm gene in the isolates either separately or in combination with *icaA*. In the same way, de Mello *et al.* (2014) and Chajęcka-Wierzchowska *et al.* (2019) detected *icaD* gene in a higher proportion than *icaA* gene in *S. epidermidis* isolated from cheese and food handlers, respectively. Unlike our results, *icaA* was the most detected gene in *S. epidermidis* isolated by Pinheiro *et al.* (2016) and by Hasanvand *et al.* (2019), while no mentioned difference between the incidence of *icaA* and *icaD* genes in *S. epidermidis* collected from milk samples was found by Abbondio *et al.* (2019).

De Silva *et al.* (2002) described an association between the *ica* operon and quantitative biofilm formation. In the current study, the emphasis is on the importance of biofilm formation in the pathogenesis of MR/VR- *S. epidermidis*, particularly the role that may be played by *icaA* and/or *icaD* genes in the phenotype of the formed biofilm. In the current study, most isolates contained *icaD* gene produced biofilm ranged from weak to strong; while the concomitant presence of both *icaA* and *icaD* was significantly associated with biofilm phenotype ($P < 0.05$) as all strains that co-expressed *icaA* with *icaD* were strong biofilm formers. Similarly, Cafiso *et al.* (2004); Pinheiro *et al.* (2016) and Chajęcka-Wierzchowska *et al.* (2019) found that the presence of *icaA* and *icaD* was significantly associated with the biofilm formation results. Unlike these findings, Ninin *et al.* (2006) could not find such a relation.

Also, it was found that only two isolates obtained from nasal swabs of patients were found to have only *icaA* gene and these isolates associated with weak biofilm phenotype. This genotypic, phenotypic correlation was explained in a study conducted by Gerke *et al.* (1998) where they found that presence of *icaA* alone revealed a weak N-acetylglucosaminyl transferase activity, while its full activity is only achieved when *icaD* is co-transcribed with *icaA*; the PIA consists of a b-1,6-linked homoglycan composed of N-acetylglucosamine, which is a substrate for the glycosyltransferase reaction in PIA biosynthesis.

Results illustrated in this study revealed that, neither *icaA* nor *icaD* could be detected in non-biofilm formers which originated from cheese and food handlers and this disagree with Los *et al.* (2010); Abu Taleb *et al.* (2012); Abbondio *et al.* (2019); Chajęcka-Wierzchowska *et al.* (2019). On the other hand, in this study six isolates obtained from patients as well as isolates

obtained by Bartoszewicz *et al.* (2004); Qin *et al.* (2007); Schommer *et al.* (2011); Darwish and Asfour (2013); Lira *et al.* (2016) and Pinheiro *et al.* (2016) showed the ability to produce biofilm while they were *ica* negative strains; contrarily, Rohde *et al.* (2004) and Salgueiro *et al.* (2017) reported that PCR positivity for *icaA/icaD* genes, can also be found in non-biofilm producers. This conflict between studies could be contributed to many theories a): Biofilm formation is largely a strain-specific trait (Mack *et al.*, 2004); b): Regulation of the *ica* operon appears to be very complex. Production of PIA is certainly subjected to on-off switching and may be involved in *S. epidermidis* phase-variation that might improve bacterial survival and growth under changing environmental conditions in vivo (Ninin *et al.*, 2006).

The microtitre plate method is considered as the golden standard method for biofilm detection as it quantified the formed biofilm in 40/51 *S. epidermidis*, while tube method identified 24/51 isolates as a non-biofilm producer which may be due to the difficulty to differentiate between weak and non-biofilm because the interpretation of the results of tube method varies according to each person observation, also differences in biofilm formation between phenotypic methods may be due to the effect of different culture media, pH, temperature, and osmotic pressure (Deka, 2014).

The high incidences of phenotypic and genotypic biofilm *S. epidermidis* producer in this study as well as other studies performed by De Silva *et al.* (2002); Cafiso *et al.* (2004); Prasad *et al.* (2012); Sahal and Bilkay (2014); Argudín *et al.* (2015) and Chajęcka-Wierzchowska *et al.* (2019) confirmed that biofilm formation and antibiotic resistance are considered to be one of the main virulence factors and are associated with each other and are regarded as important markers that differentiate commensal and pathogenic strains; also strains that are stronger biofilm formers are often multidrug resistant.

Although CoNS safety hazards associated with the presence of methicillin and vancomycin antibiotic resistance and the mutual role that played by food and food handlers in its transmission have been illustrated in a study performed by El-Zamkan *et al.* (2019), it was important to investigate some CoNS isolates (*S. epidermidis*) found in that study for their positivity for enterotoxin genes, which have been reported for its existence and for being frequent causes of food poisoning outbreaks that initiated through consumption dairy food (Rosec *et al.*, 1997; Carmo *et al.*, 2002; Veras *et al.*, 2003; Park *et al.*, 2011).

The molecular method can be considered as the routine analytical method for SE detection. Although, staphylococcal enterotoxin SEA is the most common cause of staphylococcal food poisoning worldwide, but the involvement of other staphylococcal enterotoxins has been also demonstrated (Argudín *et al.*, 2010). The results of this study revealed that *seb* gene was the most prevailing gene (30.8%) in *S. epidermidis* isolates originated from cheese samples followed by *sea* gene (7.7%), while *sec* gene could not be located in these isolates. Results obtained by Even *et al.* (2010) pointed out that 76%

Table 4. Virulence profile of *S. epidermidis* strains isolated from cheese, food handlers and hospitalized patients.

Strain No.	<i>icaA, icaD</i>	<i>Sea, seb, sec</i>	Biofilm production		Origin	
			Tube Method	MTP		
1	<i>S. epidermidis</i>	<i>icaD</i>	-	+ve	Strong	White cheese
2	<i>S. epidermidis</i>	<i>icaA, icaD</i>	<i>seb</i>	+ve	Strong	
3	<i>S. epidermidis</i>	<i>icaD</i>	-	-ve	Weak	
4	<i>S. epidermidis</i>	<i>icaA, icaD</i>	-	+ve	Strong	
5	<i>S. epidermidis</i>	-	<i>seb</i>	-ve	Non-adherent	
6	<i>S. epidermidis</i>	<i>icaA, icaD</i>	<i>seb</i>	+ve	Strong	
7	<i>S. epidermidis</i>	-	-	-ve	Non-adherent	
8	<i>S. epidermidis</i>	<i>icaD</i>	-	-ve	Weak	
9	<i>S. epidermidis</i>	-	-	-ve	Non-adherent	
10	<i>S. epidermidis</i>	<i>icaD</i>	-	+ve	Moderate	
11	<i>S. epidermidis</i>	<i>icaD</i>	<i>sea, seb</i>	+ve	Strong	
12	<i>S. epidermidis</i>	<i>icaD</i>	-	-ve	Weak	
13	<i>S. epidermidis</i>	<i>icaD</i>	-	+ve	Moderate	
14	<i>S. epidermidis</i>	<i>icaA, icaD</i>	-	+ve	Strong	Food Handlers (Nasal swab)
15	<i>S. epidermidis</i>	-	<i>seb</i>	-ve	Non-adherent	
16	<i>S. epidermidis</i>	<i>icaD</i>	<i>seb</i>	+ve	Moderate	
17	<i>S. epidermidis</i>	<i>icaD</i>	-	+ve	Moderate	
18	<i>S. epidermidis</i>	-	-	-ve	Non-adherent	
19	<i>S. epidermidis</i>	<i>icaD</i>	-	+ve	Strong	
20	<i>S. epidermidis</i>	<i>icaD</i>	-	-ve	Moderate	Food Handlers (Hand Swab)
21	<i>S. epidermidis</i>	<i>icaA, icaD</i>	<i>seb</i>	+ve	Strong	
22	<i>S. epidermidis</i>	<i>icaD</i>	-	+ve	Moderate	
23	<i>S. epidermidis</i>	<i>icaD</i>	-	+ve	Weak	
24	<i>S. epidermidis</i>	-	-	-ve	Non-adherent	
25	<i>S. epidermidis</i>	<i>icaD</i>	-	+ve	Strong	
26	<i>S. epidermidis</i>	-	<i>seb</i>	-ve	Non-adherent	
27	<i>S. epidermidis</i>	-	-	+ve	Weak	
28	<i>S. epidermidis</i>	-	-	-ve	Non-adherent	
29	<i>S. epidermidis</i>	<i>icaA, icaD</i>	-	+ve	Strong	
30	<i>S. epidermidis</i>	-	-	-ve	Weak	
31	<i>S. epidermidis</i>	-	-	-ve	Weak	
32	<i>S. epidermidis</i>	<i>icaA, icaD</i>	-	+ve	Strong	
33	<i>S. epidermidis</i>	<i>icaD</i>	-	+ve	Weak	
34	<i>S. epidermidis</i>	<i>icaD</i>	<i>seb</i>	-ve	Strong	
35	<i>S. epidermidis</i>	<i>icaA</i>	-	+ve	Weak	
36	<i>S. epidermidis</i>	-	-	-ve	Non-adherent	
37	<i>S. epidermidis</i>	<i>icaA</i>	-	+ve	Weak	
38	<i>S. epidermidis</i>	<i>icaA, icaD</i>	-	+ve	Strong	
39	<i>S. epidermidis</i>	-	<i>seb</i>	-ve	Weak	
40	<i>S. epidermidis</i>	-	-	-ve	Weak	
41	<i>S. epidermidis</i>	<i>icaD</i>	-	+ve	Moderate	
42	<i>S. epidermidis</i>	<i>icaD</i>	-	-ve	Moderate	
43	<i>S. epidermidis</i>	<i>icaA, icaD</i>	<i>seb</i>	+ve	Strong	Hospitalized patients (Diarrheal Swab)
44	<i>S. epidermidis</i>	<i>icaD</i>	<i>seb</i>	-ve	Strong	
45	<i>S. epidermidis</i>	-	-	-ve	Non-adherent	
46	<i>S. epidermidis</i>	-	<i>seb</i>	+ve	Weak	
47	<i>S. epidermidis</i>	<i>icaD</i>	-	-ve	Moderate	
48	<i>S. epidermidis</i>	-	-	-ve	Non-adherent	
49	<i>S. epidermidis</i>	<i>icaA, icaD</i>	-	+ve	Strong	
50	<i>S. epidermidis</i>	<i>icaD</i>	-	+ve	Weak	
51	<i>S. epidermidis</i>	<i>icaD</i>	<i>seb</i>	-ve	Moderate	

of the strains carrying genes encoding staphylococcal toxin belonged to *S. epidermidis* species isolated from fermented food including cheese, while, [Guimarães et al. \(2013\)](#) found that 63.2% *S. epidermidis* strains isolated from milk showed enterotoxin-encoding genes and *sea* and *seb* were the most dominating, while, [Rodrigues et al. \(2017\)](#) detected enterotoxin genes in *S. epidermidis* isolated from 3 different processing plants of fresh cheese but none of *sea*, *seb* or *sec* was amongst them. Contrarily, [Rall et al. \(2010a\)](#), [Siqueira et al. \(2017\)](#) and [Abbondio et al. \(2019\)](#) reported that none of the

enterotoxin genes (*sea*, *seb*, *sec*) could be detected in *S. epidermidis* isolated from milk.

Also, the results of this study revealed that *seb* gene only could be detected in 30.8 and 24% of *S. epidermidis* isolates obtained from food handlers and patients, respectively. This obtained result comes in parallel to data recorded by [Ahanotu et al. \(2006\)](#) and [Sospedra et al. \(2012\)](#), which proved that *seb* gene is the most common gene in coagulase negative *Staphylococci* identified from foods and food handlers. The gene was detected in closely related result in multidrug resistant *S. epi-*




Fig. 5. Dendrogram showing the relatedness between MR/VR *S. epidermidis* isolates. The dendrogram is based on the presence/absence of *seb* and biofilm-forming genes (*icaA* and *icaD*)

dermidis isolated from food handlers in a hospital as 35% by Francisco and Moreno (2018); also in a study achieved by Udo et al. (2009) 38% of *S. epidermidis* isolates produced different SEs with the majority for *seb* gene. In contrary to this result, a lower incidence of *seb* gene (17.2%) could be detected by Rall et al. (2010b). The enterotoxin production by *S. epidermidis* was also studied by several authors as Rapini et al. (2005) who observed that *S. epidermidis* isolated from food handlers produced *sea* and *sed* in 96.4%. A smaller prevalence of CoNS enterotoxin-encoding genes was observed by Crass and Bergdoll (2007), as 16.5% of CoNS strains obtained from food handlers. While the higher prevalence of *seb* gene producing *S. epidermidis* strains isolated from food handlers was observed by Santos (2003) as 63.2%. On the other hand, Çepoğlu et al., (2010) couldn't detect *seb* gene in *S. epidermidis* isolates from food handlers and detected only *see* gene.

Regarding hospitalized patients, closely related results to ours were obtained by Pinheiro et al. (2015) who recorded that 30% of *S. epidermidis* isolates obtained from patients contained *seb* gene. The production of the classical enterotoxins *sea*, *seb*, and *sec* by clinical isolates of multidrug resistant *S. epidermidis* from hospitalized patients has been also described by Cunha et al. (2007); Barretti et al. (2009); Rojas et al. (2012) and Vasconcelos et al. (2011). The differences between studies may be related to variation in the isolates studied, including the number, nature and geographic origin of the strains.

Clustering the isolates according to its detected virulence factors (The presence or absence of *icaA*, *icaD* and *seb* genes) illuminated the close relationship between samples of human handlers and dairy foods followed by their closer relationship with diarrheal samples, while the nasal swabs of patients were located in a separate arm (Fig. 5). This cluster highlights the mutual role played by food handlers in transmitting virulent *S. epidermidis* to food through cross contamination and the role of these strains in initiating foodborne diseases through food consumption. This was also observed by Doulgeraki et al. (2017) who stated that methicillin resistant *Staphylococci* could produce biofilms by contamination from human handlers rather than from food itself; yet in this study food handlers cannot be considered as standard community isolates as the hospital personnel can act as a reservoir or as a vector

for clinical *S. epidermidis* strains (Milisavljevic et al., 2005).

Conclusion

The data obtained in this study pointed out the linkage between the antibiotic resistance and biofilm formation by *S. epidermidis* and also the relation of *icaA* and *icaD* genes expression and biofilm phenotype. Additionally, the occurrence of staphylococcal enterotoxin genes in isolates of this study highlights their potentiality to cause food poisoning. Hence, Biofilm together with enterotoxin production are the most important virulence factors of *S. epidermidis*. All the above findings are dominated by the conspicuous role played by food handlers in the transmission of the virulent *S. epidermidis* to food or patients.

Acknowledgement

The authors acknowledge the contributions from the members of the hospital in collection of the samples.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Abbondio, M., Fois, I., Longheu, C., Azara, E., Tola, S., 2019. Biofilm production, quorum sensing system and analysis of virulence factors of *Staphylococcus epidermidis* collected from sheep milk samples. *Small Ruminant Research* 174, 83-87.
- Abu Taleb, A.M.F., Mohamed, M.S., Abdel-Latif, R.S., Gouda, M., 2012. The role of *ica* operon and biofilm formation in coagulase negative staphylococcal infection. *Egyptian J. Med. Microbiol* 21, 21-32.
- Ahanotu, E., Alvelo-Ceron, D., Ravita, T., Gaunt, E., 2006. Staphylococcal enterotoxin B as a biological weapon: recognition, management, and surveillance of Staphylococcal enterotoxin. *Appl. Biosaf.* 11, 120-126.
- Arciola, C.R., Baldassarri, L., Montanaro, L., 2010. In catheter infections by *Staphylococcus epidermidis* the intercellular adhesion (*ica*) locus is a molecular marker of virulent slime-producing strains. *J. Biomed. Mater. Res.* 59, 557-562.

- Argudín, M.Á., Mendoza, M.C., Rodicio, M.R., 2010. Food poisoning and *Staphylococcus aureus* enterotoxins. *Toxins* 2, 1751–1773.
- Argudín, M.A., Vanderhaeghen, W., Vandendriessche, S., Vandecandelaere, I., Denis, O., Coenye, T., Butaye, P., 2015. Biofilm formation of *ica* operon-positive *Staphylococcus epidermidis* from different sources. *APMIS Journal of Pathology, Microbiology and Immunology* 123, 1081–1089.
- Ataee, R.A., Mehrabi-Tavana, A., Izadi, M., Hosseini, S.M., Ataee, M.H., 2011. Bacterial meningitis: a new risk factor. *Journal of Medical Sciences Research* 16, 207–210.
- Bannerman, T.L., 2003. *Staphylococcus, Micrococcus*, and other catalase-positive cocci that grow aerobically. In: *Manual of Clinical Microbiology American Society Microbiology*. PR Murray, Baron EJ, Jorgensen JH, Pfaller MA, Tenover FC, Tenover FC (eds), Washington, pp. 384–404.
- Barretti, P., Montelli, A.C., Batalha, J.E., Caramori, J.C., Cunha, M.L., 2009. The role of virulence factors in the outcome of staphylococcal peritonitis in CAPD patients. *BMC Infect. Dis.* 9, 212–219.
- Bartoszewicz, M., Nowicka, J., Janikowska, E., Przondo-Mordarska, A., 2004. Adhesion dependence of coagulase-negative staphylococci to biomaterials and PIA polysaccharide synthesis on *icaADBC* operon presence. *Med. Dosw. Mikrobiol.* 56, 225–230.
- Basanisi, M.G., La Bella, G., Nobili, G., Franconieri, I., La Salandra, G., 2017. Genotyping of methicillin-resistant *Staphylococcus aureus* (MRSA) isolated from milk and dairy products in South Italy. *Food Microbiology* 62, 141–146.
- Becker, K., Heilmann, C., Peters, G., 2014. Coagulase-negative *Staphylococci*. *Clinical Microbiology Reviews* 27, 870–926.
- Becker, K., Keller, B., von Eiff, C., Bruck, M., Lubritz, G., Etienne, J., 2001. Enterotoxigenic potential of *Staphylococcus intermedius*. *Appl. Environ. Microbiol.* 67, 5551–5557.
- Blaïotta, G., Ercolini, D., Pennacchia, C., Fusco V., Casaburi, A., Pepe, O., Villani, F., 2004. PCR detection of staphylococcal enterotoxin genes in *Staphylococcus* spp. strains isolated from meat and dairy products. Evidence for new variants of *seG* and *sel* in *S. aureus* AB-8802. *J. Appl. Microbiol.* 97, 719–30.
- Breckinridge, J.C., Bergdoll, M.S., 1971. Outbreak of food borne gastroenteritis due to a coagulase-negative enterotoxin producing *Staphylococcus*. *The New England Journal of Medicine* 284, 541–543.
- Cafiso, V., Bertuccio, T., Santagati, M., Campanile, F., Amicosante, G., Perilli, M.G., Selan, L., Artini, M., Nicoletti, G., Stefani, S., 2004. Presence of the *ica* operon in clinical isolates of *Staphylococcus epidermidis* and its role in biofilm production. *Clinical Microbiology and Infection* 10, 1081–1088.
- Carmo, L.S., Dias, R.S., Linardi, V.R., Sena, M.J., Santos, D.A., Faria, M.E., Pena, E.C., Jett, M., Heneine, L.G., 2002. Food poisoning due to enterotoxigenic strains of *Staphylococcus* present in Minas cheese and raw milk in Brasil. *Food Microbiology* 19, 9–14.
- Çepoğlu, H., Vatanserver, L., Oral, N.B., 2010. Isolation of *Staphylococci* from Food Handlers and Investigation of Their Enterotoxigenicity and Susceptibility to Some Antibiotics Kafkas Univ. *Vet. Fak Derg Research Article* 16 (Suppl-A): S1–S5.
- Chajęcka-Wierzchowska, W., Zadernowska, A., Gajewska, J., 2019. *S. epidermidis* strains from artisanal cheese made from unpasteurized milk in Poland - Genetic characterization of antimicrobial resistance and virulence determinants. *International Journal of Food Microbiology* 294, 55–59.
- Chakraborty, S., Mahapatra, M., Bal, M., Roy, S., 2011. Isolation and identification of vancomycin resistant *Staphylococcus aureus* from post-operative pus sample. *Al Amen J. Med. Sci.* 2, 152–168.
- Chaves, R.D., Pradella, F., Turatti, M.A., Amaro, E.C., da Silva, A.R., Farias, A.S., Pereira, J.L., Khaneghah, A.M., 2018. Evaluation of *Staphylococcus* spp. in Food and Kitchen Premises of Campinas, Brazil. *Food Control* 84, 463–470.
- Christensen, G.D., Bisno, A.L., Simpson, W.A., Beachey, E.H., 1982. Adherence of slime producing strains of *Staphylococcus epidermidis* to smooth surfaces. *Infectious Immunity* 37, 318–326.
- Ciftci, A., Findik, A., Onuk, A., Savasan, S., 2009. Detection of methicillin resistance and slime factor production of *Staphylococcus aureus* in bovine mastitis. *Brazilian Journal of Microbiology* 40, 254–261.
- Clinical and Laboratory Standards Institute (CLSI), 2014. Performance standards for susceptibility testing, 24 h informational supplement, vol. 34: Wayne, PA: National Committee for Clinical Laboratory Standards.
- Cosgrove, S.E., 2006. The relationship between antimicrobial resistance and patient outcomes: mortality, length of hospital stay, and health care costs. *Clinical Infectious Diseases* 15, S82–S89.
- Crass, B., Bergdoll, M.S., 2007. Involvement of coagulase-negative staphylococci in toxic shock syndrome. *Journal of Clinical Microbiology* 23, 43–45.
- Cunha, M.L.R.S., Calsolari, R.A.O., Araújo-Júnior, J.P., 2007. Detection of enterotoxin and toxic shock syndrome toxin 1 genes in *Staphylococcus*, with emphasis on coagulase-negative staphylococci *Microbiol. Immunol.* 51, 381–390.
- Da Cunha, M., Calsolari, R.A., Júnior, J.P., 2007. Detection of enterotoxin and toxic shock syndrome toxin 1 genes in *Staphylococcus*, with emphasis on coagulase-negative *Staphylococci*. *Microbiology and Immunology* 51, 381–390.
- Darwish, S.F., Asfour, H.A., 2013. Investigation of biofilm forming ability in *Staphylococci* causing bovine mastitis using phenotypic and genotypic assays. *The Scientific World Journal* 2013, Article ID 378492, 9 pages.
- De Freitas, G.F., Nóbrega, D.B., Richini-Pereira, V.B., Marson, P.M., de Figueiredo, J.C., Langoni, H., 2013. Enterotoxin genes in coagulase-negative and coagulase-positive staphylococci isolated from bovine milk. *J. Dairy Sci.* 96, 2866–2872.
- de Mello, J.F., da Rocha, L.B., Lopes, L.S., Frazzton, J., da Costa, M., 2014. Sanitary quality, occurrence and identification of *Staphylococcus* spp. in food services. *Braz. J. Microbiol.* 45, 211–216.
- De Silva, G.D.I., Justice, A., Wilkinson, A.R., Buttery, J., Herbert, M., Day, N.P.J., Peacock, S.J., 2001. Genetic population structure of coagulase-negative *Staphylococci* associated with carriage and disease in preterm infants. *Clin. Infect. Dis.* 33, 1520–1528.
- De Silva, G.D.I., Kantzanou, M., Justice, A., Massey, R.C., Wilkinson, A.R., Day, N.P.J., Peacock, S.J., 2002. The *ica*-operon and biofilm production in coagulase-negative *Staphylococci* associated with carriage and disease in a neonatal intensive care unit. *Journal of Clinical Microbiology* 40, 382–388.
- Deka, N., 2014. Comparison of tissue culture plate method, tube method and Congo red agar method for the detection of biofilm formation by coagulase negative *Staphylococcus* isolated from non-clinical isolates. *Int. J. Curr. Microbiol. App. Sci.* 3, 810–815.
- Depardieu, F., Perichon, B., Courvalin, P., 2004. Detection of the van Alphabet and Identification of Enterococci and *Staphylococci* at the Species Level by Multiplex PCR. *J. Clin. Microbiol.* 42, 5857–5860.
- Doulgeraki, A.I., DiCiccio, P., Ianieri, A., Nychas, G.J.E., 2017. Methicillin resistant food-related *Staphylococcus aureus*: a review of current knowledge and biofilm formation for future studies and applications. *Res. Microbiol.* 168, 1–15.
- Ebrahimi, A., Ghasemi, M., Ghasemi, B., 2014. Some virulence factors of staphylococci isolated from wound and skin infections in Shahrekord, IR Iran. *Jundishapur J. Microbiol.* 7, e9225.
- El-Zamkan, M.A., Mubarak, A.G., Ali, A.O., 2019. Prevalence and phylogenetic relationship among methicillin and vancomycin resistant *Staphylococci* isolated from hospital's dairy food, food handlers, and patients. *Journal of Advanced Veterinary and Animal Research.* 6, 463–471.
- Even, S., Leroy, S., Chalier, C., Zakour, N.B., Chacornac, J., Lebert, I., Jamet, E., Desmonts, M., Coton, E., Pochets, S., Donnio, P., Gaudier, M., Talon, R., Le Loir, Y., 2010. Low occurrence of safety hazards in coagulase negative *Staphylococci* isolated from fermented foodstuffs. *Int. J. Food Microbiol.* 139, 87–95.
- Ferreira, F.A., Souza, R.R., Bonelli, R.R., Américo, M.A., Fracalanza, S.E.L., Figueiredo, A.M.S., 2012. Comparison of invitro and in vivo systems to study *ica*-independent *Staphylococcus aureus* biofilms. *J. Microbiol. Meth.* 88, 393–398.
- Fey P.D., Olson, M.E., 2010. Current concepts in biofilm formation of *Staphylococcus epidermidis*. *Future Microbiology* 5, 917–933.
- Francisco, J.J., Moreno, B., 2018. Importance of food handlers as a source of enterotoxigenic coagulase negative *Staphylococci*. *Zentralbl Bakteriell Mikrobiol Hygiene* 181, 464–474.
- Gad, G.F., El-Feky, M.A., El-Rehewy, M.S., Hassan, M.A., Abolella, H., El-Baky, R.M., 2009. Detection of *icaA*, *icaD* genes and biofilm production by *Staphylococcus aureus* and *Staphylococcus epidermidis* isolated from urinary tract catheterized patients. *J. Infect. Dev. Ctries.* 3, 342–351.
- Gerke, C., Kraft, A., Süßmuth, A., Schweitzer, O., Götz, F., 1998. Characterization of the N-Acetylglucosaminyltransferase activity

- involved in the biosynthesis of the *Staphylococcus epidermidis* Polysaccharide Intercellular Adhesin. *J. Biol. Chem.* 273, 18586–18593.
- Guimarães, F. F., Nóbrega, D.B., Richini-Pereira, V.B., Marson, P.M., Pan-toja, J.C.F., Langoni, H., 2013. Enterotoxin genes in coagulase-negative and coagulase-positive *Staphylococci* isolated from bovine milk. *J. Dairy Sci.* 96, 2866–2872.
- Hanke, M.L., Heim, C.E., Angle, A., Sanderson S.D., Kielian T., 2013. Targeting macrophage activation for the prevention and treatment of *Staphylococcus aureus* biofilm infections. *J. Immunol.* 190, 2159–2168.
- Hasanvand, H., Teymouri, F., Ohadi, E., Azadegan, A., Kalani, B.S., 2019. Biofilm Formation in *Staphylococcus epidermidis* Isolated from Hospitalized Patients. *Arch. Clin. Infect. Dis.* 14, e64496.
- Irlinger, F., 2008. Safety assessment of dairy microorganisms: Coagulase-negative *Staphylococci*. *Int. J. of Food Microbiol.* 126, 302–310.
- Jaglic, Z., Michu, E., Holasova, M., Vlkova, H., Babak, V., Kolar, M., Bardon, J., Schlegelova, J., 2010. Epidemiology and characterization of *Staphylococcus epidermidis* isolates from humans, raw bovine milk and a dairy plant. *Epidemiol. Infect.* 138, 772–782.
- Jean-Baptiste, N., Benjamin, D., Cohen-Wolkowicz, M., Fowler, V.J., Laughon, M., Clark, R.H., Smith, P.B., 2011. Coagulase-negative *Staphylococcal* infections in the neonatal intensive care unit. *Infect. Control. Hosp. Epidemiol.* 32, 679–686.
- Kloos, W.E., Bannerman, T.L., 1994. Update on clinical significance of coagulase-negative *Staphylococci*. *Clin. Microbiol. Rev.* 7, 117–140.
- Kloos, W.E., Schleifer, K.H., 1975. Simplified scheme for routine identification of human *Staphylococcus* species. *J. Clin. Microbiol.* 1, 82–88.
- Kord, M., Ardebili, A., Jamal, M., Jahanbakhsh, R., Behnampour, N., Ghaemi, E.A., 2018. Evaluation of Biofilm Formation and Presence of Ica Genes in *Staphylococcus epidermidis* Clinical Isolates. *Osong Public Health Res. Perspect.* 9, 160–166.
- Kot, B., Wierzchowska, K., Gruzewska, A., Lohinau, D., 2017. The effects of selected phytochemicals on biofilm formed by five methicillin-resistant *Staphylococcus aureus*. *Natural Product Research* 32, 1299–1302.
- Liberto, M.C., Matera, G., Quirino, A., 2009. Phenotypic and genotypic evaluation of slime production by conventional and molecular microbiological techniques. *Microbiol. Res.* 164, 522–528.
- Lira, M.C., Givisiez, P.E.N., de Sousa, F.G.C., Magnani, M., de Souza E.L., Spricigo, D.A., Gebreyes, W.A., de Oliveira, J.P., 2016. Biofilm-forming and antimicrobial resistance traits of *Staphylococci* isolated from goat dairy plants. *J. Infect. Dev. Ctries.* 10, 932–938.
- Los, R., Sawicki, R., Juda, M., 2010. A comparative analysis of phenotypic and genotypic methods for the determination of the biofilm-forming abilities of *Staphylococcus epidermidis*. *FEMS Microbiol. Lett.* 10, 97–103.
- Mack, D., Becker, P., Chatterjee, I., Dobinsky, S., Knobloch, J.K., Peters, G., Rohde, H., Herrmann, M., 2004. Mechanisms of biofilm formation in *Staphylococcus epidermidis* and *Staphylococcus aureus*: functional molecules, regulatory circuits, and adaptive responses. *Int. J. Med. Microbiol.* 294, 203–212.
- Madhusoodanan, J., Seo, K.S., Remortel, B., Park, J.Y., Hwang, S.Y., Fox, L.K., Park, Y.H., Deobald, C.F., Wang, D., Liu, S., Daugherty, S.C., Gill, A.L., Bohach, G.A., Gill, S.R., 2011. An enterotoxin-bearing pathogenicity island in *Staphylococcus epidermidis*. *J. Bacteriol.* 193, 1854–1862.
- Marín, M.E., de la Rosa, M.C., Cornejo, I., 1992. Enterotoxigenicity of *Staphylococcus* strains isolated from Spanish dry-cured hams. *Appl. Environ. Microbiol.* 58, 1067–1069.
- McClure, J.A., Conly, J.M., Lau, V., Elsayed, S., Louie, T., Hutchins, W., Zhang, K., 2006. Novel multiplex PCR assay for detection of the staphylococcal virulence marker Panton-Valentine leukocidin genes and simultaneous discrimination of methicillin-susceptible from resistant *Staphylococci*. *J. Clin. Microbiol.* 44, 1141–1146.
- Mehrotra, M., Wang, G., Johnson, W.M., 2000. Multiplex PCR for Detection of Genes for *Staphylococcus aureus* Enterotoxins, Exfoliative Toxins, Toxic Shock Syndrome Toxin 1, and Methicillin Resistance. *Journal of Clinical Microbiology* 38, 214–219.
- Milislavljevic, V., Wu, F., Cimmotti, J., Haas, J., Della-Latta, P., Larson, E., Saiman, L., 2005. Genetic relatedness of *Staphylococcus epidermidis* from infected infants and staff in the neonatal intensive care unit. *Am. J. Infect. Control* 33, 341–347.
- Morgenstern, M., Erichsen, C., von Ruden, C., Metsemakers, W.J., Kates, S.L., Moriarty, T.F., 2016. Staphylococcal orthopaedic device-related infections in older patients. *Injury* 47, 1427–1434.
- Ninin, E., Caroff, N., Espaze, E., 2006. Assessment of *ica* operon carriage and biofilm production in *Staphylococcus epidermidis* isolates causing bacteraemia in bone marrow transplant recipients. *Clin. Microbiol. Infect.* 12, 446–452.
- Oliveira, A., de Cunha, M.L., 2010. Comparison of methods for the detection of biofilm production in coagulase-negative staphylococci. *BMC Res. Notes* 3, 260–264.
- Oliveira, A.M., Padovani, C., Miya, N., Sant'Ana, A.S., Pereira, J.L., 2011. Incidence of enterotoxin D producing *Staphylococcus* spp in Brazilian cow's raw milk and its relation with coagulase and thermonuclease enzymes. *Foodborne Pathog. Dis.* 8, 159–163.
- Omori, G., Kato, Y., 1959. A staphylococcal food-poisoning caused by a coagulase negative strain. *Bilken's Journal* 2, 92.
- Otto, M., 2009. *Staphylococcus epidermidis* –the 'accidental' pathogen. *Nat. Rev. Microbiol.* 7, 555–567.
- Park, J.Y., Fox, L.K., Seo, K.S., McGuire, M.A., Park, Y.H., Rurangirwa, F.R., Sischo, W.M., Bohach, G.A., 2011. Detection of classical and newly described staphylococcal superantigen genes in coagulase-negative staphylococci isolated from bovine intramammary infections. *Vet. Microbiol.* 147, 149–154.
- Piechota, M., Kot, B., Frankowska-Maciejewska, A., Woźniak-Kosek, A., 2018. Biofilm Formation by Methicillin-Resistant and Methicillin-Sensitive *Staphylococci* Strains from Hospitalized Patients in Poland. *BioMed. Res. Int.* Volume 2018, Article ID 4657396, 7 pages.
- Pinheiro, L., Brito, C.I., de Oliveira, A., Martins, P.Y.F., Pereira, V.C., da-Cunha, M.R., 2015. *Staphylococcus epidermidis* and *Staphylococcus haemolyticus*: Molecular Detection of Cytotoxin and Enterotoxin Genes. *Toxins (Basel)* 7, 3688–3699.
- Pinheiro, L., Brito, C.I., de Oliveira, A., Pereira, V.C., da Cunha, M.L.R.S., 2016. *Staphylococcus epidermidis* and *Staphylococcus haemolyticus*: detection of biofilm genes and biofilm formation in blood culture isolates from patients in a Brazilian teaching hospital. *Diagn. Microbiol. Infect. Dis.* 86, 11–14.
- Podkowik, M., Park, J.Y., Seo, K.S., Bystron, J., Bania, J., 2013. Enterotoxigenic potential of coagulase-negative *Staphylococci*. *Int. J. Food Microbiol.* 163, 34–40.
- Prakash, P.H., Rajan, V., Gopal, S., 2016. Predominance of SCCmec type IV and V among biofilm producing device-associated *Staphylococcus aureus* strains isolated from tertiary care hospitals in Mysuru, India. *Enferm. Infec. Microbiol. Clin.* 35, 229–235.
- Prasad, S., Nayak, N., Satpathy, G., Nag, H.L., Venkatesh, P., Ramakrishnan, S., Ghose, Supriyo, Nag, T.C., 2012. Molecular & phenotypic characterization of *Staphylococcus epidermidis* in implant related infections. *Indian J. Med. Res.* 136, 483–490.
- Qin, Z., Yang, X., Yang, L., Jiang, J., Ou, Y., Molin, S., Qu, D., 2007. Formation and properties of in vitro biofilms of *ica*-negative *Staphylococcus epidermidis* clinical isolates. *J. Med. Microbiol.* 56, 83–93.
- Rall, V.L., Sforzin, J.M., de Deus, M.F., de Sousa, D.C., Camargo, C.H., Godinho, N.C., Galindo, L.A., Soares, T.C., Araújo Júnior, J.P., 2010a. Polymerase chain reaction detection of enterotoxins genes in coagulase-negative *Staphylococci* isolated from Brazilian Minas cheese. *Foodborne Pathog. Dis.* 7, 1121–1123.
- Rall, V.L.M.I., Sforzin, J.M.I., Augustini, V.C.M.I., Watanabe, M.T.I., Fernandes, A.I., Rall, R., Silva, M.G., Araújo, J.P. 2010b. Detection of enterotoxin genes of *Staphylococcus* sp. isolated from nasal cavities and hands of food handlers. *Braz. J. Microbiol.* 41, 59–65.
- Rapini, L.S., Cerqueira, M.M., Carmo, L.S., Veras, J.F., Souza, M.R., 2005. Presença de *Staphylococcus* spp. produtores de enterotoxinas e da toxina da síndrome do choque tóxico em manipuladores de queijo de cabra. *Arq. Bras. Med. Vet. Zootec* 57, 825–829.
- Rodrigues, M. X., Silva, N.C.C., Trevilin, J.H.T., Cruzado, M.M.B., Mui, T.S., Duarte, F.R.S., Castillo, C.J.C., Canniatti-Brazaca, S.G., Porto, E., 2017. Molecular characterization and antibiotic resistance of *Staphylococcus* spp. isolated from cheese processing plants. *J. Dairy Sci.* 100, 5167–5175.
- Rodríguez, M., Núñez, F., Córdoba, J.J., Bermúdez, E., Asensio, M.A., 1996. Gram-positive, catalase-positive cocci from dry cured Iberian ham and their enterotoxigenic potential. *Appl. Environ. Microbiol.* 6, 1897–1902.
- Rohde, H., Kalitzky, M., Kroger, N., Scherpe, S., Horstkotte, M.A.,

- Knobloch, J.K.M., Zander, A.R., Mack, D., 2004. Detection of virulence-associated genes not useful for discriminating between invasive and commensal *Staphylococcus epidermidis* strains from a bone marrow transplant unit. *J. Clin. Microbiol.* 42, 5614-5619.
- Rojas, M.B., Antonelli, C.M., Pereira, E.P.L., Cunha, M.L., 2012. Detection of enterotoxin a in coagulase-negative staphylococci isolated from nutrition students. *Archives of Clinical Microbiology* 3, 6.
- Rosec, J.P., Guiraud, J.P., Dalet, C., Richard, N., 1997. Enterotoxin production by *Staphylococci* isolated from foods in France. *Int. J. Food Microbiol.* 35, 213-221.
- Sahal, G., Bilkay, I.S., 2014. Multi drug resistance in strong biofilm forming clinical isolates of *Staphylococcus epidermidis*. *Braz. J. Microbiol.* 45, 539-544.
- Saising, J., Singdam, S., Ongsakul, M., Voravuthikunchai, S.P., 2012. Lipase, protease, and biofilm as the major virulence factors in *Staphylococci* isolated from acne lesions. *Biosci. Trends.* 6, 160-164.
- Salgueiro, V.C., Iorio, N.L., Ferreira, M.C., Chamon, R., dos Santos, K.R., 2017. Methicillin resistance and virulence genes in invasive and nasal *Staphylococcus epidermidis* isolates from neonates. *BMC Microbiol.* 17, 15-25.
- Santos, D.A., 2003. O papel do manipulador de alimentos em surtos de intoxicação alimentar causados por espécies de *Staphylococcus* ocorridos em quatro cidades do Estado de Minas Gerais, Brasil. 85f. Dissertação (Mestrado em Ciências Biológicas) - Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte.
- Schlegelova, J., Babak, V., Holasova, M., Dendis, M., 2008. The biofilm positive *Staphylococcus epidermidis* isolates in raw materials, foodstuffs and on contact surfaces in processing plants. *Folia Microbiol.* 53, 500-504.
- Schommer, N.N., Christner, M., Hentschke, M., Ruckdeschel, K., Aepfelbacher, M., Rohde, H., 2011. *Staphylococcus epidermidis* uses distinct mechanisms of biofilm formation to interfere with phagocytosis and activation of mouse macrophage-like cells 774A.1. *Infect. Immun.* 79, 2267-2276.
- Siqueira, A. K., Salerno, T., Lara, G.H.B., Condas, L.A.Z., Pereira, V.C., Riboli, D.F.M., LISTONI, F.J.P., Da Silva, A.V., Leite, D.S., Da Cunha, M.L.R.S., Ribeiro, M.G., 2017. Enterotoxin Genes, Multidrug Resistance, And Molecular Typing of *Staphylococcus* spp. Isolated from Organic Bovine Milk. *Braz. J. Vet. Res. Anim. Sci.* 54, 81-87.
- Smith, K., Ramage, G., Lappin, D., Lang, S., 2008. Biofilm formation by Scottish clinical isolates of *Staphylococcus* species. *J. Med. Microbiol.* 57, 1018-1023.
- Sospedra, I., Marín, R., Mañes, J., Soriano, J.M., 2012. Rapid whole protein quantification of staphylococcal enterotoxin B by liquid chromatography. *Food. Chem.* 133, 163-166.
- Spiliopoulou, A.I., Kolonitsiou, F., Krevvata, M.I., Leontsinidis, M., Wilkinson, T.S., Mack, D., 2012. Bacterial adhesion, intracellular survival and cytokine induction upon stimulation of mononuclear cells with planktonic or biofilm phase *Staphylococcus epidermidis*. *FEMS Microbiol. Lett.* 330, 56-65.
- Stepanović, S., Vuković, D., Dakic, I., Savić, B., Vlahović, M.S., 2000. A modified microtiter-plate test for quantification of staphylococcal biofilm formation. *Journal of Microbiological Methods* 40, 175-179.
- Udo, E.E., Al-Bustan, M.A., Jacob, L.E., Chugh, T.D., 2009. Enterotoxin production by coagulase negative *Staphylococci* in restaurant workers from Kuwait City may be a potential cause of food poisoning. *Journal of Medical Microbiology* 48, 819-823.
- Valle, J., Gomez-Lucia, E., Piriz, S., Goyache, J., Orden, J.A., Vadillo, S., 1990. Enterotoxin production by *Staphylococci* isolated from healthy goats. *Appl. Environ. Microbiol.* 56, 1323-1326
- Vasconcelos, N.G., Pereira, V.C., Araújo Júnior, J.P., Da Cunha Mde, L., 2011. Molecular detection of enterotoxins E, G, H and I in *Staphylococcus aureus* and coagulase negative *Staphylococci* isolated from clinical samples of newborns in Brazil. *J. Appl. Microbiol.* 111, 749-762.
- Veras, J.F., Santos, D.A., Carmo, L.S., 2003. Levantamento de surtos de toxinfecção alimentar envolvendo leite e produtos derivados no estado de Minas Gerais, Brasil. *Higiene Alimentar* 17, 218-222.
- Vuong, C., Otto, M., 2002. *Staphylococcus epidermidis* infections. *Microbes Infect.* 4, 481-489.
- Weir, D., Jonem, C., Ammerman, L., Dybdahl, K., Tomlinson, S., 2007. Report of a strain of *Staphylococcus caprae* with the genes for enterotoxin A and enterotoxin-like toxin type P. *J. Clin. Microbiol.* 45, 3476-3477.
- Zell, C., Resch, M., Rosenstein, R., Albrecht, T., Hertel, C., Götz, F., 2008. Characterization of toxin production of coagulase-negative *Staphylococci* isolated from food and starter cultures. *Int. J. Food Microbiol.* 127, 246-251.