



## Occurrence of Quadruplets in Nelore Cattle: A Case Report

Gregório Miguel Ferreira de Camargo<sup>1\*</sup>, Lucas Henrique Brito Sales<sup>2</sup>, Raphael Bernal Costa<sup>1</sup>

<sup>1</sup>Avenida Adhemar de Barros, n 500, Ondina, Salvador-BA, CEP:40170-110, Escola de Medicina Veterinária e Zootecnia, Universidade Federal da Bahia (UFBA), Brazil.

<sup>2</sup>CRV Lagoa da Serra, Sertãozinho-SP, Brazil

### ARTICLE INFO

#### Case Report

#### Received:

09 June 2020

#### Accepted:

02 July 2020

#### Keywords:

Multiple birth,  
*Bos taurus indicus*

### ABSTRACT

Cattle are monotocous species and cow usually deliver one calf per birth. The present report has the aim to notify the birth occurrence of quadruplets of a Nelore cow: first time in scientific literature for this breed. The report presents negative and positive consequences of event.

*J. Adv. Vet. Res. (2020), 10 (3), 177-178*

### Introduction

Cattle are monotocous species, with one or, occasionally, two newborns. There are few reports in literature for quadruplets in cattle (Basrur and Kanagawa, 1969; Andreu-Vázquez *et al.*, 2012; Biswas *et al.*, 2015), being a very rare event. Prediction studies indicated that quadruplets rates in cattle would be 0.1% if multiple birth rate of herd would be 55% (Bennet *et al.*, 1998). Multiple births are undesired in dairy cattle due to detriments caused and abnormal karyotype (Andreu-Vázquez *et al.*, 2012). However, in beef cattle, the trait has economic potential for selection (Kirkpatrick, 2002; Komisarek and Dorynek, 2002).

### Case report

The present report was aimed to notice, for first time in the available scientific literature, the occurrence of birth of quadruplets of a Nelore cow (*Bos taurus indicus*). No ethical

statement was needed since no handling or sampling of animals was done. It is just a notification. A Nelore cow, in its second parturition, at four years and seven months of age, calved quadruplets, being three male calves and one female calf (Fig. 1). It is a rare event. The pregnancy resulted from artificial insemination of the cow by an Angus bull. One of the male calves died after birth. The other three are alive and healthy. Curiously, the cow calved twins in its first parturition. We tried to diagnose calves for freemartinism. However, farm was located far from the laboratory and it did not allow sampling fresh blood for testing

### Discussion

Twinning rates or multiple births are unknown, in scientific literature, for Nelore cattle, but occurrence of twins in commercial herds was noticed by breeders. There are reports of quadruplets in other cattle breeds (indicine and taurine) by Basrur and Kanagawa (1969); Andreu-Vázquez *et al.* (2012) and Biswas *et al.* (2015). It is worth to know that twinning rates in cattle are low, varying from 0.5% to 5% (Vinet *et al.*, 2012).

Some negative consequences come with the occurrence of multiple births such as: smaller birth weight, increase in calf

\*Corresponding author: Gregório Miguel Ferreira de Camargo  
E-mail address: [gregorio.camargo@ufba.br](mailto:gregorio.camargo@ufba.br)



Fig. 1. Nelore cow with three crossbred calves alive and healthy.

mortality, dystocia, stillbirth, calf rejection by dam, retention of placenta, increase in open days and freemartinism (Kirkpatrick, 2002; López-Gatius et al., 2017). Moreover, it is recommended to remove one of the embryos in dairy cattle (Andreu-Vázquez et al., 2012; López-Gatius et al., 2017). However, Kirkpatrick (2002) proposed alternative cow and twin calf management to avoid drawbacks of twinning in beef production systems.

In beef cattle production, the increase in number of calves per parturition improves economic return (Guerra-Martinez et al., 1990; Komisarek and Dorynek, 2002). In production systems with taurine breeds, in which calves are heavier at birth, selection for twins is potentially of interest (Moioli et al., 2017).

Heritability estimates for twinning rates are low (Moioli et al., 2017; Lett and Kirkpatrick, 2018), indicating low additive genetic contribution. The fact that dam, herein reported, had multiple births in both parturitions might be an indication of additive genetic contribution, since other contemporary dams were in same environment and calved singletons. It might also be a contribution of non-additive genetic, since sire was from another breed. Although heritability estimates were low, Van Vleck and Gregory (1996) reported increase in twinning rates by selection. Moioli et al. (2017) reported great increase in heritability estimates when genomic data were incorporated, being interesting when trait is selection criteria.

Nelore breed is raised in tropical environment and mainly in semi-extensive or extensive production systems. Calves are born with average weight of 30 kg. In this situation, selection for multiple births would not be advantageous. However, in some farms, production is intensified by using new technologies. Therefore, it might be a trait interesting to be selected due to its economic return.

The report of animals with superior phenotype, as it is in the present case, are worth to be done. We notice variability within breed. It may be taken into account for further studies of reproductive physiology and biotechnologies and also propositions of phenotypes of interest.

## Acknowledgement

The authors wish to thank the beef cattle breeder for the notification of occurrence and all information that allowed the report.

## Conflict of interest

None of the authors has any conflict of interest to declare.

## References

- Andreu-Vázquez, C., Garcia-Ispuerto, I., López-Gatius, F., 2012. Manual rupture versus transvaginal ultrasound-guided aspiration of allanto-amniotic fluid in multiple pregnancies: a clinical approach to embryo reduction in dairy cattle. *Journal of Reproduction and Development* 58, 420-424.
- Biswas, J., Biswas, S. Pan., S., Mandal, A., 2015. A cytogenetic study of heterosexual quadruplets of cattle (*Bos indicus*) - a case report. *Veterinarski Arhiv* 85, 105-110.
- Basrur, P.K., Kanagawa, H., 1969. Parallelism in chimeric ratios in heterosexual cattle twins. *Genetics* 63, 419-425.
- Bennet, G.L., Echternkamp, S.E., Gregory, K., 1998. A Model of Litter Size Distribution in Cattle. *Journal of Animal Science* 76, 1789-93.
- Guerra-Martinez, P., Dickerson, G.E., Anderson, G.B., Green, R.D., 1990. Embryo-transfer twinning and performance efficiency in beef production. *Journal of Animal Science* 68, 4039-4050.
- Kirkpatrick, B.W., 2002. Management of twinning cow herds. *Journal of Animal Science* 80, E14-E18.
- Komisarek, J., Dorynek, Z., 2002. Genetic aspects of twinning in cattle. *Journal of Applied Genetics* 43, 55-68.
- Lett, B. M., Kirkpatrick, B.W., 2018. Short communication: Heritability of twinning rate in Holstein cattle. *Journal of Dairy Science* 101, 4307-4311.
- López-Gatius, F., Andreu-Vázquez, F.C., Mur-Navales, R., Cabrera, V.E., Hunter R.H.F., 2017. The dilemma of twin pregnancies in dairy cattle. A review of practical prospects. *Livestock Science* 197, 12-16.
- Moioli, B., Steri, R., Marchitelli, C., Catillo, G., Buttazzoni, L., 2017. Genetic parameters and genome-wide associations of twinning rate in a local breed, the Maremmana cattle. *Animal* 11, 1660-1666.
- Van Vleck, L.D., Gregory, K.E., 1996. Genetic trend and environmental effects in a population of cattle selected for twinning. *Journal of Animal Science* 74, 522-528.
- Vinet, A., Drouilhet, L., Bodin, L., Mulsant, P., Fabre, S., Phocas, F., 2012. Genetic control of multiple births in low ovulating mammalian species. *Mammalian Genome* 23, 727-740.