

Phytosociology and Biodiversity Indices of the Woody plants of Alfula Area, West Kordofan, Sudan

Ismail Mirghani ISMAIL

Forestry and Gum Arabic Research Centre, Khartoum, Sudan

*Corresponding author, email: ismail.mirghani@yahoo.com

ARTICLE INFO

Article history:

Received: 22 January 2020

Accepted after corrections

28 February 2020

Keywords:

Species diversity,
distribution, similarity,
vegetation

ABSTRACT

This study was conducted in Alfula area, west Kordofan state. This study aimed at determining the floristic composition, phytosociology, and species diversity of woody species and to produce a checklist of the encountered woody flora. Qualitative and quantitative data collected from forty circular sample plots of 0.1ha (17.84m radius), The Shannon-Wiever and Pielous indices were used to assess species diversity, also similarity between sampled sites and distribution pattern of species was assessed. Fifty-eight species belongs to 16 families were recorded. The diversity indices values varying along studied sites. The total highest density of all trees and shrubs was 866 stem/ha. most dominant species is *Geuara senegalensis*. Most species were aggregately distributed.

1. Introduction

Sudan is one of largest African countries, with an area of 1882000 km². Ecologically the vegetation of Sudan was classified into five zones, these zones are: desert, semi-desert, low rainfall woodland savanna, riverian forest, mountain vegetation. The area of this study located in low rainfall woodland savanna (Harrison and Jackson, 1958). Previously several regional studies regarding vegetation and flora has conducted in Sudan in general, while few study has been conducted in the Alfula area, these studies including (Doka and Yagi, 2009; Ibrahiem, 2003). The diversification and highly richness of vegetation cover are the important reasons led to selection of this area for the present study. In-addition to that, the area is considered as an important source of animal wealth. This study aimed at determining the floristic composition, phytosociology, and species diversity of woody species and to produce a checklist of the existing woody flora.

2. Methods

2.1. Study area

The study was carried out in Alfula area, western Kordofan State (Fig.1). The study area is generally flat and dominated by Gardoud soil with exception of limited areas characterized by clay soil around seasonal water courses (Wadies) (table 1). The area of this study was classified as low rainfall woodland savanna where the annual precipitation rate ranging from 250-1000 mm (Harreison and Jackson, 1958; Sahni, 1968).

Table 1. Characteristic feature of selected sites

Site	Location	Latitude (N)	Longitude (E)	Altitude (M)	Habitat
1	Khoe Gaghmni	11 41 061	028 17 882	1639	Khor bank (Clay)
2	Alwadi Alazrag	11 33 505	028 17310	1559	Khor bank (Clay)
3	Shag Alganna	11 40 826	2813383	1736.8	Gardood
4	Tirgado	1147330	2818833	1873.4	Gardood
5	Alodaya road	1150966	2818860	1863.6	Gardood
6	Baboya	1146572	028 27 597	1783.4	Gardood
7	Kadam	12 03 891	028 47 649	1743.8	Gardood
8	Dahia	12 10 402	028 56310	1757.5	Khor bank (Clay)




Figure 1. Study area: Alfula area, West Kordofan state, Sudan

2.2. Data collection

The vegetation data collected from 40 systematic circular sample plots of 0.1 ha (radius = 17.84 m) with intervals 0.15 mile between each plots, along 8 line.

2.3. Data analysis

Phytosociological analysis

In order to assess the dominance of species in the vegetation communities, density (D), abundance (A), frequency (F %), relative density (RD %), relative.

Density (D) = $\frac{\text{number of plant of a certain species}}{\text{Total area sampled}}$

Relative density (RD %) = $\frac{\text{Density of species X100}}{\text{Total density of all species}}$

Abundance (A) = $\frac{\text{Total number of individual x 100}}{\text{No. of quadrate where species occur}}$

Relative abundance (RA %) = $\frac{\text{Abundance of species x 100}}{\text{Total abundance of all species}}$

Frequency (F %) = $\frac{\text{No. of quadrate where species occur x 100}}{\text{Total numbers of quadrate}}$

$$\text{Relative frequency (RF\%)} = \frac{\text{frequency of species} \times 100}{\text{Total frequency of all species}}$$

$$\text{Importance value index (IVI)} = \text{RD\%} + \text{RA\%} + \text{RF\%}$$

Diversity indices

Species richness was determined as the total number of species present in the studied site. The Shannon diversity index applied to estimate woody plant species diversity along the study area (Shannon and Weaver, 1963). This index was calculated by the equation $H_s = -\sum p_i \ln p_i$. Where, p_i is the proportion of individuals found in the i th species and 'ln' denotes the natural logarithm. Pielou index used for estimation of species evenness (E) after (Pielou, 1966). This index was calculated by the equation $E = H/\ln S$. Where: 'H' is the Shannon-Wiener diversity measure, 'S' is Number of species. Species distribution test and Comparisons of woody plant species composition between different plots were estimated using single linkage cluster analysis based on Jaccard similarity, Biodiversity Pro version 2 (McAleece, 1998).

3. Results and discussion

3.1. Common flora

Botanical account on encountered flora during vegetation survey is given as a checklist. 58 plant species belongs to 16 families. The family *Fabaceae* (through its 3 subfamily) is the highest represent with 19 species, followed by *Combretaceae* which represented by 6 species and *Poaceae* which represented by 5 species; while the other families represented by number of species ranging between 1 to 4 (Tab. 2).

Table 2. A checklist of encountered common flora of AlFula area

Family	plant species	Vernacular name	Habit
Asparagaceae subfamily Asparagoideae	<i>Asparagus flagellaris</i> (Kunth) Baker.	Dayog	Herb
Poaceae subfamily Chloridoideae	<i>Eragrostis tremula</i> Steud.	Bannu	Grass
	<i>Schoenefeldia gracilis</i> Kunth.	Danab Alnaga	Grass
Poaceae subfamily Panicoideae	<i>Pennisetum pedicellatum</i> Tan.	Danab Elkadees	Grass
	<i>Cenchrus biflorus</i> Roxb.	Haskanit	Grass
	<i>Cenchrus ciliaris</i> L., Mant. Alt.	Haskanit	Grass
Vitaceae subfamily Vitoideae	<i>Cissus quadrangularis</i> L.	Salala	Herb
Zygophyllaceae Subfamily Tribuloideae	<i>Balanites aegyptiaca</i> (L.) Delile	Heglig	Tree
Fabaceae Subfam. Caesalpinoideae	<i>Bauhinia rufescens</i> Lam.	Kulkul	Tree
	<i>Cassia seiberiana</i>	Um Kashau, Um Masareen	Tree
	<i>Piliostigma reticulatum</i> (DC.) Hochst.	Kharoub	Tree
	<i>Senna obtusifolia</i> (L.) Irwin & Barneb.	Kawal	Herb
	<i>Tamarindus indica</i> L.	Aradeib	Tree
Fabaceae subfam. Mimosoideae	<i>Acacia mellifera</i> (Vahl) Benth.	Titir	Shrub
	<i>Acacia niloticasubsp.adstringens</i> (Schum. & Thonn.) Roberty.	Sunot	Tree
	<i>Acacia nilotica subsp. tomentosa</i>	Sunot	Tree
	<i>Acacia oerfota</i> (Forssk.) Schweinf.	Laot	Shrub
	<i>Acacia sieberiana</i> DC.	Kuk	Tree
	<i>Acacia senegal</i> (L.) Willd.	Hashab	Tree
		Talih	Tree
	<i>Albizia amara</i> (Roxb.) Boiv.	Arad	Tree
	<i>Albizia anthelmintica</i> Brongn.	Um Takirma	Tree
	<i>Dichrostachys cinerea</i> (L.)White & Arn.	Kadad	Shrub
	<i>Entada africana</i> Guill. & Perr.	Seisban. Entada	Tree
<i>Faidherbia albida</i> (DeL.) Chev.	Haraz	Tree	
Fabaceae sub-fam. Faboideae	<i>Dalbergia melanoxylo</i> n Guill. & Perr.	Abanus	Tree
	<i>Lonchocarpus laxiflorus</i> Guill. & Perr.	Khashkhash Azrag	Tree
	<i>Zornia glochidiata</i> Reichb. ex DC.	Sheleni	Herb
Rhamnaceae	<i>Ziziphus spina-christi</i> (L.) Desf.	Sidir	Tree
	<i>Ctenolepis cerasiformis</i> (Stocks) C. B. Clarke.	Sim Elter	Herb
Euphorbiaceae subfamily Euphobioideae	<i>Euphorbia venenifica</i> Tremaux ex Kotschy		Shrub

Combretaceae	<i>Anogeissus leiocarpus</i> (DC.) Guill. & Perr.	Sahab	Tree
	<i>Combretum aculeatum</i> Vent.	Habeel	Tree
	<i>Combretum glutinosum</i> Perr. ex DC.	Habeel	Tree
	<i>Guiera senegalensis</i> J. F. Gmel.	Ghibeish	Tree
	<i>Terminalia brownii</i> Fresen	Subagh	Tree
Anacardiaceae	<i>Terminalia laxiflora</i> Engl. & Diels Monogr	Darout	Tree
	<i>Lannea humilis</i> (Oliv.) Engl.	Layoun	Tree
	<i>Lannea fruticosa</i> (Hochst. ex A. Rich.) Engl.	Layoun	Tree
Malvaceae subfamily Bombacoideae	<i>Sclerocarya birrea</i> (A. Rich.) Hochst.	Himeid	Tree
Malvaceae Subfamily Grewioideae	<i>Adansonia digitata</i> Linn.	Tabaldi	Tree
	<i>Grewia flavescens</i> Juss.	Khlekhsan	Shrub
	<i>Grewia tenax</i> (Forsk.) Fiori.	Gudiem	Shrub
Malvaceae subfamily Malvoideae	<i>Grewia villosa</i> Willd.	Tikko	Shrub
	<i>Sida alba</i> Linn.	Um Shadeda	Herb
Malvaceae subfamily Sterculioideae	<i>Sida cordifolia</i> L.	Nayada	Herb
	<i>Sterculia setigera</i> Del.	Tartar	Tree
Capparaceae	<i>Boscia angustifolia</i> A. Rich.	Sareh	Tree
	<i>Boscia senegalensis</i> Lam.	Mikheit	Tree
	<i>Capparis decidua</i> (Forsk.) Edgew.	Tundub	Small tree
	<i>Capparis tomentosa</i> Lam.		Shrub
Rubiaceae subfamily Ixoroideae	<i>Catunaregam nilotica</i> (Stapf.) Tirveng.	Shagart Elmarfaein	Tree
	<i>Feretia apodanthera</i> Del.	Shai Elbitera	Tree
Apocynaceae subfamily Apocynoideae	<i>Adenium obesum</i> (Forssk.) Roem & Schult.	Shagart Alsim	Tree
Apocynaceae subfamily Asclepiadoideae	<i>Leptadenia pyrotechnica</i> (Forsk.) Decne.	Marikh	Shrub
Boraginaceae	<i>Cordia sinensis</i> Lam.	Andrab	Tree
Acanthaceae Subfamily Acanthoideae	<i>Hygrophila auriculata</i> (Schumach.) Heine.	Abu Shweika	Herb
	<i>Peristrophe paniculata</i> (Forssk.) Brummitt.	Abu Rukab	Herb
Bignoniaceae	<i>Stereospermum kunthianum</i> Cham.	Khashkhash abiad	Tree

3.2. Density

The total density of all woody plants was varying along different sites. Baboya (Site 6) recorded (866 plant/ha as a highest density of woody plants, while Dahia (Site 8) has recorded (182 plant/ha as a lowest density. *Guiera senegalensis* recorded highest values of relative density (92.61%), (60.74 %) and (47.13 %) in Baboya (Site 6), Tirgado (Site 4) and Alodaya road (Site 5) respectively; *Grewia flavescens* was a second species that recorded high relative density (41.67%) in Alodaya road (Site 5) (Fig. 2, Tab. 3). The highest relative density which recorded by above mentioned species may attributed to those species are not palatable enough for livestock, especially in regeneration stages.




Figure 2. Density along different studied sites

Table 3. density, relative density(RD), relative abundance (RA), relative frequency (RF), importance value index (IVI) and distribution pattern of the plant species in the study area

Species	site 1 (Khor Jaghmi)		site 2 (Alwadi Alazrag)		site 3 (Shag AlGanna)		site4 (Tirgado)		site 5 (Alodya Road)		site 6 (Baboya)		site 7 (Kadam)		site 8 (Dahia)		Aggregation
	RD	IVI	RD	IVI	RD	IVI	RD	IVI	RD	IVI	RD	IVI	RD	IVI	RD	IVI	
<i>Acacia mellifera</i>	0	0	1.32	10.65	0	0	0.37	4.79	0	0	0	0	0	0	0	0	Random
<i>Acacia nilotica</i> subsp <i>adstringens</i>	16.6	43.27	0	0	5.75	21.61	4.07	20.13	0	0	0	0	0	0	20.88	50.35	Aggregated
<i>Acacia nilotica</i> subsp. <i>tomentosa</i>	0	0	33.55	69.9	0	0	0	0	0	0	0	0	0	0	0	0	Aggregated
<i>Acacia oerfota</i>	0	0	0	0	0	0	0	0	0	0	0	0	5.21	19.09	3.3	15.55	Aggregated
<i>Acacia seiberiana</i>	0.72	6.97	0	0	1.77	11.78	0	0	0	0	0	0			0	0	Random
<i>Acacia senegal</i>	8.63	35.65	0	0	1.33	8.93	0	0	0	0	0	0	5.21	21.13	1.1	7.96	Aggregated
<i>Acacia seyal</i> subsp. <i>seyal</i>	20.9	51	25	57.37	0	0	0	0	0.29	5.02	0	0	0	0	0	0	Aggregated
<i>Adenium obesum</i>	0	0	0	0	0	0	0.37	4.79	0	0	0.23	5.66	0	0	0	0	Random
<i>Albizia amara</i>	4.32	19.1	0	0	8.85	32.37	1.48	12.58	4.6	23.53	1.56	16.01	7.29	27.41	8.33	18.12	Aggregated
<i>Albizia anthelmenthica</i>	0	0	0	0	0.44	5.45	0	0	0	0	0.23	5.66	1.04	7.4	0	0	Random
<i>Anogeissus leiocarpa</i>	0.72	6.97	5.26	19.53	0	0	0	0	0	0	0.69	14.82	3.13	14.19	0	0	Aggregated
<i>Balanites aegyptiaca</i>	20.9	51.42	7.24	33.7	1.77	10.67	0.74	6.45	0	0	0	0	22.92	53.27	6.59	24.49	Aggregated
<i>Bauhenia reticulata</i>	0	0	0	0	0	0	0.37	4.79	0.29	5.02	0	0	8.33	26.62	6.59	23.02	Aggregated
<i>Bauhenia rufescens</i>	0	0	0	0	0	0	0.37	4.79	0	0	0	0	0	0	2.2	11.76	Random
<i>Boscia angustifolia</i>	0	0	0	0	0	0	1.11	11.78	0.86	9.69	0	0	0	0	0	0	Aggregated
<i>Boscia senegalensis</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	38.46	82.43	Aggregated
<i>Cassia seiberiana</i>	0	0	0	0	0	0	0.74	8.28	0.86	9.69	0	0	0	0	0	0	Aggregated
<i>Catonaregam nilotica</i>	0	0	0.66	6.15	0	0	0	0	0	0	0	0	0	0	0	0	Random
<i>Combretum aculeatum</i>	6.48	23.25	0	0	3.98	19.39	0	0	0	0	0	0	4.17	17.58	0	0	Aggregated
<i>Combretum glotinosum</i>	0	0	0	0	1.33	8.93	0.37	4.79	0	0	0.23	5.66	2.08	10.79	0	0	Random
<i>Cordia sinensis</i>	0	0	0.66	6.15	0	0	0	0	0	0	0	0	0	0	0	0	Random
<i>Dalbergia melanoxylon</i>	0	0	4.61	19.96	3.54	18.12	1.48	10.32	1.72	14.64	2.31	26.22	0	0	0	0	Aggregated
<i>Dicrostachys cinerea</i>	0	0	0.66	6.15	0	0	0	0	1.72	14.64	0	0	0	0	0	0	Aggregated
<i>Feidherbia albida</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1.1	12.13	Random
<i>Feretia apodanthera</i>	0	0	14.47	40.25	0	0	0	0	0	0	0	0	0	0	0	0	Aggregated
<i>Guiera senegalensis</i>	0	0	0	0	38.5	81.59	60.74	126.3	47.13	106.53	92.61	201.33	31.25	74.55	0	0	Aggregated
<i>Grewia flavescens</i>	0	0	0	0	31.86	70.07	25.56	60.36	41.67	96.18	2.31	18.97	0	0	0	0	Aggregated
<i>Grewia villosa</i>	5.76	21.67	0	0	0	0	0	0	0	0	0	0	0	0	0	0	Aggregated
<i>Lannea fruticosa</i>	0	0	0	0	0	0	1.48	10.32	0	0	0	0	0	0	0	0	Aggregated
<i>Lannea humilis</i>	0	0	0	0	0	0	0	0	0	0	0.23	5.66	0	0	0	0	Random
<i>Lonchocarpus laxiflorus</i>	0	0	0	0	0	0	0.37	4.79	0	0	0	0	0	0	0	0	Random
<i>Sclerocarya birrea</i>	0	0	0	0	0	0	0.37	4.79	0.29	5.02	0	0	0	0	0	0	Random
<i>Asparagus flagellaris</i>	0	0	0	0	0.44	5.45	0	0	0	0	0	0	0	0	0	0	Random
<i>Streospermum kunthianum</i>	0	0	0	0	0	0	0	0	0.29	5.02	0	0	0	0	0	0	Random
<i>Tamarindus indica</i>	0	0	4.61	18.05	0	0	0	0	0	0	0	0	0	0	1.1	7.96	Aggregated
<i>Terminalia brownii</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2.2	11.76	Random
<i>Terminalia laxiflora</i>	0	0	0	0	0	0	0	0	0.29	5.02	0	0	0	0	0	0	Random
<i>Ziziphus spina christi</i>	15.1	40.69	1.97	12.13	0.44	5.45	0	0	0	0	0	0	9.38	27.97	12.09	34.48	Aggregated

3.3. Dominance

The dominance of the plant species was determined using the Importance Value Index (IVI) of assessed species. From the analysis results the most dominant species is *Geuara senegalensis* with values of IVI (201.33),(126.3) (106.53), (81.59) and (74.55) in site 6 (Baboya), site 4 (Tirgado), site 5 (Alodya Road), site 3 (Shag AlGanna) and site 7 (Kadam) respectively, and followed by *Grewia flavescens* with values of IVI (96.18), (70.07) and (60.36) in site 5 (Alodya Road), site 3 (Shag AlGanna) and site4 (Tirgado) respectively, and *Acacia nilotica* subsp. *tomentosa* with values of IVI (69.9) in site 2 (Alwadi Alazrag); while the other species have an IVI ranging between 51 - 4.79. The dominance of *Geuara senegalensis* may attribute to the suitability of environmental factors such as precipitation amount, temperature and soil type especially all those sites in which this species is dominant are Gardoud soil (Tab. 3).

3.4. Distribution Patterns

Distribution pattern of plant were assessed and results showed that 15 plants were randomly and 22 were aggregated (Tab. 3). It is clear from above mentioned results that most of plant species were aggregate distributed and this indicates that the habitat is suitable for these species.

3.5. Diversity indices

The results revealed that richness was highest at site 4 (Tirgado) (16), followed by site 3 (Shag Alghana) with richness value (13). The highest Shannon diversity index was 0.899 in site 7 (Kadam), followed by 0.867 in site 1 (Khor Jaghmi), whereas the least Shannon diversity index was 0.161 in site 6 (Baboya). The highest species evenness index was recorded in site 1(Khor Jaghmi) (0.867), whereas the least evenness index was 0.179 in 6 (Baboya) (Fig.3). These values are relatively small when compared to the values that calculated in other sites in Sudan such as Jebel ELDair (Ismail and Mahmoud, 2010), Nuba mountains (Ismail and Alawad, 2017), and Jebel AlGerri (Ismail and ElSheikh, 2016). The decrease of diversity richness may due to that the area of this study is considered as one of important animal routes in Sudan, where as the animal browsing the trees and removing the regeneration cover.




Figure 3. Species richness and diversity indices of woody plants

3.6. Site similarity

The similarity measures indicated that the higher percentage of similarity was found between Site 3 (Shag AlGanna) and Site 7 (Kadam), where as (44.4%) of species were shared between the two sites, followed by (38.09%) between site 3 (Shag AlGanna) and site 4 (Tirgado) (Fig. 4), this might be explained by the fact that these plots have similar altitudes and environment characteristics specially these sites are located in Gardood soil..




Figure 4. Dendrogram showing percentage of similarity between studied sites

4. Conclusion

The results of the study revealed that Alfula area was varying in richness of plant diversity of the woody species within different sites. In addition, the study showed that the density is ranging from 866 stem/ha as a highest value density to 182 stem/ha as a lowest value density and most of the species were aggregately distributed with higher percentage of similarity for most studied sites. the study area is characterized by relatively low diversity of woody plant when compared with the areas that located in the same ecological zone, A checklist of trees and shrubs that were existing in the study area is produced and the species are listed, (Tab.2).

References

1. Doka I.G. and Yagi S. M., 2009. Ethnobotanical Survey of Medicinal Plants in West Kordofan (Western Sudan). *Ethnobotanical Leaflets* 13: 1409-1416.
2. Harrison, M.N. and J.K. Jackson. 1958. Ecological classification of the vegetation of the Sudan. Agricultural Publication Committee, Khartoum. 44 pp
3. Ibrahim M. A. M. 2003. A Study of the Vegetation of El Fula Area, Western Kordofan State. A thesis submitted in fulfillment of the requirement for the award of the degree of Ph.D. (Forestry), U of K.
4. Ismail and ElSheikh, 2016. Floristic composition, Phytosociology and species diversity of Woody vegetation of Jebel AlGerri, Blue Nile State, Sudan. *Jour. of Nat. Ressour. & Environ. STU.* , 4 (2): 24-34
5. Ismail I. M. and Alawad, A. A. 2017. Changes in Plant Species Richness and Density along Altitudinal Gradient in Rashad and Alabassia Localities, South Kordofan, Sudan. *Neelain Journal of Science and Technology*. 1 (1): 21–28.
6. Ismail I. M. and Mahmoud A. E. 2010. Floristic Composition and Species Diversity of Woody Vegetation of Jabal Eldair (North Kordofan State-Sudan). *Sudan Silva*, 14 (1):49-60.
7. McAleece N., 1998. Bio Diversity Professional Beta. The Natural History Museum and the Scottish Association for Marine Science.
8. Pielou E.C., 1966. The measurement of diversity in different types of biological collections. *Journal of Theoretical Biology* 13: 131-144.
9. Sahni, K.C. 1968. Important Trees of the Northern Sudan, Published for UNDP and FAO by University of Khartoum Press, Khartoum, Sudan.
10. Shannon C. E. and Weaver W., 1963. The Mathematical theory of Communication. University of Illinois Press, Urbana, USA

Please cite this Article as:

Ismail I. M., 2020. Phytosociology and Biodiversity Indices of the Woody plants of Alfula Area, West Kordofan, Sudan. *Agric. For. J.*, 4(1): 40-46.

DOI: <https://doi.org/10.5281/zenodo.3837661>