

## Functional brain networks involved in gaze and emotional processing

Journal:	<i>European Journal of Neuroscience</i>
Manuscript ID	Draft
Manuscript Type:	Research Report
Date Submitted by the Author:	n/a
Complete List of Authors:	Ziaei, Maryam; University of Queensland, Centre for Advanced Imaging Ebner, Natalie; University of Florida, Psychology Burianová, Hana; University of Queensland, Centre for advanced imaging
Key Words:	amygdala, emotional expression, eye gaze, Functional connectivity, multivariate

## Functional brain networks involved in gaze and emotional processing

Maryam Ziaei<sup>1,2</sup>, Natalie C. Ebner<sup>3,4</sup>, Hana Burianová<sup>2,5</sup>

<sup>1</sup>School of Psychology, University of Queensland, Brisbane, Australia

<sup>2</sup>Centre for Advanced Imaging, University of Queensland, Brisbane, Australia

<sup>3</sup>Department of Psychology, University of Florida, Gainesville, Florida, USA

<sup>4</sup>Department of Aging and Geriatric Research, Cognitive Aging and Memory Program, Clinical Translational Research Program (CAM-CTRP), Institute on Aging, University of Florida, Gainesville, Florida, USA

<sup>5</sup>Department of Psychology, Swansea University, Swansea, United Kingdom

## Abstract

Eye-gaze direction plays a fundamental role in the perception of facial features and particularly the processing of emotional facial expressions. Yet, the neural underpinnings of the integration of eye gaze and emotional facial cues are not well understood. The primary aim of this study was to delineate the functional networks that subservise the recognition of emotional expressions as a function of eye gaze. Participants were asked to identify happy, angry, or neutral faces, displayed with direct or averted gaze, whilst their neural responses were measured with fMRI. The results show that recognition of happy expressions, irrespective of eye-gaze direction, engages the critical nodes of the default mode network. Recognition of angry faces, on the other hand, is gaze-dependent, engaging the critical nodes of the salience network when presented with direct gaze, but fronto-parietal areas when presented with averted gaze. Functional connectivity analysis further shows gaze-dependent engagement of a large-scale network connected to bilateral amygdala during the recognition of angry expression. The study provides novel insights into the functional connectivity between the amygdala and other critical social-cognitive brain nodes, which are essential in processing of ambiguous, potentially threatening signals. These findings have important implications for psychiatric disorders, such as post-traumatic stress disorder, which are characterized by aberrant limbic connectivity.

**Keywords:** Amygdala, emotional expression, eye gaze, functional connectivity, multivariate

## Introduction

Eye-gaze perception plays a fundamental role in social and non-verbal communication, signaling one's intention to approach (direct gaze) or avoid (averted gaze) a person. Together with facial emotional cues, eye gaze carries important information about the underlying emotions and thus can enhance or disrupt perception of the expressed emotion. According to shared signal hypothesis (Adams & Kleck, 2005), when eye gaze matches the underlying emotion (e.g., angry expression with direct gaze), perception of that emotion would be enhanced. However, when eye gaze and emotion convey discordant information (e.g., angry expression with averted gaze), emotion perception would be diminished, possibly due to an increase in the ambiguity of social signaling. The ability to integrate the different facial cues to determine others' intentions and affective or mental state is thus crucial to one's everyday social communication. Thus, understanding the functional networks of such highly complex processes will provide insights into the underlying mechanisms involved in social-cognitive or social functioning impairments among various psychiatric, neurological, and neurodegenerative illnesses (Burns, 2006; Kennedy & Adolphs, 2012; Yu & Wu, 2013).

A number of accounts have been proposed to explain the mechanisms involved in the processing of concomitant eye gaze and emotional expressions. According to the shared signal hypothesis, eye gaze and emotional cues share the congruent values of approach or avoidance tendencies and, therefore, should be processed more efficiently when they are both approach/avoidance congruent (Adams & Kleck, 2005). Alternatively, the proponents of the self-relevance appraisal hypothesis argue that facial cues are appraised according to their relevance to the observers' needs, goals, and well-being and thus should be processed more efficiently when the cues are perceived as more self-relevant (Sander et al., 2007). Neuroimaging and lesion-based studies have provided support for both of these accounts and highlighted the importance of the amygdala in the integration of emotional cues with eye-gaze cues. Although some studies have reported the role of the amygdala in the recognition of

angry emotion with direct eye gaze (Cristinzio et al., 2010; N'Diaye et al., 2009; Sander et al., 2007; Sato et al., 2010), others have found the opposite results and showed enhanced activity in the amygdala when presented angry emotion with averted eye gaze (Adams et al., 2012; Adams et al., 2003; Adams & Kleck, 2005). Besides these inconsistencies, only a few studies assessed such emotion-gaze interactions for happy expressions, with disparate results. Adams and Kleck (2005) found enhanced recognition of happiness with direct gaze; however, Cristinzio et al. (2010) and Sander et al. (2007) did not find any significant differences in the intensity rating of happy facial expressions as a function of eye-gaze orientation. Although parts of discrepancies reflect differences in the paradigm used in abovementioned studies, the underlying neural circuitry of emotion-gaze integration is still under investigated. Therefore, the primary aim of this study was to examine whole-brain activity during the recognition of happy and angry facial expressions as a function of eye gaze. Given that gazes are used as indicators of expresser's attentional orientation (Sander et al., 2007), we treated the eye-gaze cues as means of conveying signals by a target face and not what the observers felt, similar to Adams and Kleck (2005).

Although amygdala has been considered a major hub for different social processes, e.g., social perception or social attribution (Bickart et al., 2014), it is still unclear to which brain regions is amygdala functionally connected when processing socially-relevant and communicative signals. As the ability to understand and integrate socially-relevant cues is essential for social cognition, these processes undoubtedly rely on a large number of brain structures and their connections (Kennedy & Adolphs, 2012). In other words, given the complexity of the underlying cognitive integration of emotion and eye gaze, it is reasonable to suggest that these processes would be supported by a large-scale, distributed functional network. However, to our knowledge, no existing empirical research has examined functional connectivity with the amygdala during recognition of eye-gaze and emotional expressions

cues. Thus, the second aim of this study was to delineate a task-related network that is functionally connected to bilateral amygdala and to assess the strength of connectivity within this network as a function of eye gaze. Delineating the amygdala network underlying critical social-cognitive processes may aid our understanding of the markers of proper social functioning and, in turn, of specific disruptions in the functional circuitry underlying emotion-related psychiatric or neurological disorders.

## **Methods**

### ***Participants***

Twenty-one healthy young adults (age 17-27 years,  $M = 20.65$ ,  $SD = 2.66$ , 10 males) participated in this study. One participant was excluded from the whole-brain analysis due to extensive movement and two participants were removed from the connectivity analysis due to outlier nature of the brain signals. All participants were undergraduate students recruited from the University of Queensland in exchange for course credit or \$15 AUD per hour. Participants were screened for claustrophobia, neurological and psychiatric disorders, and magnetic resonance imaging (MRI) compatibility. All participants were right-handed English speakers, had normal or corrected-to-normal vision, and had no history of neurological impairment or psychiatric illnesses. They took part in two separate testing sessions: neuropsychological assessment and functional MRI (fMRI) scanning session. They were provided with a written consent as approved by the Human Research Ethics Committee at the University of Queensland and were debriefed upon the completion of the second session.

### ***Materials***

The stimuli consisted of color, front-view faces selected from the FACES database (Ebner et al., 2010) and included happy, angry, and neutral expressions. The gazes of the posers were photoshopped toward either right or left side. Eight lists of 60 faces were created

using MATLAB, based on three selection criteria: gender of the poser (male/female), gaze (direct/averted), and emotional expression (happy/angry/neutral). Each face identity was presented once with only one emotional expression displayed within each run. Each of these lists consisted of equal numbers of male and female posers (30), direct and averted gaze directions (30), and emotional expressions (20). Finally, the faces in each list were matched based on the independent ratings of attractiveness ( $M = 41.66$ ,  $SD = 13.08$ ; Ebner et al. (2010)). Each participant was presented with five of the lists (300 trials in total) and the presentation order of the lists was counterbalanced across participants in the scanner. In order to avoid habituation toward the faces, no more than two faces of each category (age of the face, facial expressions, and gaze direction) were repeated in a row. The faces were presented in 600 x 450 pixels, which were adjusted for the presentation in the scanner and presented against gray background, using E-prime software.

### ***Experimental design***

The scanner session lasted for 50 minutes and consisted of 2 components: structural magnetic resonance imaging (sMRI) and functional magnetic resonance imaging (fMRI) of emotion recognition task. Prior to the scanning, participants were verbally and visually instructed about the task and practiced until they were familiarized with the instructions. During the emotion recognition task in the scanner, participants were asked to identify, as fast and accurate as possible, whether the faces displayed happy, angry, or neutral expression by pressing the relevant buttons on an MRI-compatible response box. Each face was presented, one at a time, for 3.5 seconds, followed by a fixation cross, which was randomly jittered using three time intervals: 0.5 seconds (20 trials), 1 seconds (20 trials), and 1.5 seconds (20 trials). The jittered ITI allowed for an independent estimation of the BOLD response on a trial-by-trial basis. The task consisted of five runs of the emotion recognition task; each run

lasted for 4.5 minutes. Participants performed two runs of scanner task, which was followed by an acquisition of sMRI, then performed three runs of the emotion recognition task.

### ***Image Acquisition, Preprocessing, and Analysis***

Functional images were acquired at the Centre for Advanced Imaging using a 3-T Siemens scanner with a 32-channel head coil. The functional images were obtained using a whole-head T2\*-weighted echo-planar image (EPI) sequence (93 slices, repetition time (TR) = 3000ms, echo time (TE) = 45ms, flip angle = 90°, field of view (FOV) = 192mm, voxel size = 2mm<sup>3</sup>). High-resolution T1-weighted images were acquired with a MPRAG sequence (126 slices with 1mm thickness, TR = 1900ms, TE = 2.3ms, TI = 900ms, FOV = 230ms, voxel size = 0.9mm<sup>3</sup>). The tasks were presented to participants on a computer screen through a mirror mounted on top of the head coil. Participants were provided with earplugs and cushions inside the head coil to dampen noise and minimize head movement.

For functional analysis, T2\*-weighted images were pre-processed with Statistical Parametric Mapping Software (SPM8; <http://www.fil.ion.ucl.ac.uk/spm>) implemented in MATLAB 2010b (Mathworks Inc., MA). Following the realignment to a mean image for head-motion correction, images were segmented to gray and white matter. Then, images were spatially normalized into a standard stereotaxic space with voxel size of 2mm<sup>3</sup>, using the Montreal Neurological Institute (MNI) template, and spatially smoothed with a 6-mm Gaussian Kernel.

The procedure of the fMRI analysis was twofold. First, we examined the whole-brain activity during emotion recognition of faces displayed with direct or averted gaze. For this purpose, we conducted a whole-brain analysis in which the BOLD response for the whole brain was measured across the experimental conditions. Second, we examined the connectivity of the functional network underlying emotion recognition of faces with direct



and averted gaze. For this purpose, we selected bilateral amygdala as the seed region and correlated its BOLD intensity with that of the rest of the brain.

The fMRI data were statistically analyzed using a multivariate analytical technique Partial Least Squares (PLS; McIntosh et al. (1996); McIntosh et al. (2004)); for a detailed tutorial and review of PLS, see Krishnan et al. (2011), as implemented in PLS software ([http://research.baycrest.org/pls\\_software](http://research.baycrest.org/pls_software)) running on MATLAB 2010b (The MathWorks Inc., MA). PLS analysis uses singular value decomposition (SVD) of a single matrix that contains all participants' data to find a set of orthogonal latent variables (LVs), which represent linear combinations of the original variables. Therefore, PLS enables differentiation of the degree of contribution of different brain regions associated with task demands, behavioral or anatomical covariates, or functional seed activity. The first LV usually accounts for the largest covariance in the data, with progressively smaller amount for subsequent LVs. Each LV delineates cohesive patterns of brain activity related to experimental conditions. Additionally, brain scores are calculated as the dot product of a subject's image volume of each LV. The brain score reflects how strongly each subject contributes to the pattern expressed in each LV. Each LV consists of a singular image of voxel saliences (*i.e.*, a spatiotemporal pattern of brain activity), a singular profile of task saliences (*i.e.*, a set of weights that indicate how brain activity in the singular image is related to the experimental conditions, functional seeds, or behavioral/anatomical covariates), and a singular value (*i.e.*, the amount of covariance accounted for by the LV). Given that the task was event-related, therefore, the analysis was conducted on the 15-sec period (5 TRs), starting at the onset of the faces, and activity at each time point in the analysis was normalized to activity in the first TR (Labeled 0 in the Figure 3). The PLS analysis for the event-related data reveals a set of brain regions related to the task for each TR on each LV. For each TR, the pattern of brain activity

identified for that TR is calculated for each participant. Mean brain scores across participants and across the entire brain are then plotted across the 5 TRs used in the analysis.

The statistical significance of each LV is assessed using permutation test, which determines that the probability of a singular value from 500 random reordering and resampling is larger than initial obtained value (McIntosh et al., 1996). In addition to the permutation test, to determine the reliability of the salience for each brain voxel, a standard error of each voxel's salience on each LV is estimated by 100 bootstrap resampling steps (Efron & Tibshirani, 1985). Peak voxels with a bootstrap ratio (BSR; *i.e.*, salience/standard error) > 2.5 were considered to be reliable, as these approximate  $p < 0.01$  (Sampson et al., 1989). As the activation patterns identified by PLS and corresponding brain responses is done in one single step, therefore, there is no need for multiple comparison correction.

#### *Whole-Brain Analysis*

We assessed whether emotion recognition is modulated by eye gaze and identified the specific functional loci for a priori selected anatomical region (amygdala) by examining whole-brain activations during two emotional expressions (angry and happy) and two eye-gaze directions (averted and direct). Neutral faces were utilized in the experimental design as a control condition, in order to remove the effect of visual perception (for a review see Sabatinelli et al. (2011)). A separate set of analysis included neutral conditions and revealed two main findings. First, the brain networks involved for happy and angry expressions did not change as a matter of including neutral conditions in the analysis. Second, salience network, including anterior cingulate gyrus and bilateral insula, was involved during recognition of neutral expressions irrespective of the eye gaze. However, given that previous works also found that the ambiguity of neutral faces may lead to uncertainty and heightened vigilance, which, in turn, may increase amygdala activity (Blasi et al., 2009), all of the analyses in the results section were reported only for happy and angry facial expressions.

### *Functional Connectivity*

We examined task-related functional connectivity during angry emotional expressions for direct and averted gaze by correlating activity in bilateral amygdala with activity in the rest of the brain during angry emotion recognition. Although amygdala activity has been reported in processing of happy facial expressions (Canli et al., 2002), we did not find any amygdala activity in the whole-brain findings during happy facial recognition; thus, we conducted the functional connectivity analyses on the angry expression conditions only.

The selection of bilateral amygdala was based on two criteria: first, theoretical – previous studies have highlighted the critical role of bilateral amygdala in gaze and emotional processing (Calder & Young, 2005; Carlin & Calder, 2013; Itier & Batty, 2009; Shepherd, 2010); and second, data-driven – in the whole-brain analysis we identified the functional loci for the a priori amygdala regions, left (-18 -4 -12) and right (20 -8 -12) during recognition of angry expressions. To delineate the functional network involved during gaze and emotional processing, we extracted the BOLD values from the peak voxels of the seed regions for the angry conditions and correlated them with activity in the rest of the brain across all participants. These correlations were then combined into a matrix and decomposed with singular value decomposition, resulting in a set of LVs characterizing the set of regions where activity was correlated with seed activity during direct or averted gaze conditions. The significance and reliability of the analysis were determined by permutation test and bootstrap sampling, as described above.

## **Results**

### ***Behavioral Results***

A 2 (eye-gaze direction) by 2 (emotions: happy and angry) repeated measures ANOVA on accuracy revealed a significant main effect of emotion,  $F = (1, 18) = 13.01$ ,  $p < .01$ ,  $\eta_p^2 =$

.42, with higher accuracy for happy than angry faces. A similar analysis was conducted for the response times. Due to the long RT (+3 SD more than the group mean), one participant was excluded from the analysis performed on RTs. A significant main effect of emotion,  $F = (1,17) = 34.47, p < .001, \eta_p^2 = .67$ , suggests that happy faces were recognized faster than angry faces. No significant main effect of gaze or interactions between emotion and eye-gaze directions were found for RTs or accuracy (all  $F_s < 1$ ).

[Insert Figure 1 about here]

### ***Behavioral eye-tracker Results***

In addition to the fMRI session, participants undertook a separate behavioral session in which various cognitive and emotional background measures were collected (Table 1). In order to examine participants' eye-tracker patterns for different emotional conditions, they have performed an eye-tracker task in which they were presented with same faces from the scanner task intermixed with not-previously seen faces. Fixation points and fixation times were recorded using a chin-rest SR EyeLink 1000 eye tracker (SR Research, Ontario, Canada). Prior to the onset of the faces, participants performed a 9-point eye calibration and validation procedure to ensure the accurate recording of the eye positions from different points on the screen. Participants were instructed to focus on each fixation point on the screen until it disappears and then move to the next point. The task included 60 number of happy, 60 number of angry and 60 number of neutral faces with equal number of gaze directions (90 direct and 90 avert in total). Each face was presented on the screen for 3.5 sec. and 1 sec. fixation cross was included between faces to ensure that participants' focus was on the center of the screen prior to the start of the next face. Two region of interest (ROIs) were selected, one for the mouth and one for the eye regions for each face separately. Due to the technical problems during eye-tracker recording, we only obtained clean data from 17 participants. Fixation duration was included in the analyses if they were within each ROI. A two (ROIs;

mouth or eye ROI) by two (emotions; angry and happy) repeated measure ANOVA on fixation durations revealed no significant main effect of emotion ( $F(1,16) = .26, p > .05, \eta_p^2 = .01$ ), main effect of ROI ( $F(1,16) = 1.64, p > .05, \eta_p^2 = .09$ ), main effect of ROI ( $F(1,16) = .84, p > .05, \eta_p^2 = .05$ ), or interaction between ROIs and emotions ( $F(1,16) = 1.51, p > .05, \eta_p^2 = .08$ ), suggesting that fixation times spent on each faces' regions were equal when recognizing happy or angry expressions. Although happy expressions seem to have salient facial feature, open mouth, that does not seem to have any interference on the amount of time participants spent on different regions of faces.

### ***Whole-brain Results***

The results from whole-brain analyses delineated two significant LVs. LV1 accounted for 48% of covariance in the data and revealed a set of brain regions, which were engaged during the processing of angry averted conditions relative to the other conditions. In line with our first prediction, this set of regions included bilateral amygdala as well as bilateral inferior frontal gyrus (IFG), right middle frontal gyrus, bilateral superior frontal gyrus, medial frontal gyrus, left cingulate gyrus, bilateral inferior parietal lobe (IPL), bilateral insula, left superior temporal gyrus (STG), putamen, bilateral thalamus, and bilateral cuneus (Fig. 2, Panel A & Table 2). LV2 accounted for 33% of covariance in the data, revealing a set of regions with increased activity during recognition of angry direct faces relative to the other conditions (Fig. 2, Panel B & Table 3). These areas included right superior frontal gyrus, right cingulate gyrus, right middle temporal gyrus, right superior parietal lobe, bilateral occipital gyrus, bilateral insula, bilateral putamen, and left amygdala.

Happy facial expressions with both direct and averted gaze directions, on the other hand, activated bilateral anterior cingulate gyrus, left medial frontal gyrus, bilateral superior frontal gyrus, bilateral middle temporal gyrus, left IPL, bilateral superior parietal lobe, left precuneus, and left posterior cingulate cortex (PCC; Fig. 2, Panel C & Table 3).

[Insert Figure 2 and Tables 2&3 here]

Furthermore, we extracted and compared the time courses of the amygdala during recognition of angry with averted relative to direct gaze conditions. During the angry expression with averted gaze condition, activity in left amygdala peaked around 6 seconds, whereas activity in right amygdala showed a more sustained activation during recognition of angry averted condition relative to the angry direct condition (Fig. 3). A series of independent *t*-tests showed significant differences between signal intensity of right and left amygdala at time points 3, 6, and 9-sec after stimulus onset during recognition of angry expressions with averted gaze relative to the angry expression with direct gaze, all  $ps < .05$ .

[Insert Figure 3 here]

### ***Functional Connectivity Results***

The results from the seed PLS analysis revealed one significant LV, which explained 67% of covariance in the data and delineated a functional network connected to bilateral amygdala. This functional network was engaged significantly more strongly during recognition of angry emotion with averted gaze than it was during recognition of angry emotion with direct gaze (Fig. 4 & Table 4). This network included bilateral middle frontal gyrus, bilateral superior frontal gyrus, right anterior cingulate gyrus, right inferior frontal gyrus, bilateral STG, bilateral PCC, left IPL, precuneus, and bilateral thalamus.

[Insert Figure 4 and Table 4 here]

### **Discussion**

The aims of the study were to examine whole-brain activity and functional connectivity during emotion recognition of faces displayed with direct or averted eye gaze. Three primary findings emerged: i) although participants did not show any modulation of eye gaze for happy

expressions, recognition of angry expressions was modulated by the direction of eye gaze; ii) in line with some previous works (Adams et al., 2012; Adams et al., 2003; Adams & Kleck, 2005), bilateral amygdala was involved significantly more strongly during the recognition of angry faces with *averted* gaze than angry faces with direct gaze; and iii) functional connectivity results revealed a social-cognitive network, which was connected to bilateral amygdala significantly more strongly during the recognition of angry faces with averted gaze than angry faces with direct gaze. These findings show that the discriminability of facial expressions plays a critical role in the processing of concomitant eye gaze and emotion expressions, and provide novel evidence for a functional amygdala network, which integrates information of eye gaze and emotion of particularly ambiguous stimuli.

During the recognition of angry expressions with direct gaze, the whole-brain analysis showed activity in the insula and dorsal ACC, critical nodes of the salience network. The salience network is known to be important in orienting and allocating cognitive control resources toward subsequent stimulus processing (Barrett & Satpute, 2013) and orienting attention towards them in order to adaptively guide behavior (Menon, 2015). The engagement of the salience network during the recognition of angry expressions suggests that these regions are essential in orienting cognitive resources towards threatening stimuli. Moreover, the engagement of anterior insula during the processing of angry expressions with direct gaze is in line with previous studies that show the involvement of this region in a wide range of cognitive (Desimone & Duncan, 1995; Hopfinger et al., 2000; Menon & Uddin, 2010) and emotional (Lindquist et al., 2012; Ochsner & Gross, 2005) tasks. Anterior insula constitutes a hub of the ventral attentional network, which communicates salient information to other cortical and subcortical networks in order to evaluate and switch between cognitive networks (Menon & Uddin, 2010; Sridharan et al., 2008). It is thus not surprising that the anterior

insula, and in general, the salience network, is engaged more strongly during the processing of angry direct faces in order to orient attentional resources toward a threatening stimulus.

In contrast to recognition of angry emotion with direct gaze, recognition of angry facial expression with averted gaze engaged frontal and parietal regions, as well as bilateral amygdala. This finding is in line with previous findings showing amygdala activity during angry expressions with averted gaze (Adams et al., 2003), but is in contradiction with other studies that showed increased activity of amygdala in response to angry faces with direct gaze (N'Diaye et al., 2009; Sato et al., 2004). It must be acknowledged, however, that the differences in stimulus presentation duration and stimulus set across studies might be contributing to such discrepancies. In order to reconcile these differences across discrepant studies, Adams et al. (2012) conducted several experiments in which different stimulus sets (Ekman faces and NimStim faces) and different presentation durations were employed (1-sec vs. 300-msec), with participants passively viewing the stimuli. Their findings demonstrate that amygdala shows an early, reflexive response toward a clear threat (angry direct gaze), but is more tuned toward ambiguous threat (angry averted gaze) at a later, reflective response. In addition, our study lends support to the notion that task instructions in emotion recognition research are critically important. Using an explicit emotion recognition task as in the present study, we found amygdala to be engaged during recognition of angry averted gaze. This finding supports the idea that amygdala subserves the processing of highly ambiguous signals as conveyed by the combination of angry facial expressions and averted gaze using naturalistic stimuli, such as those from the FACE database.

In addition to the gaze-dependent differentiation of regional activations during the recognition of angry facial expressions, functional connectivity results revealed a large-scale network whose connectivity was significantly stronger during the recognition of angry averted faces than angry direct faces. In addition to bilateral amygdala, this network included



IPL, STS, and medial PFC (mPFC), the critical nodes of social brain network. Activity in STS and mPFC has been reported in a variety of tasks, such as social cognition (Allison et al., 2000), emotion processing and eye gaze (N'Diaye et al., 2009; Pourtois et al., 2004), biological motion perception (Pelphrey & Morris, 2006; Vander Wyk et al., 2009), as well as perspective taking (Gallagher & Frith, 2003; Mitchell et al., 2005; Saxe & Powell, 2006). Thus, we interpret the strong connectivity of STS and mPFC with bilateral amygdala during the recognition of angry averted gaze in line with the idea that averted gaze is ambiguous and may require significantly more inference of the mental state of others than direct gaze does. Therefore, recognition of angry emotions with averted gaze relies on distributed social brain network, which is functionally connected to the amygdala. The identified functional network for angry averted gaze resembles the subnetwork of social brain that has been shown to be involved in detecting socially salient stimuli (Kennedy & Adolphs, 2012). Our results extend these findings and suggest that the functional network connected to the amygdala is strongly involved during recognition of salient, ambiguous, and socially-communicative cues. The connection between brain regions from core (e.g., STS and fusiform gyrus) as well as extended systems (e.g., mPFC, IPL, insula, precuneus, and striatum) and the amygdala indicates the integration of these two systems at higher social-cognitive processes (Haxby & Gobbini, 2011). Therefore, our results extend findings from previous literature by showing that recognizing threat in an ambiguous situation from facial cues relies strongly on the functional network of amygdala. Further studies are required to provide further insight into the changes occur in the functional network of amygdala among psychiatric and neurological illnesses and whether changes in this network are associated with deficits in social functioning among these patients.

Recognition of happy expressions, however, was not modulated by eye-gaze directions at either behavioral or neural levels, in line with previous behavioral studies, which show that

happy facial expressions are insensitive to gaze modulation (N'Diaye et al., 2009; Sander et al., 2007). This finding could be explained in line with the speed-of-processing hypothesis, which states that the distinguished features of happy facial expressions – e.g., teeth showing – make the recognition of happy expression easier and could prevent the interference from the eye regions (Graham & Labar, 2012). Regardless of gaze, we show that recognition of happy expressions engages the critical nodes of the default mode network (DMN; e.g., vmPFC, PCC, precuneus, and STS; Raichle et al. (2001)). DMN is involved in perspective-taking of desire, beliefs, and intentions of others, i.e., processes that are self-referential in nature (Buckner et al., 2008). This network has an extensive connectivity with regions involved in emotion processing (Grimm et al., 2009; Sheline et al., 2009) and is mainly involved when a task demand decreases (Buckner et al., 2008; Mckiernan et al., 2003). This network is also implicated in social function (Kennedy & Adolphs, 2012; Mars et al., 2012); thus, we suggest that recognition of happy expressions may be easier and thus impose lower demands on cognitive resources relative to other conditions, and as a result, may rely more heavily on self-referencing processes subserved by the DMN. Angry expressions however, may require more cognitive effort to a greater extent than happy expressions.

One potential argument for the lack of eye-gaze effect for happy expressions is the distinguished feature of the happy faces, e.g., teeth showing. Such dominant feature potentially could capture participants' attention toward mouth areas relative to eye regions and subsequently result in lack of eye-gaze modulation for happy expression. Although we cannot confirm whether participants were paying more attention to the eye or mouth areas while performing emotion recognition task in the scanner, our eye-tracker data, outside the scanner, showed that there were no significant differences between eye and mouth areas. Therefore, the lack of sensitivity to eye gaze for happy expressions does not seem to be attributable to the amount of time they lingered on eye vs. mouth regions. These findings are

in line with previous work suggesting lack of differences between fixation changes between mouth or eye regions for happy and angry expressions, but not fear (Gamer & Buchel, 2009).

There is a methodological consideration that has to be highlighted here. Participants in this study were asked to identify the emotional expressions of the face rather than gender or intensity ratings. Previous studies that did not find any effect of gaze modulation for happy expressions were used intensity ratings (Cristinzio et al., 2010; Sander et al., 2007) for instance. Task instruction might have an impact on the interplay between eye gaze and emotional expressions. We speculate that asking participants to focus on variant or invariant features of the faces might have differential impact on the recruitment and interaction between core and extended systems (Haxby & Gobbini, 2007). Therefore, future research is required to investigate the impact of different task instruction on the interplay between eye gaze and emotional expressions.

In conclusion, the current study examined the underlying neural mechanisms involved in the recognition of emotional expressions displayed with direct or averted gaze. The findings suggest that the brain activity involved in the recognition of angry expressions is modulated by eye-gaze direction, whereas recognition of happy expressions is not influenced by eye gaze. The results imply that the valence and discriminability of stimuli are critical factors in understanding eye gaze and emotion interaction. Moreover, for the first time, we identified a functional network, which comprises bilateral amygdala and the main nodes of the social-cognitive network, which seem critical to the processing of ambiguous and potentially threatening social signals. These findings provide critical insights into the underlying brain networks involved in processing socially communicative signals, which can be used as biomarkers for further diagnosis of psychiatric and neurological illnesses.

### **Acknowledgment**

Authors would like to acknowledge the valuable practical support provided by the imaging staff at the Centre for Advanced Imaging and valuable comments from the fMRI lab members on the results. This work was supported by Australian Research Council DP1093234 and 5021230-01-414-21-017115. Authors declare no conflict of interest.

For Peer Review

## References

- Adams, R. B., Jr., Franklin, R. G., Jr., Kveraga, K., Ambady, N., Kleck, R. E., Whalen, P. J., Hadjikhani, N., & Nelson, A. J., 2012. Amygdala responses to averted vs direct gaze fear vary as a function of presentation speed. *Soc Cogn Affect Neurosci* 7, 568-577.
- Adams, R. B., Jr., Gordon, H. L., Baird, A. A., Ambady, N., & Kleck, R. E., 2003. Effect of gaze on amygdala sensitivity to anger and fear faces. *Science* 300.
- Adams, R. B., Jr., & Kleck, R. E., 2005. Effects of direct and averted gaze on the perception of facially communicated emotion. *Emotion* 5, 3-11.
- Allison, T., Puce, A., & McCarthy, G., 2000. Social perception from visual cues: role of the STS region. *Trends in Cognitive Science* 4, 267-278.
- Barrett, L. F., & Satpute, A. B., 2013. Large-scale brain networks in affective and social neuroscience: towards an integrative functional architecture of the brain. *Current Opinion in Neurobiology* 23, 361-372.
- Bickart, K. C., Dickerson, B. C., & Barrett, L. F., 2014. The amygdala as a hub in brain networks that support social life. *Neuropsychologia* 63, 235-248.
- Blasi, G., Hariri, A. R., Alce, G., Taurisano, P., Sambataro, F., Das, S., Bertolino, A., Weinberger, D. R., & Mattay, V. S., 2009. Preferential amygdala reactivity to the negative assessment of neutral faces. *Biological psychiatry* 66, 847-853.
- Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L., 2008. The brain's default network: anatomy, function, and relevance to disease. *Ann. N. Y. Acad. Sci.* 1124, 1-38.
- Burns, J., 2006. The social brain hypothesis of schizophrenia. *World psychiatry* 5, 77-81.
- Calder, A. J., & Young, A. W., 2005. Understanding the recognition of facial identity and facial expression. *Nat Rev Neurosci* 6, 641-651.
- Canli, T., Sivers, H., Whitfield, S. L., Gotlib, I. H., & Gabrieli, J. D., 2002. Amygdala response to happy faces as a function of extraversion. *Science* 296, 2191.
- Carlin, J. D., & Calder, A. J., 2013. The neural basis of eye gaze processing. *Current Opinion in Neurobiology* 23, 450-455.
- Cristinzio, C., N'Diaye, K., Seeck, M., Vuilleumier, P., & Sander, D., 2010. Integration of gaze direction and facial expression in patients with unilateral amygdala damage. *Brain* 133, 248-261.
- Desimone, R., & Duncan, J., 1995. Neural mechanisms of selective visual attention. *Annual review of neuroscience* 18, 193-222.
- Ebner, N. C., Riediger, M., & Lindenberger, U., 2010. FACES--a database of facial expressions in young, middle-aged, and older women and men: development and validation. *Behav Res Methods* 42, 351-362.
- Gallagher, H. L., & Frith, C. D., 2003. Functional imaging of 'theory of mind'. *Trends in Cognitive Science* 7, 77-83.
- Gamer, M., & Buchel, C., 2009. Amygdala activation predicts gaze toward fearful eyes. *J Neurosci* 29, 9123-9126.
- Graham, R., & Labar, K. S., 2012. Neurocognitive mechanisms of gaze-expression interactions in face processing and social attention. *Neuropsychologia* 50, 553-566.
- Grimm, S., Boesiger, P., Beck, J., Schuepbach, D., Bermpohl, F., Walter, M., Ernst, J., Hell, D., Boeker, H., & Northoff, G., 2009. Altered negative BOLD responses in the default-mode network during emotion processing in depressed subjects. *Neuropsychopharmacology* 34, 932-843.
- Haxby, J. V., & Gobbini, M. I. (2007). *Distributed neural systems for face perception*.
- Haxby, J. V., & Gobbini, M. I. (2011). *Distributed neural systems for face perception: The Oxford Handbook of Face Perception*.

- Hopfinger, J. B., Buonocore, M. H., & Mangun, G. R., 2000. The neural mechanisms of top-down attentional control. *Nature neuroscience* 3, 284-291.
- Itier, R. J., & Batty, M., 2009. Neural bases of eye and gaze processing: the core of social cognition. *Neurosci Biobehav Rev* 33, 843-863.
- Kennedy, D. P., & Adolphs, R., 2012. The social brain in psychiatric and neurological disorders. *Trends Cogn Sci* 16, 559-572.
- Krishnan, A., Williams, L. J., McIntosh, A. R., & Abdi, H., 2011. Partial Least Squares (PLS) methods for neuroimaging: A tutorial and review. *Neuroimage* 56, 455-475.
- Lindquist, K. A., Wager, T. D., Kober, H., Bliss-Moreau, E., & Barrett, L. F., 2012. The brain basis of emotion: a meta-analytic review. *Behavioral and Brain Sciences* 35, 121-143.
- Mars, R. B., Neubert, F. X., Noonan, M. P., Sallet, J., Toni, I., & Rushworth, M. F., 2012. On the relationship between the "default mode network" and the "social brain". *Front Hum Neurosci* 6, 189.
- McIntosh, A. R., Bookstein, F. L., Haxby, J. V., & Grady, C. L., 1996. Spatial Pattern Analysis of Functional Brain Images Using Partial Least Squares. *Neuroimage* 3, 143-157.
- McIntosh, A. R., Chau, W. K., & Protzner, A. B., 2004. Spatiotemporal analysis of event-related fMRI data using partial least squares. *Neuroimage* 23, 764-775.
- Mckiernan, K. A., Kaufman, J. N., Kucera-Thompson, J., & Binder, J. R., 2003. A parametric manipulation of factors affecting task-induced deactivation in functional neuroimaging. *Journal of Cognitive Neuroscience* 15, 394-408.
- Menon, V., 2015. Saliency Network. 597-611.
- Menon, V., & Uddin, L. Q., 2010. Saliency, switching, attention and control: a network model of insula function. *Brain Structure and Function* 214, 655-667.
- Mitchell, J. P., Banaji, M. R., & MacRae, C. N., 2005. The link between social cognition and self-referential thought in the medial prefrontal cortex. *Journal of Cognitive Neuroscience* 17, 1306-1315.
- N'Diaye, K., Sander, D., & Vuilleumier, P., 2009. Self-relevance processing in the human amygdala: gaze direction, facial expression, and emotion intensity. *Emotion* 9, 798-806.
- Ochsner, K. N., & Gross, J. J., 2005. The cognitive control of emotion. *Trends in Cognitive Science* 9, 242-249.
- Pelphrey, K. A., & Morris, J. P., 2006. Brain Mechanisms for Interpreting the Actions of Others From Biological-Motion Cues. *Curr Dir Psychol Sci* 15, 136-140.
- Pourtois, G., Sander, D., Andres, M., Grandjean, D., Reveret, L., Olivier, E., & Vuilleumier, P., 2004. Dissociable roles of the human somatosensory and superior temporal cortices for processing social face signals. *Eur J Neurosci* 20, 3507-3515.
- Raichle, M. E., MacLeod, A. M., Snyder, A. Z., Powers, W. J., Gusnard, D. A., & Shulman, G. L., 2001. A default mode of brain function. *PNAS* 98, 676-682.
- Sabatinelli, D., Fortune, E. E., Li, Q., Siddiqui, A., Krafft, C., Oliver, W. T., Beck, S., & Jeffries, J., 2011. Emotional perception: meta-analyses of face and natural scene processing. *Neuroimage* 54, 2524-2533.
- Sampson, P. D., Streissguth, A. P., Barr, H. M., & Bookstein, F. L., 1989. Neurobehavioral effects of prenatal alcohol: Part II. Partial Least Squares analysis. *Neurotoxicology and Teratology* 11, 477-491.
- Sander, D., Grandjean, D., Kaiser, S., Wehrle, T., & Scherer, K. R., 2007. Interaction effects of perceived gaze direction and dynamic facial expression: Evidence for appraisal theories of emotion. *European Journal of Cognitive Psychology* 19, 470-480.

- Sato, W., Kochiyama, T., Uono, S., & Yoshikawa, S., 2010. Amygdala integrates emotional expression and gaze direction in response to dynamic facial expressions. *Neuroimage* 50, 1658-1665.
- Sato, W., Yoshikawa, S., Kochiyama, T., & Matsumura, M., 2004. The amygdala processes the emotional significance of facial expressions: an fMRI investigation using the interaction between expression and face direction. *Neuroimage* 22, 1006-1013.
- Saxe, R., & Powell, L. J., 2006. It's the thought that counts: specific brain regions for one component of theory of mind. *Psychological Science* 17, 692-699.
- Sheline, Y. I., Barch, D. M., Price, J. L., Rundle, M. M., Vaishnavi, S. N., Snyder, A. Z., Mintun, M. A., Wang, S., Coalson, R. S., & Raichle, M. E., 2009. The default mode network and self-referential processes in depression. *Proceedings of the National Academy of Sciences* 106, 1942-1947.
- Shepherd, S. V., 2010. Following gaze: gaze-following behavior as a window into social cognition. *Front Integr Neurosci* 4, 5.
- Sridharan, D., Levitin, D. J., & Menon, V., 2008. A critical role for the right fronto-insular cortex in switching between central-executive and default-mode networks. *Proceedings of the National Academy of Sciences* 105, 12569-12574.
- Vander Wyk, B. C., Hudac, C. M., Carter, E. J., Sobel, D. M., & Pelphrey, K. A., 2009. Action understanding in the superior temporal sulcus region. *Psychological Science* 20, 771-777.
- Yu, R. L., & Wu, R. M., 2013. Social brain dysfunctions in patients with Parkinson's disease: a review of theory of mind studies. *Transl Neurodegener* 2, 7.

**Table 1**

Descriptive statistics for the background cognitive measures

<i>Measure</i>	<i>M</i>	<i>SD</i>
<b>NART FSIQ</b>	113.75	3.84
<b>Age</b>	20.65	2.66
<b>RMET</b>	27.47	1.94
<b>Ekman emotion recognition</b>		
<i>Sadness</i>	7.78	1.81
<i>Disgust</i>	7.68	1.56
<i>Happiness</i>	9.60	0.58
<i>Surprise</i>	9.15	1.06
<i>Fear</i>	7.21	2.55
<i>Anger</i>	7.36	1.64
<b>PRSF</b>		
<i>Social Inappropriateness</i>	19.73	4.90
<i>Social Appropriateness</i>	58.10	6.90
<i>Prejudice</i>	6.84	1.06
<b>Empathy Quotient</b>	42.16	10.35
<b>Big Five Inventory</b>		
<i>Extraversion</i>	27.89	6.05
<i>Agreeableness</i>	31.31	3.41
<i>Conscientiousness</i>	30.78	5.66
<i>Neuroticism</i>	21.10	6.17
<i>Openness</i>	33.36	6.29
<b>Eye-tracker task</b>		
<i>Angry_ Eye region</i>	340.87	79.03
<i>Angry_ Mouth region</i>	355.98	145.96
<i>Happy_ Eye region</i>	306.64	53.33
<i>Happy_ Mouth region</i>	388.24	196.20



---

*Note.* NART FSIQ = National Adult Reading Test Full-Scale Intelligence Quotient, RMET = Reading the Mind in the Eye Test, PRSF = Peer-Report Social Functioning Scale.

For Peer Review

**Table 2**

Regions from LV1 of whole-brain analysis showing increased activity for angry facial expression with averted gaze vs. all other conditions

Regions	Hem	BA	MNI coordinates		BSR
			XYZ		
Medial Frontal Gyrus	L	6	[0 2 56]		6.22
Superior Frontal Gyrus	L	9	[-36 50 24]		5.03
	R	9	[38 50 26]		3.74
Inferior Frontal Gyrus	R	46	[52 36 10]		7.40
	L	9	[-60 10 24]		6.67
Anterior Cingulate Gyrus	L	32	[-2 14 40]		4.92
Superior Temporal Gyrus	L	22	[-56 6 2]		6.67
Inferior Parietal Lobe	L	40	[-56 -16 26]		5.52
	R	40	[64 -32 26]		4.02
Precentral Gyrus	L	43	[-54 -2 10]		6.13
	R	44	[56 10 0]		6.04
Postcentral Gyrus	R	3	[58 -12 28]		6.25
	L	3	[-44 -14 58]		6.82
Posterior Cingulate Gyrus	L	23	[-2 28 28]		4.59
Posterior Cingulate Gyrus	R	30	[12 -60 6]		5.95
	L	30	[-12 -68 8]		4.58
Middle Occipital Gyrus	R	18	[32 -86 -2]		8.39
	L	19	[-34 -88 4]		6.03
Cuneus	L	23	[-6 -72 12]		4.97
Insula	L	13	[-46 -2 4]		6.35
	R	13	[48 6 0]		6.1
Putamen	L		[-28 -2 10]		5.4
Thalamus	L		[-8 -20 10]		4.93
	R		[10 -14 10]		4.22
Amygdala	L		[-18 -4 -12]		5.34
	R		[20 -8 -12]		3.19
Cerebellum	L		[-25 -70 -15]		4.57

R

[36 -55 -15]

5.88

---

*Note.* BSR = Bootstrap Ratio,  $BSR > 2.5$ ,  $p < .005$ ; Hem = Hemisphere; R = right; L = left; BA = Brodmann Areas; x coordinate = right/ left; y coordinate = anterior/posterior; z coordinate = superior/inferior.

For Peer Review

**Table 3**

Regions from LV2 of whole-brain analysis showing increased activity for angry facial expressions with *direct* gaze and happy facial expressions with both direct and averted gaze relative to the other conditions

Regions	Hem	BA	MNI coordinates	BSR
			XYZ	
<b>Angry facial expression (direct gaze) &gt; happy facial expressions</b>				
Superior Frontal Gyrus	R	6	[4 20 46]	4.99
Anterior Cingulate Gyrus	R	32	[0 22 39]	3.94
Middle Temporal Gyrus	R	37	[46 -62 0]	4.69
Superior Parietal Lobe	R	7	[26 -58 47]	6.32
Occipital Gyrus	R	19	[36 -80 0]	3.7
	L	19	[-46 -76 -4]	4.8
Insula	R	13	[44 20 2]	4.6
	L	13	[-42 14 2]	4.8
Amygdala	L		[-24 -12 -15]	3.90
Putamen	R		[24 6 6]	3.64
	L		[-26 2 6]	3.73
Cerebellum	R		[40 -68 -8]	4.65
	L		[-46 -76 -6]	5.56
<b>Happy facial expression (direct &amp; averted gaze) &gt; angry facial expressions</b>				
Middle Frontal Gyrus	L	6	[-26 2 48]	5.31
Superior Frontal Gyrus	R	8	[26 30 46]	3.68
	L	8	[-24 34 46]	3.7
Medial Frontal Gyrus	L	32	[-6 16 48]	5.82
Anterior Cingulate Gyrus	R	24	[2 30 -14]	4.51
	L	32	[-10 42 -6]	5.78
Middle Temporal Gyrus	R	39	[46 -70 30]	3.65
	L	39	[-48 -70 30]	4.23
Middle Temporal Gyrus	L	21/	[-58 -32 2]	8.51

22

Superior Parietal Lobe	R	7	[28 -56 62]	5.18
	L	7	[-26 -64 56]	5.56
Inferior Parietal Lobe	L	7	[-32 -52 48]	4.73
	L	40	[-50 -36 48]	5.18
Posterior Cingulate Gyrus	L	31	[-8 -34 46]	3.77
Precuneus	L	31	[-12 -62 24]	4.58

---

*Note.* BSR = Bootstrap Ratio,  $BSR > 2.5$ ,  $p < .005$ ; Hem = Hemisphere; R = right; L = left;

BA = Brodmann Areas; x coordinate = right/ left; y coordinate = anterior/posterior; z coordinate = superior/inferior.

**Table 4**

Regions from functional connectivity with bilateral amygdala for angry facial expressions

Regions	Hem	BA	MNI coordinates	BSR
			XYZ	
Middle Frontal Gyrus	R	46	[50 42 10]	7.68
	L	10/46	[-42 48 12]	5.17
Inferior Frontal Gyrus	R	44	[62 12 18]	5.92
Superior Frontal Gyrus	L	8	[-2 42 44]	5.71
	R	6	[5 12 56]	8.55
Middle Frontal Gyrus	L	9	[-43 26 27]	8.26
	L	8	[-26 26 46]	5.13
Anterior Cingulate Gyrus	R	24	[22 16 48]	9.22
Precentral Gyrus	R	6	[62 2 10]	7.33
	L	4	[-56 -2 18]	10.91
Superior Temporal Gyrus	R	22	[38 -52 16]	6.44
	L	38	[-43 10 -31]	8.55
Inferior Parietal Lobe	L	40	[-40 -50 54]	9.57
Postcentral Gyrus	R	40/43	[60 -18 18]	4.39
Posterior Cingulate Gyrus	L	23	[-2 28 28]	4.59
	R	31	[12 -62 18]	5.93
Fusiform Gyrus	R	37	[41 -59 -15]	5.47
Cuneus	R	18	[18 -68 18]	6.72
Precuneus	L	31	[-14 -66 18]	5.31
Caudate	L		[14 4 18]	9.33
Putamen	R		[-28 -2 10]	5.87
Thalamus	L		[-14 -28 12]	7.03
	R		[10 -14 10]	4.16
Cerebellum	L		[-16 -68 -15]	7.96
	R		[10 -63 -15]	6.26
Amygdala	R		[20 -8 -14]	19.33
	L		[-18 -4 -12]	12.33

*Note.* BSR = Bootstrap Ratio,  $BSR > 2.5$ ,  $p < .005$ ; Hem = Hemisphere; R = right; L = left; BA = Brodmann Areas; x coordinate = right/ left; y coordinate = anterior/posterior; z coordinate = superior/inferior.

For Peer Review

## Figure Legends

**Fig. 1. Behavioral Results.** Behavioral results from emotion recognition task in the scanner. Participants were faster and more accurate for recognizing happy expressions relative to angry expressions. Bars represent 1 standard error of the mean (SEM).

**Fig. 2. Whole-Brain Results.** Patterns of whole-brain activity during the recognition of angry expressions with averted gaze (A), angry expression with direct gaze (B), and happy expression with direct and averted gaze (C), relative to the other conditions. Error bars denote 95% confidence intervals for the correlations calculated from the bootstrap procedure. All reported regions have  $BSR \geq 2.5$  and cluster size  $\geq 100$  voxels. L = left hemisphere, R = right hemisphere.

**Fig. 3. BOLD Signal Intensity in Bilateral Amygdala.** Peak voxel intensity of left (-18 -4 -12) and right (20 -8 -12) amygdala during the four experimental conditions within 12-sec after stimulus onset.

**Fig. 4. Functional Connectivity Results.** (A) The functional network connected to bilateral amygdala during the angry conditions. (B) Correlations between activity in bilateral amygdala and the functional network during the angry conditions. Error bars denote 95% confidence intervals for the correlations calculated from the bootstrap procedure. Brain/correlation scores were considered unreliable when CIs crossing zero and considered significantly different if CIs do not overlap. All reported regions have  $BSR \geq 2.5$  and cluster size  $\geq 100$  voxels.



Fig. 1




Fig. 2




Fig. 3




Fig. 4

