



ISSN: 2231-6302

# A comprehensive review of ethnobotanical plants used by the people of Pir Panjal Range in (Jammu Division) Union Territory Himalaya of Jammu and Kashmir- India

Tahir Mahmood\*

Centre for Biodiversity Studies, Baba Ghulam Shah Badshah University Rajouri 185234

## ABSTRACT

The current paper provides a taxonomic inventory of the medicinal plant species collected by the author during the last one decade from Pir Panjal range in (Jammu Division) Himalaya of Union territory Jammu & Kashmir- India. The inventory records a total no of 76 medicinal species belonging to 45 families of the total taxa were recorded for the medicinal Purposes. The inventory is expected to provide baseline scientific data for further studies on plant diversity in Jammu division and can be used to facilitate the long-term conservation and sustainable use of medicinal plant resources in the Himalaya region, and among all the families Cucurbitaceae and Euphorbiaceae were found to be most dominant families in term of the species in the areas with 06 species, followed by Polygonaceae and Rosaceae.

**KEYWORDS:** Ethnobotanical, Medicinal Plants, Jammu division, Pir Panjal, Himalaya, J&K-India

**Received:** May 08, 2020  
**Accepted:** June 17, 2020  
**Published:** June 23, 2020

**\*Corresponding Author:**  
Tahir Mahmood,  
Email: [tahirchoudhary786@gmail.com](mailto:tahirchoudhary786@gmail.com)

## INTRODUCTION

Plants are remarkable source of valuable substances for human beings. These are showing variation in their habitat as well as their habit. As per climatic condition, the plants are showing their presence in different sites. Plants are essential for healthier life because they provide us medicines, which are both effective and safe, without any side effect. Some pathological conditions in human being that could not be fully treated by conventional pharmaceutical are numerous [1] for this reason, there is a growing tendency in use of herbal preparations. The world health organization (WHO) estimates that 80% of the world population depends on plants remedies for its primary health care needs [2]. The local peoples of the rural areas have good knowledge about the uses of plants and they prefer medicinal plants due to their easy availability and cheap therapy as compared to costly pharmaceuticals. The traditional Practioners are playing an important role in providing health coverage to 75% of the population residing in villages and rural areas. Maximum 76% rural peoples depend on forest products for fulfilling their daily needs. India ethnobotanical work has been done in the past [3-5]. while in all these studies qualitative approaches have been adapted to document ethnobotanical information [6]. Ethnobotany of Jammu division Union

territory, Jammu & Kashmir is getting various studies have been reported from various parts of the areas [7-15]. While in contrast, ethnobotanical research has been somewhat neglected in the south foot hilly areas of district rajouri province Jammu particularly. In province Jammu, few studies were carried out by some Scholars and Scientists in the past [16] conducted research Flora of Jammu and Plants of Neighborhood Bishen, Flora of upper Liddar Valleys of Kashmir Himalaya. Ethnobotanical study of useful climbers creepers and twiners of Baba Ghulam Shah Badshah University campus and adjoining areas of district rajouri Jammu and Kashmir [10]. Ethnobotany of medicinal plants in district Mastung of Balochistan province –Pakistan [17]. The present study can be considered as the first time and one which deals with an ethnobotanical study on medicinal plants in this region. Jammu division has also got importance for its topography as well, inside having high mountains, with desert habitats and having high rich diversity of medicinal plants. The rural areas of the Jammu division still depend on these wild plants for cure the disease and having a good ethnobotanical knowledge about medicinal plants. but currently the ethnobotanical knowledge is disappearing very fast from the urban areas of the Jammu division because of being closer to and bounded with the capital city of province Jammu' having health and other facilities.

Copyright: © The authors. This article is open access and licensed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted, use, distribution and reproduction in any medium, or format for any purpose, even commercially provided the work is properly cited. Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.

The aim of the current research is to highlight the key of medicinal plants in Jammu division of Pir Panjal range Himalaya of Union territory Jammu & Kashmir- India.

The aim of this study was to document ethnomedicinal uses of plants and analyzed ethnobotanical information using quantitative indices of information consent factor (ICF), fidelity level (FL), use value (UV), use report (UR) frequency citation (FC) and relative frequency citation (RFC).

## MATERIAL AND METHODS

### Jammu Division Geo-ethnographical Overview

Jammu division has an area of 26.64 km<sup>2</sup> with ten districts. Jammu, Doda, Kathua, Ramban, Reasi, Kishtwar, Poonch, Rajouri, Udhampur and Samba, Union territory: Jammu and Kashmir (Figure 1). According to the census 2011, the total population of Jammu Division is 5,350,811. Its lie between... 18', East longitude and 32 degree 50' and 33 degree 30' North latitude. The Jammu division presents a composite culture Pahari, Gojri, Dogri and Kashmiri. Irrespective of ethnic groups all speak the pahari language with easily. The climate varies from semi-tropical in the souther part to temperate in the mountainous northen part. The sub-tropical region receives regular monsoons whereas the northen part prone to hailstorms experiences excessive rains. The Jammu division is drained by small and big rivers. Some of the tribal peoples annually migrate during winter from higher altitudes to lower, During the summer from lower to higher altitudes with their families along with Cattles ( Sheeps, Goats, Horses etc.,) Migration to other countries is 14.9% for their bread and butter of all migrants. Migration starts in April ending and continues till June. The migrants return from September and continue till November.

### Socio-economic Condition of the Area

Jammu division is the major earning means of the peoples in the region. Nearly 57% of the population of Jammu division depends on agricultures. Important cultivated plants are wheat, maize, potato, onion, and other vegetables. Some of the local inhabitants collect medicinal plants from forests, deserts, mountains and plains and sell them to the local traditional herbs sellers in very cheap prices. Local traditional herbs sellers then supply these plants to the pharmaceutical companies in good prices. The Jammu division has been released with diverse flora included a great numbers of medicinal plants. The rural areas of the division are still dependent on medicinal plants for their health care because of lack health centers in the area. If the sustainable use of wild flora and cultivation of medicinal plants are promoted in the area, this will strongly affect on the socio-economic condition of the local inhabitants.

### Field Interviews

For the study and documentation of medicinal plants, intensive exploration trips were conducted about twelve months from January 2019 to February 2020. The questionnaire was mainly focused on the ethnobotanical claims and traditional believes of local communities and nearby peoples. The interviews were conducted using the local languages that are Phari, Gojri, Urdu, as the first author is a local person of the region. for the ethnobotanical information, a total no 197 inhabitants of the area were interviewed. 86 women, 99 men, and 12, traditional healers were interviewed. The informants were divided into three different age of groups i.e. 22-42-43-63-64-84- years old. All the informal meetings were held 26 different villages of the district rajouri province Jammu i.e. Kurhad, Prori, darhal, Khorwalli, Palma, Bakori, Budhal, Soaker. Saj. DK, Dodaj, Rehan, Hobby, Kandi, Jari-wali, Basholi. Udhampur. Reasi.




Figure 1: Map of India showing Union Territory Jammu, Kashmir Ladakh. Map showing Jammu division Union territory of Jammu & Kashmir, India

## Collection, Identification and Deposition of Medicinal Plants

The plants were collected during twelve months (January-2019-to Faburary-2020).The Jammu division covering almost all the seasons of the year and from all the parts of districts. The collected plants specimens were dired and preserved processed as per routine herbarium techniques recommended by Jain and Rao [18] for reconfirmation of plants identification, the flora of Flowers of the Himalaya [19]. Exotic Ornamental Flora of Kashmir [20] Flora of British India [21] and Flora of Jammu and Kashmir. Vouchers specimens were deposited in the herbarium, Centre for Biodiversity Baba Ghulam Shah Badshah University rajouri for futures references.

## Quantitative Analysis of Ethnobotanical Results

The data collected was analysis using quantitative value indices.

### Information consensus factor (ICF)

Information consensus factor (ICF) was obtained [22] using the following formula;

$$ICF = (N_{ur} - N_t) / (N_{ur} - 1)$$

Where  $N_{ur}$  refers to the total number of uses reports for each disease category and  $N_t$  it is the number of taxa used in that category. It used to test the homogeneity of knowledge on the uses of species in the illness categories between the populations. The ICF provides a range of (0-1). High ICF shows that there is a narrow well-defined group of species used to cure a particular ailment category and/or that information is exchanged between informants and low ICF values (close to zero) indicate that informants disagree over which plant to use due to random choosing or lack of exchange of information about the use among informants [23].

### Fidelity level (FL)

Fidelity level (FL) index was calculated by using the following formula as described by Friedman *et al.* [24] to determine the most preferred species used in the treatment in the same category:

$$FL = (N_p / N) 100$$

Where  $N_p$  is the number of informants citing the use of the plant for a particular illness and  $N$  is the total numbers of informants citing the species for any illness. High FL value indicates high frequency of use of the plant species for treating a particular ailment category by the informants of the study area.

### Frequency citation (FC) and relative frequency citation (RFC)

The FC of the species of plants being utilized was evaluated using the formula:  $FC = (\text{Number of times a particular species was mentioned} / \text{total number of times that all the species were$

mentioned) 100 and the relative frequency citation (RFC) index by using the following formula:

$$RFC = FC / N \quad (0 \text{ RFC } 1)$$

The index is obtained by dividing the number of informants mentioning a useful species FC or frequency of citation by the total number of informants in the survey (N). RFC value varies from 0 (when nobody refers to plants as a useful one). to 1 (when all the informants mention it as useful). RFC index, which does not consider the use-category (UR or use-report it is a single record for use of a plant mentioned by an individual).

### Use value (UV) and use report (UR)

$$UV = \sum U/n$$

Where UV is the use value of a species, 'U' is the number of use reports cited by each informants for a given plant species and 'n' is the total number of informants interviewed for a given plant. The UV is applied in determining the plants with the highest use (most frequently indicated) in the treatment of an ailment, while use report (UR) is the use recorded for every species,

## RESULTS AND DISCUSSION

### Use of Plants and Demography

A total no of 197 inhabitants of the Jammu division (40% women, (30%) men and (09%) men traditional healers of different age of groups were interviewed. The informants were divided into three different ages of groups. Most of the informants above belonged to the age of 60 year (Fig.2) and many informants were categorized (Table 1) in total 45 families and 76 species with local name of the plants, family




Table 1: All informants are categorized

S.no	Category	No of peoples Interviewed
1.	Men	86
2.	Women	99
3.	Traditional healers	12
Total		197

Table 2: List of medicinal plant used by local people of Jammu division Union territory Jammu &amp; Kashmir

S.no	Botanical name	Local name	Voucher number	Family	Life form	Part used	Disease treated	Preparation mode(s)	FC*	RFC*	UR*	UV*
1	<i>Abelmoschus moschatus</i> Medik	Ban-bar	CBS-117	Malvaceae	Herb	Roots	Chest pain,	Juice-Raw	12	0.05	3	0.11
2	<i>Allium cepa</i> L.	Payaz	CBS-119	Alliaceae	Herb	Bulb	Pimples, Skin infection	Roasted	14	0.068	3	0.26
3	<i>Aloe vera</i> L	<i>Aloe vera</i>	CBS-121	Liliaceae	Herb	Leaves	Jaundice	Juice	9	0.04	1	0.11
4	<i>Ajuga bracteosa</i> L	Kauri booti	CBS-122	Lamiaceae	Herb	Leaves	Jaundice	Decoction	12	0.054	2	0.16
5	<i>Adiantum Capillus veneris</i> L.	Gauthier	CBS-1.23	Adiantaceae	Herb	Whole plant	Herpes	Sap	4	0.017	2	
6	<i>Asparagus racemosus</i> Willd	Sanspai	CBS-124	Liliaceae.	Climber.	Roots	Constipation, Stomachache	Decoction/ Infusion	21	0.1	2	0.08
7	<i>Achryanthus aspera</i> L	Phut kando	CBS-125	Amaranthaceae	Herb	Root	Jaundice, Constipation	Powder	8	0.04	1	0.12
8	<i>Bergenia ciliata</i> Haw	Zakham-ayath	CBS-126	Saxifragaceae	Herb	Root, Leaves	Wound healing	Decoction/ Paste	0.1	2	0.09	
9	<i>Berberis lycium</i> Royle	Simloo	CBS-127	Berberidaceae	Shrub	Roots	Jaundice, Wounds, Back pain	Powder / Decoction	39	0.173	5	0.12
10	<i>Buddleja asiatica</i> Lour	Batti	CBS-128	Scrophulariaceae	Shrub	Leaves	Skin disease	Juice	17	0.08	3	0.18
11	<i>Brassica campestris</i> L	Sarson	CBS-129	Brassicaceae	Herb	Seed Leaves	Skin,	Cooked/Oil	16	0.072	1	0.06
12	<i>Cannabis Sativa</i> L	Bhang	CBS-130	Cannabaceae	Herb	Seeds, Leaves	Piles, Hallucination.	Powder / Decoction	0.05	2	0.23	
13	<i>Cardiospermum halicacabum</i> L.	Qulqul	CBS-131	Sapindaceae	Climber	Leaves, Stem	Swelling, Snakebite, Rheumatism	Juice/Powder	7	0.032	2	0.27
14	<i>Coriandrum sativum</i> L	Dhania	CBS-132	Apiaceae	Herb	Whole plant	Spice, Diabetes, Ulcers	Powder, Decoction	38	0.172	4	0.11
15	<i>Calotropis procera</i> Aiton. D	Aak	CBS-212	Apocynaceae	Shrub	Leaves	Chest pain, Fever	Powder	9	0.04	1	0.11
16	<i>Cuscuta reflexa</i> Roxb	Neela dhari	CBS-213	Cuscutaceae	Climber	Stem, Leaves	Rheumatic pain, Dandruff	Decoction	25	0.11	2	0.08
17	<i>Cinnamomum tamala</i> Buch	Dalchini.	CBS-214	Lauraceae	Tree	Whole plant	Constant weight, Swelling	Decoction / Powder	10	0.05	2	0.23
18	<i>Cucumis Sativus</i> L	Kakri-kheera	CBS-215	Curcubitaceae	Climber	Fruit	Diuretic, haemostatic	Infusion	9	0.04	1	0.11
19	<i>Carica papaya</i> L	Pappetaa	CBS-216	Caricaceae	Tree,	Fruit	Worm infestation, constipation	Juice/ Powder.	10	0.05	2	0.23
20	<i>Cedrus deodara</i> Roxb	Deodar	CBS-217	Pinaceae	Tree		Wounds		22	0.1	3	0.14
21	<i>Cynodon dactylon</i> Linn	Khabbal	CBS-218	Poaceae	Herb	Whole plant	Swelling, Sprains	Raw. Paste	13	0.06	2	0.15
22	<i>Catharanthus roseus</i> L	Sada bahar	CBS-219	Apocynaceae	Shrub	Leaves	Diabetes, Malaria	Juice	18	0.08	2	0.11
23	<i>Cassia fistula</i> L	Amaltas	CBS-220	Fabaceae	Tree	Whole plant	Pulp of pods	Powder	13	0.06	2	0.15
24	<i>Cyperus rotundus</i> L.	Nut grass	CBS-221	Cyperaceae	Herb	Root	Nausea, Fever, Inflammation	Juice/Decoction	15	0.07	2	0.13
25	<i>Daucus carota</i> L	Gaajir	CBS-222	Apiaceae	Herb	Root	Fairness of skin	Roots juice	11	0.05	1	0.09
26	<i>Equisetum Debile</i> Roxb	Tarutkaah	CBS-307	Equisetaceae	Herb	Whole plant	Diuretic, hand burning	Juice powder	22	0.1	2	0.09
27	<i>Euphorbia wallichii</i>	Hirbi	CBS-308	Euphorbiaceae	Herb	Milk	Severe constipation	Juice	4	0.02	1	0.25
28	<i>Elaeagnus umbellata</i> Burn	Kankoli	CBS-309	Elaeagnaceae	Shrub	Fruit	Mouth sores	Fruit	12	0.05	2	0.17
29	<i>Euphorbia helioscopia</i> L	Doudal	CBS-310	Euphorbiaceae	Shrub	Leaves	Anti-allergic	Milk/Paste	8	0.04	1	0.12
30	<i>Euphorbia hirta</i> L	Cat hair	CBS-311	Euphorbiaceae	Shrub	Leaves	Pathogenic bacteria	Powder/Infusion	19	0.09	2	0.10
31	<i>Foeniculum vulgare</i> Mill	Sounf	CBS-312	Apiaceae	Herb	Fruit Leaves	Hypertension, Breast tissues	Powder	13	0.06	2	0.15

(Contd...)

Table 2: (Continued)


S.no	Botanical name	Local name	Voucher number	Family	Life form	Part used	Disease treated	Preparation mode(s)	FC*	RFC*	UR*	UV*
32	<i>Iris domestica</i> L	Reach jaari	CBS-313	Iridaceae	Herb	Fruit	Asthma, Throat troubles	Powder/Decoction	10	0.05	1	0.1
33	<i>Justicia adhatoda</i> L	Baykar	CBS-314	Acanthaceae	Shrub	Leaves	Swelling	Juice	7	0.032	1	0.14
34	<i>Jasminum officinal</i> Linn	Chameli	CBS-315	Oleaceae	Climber	Fruit	Ringworm, Narcotic	Decoction	8	0.04	1	0.12
35	<i>Luffa cylindrica</i> L	Jungli lokii	CBS-316	Curcubitaceae	Climber	Fruits	Diuretic, Splenopathy	Cooked/Juice	29	0.13	2	0.06
36	<i>Litsea glutinosa</i> Lour	Medha sak	CBS-317	Lauraceae	Tree	Bark	Aphrodisiac, Sprains, Fracture.	Bark, Powder	23	0.1	1	0.04
37	<i>Lathyrus aphaca</i> L.	Jungli mutter	CBS-318	Fabaceae	Herb	Seeds	Toothache, Narcotic	Powder. Cooked	28	0.12	1	0.03
38	<i>Lathyrus sativus</i> L	Phali	CBS-319	Fabaceae	Climber	Seeds	Produces protein	Cooked	20	0.09	2	0.1
39	<i>Mallotus philippensis</i> L	Kamila	CBS-320	Euphorbiaceae	Shrub	Fruit, Bark	Dyeing silk, Wool	Powder	16	0.072	4	0.25
40	<i>Momordica chararita</i>	Kerala	CBS-107	Cucurbitaceae	Climber	Fruit	Ulcers, Diabetes.	Juice/Decoction	19	0.09	2	0.10
41	<i>Marsilea quadrifolia</i> L	Khatri	CBS-108	Marsileaceae	Climber	Leaves	Stomach worms Snake bites, Skin injuries	Milk, Decoction	7	0.032	1	0.14
42	<i>Mentha arvensis</i> L	Podina	CBS-109	Lamiaceae	Herb	Whole plant	Stomach pain	Powder	38	0.17	5	0.13
43	<i>Morus alba</i> L	Thooth	CBS-110	Moraceae	Tree	Leaves Fruit	Purgative, Toothache	Decoction	9	0.04	1	0.11
44	<i>Melia azedaranch</i> L	Deralik	CBS-111	Meliaceae	Tree	Leaves Fruit	Wound, burning of hands and feet	Infusion Paste	27	0.12	3	0.11
45	<i>Musa paradisiaca</i> L Stones, Flushing urinary blocks	Kaila	CBS-112	Musaceae	Shrub	Fruit	Dissolving kidney, Urinary bladder	Food	28	0.12	2	0.07
46	<i>Nerium indicum</i> Mill	Gandilo	CBS-401	Apocynaceae	Shrub	Flowers, Root	Scabies, Ringworm	paste /Juice	11	0.05	3	0.27
47	<i>Oxalis corniculata</i> L	Desi Shutal	CBS-402	Oxalidaceae	Climber	Leaves	Fractured bones, Purify blood	Raw leaves	11	0.05	3	0.27
48	<i>Olea ferruginea</i> Royle	Khahoo	CBS-403	Oleaceae	Tree	leaves	Mouth, Toothache,	Decoction	10	0.05	1	0.1
49	<i>Punica granatum</i> L	Daruna	CBS-404	Puniaceae	Tree	Fruits Bark	Jaundice, Diabetic, Syphillis.	Juice/powder	22	0.1	3	0.14
50	<i>Persicaria amplexicaulis</i> (D. Don)	Masloon	CBS-405	Polygonaceae	Herb	Root	fever, Pain.	Decoction, Juice	23	0.1	1	0.04
51	<i>Phyllanthus emblica</i> L	Aamia	CBS-406	Phyllanthaceae	Tree	Fruits	Bleeding / Cough.	Fruits / Powder	15	0.07	1	0.06
52	<i>Pyrus pashia</i> Buch	Batangi	CBS-407	Rosaceae	Shrub	Fruit	Diarrhoea, Constipation	Juice	22	0.1	2	0.09
53	<i>Rosa moschata</i> Herrm	Phalwari	CBS-408	Rosaceae	Climber	Roots, Flower	Aphrodisiac, Digestive	Cooked/Powder	8	0.04	1	0.12
54	<i>Rubus ellipticus</i> Sm	Gurcho	CBS-409	Rosaceae	Shrub	Fruit	Cooling effect	Raw Fruits	8	0.03	1	0.12
55	<i>Rosa indica</i> L	Jungli gulab	CBS-410	Rosaceae	Shrub	Flower	Eye infection, Constipation	Raw petals, Decoction	18	0.08	4	0.22
56	<i>Rumex patentia</i> L	Hulla	CBS-411	Polygonaceae	Herb	Leaves-Bark	Constipation, Tumors	Paste/ Roasted	13	0.06	2	0.15
57	<i>Rumex hastatus</i> D. Don	Khatti buti	CBS-412	Polygonaceae	Herb	Leaves	Wounds	Paste/ Decoction	16	0.072	1	0.06
58	<i>Ricinus communis</i> L	Daalda butoe	CBS-413	Euphorbiaceae	Herb	Seeds	Breast tumours,	Juice /Infusion	14	0.063	1	0.07
56	<i>Raphanus Sativus</i> L	Mulli	CBS-414	Brassicaceae	Herb	leaves	Diuretic, Digestive	Juice	18	0.08	3	0.17
60	<i>Solanum melongena</i> L	Pathaa	CBS-415	Solanaceae	Herb	Roots, Leaves	Ulcer ,Nose, Narcotic	Juice/Paste	9	0.04	2	0.22
61	<i>Solanum surattense</i> Dunal	Mookri	CBS-416	Solanaceae	Herb	Leaves	Warms, Dandruff.	Sap/	12	0.05	2	0.15

(Contd...)



Table 2: (Continued)

S.no	Botanical name	Local name	Voucher number	Family	Life form	Part used	Disease treated	Preparation mode(s)	FC*	RFC*	UR*	UV*
62	<i>Skimmia lauroala</i> Franch	Patlo	CBS-417	Rutaceae	Herb	Leaves	Purify the air, Aeromatic	Fog	15	0.07	1	0.06
63	<i>Solanum Nigrum</i> L	Kaach maach	CBS-418	Solanaceae	Herb	Seeds, Leaves	Throat pain, Toothache	Juice /Powder	25	0.11	2	0.08
64	<i>Solena amplexicaulis</i> (Lam)Gandhi	Bun kereli	CBS-419	Cucurbitaceae	Creeper	Roots,Leaves	Invigorating, Astringent,	Cooked	29	0.13	1	0.03
65	<i>Trichosanthes cucumerina</i> L	Khaakri	CBS-420	Cucurbitaceae	Creeper	Fruits	Jaundice/ Liver, Digestive	Cooked/Juice	12	0.05	1	0.08
66	<i>Taraxacum officinale</i> Willd	Hund	CBS-503	Asteraceae	Herb	Whole plant	Delivery, Dandelion wine	Cooked/	23	0.1	1	0.04
67	<i>Tinospora cordifolia</i> Willd	Gulanacha	CBS-504	Menispermac	Climber		Diabetes,Allergic rhinitis,Cancer	Cooked	13	0.06	2	0.15
68	<i>Typha latifolia</i> L	Cat-tail	CBS-505	Typhaceae	Herb	Leaves	Boils, Burns, Wounds	Decoction	7	0.032	1	0.14
69	<i>Vitex Negundo</i> L	Banna	CBS-506	Verbenaceae	Shrub	Leaves	Earache,Wound	Decoction	15	0.07	1	0.06
70	<i>Vitis jacquemontii</i> L	Daakh	CBS-509	Vitaceae	Climber	Leaves	Skin disease, Chest Pan.	Sap	18	0.08	2	0.11
71	<i>Sapindus mukorossi</i> L	Raetha	CBS-511	Sapindaceae	Tree	Fruit, Leaves	Asthma, Diarrhea, Cholera	Infusion	9	0.04	1	0.11
72	<i>Zanthoxylem armatum</i> DC	Timber	CBS-512	Rutaceae	Shrub	Fruit, Bark	Blood pressure, Stimulation	Gum	23	0.1	1	0.04
73	<i>Ziziphus mauritiana</i> Lam	Beri	CBS-513	Rhamnaceae	Tree	Fruits, Seeds	Fever, Ulcers, Cephalalgia	Decoction/Powder	8	0.03	1	0.13
74	<i>Ziziphus oxyphylla</i> Edgew	Cocon beri	CBS-514	Rhamnaceae	Tree	Fruits, Seeds	Constipation, Fever	Decoction/Powder	8	0.03	1	0.13
75	<i>Zea mays</i> L	Maak	CBS-515	Poaceae	Herb	Maize starch	kidney stones	Juice	29	0.13	2	0.06
76	<i>Zingiber officinale</i> Roscoe	Adrak	CBS-069	Zingiberaceae	Herb	Tuber	Common Spice	Powder	19	0.09	2	0.1



**Figure 2:** Distribution of gender, age and number of informants interviewed


name their uses and parts of the plants used for their medicinal values, use report (UR) use value, (UV) frequency citation (FC) and relative frequency citation (RFC) are listed in Table 2. The best represented used families in terms of the number of species are Cucurbitaceae (6 species), Euphorbiaceae, Rosaceae, Polygonaceae, Apiaceae, Apocynaceae, with 7 species each (Table 3) the most common part of the plants used are their leaves and whole plant (19%) each (Fig. 3) plant are often used as decoction (29%) and a small portion is also used roasted, juice and soups. Highest plants species are used in the treatment of gastrointestinal disease (21 species) Moreover a single plant is used for the more than one disease for example, *Mentha arvensis* (Stomach pain) *Luffa cylindrica* (Diuretic, Splenopathy) *Zanthoxylum armatum* (Blood pressure, Stimulation) *Berberis lyceum* Royle (Jaundice, Wounds, Back pain) Highest ICF value (1) was recorded for antidote category. 100% fidelity level was found for four plant species i.e. *Zea mays*, *Pyrus pashia*, *Musa paradisiacal*, and *Momordica charanita*. The highest use value was reported for the *Litsea glutinosa* (0.6). Highest RFC value was calculated for *Berberis lyceum*, *Coriandrum sativum*, (0.23). and other five uses reports for each in Table 3.

The results of the study showed that Cucurbitaceae is the largest medicinal plant family. The values and characteristics of family, Cucurbitaceae as a Predominant in this area, among all the families Cucurbitaceae and Euphorbiaceae were found to be most dominant families in term of the species in the area with 06 species, followed by Polygonaceae and Rosaceae.

### Herbal Drug Preparation Method

Among herbal drug preparation, decoction (21%) with 29 species). And infusion (17% with 28 species) (Fig. 3) are highly used in the area.

The result of wide spread use of decoction and infusion agree with the results of Gurdal and Kultur [25] and Ahmed *et al.* [26] who reported that decoction was the most commonly used



**Table: 3** Most used families of the study area

Family name	Number of taxa
Cucurbitaceae	6
Euphorbiaceae	5
Rosaceae	4
Polygonaceae	3
Apiaceae	3
Apocynaceae	3
Total	24

preparation method. Followed by infusion. In the study eight internal application methods were recorded i.e. decoction, infusion, powder, raw, cooked, tea, soup and juice have been used internally. The gum was used as chewing gum and 4 direct external application methods like milk/sap, steam, smoke, and roasted were also recorded.

### Plant Part used for Medicinal Purposes and their Life Form

Among the different parts of the plants used in therapy, the whole plant and leaves are used frequently (21% of each) (Fig 4). In literature, it was also noted that the leaves are more accessible or available in nature and are relatively more abundant as compared to other plant parts which may explain why they are used, while the frequent use of whole plant in the region may be that the area is mountainous and very less rain falls in the region, mostly plants are herbaceous and wild bushes (Fig. 5.) due to this the people collect the aerial part of the plants and use their decoction and infusion commonly. The herbaceous habit is not only dominant life form in our study but it is a common and widespread ecological phenomenon around the world. That for the preparation of remedies from the whole plants is very commonly used (23.13%) followed by leaves (19.28%). It is also noticed, that if only one plant part is required e.g. leaf, flower or fruit for the need is local people collects the whole plants instead of single part, the practice of plant parts collections has adversely affected the population size. The other plants used by the local people were seeds (20%), fruits (10%) and other (Fig. 4) due to extensive use of seeds and whole plant, The pressure on the survival of such wild populations has increased. The least

used parts are tuber and roots, probably due to their low level of approach that very few plants have tubers in the area and the roots of shrub and tree are very difficult to get.

### Quantitative Analysis

#### Informants consensus factor (ICF) and fidelity level (FL)

The informant consensus factor (ICF) of medicinal plants in our study ranges from (0-1.0) (Table 4). Antidote category has highest ICF Value (1.0) in which only one species *Calotropis procera* is used for snake bite and scorpion stung. The second highest value observed is for respiratory disease (0.39). the least agreement between the informants was observed for plants used for nose, ear and throat disease (ENT) (Earache, throat inflammation) and eye disease both having the zero ICF. Similar result were reported by Jamila and Mostafa [27], who reported the second highest ICF for respiratory disease (ICF: 0.81) and least ICF for eye and vision problems (ICF: 0.21). Fidelity level (FL) of 21 plant species was found against a given ailment category (Table 5) 100% fidelity level was calculated for three plant species. According to our findings, we suggest that high FL indicates the prevalence of

specific disease in the area that are treated with the medicinal plants with the high FL values.

### Threats to Medicinal Plants and Indigenous Knowledge in the Area

Majority of the people of the areas are educated but especially in the rural areas are 56% illiterate of the division and the earning sources of the locals are only agriculture and livestock. Some of the local inhabitants collect medicinal plants-- *Momordica charanita*, *Punica granatum*, *Phyllanthus emblica*, *Raphanus Sativus*, *Zanthoxylem armatum*, *Zingiber officinale* *Mentha arvensis*, *Litsea glutinosa*, *Lathyrus aphaca* and sell them to the local herb sellers in very cheap prices and these species are traded to the pharmaceutical companies in good prices. Over grazing point, urbanization, and uprooting of medicinal plants and serious threat in the areas, These threat increase the risk of their extinction and calls for a strict control over their protection by the authorities. The sustainable use of wild flora for cultivation of medicinal plants should be promoted in the area, This will strongly improve the socioeconomic condition of the local inhabitants.




Figure 3: Percentage of herbal drug preparation




Figure 4: Percentage of plants used



Table 4: Percentage of species and citations in each medicinal use category

S no.	Disease category	No of use reports	%age of reports	No of species	%age of taxa used	Informants consensus factor (ICF)
1	Jaundice, Wounds, Back pain, Skin disease,	2	1	1	1	1.0
2	Dissolving kidney, Urinary bladder Food, Stones, Flushing urinary blocks	34	18	18	12	0.39
3	Delivery, Dandelion wine, Constipation, Fever.	17	16	9	11	0.07
4	Worm infestation, constipation, Diuretic, hand burning. Fever, Skin allergy	14	13	7	9	0.08
5	Ear, nose and throat disease (ENT) Earache, throat inflammation,	4	3	2	2	0
6	Jaundice/ Liver, Digestive, Fever, Ulcers, Cephalalgia	18	16	9	11	0.12
7	Blood pressure, Stimulation, Aphrodisiac, Digestive	7	5	4	3	0.34
8	Infectious disease (Malarial fever, typhoid, measles	15	14	8	7	0.09
9	Bones fracture, dislocation, joints pain	12	7	5	5	0.3
10	Skin disease, Chest Pan. hands burning	9	8	6	6	0.14
11	Produces protein, Toothache, Narcotic, Aphrodisiac, Sprains, Fracture	35	19	19	13	0.52

Table 5: Fidelity level (FL) of medicinal plants of the study area

S.no	Plants name	No of informants repoted the taxa	number of aliments treated	No. of use frequently determined by informant	FL
1	Berberis lyceum	16	4	16	100
2	Bergenia ciliata	34	5	38	86.48
3	Coriandrum sativum	25	4	25	100
4	Momordica charanita	36	5	36	100
5	Cuscuta reflexa	10	3	18	55.56
5	Cedrus deodara	17	3	22	77.28
7	Cinnamomum tamala	25	2	25	100
8	Equisetum debile	13	3	13	93.67
9	Luffa cylindrica	20	4	29	98.97
10	Lathyrus aphaca	26	2	18	69.24
11	Musa paradisiacal	24	3	24	100
12	Punica granatum	33	1	29	87.88
13	Persicaria amplexicaulis	21	2	16	76.19
14	Phyllanthus emblica	28	1	26	92.86
15	Rosa indica	20	2	18	90
16	Ricinum communis	29	2	24	82.76
17	Trichosanthes cucumerina	12	2	11	91.67
18	Taraxacum officinal	24	2	21	87.17
19	Vitis Jacquemontii	15	1	8	53.34
20	Zanthoxylem armatum	26	3	26	100
21	Zee mays	28	2	28	100




Figure 5: Percentage of plant life form

## CONCLUSION

This study first documented the information about the traditional medicinal plants in Pir Panjal range in Jammu division Himalaya

of Union territory Jammu & Kashmir-India. The area is rich in medicinal plants and these plants are still commonly used for medicinal purpose of people in their daily lives. There is a gradual loss of traditional knowledge about these medicinal plants in new generation. Thus it is felt important to document and reconstitute the remainders of the ancient medical practice which exist in the area as well as other part of the region and Preserve this knowledge for future generation. This data matches with that of Singh and Kim. [28] provide a list of some alpine plants of Poonch; Kim [28] presented a brief account of some medicinal plants of Pir Panjal range: Singh [29] gave an introductory account of some wild flowering plants of Rajouri; Vir Jee *et al.* [30] reported their concise taxa-ethnobotanical observation made in some rural areas of Rajouri. Thus, such type of study may also bring to light some new source of drugs for control the disease. This study also provides basic for the conservation of the local flora; It will also provide various socio-economic dimensions associated with the common people.

## ACKNOWLEDGEMENTS

This research was carried out in Centre for Biodiversity Studies, School of Biosciences and Biotechnology Baba Ghulam

Shah Badshah University, Rajouri. I am thankful to the Vice Chancellor, Professor Javed Musarrat for believing me and giving me opportunity to serve as Extension Scientist in Centre for Biodiversity Studies. I would like to place on record my sincere gratitude to my mentor, Professor, Iqbal Parwez Dean School of Biosciences and Biotechnology for his guidance, constant support and motivations, Without his help this work would not have been possible.

## REFERENCES

- Redzić, S. S. 2007. The ecological aspect of ethnobotany and ethnopharmacology of population in Bosnia and Herzegovina. *Collegium Antropologicum*, 31; 869-890.
- Ullah, R., Hussain, Z., Iqbal, Z., Hussain, J., Khan, N., Muhammad, Z., Ayaz, Z., Ahmad, S., 2010. Traditional uses of medicinal plants in Dara Adam Khel NWFP Pakistan. *Journal of Medicinal Research*, 17: 1815-1821.
- Safa, O., Soltanipoor, M.A., Rastegar, S., Kazami, M., Dehkord, K.N., Ghannadi, A., 2012. An ethnobotanical survey on Harmozgan Province, Iran. *Avicenna Journal of Phytomedicine*, 3(1): 64-81.
- Nasab, K.F., Khosravi, A.R. 2014. Ethnobotanical study of medicinal plants of Sirjan in Kerman Province, Iran. *Journal of Ethnopharmacology*, 154; 190-197
- Singh, H., Husain, T., Agnihotri, P., Pande, P.C., Khaton, S. 2014. An ethnobotanical study of medicinal plants used in sacred groves of Kumaon Himalaya. Uttarakhand. *Iran Journal of Ethnopharmacology*, 154; 98-108.
- Hamayun, M., Khan, M.A. and Hayat, T., 2005. Ethnobotanical profile of Utror and Gabral Valleys, District Swat, Pakistan. [www.ethnoleaflets.com/leaflets/swat.htm](http://www.ethnoleaflets.com/leaflets/swat.htm).
- Shah, A., Abass, G., Sharma MP. 2012. Ethnomedicinal study of Some Medicinal plants from Tehsil Budhal, District Rajouri Jammu and Kashmir. *International Multidisciplinary Research Journal*, 2(6); 05-06
- Mahmood, T., Shah, A., 2012. Medicinal plants used by traditional healers in Poonch district of Jammu and Kashmir, *Life Science Leaflets*, 5: 53-60.
- Mahmood, T., Kadam, D., 2012. Some medicinal Plants used for the treatment of Jaundice and Hepatitis based on Tribal and rural people of Poonch & Rajouri Jammu and Kashmir. *Environment and Ecology*, 3:449-454
- Mahmood, T., 2019. A report on ethnomedicinal Plants used by Gujjar - Bakarwal tribes of Some Parts of Pir Panjal Himalayas of District Rajouri Jammu and Kashmir India. *Environment and Ecology*, 37(1A); 293-303.
- Mahmood, T., 2019. Ethnomedicinal study of useful climbers creepers and twiners of Baba Ghulam Shah Badshah University campus and adjoining areas of District Rajouri (Jammu and Kashmir). *Med. Aromatic Plants (Loss Angles)*, Vol.8 Iss.6 No.8:340. doi:10.35248 2167-0412-19. 8.340
- Pant, S., Verma, S., 2008. Ethnomedicinal notes on Tree Species of Pir Panjal Biodiversity Park of Baba Ghulam Shah Badshah University Rajouri, Jammu and Kashmir, India. *Ethnomedicinal Leaflets*, 12:404-412.
- Malik, A.H, Khuroo, A.A., Dar, G.H., Khan, Z., S. 2010. The woody flora of Jammu and Kashmir India. An updated Checklist. *Journal of Economic and Taxonomic Botany*, 34(2): 274-297.
- Shah, A., Bharati, K.A., Ahmad, J., Sharma, M.P. 2015. New ethnomedicinal claims from Gujjar-Bakarwal tribes of rajouri and poonch districts of Jammu and Kashmir, India. *Journal of Ethnopharmacology*, 166: 119-28.
- Dar, G.H. Akhter, H, Malik and Anzar, A. Khuroo. 2014. A contribution flora of Rajouri and Poonch District in the Pir Panjal Himalaya Jammu and Kashmir, India. [www.checklist.org.br](http://www.checklist.org.br)
- Sharma, B.M and Kachroo, P. 1981-82. Flora of jammu and plants of Neighbourhood. Vols. I-II. Bishen Singh Mahendra Pal Singh, Dehra Dun, India
- Bibi, T., Ahmad, M., Tareen, B.R., Tareen, M.N., Jabeen, R., Rehman, Ur R., Sultana, S., Zafar, M., Yaseen, G., (2014) Ethnobotany of medicinal plants in district Mastung of Balochistan province – Pakistan. *Journal of Ethnopharmacology*, 79-89.
- Jain, S.K., Rao, R.R., 1977. A Handbook of field and herbarium Methods. Today and Tomorrow Printers and Publishers, New Delhi.
- Polunin, O., and A. Stainton. 1986. Flowers of the Himalaya, 2<sup>nd</sup> reprint. Oxford University Press, Delhi, Bombay, Calcutta & Madras.
- Shabana A., G. H. Dar and A. Q. Jhon. 2013. Exotic Ornamental Flora of Kashmir: A Field Guide. Dominant Publishers & Distributors Pvt. Ltd., New Delhi, India.
- Hooker, J.D. 1872-97. The flora of British India. Vols I-VII. London: L. Reeve and Co.
- Trotter, R.T., Logan, M.H., 1986. Informants consensus; a new approach for identifying potentially effective medicinal plants, In: Etkin, N.L. (Ed.), Plants in indigenous Medicine and Diet. Redgrave Publishing Company, Bedford Hills, New York, pp. 91-112
- Gazzaneo, L., R.S., Lucena, R.F.P. Albuquerque. U.P., 2005. Knowledge and use of medicinal plants by the local specialists in a region of Atlantic forest in the state of Pernambuco (North eastern Brazil). *Journal of Ethnomedicines*, 1; 1-11.
- Friedman, J., Yaniv, Z., Dafni, A., Palewitch, D., 1986. A preliminary classification of the healing potential of medicinal plants, based on the rational analysis of an ethnopharmacology field survey among Bedouins in Negev Desert, Israel. *Journal of Ethnopharmacology*, 16: 275-287.
- Gurdal, B., Kultur, S., 2013. An ethnobotanical study of medicinal plants in Marmaris (Mugla, Turkey). *Journal of Ethnopharmacology*, 146.113-126.
- Ahmad, M., Shazia, S., Fazl., Hadi, S., Hadda, T.B., Rashid, S., Zafar, M., Khan, M.A., Khan, M.P.Z., Yaseen, G. 2014. An ethnobotanical study of medicinal plants in high mountainous region of Chail valley (District Swat-Pakistan). *Journal of Ethnobiology and Ethnomedicines*, 10: 1-36.
- Jamila, F., Mostafa, E. 2014. Ethnobotanical survey of medicinal plants used by people in Oriental Morocco to manage various ailments. *Journal of Ethnopharmacology*, 154:76-87.
- Kirn., H.S. 1992. Pir Panjal Range, a paradise of medicinal plants; pp.63-65, in: M. P. Sharma (ed.), Bahar-e-Rajouri-92. Nehru Yuva Kendra, Rajouri
- Singh, H., 1992. Wild flowers of Rajouri mountains; PP60-62. in: M.P. Sharma (ed.), Bahar-e-Rajouri-92. Nehru yuva Kendra, Rajouri
- Vir Jee, Dar G.H, P. Kachroo and G. M. Bhat. 1984. Taxo-ethnobotanical Studies of the rural areas in district Rajouri (Jammu). *Journal of Economic and Taxonomic Botany* 5 (4): 831-838.