

ROMANIAN NEUROSURGERY

Vol. XXXIV | No. 2 June 2020

Outcome analysis of upper and lower limb motor functions after anterior cervical discectomy and fusion for degenerative cervical disc disease

Ninad N. Srikhande, V.A. Kiran Kumar, N.A. Sai Kiran, Amrita Ghosh, Ranabir Pal, Luis Rafael Moscote-Salazar, V. Anil Kumar, Vishnu Vardhan Reddy, Amit Agrawal



Outcome analysis of upper and lower limb motor functions after anterior cervical discectomy and fusion for degenerative cervical disc disease

Ninad N. Srikhanda¹, V.A. Kiran Kumar¹, N.A. Sai Kiran¹, Amrita Ghosh², Ranabir Pal³, Luis Rafael Moscote-Salazar⁴, V. Anil Kumar⁵, Vishnu Vardhan Reddy¹, Amit Agrawal¹

¹ Department of Neurosurgery, Narayana Medical College Hospital, Chinthareddypalem, Nellore, Andhra Pradesh, INDIA

² Department of Biochemistry, Calcutta Medical College, 88, College street, Kolkata-700073, INDIA

³ Department of Community Medicine, MGM Medical College and LSK Hospital, Kishanganj -855107, Bihar, INDIA

⁴ Neurosurgery-Critical Care, Red Latino, Organización Latinoamericana de Trauma y cuidado, Neurointensivo, Bogota, COLOMBIA

⁵ Department of Anaesthesia, Narayana Medical College Hospital, Chinthareddypalem, Nellore, Andhra Pradesh, INDIA

ABSTRACT

Background: Anterior cervical discectomy and fusion (ACDF) is the most commonly performed surgical procedure for symptomatic cervical disc disease. In this study, we analysed the upper and lower limb motor functions after ACDF for disc prolapse in patients with degenerative cervical disc disease.

Methods: One hundred consecutive adult patients who underwent ACDF for single or two-level cervical disc prolapse during the study period (October 2015 to October 2017) were included in the study.

Results: Preoperative motor deficits in limbs were noted in 73% (73/100) of the patients. Enhance recovery of motor deficits was noted in 72.6% (53/73) of these patients and persisting motor deficits in the remaining patients (20/73- 27.4%). Five patients (5/27- 18.5%) without any preoperative motor deficits developed motor deficits after ACDF. Detailed pre and postoperative (at the time of discharge) motor power (graded by MRC grade) in all 4 limbs (Shoulder abduction / adduction / flexion / extension, elbow flexion / extension, wrist flexion / extension, hip abduction / adduction / flexion / extension, knee flexion/extension, ankle flexion/extension) was recorded. Statistically significant improvement in motor power (as recorded at the time of discharge) was noted in all the tested muscle groups after ACDF.

Conclusion: Early improvement in preoperative motor deficits can be expected in the majority of the patients with cervical PIVD following ACDF.

Keywords

anterior cervical discectomy and fusion, outcome, cervical disc degeneration



Corresponding author:
V. A. Kiran Kumar

Assistant Professor of Neurosurgery,
Narayana Medical College Hospital
Chinthareddypalem, Nellore, Andhra
Pradesh, India

drananth21@gmail.com

Copyright and usage. This is an Open Access article, distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>) which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited.

The written permission of the Romanian Society of Neurosurgery must be obtained for commercial re-use or in order to create a derivative work.

ISSN online 2344-4959
© Romanian Society of
Neurosurgery



First published
June 2020 by
London Academic Publishing
www.lapub.co.uk

INTRODUCTION

ACDF is most commonly done to treat a symptomatic cervical PIVD. (1, 16, 20, 22, 23, 25) ACDF is a safe procedure and is rarely associated with post-operative complications. (2, 7, 9, 12, 17) Significant proportion of patients shows remarkable recovery in motor deficits following ACDF. (11, 16) Authors in this publication analysed in detail the early recovery of motor deficits following ACDF for single or two level degenerative cervical PIVD.

MATERIAL AND METHODS

One hundred consecutive adult patients who underwent ACDF for degenerative cervical PIVD during the study period (October 2015 to October 2017) were included in the study. Patients with traumatic PIVD were excluded. Approval from the institutional ethical committee was taken for this study. Detailed pre and postoperative (At the time of discharge) motor power (graded by MRC grade except hand grip which was subjectively graded from 0-100%) in all 4 limbs (Shoulder abduction / adduction / flexion / extension, elbow flexion / extension, wrist flexion / extension, hip abduction / adduction / flexion / extension, knee flexion / extension, ankle flexion / extension) were analysed.

Statistical analysis

Data analysis was done by using statistical software SPSS Statistics Version 24.0. Descriptive statistics including mean and standard deviation for continuous variables, and frequency and percentage for categorical variables were used for data expression. Appropriate tests like Chi-square test, Wilcoxon signed rank test etc. were used for checking statistically significant correlation. A probability (P) value of <0.05 was considered significant.

RESULTS

Preoperative motor deficits in limbs were noted in 73% (73/100) of the patients. At the time of discharge, enhance recovery of motor deficits was noted in 72.6% (53/73) of these patients and persisting motor deficits in the remaining patients (20/73- 27.4%). Five patients (5/27- 18.5%) without any preoperative motor deficits developed motor deficits after ACDF. Pre and postoperative (At the time of discharge) motor power in all 4 limbs is compared in Tables 1-4. Statistically significant improvement in motor power at the time of discharge was recorded in all the tested muscle groups after ACDF.

Right upper limb	Preoperative (n=100)				Postoperative (n=100)				P-value	
	Mean [#]	Median [#]	Range [#]	IQR	Mean [#]	Median [#]	Range [#]	IQR		
Shoulder										
Shoulder abduction	3.80±1.4	4	0-5	2	4.15±1.29	5	0-5	1	0.001 (S)*	
Shoulder adduction	3.80±1.4	4	0-5	2	4.16±1.29	5	0-5	1	0.001 (S)*	
Shoulder flexion	3.81±1.4	4	0-5	2	4.18±1.29	5	0-5	1	0.001 (S)*	
Shoulder extension	3.80±1.4	4	0-5	2	4.17±1.32	5	0-5	1	0.001 (S)*	

Elbow										
Elbow flexion	3.73±1.5	4	0-5	1		4.17±1.32	5	0-5	1	0.001 (S)*
Elbow extension	3.71±1.5	4	0-5	2		4.16±1.33	5	0-5	1	0.001 (S)*
Wrist										
Wrist flexion	3.71±1.6	4	0-5	1		4.14±1.27	5	0-5	1	0.001 (S)*
Wrist extension	3.72±1.6	4	0-5	2		4.14±1.29	5	0-5	1	0.001 (S)*
Right hand grip	Preoperative (Number of patients)				Postoperative (Number of patients)				P-value	
0-25%	11				4				0.001 (S)†	
25-50%	11				13					
50-75%	29				22					
75-100%	49				61					

Table 1. Comparison of preoperative and postoperative motor power in right upper limb.

#Power graded according to MRC grade.

*Obtained Using Wilcoxon signed rank test; † Obtained using Chi square test; S: Significant.

Left upper limb	Preoperative (n=100)				Postoperative (n=100)				P-value*	
	Mean#	Median#	Range#	IQR	Mean#	Median#	Range#	IQR		
Shoulder										
Shoulder abduction	3.75±1.41	4	0-5	2	4.12±1.36	5	0-5	1	0.001 (S)*	
Shoulder adduction	3.74±1.41	4	0-5	2	4.14±1.34	5	0-5	1	0.001 (S)*	
Shoulder flexion	3.72±1.42	4	0-5	2	4.13±1.34	5	0-5	1	0.001 (S)*	
Shoulder extension	3.72±1.45	4	0-5	2	4.13±1.32	5	0-5	1	0.001 (S)*	
Elbow										

Knee flexion	3.40±1.86	4	0-5	2	3.90±1.60	5	0-5	2	0.001(S)
Knee extension	3.38±1.86	4	0-5	2	3.90±1.59	5	0-5	2	0.001(S)
Ankle									
Ankle flexion	3.33±1.85	4	0-5	2	3.84±1.58	4.5	0-5	2	0.001(S)
Ankle extension	3.35±1.86	4	0-5	2	3.85±1.58	5	0-5	2	0.001(S)

Table 3. Comparison of preoperative and postoperative motor power in right lower limb.

#Power graded according to MRC grade

*Obtained Using Wilcoxon signed rank test; S: Significant

Left lower limb	Preoperative (n=100)				Postoperative (n=100)				P-value*
	Mean	Median	Range	IQR	Mean	Median	Range	IQR	
Hip									
Hip abduction	3.32±1.86	4	0-5	2	3.98±1.53	5	0-5	1	0.001 (S)
Hip adduction	3.33±1.87	4	0-5	2	3.99±1.52	5	0-5	1	0.001 (S)
Hip flexion	3.32±1.86	4	0-5	2	4.00±1.49	5	0-5	1	0.001 (S)
Hip extension	3.29±1.84	4	0-5	2	3.98±1.49	5	0-5	1	0.001 (S)
Knee									
Knee flexion	3.27±1.83	4	0-5	2	3.95±1.52	5	0-5	1	0.001 (S)
Knee extension	3.27±1.84	4	0-5	2	3.94±1.51	5	0-5	1	0.001 (S)
Ankle									
Ankle flexion	3.22±1.89	4	0-5	3	3.88±1.55	5	0-5	2	0.001 (S)
Ankle extension	3.20±1.89	4	0-5	3	3.89±1.56	5	0-5	2	0.001 (S)

Table 4. Comparison of preoperative and postoperative motor power in right lower limb.

#Power graded according to MRC grade

*Obtained Using Wilcoxon signed rank test; S: Significant

DISCUSSION

Cervical PIVD is a common degenerative disc disease affecting millions of people. (24) Cervical disc herniation can occur as a result of ageing, wear and tear, or sudden stress from an accident.(5) Majority of these patients present with neck pain radiating to upper limbs. (24) Other presenting symptoms include motor deficits, stiffness in limbs, sensory deficits, paresthasias in limbs etc. (16, 24) Majority of the patients presenting with only neck pain or radicular pain can be managed with medication as

and conservative measures like physiotherapy, cervical collar etc. Patients with significant pain not responding to conservative measures and patients with neurological deficits like sensory/motor deficits and bladder symptoms respond well to surgery. (10, 16, 21)

ACDF is a common surgical procedure performed for symptomatic degenerative cervical disc disease. (6, 16) It helps to relieve the pressure on nerve roots and/or on the spinal cord, (14) thus resulting in improvement in various clinical symptoms including

neck pain, radicular pain, motor weakness, sensory symptoms, tightness in limbs and bladder disturbances. (10, 13, 16) Various complications reported with ACDF include dysphagia, hoarseness of voice, wound hematoma, graft migration, pseudoarthrosis, wound infection etc. (13) Patients can rarely have sensory or motor deficits after ACDF due to small risk of damage to the spinal cord, nerve roots or both. (10, 13, 16)

Improvement in neurological deficits ranging from 36-93% has been reported in various series after ACDF. (3, 13, 15, 16, 18) Lehman et al (16) reported preoperative motor deficits in 55% of the patients and reported recovery of these deficits in 95% of them at 1 year. Chiles et al (4) reported strength improvement rates ranging from 79.1% to 90.9% in various individual muscle groups of upper and lower limbs following ACDF. In the present study very high proportion of patients (73%) presented with motor deficits and early complete recovery of these deficits were noted in 72.6% (53/73) of these patients.

Majority of the studies on ACDF have graded neurological deficits using various scores like Nurick's grade, JOA, modified JOA scores etc. which combine both sensory and motor deficits. (4, 8, 16, 19) Detailed assessment of motor deficits with grading of motor power for various muscle groups has not been done in most of the studies on ACDF. (8, 16) In the present study we compared the preoperative motor power and early postoperative (at the time of discharge) motor power following ACDF in all major groups of muscles of lower and upper extremity and found that significant improvement in motor power in early postoperative period. Long term follow-up studies in patients following ACDF is required as they can develop new deficits secondary to adjacent segment disease. (16)

CONCLUSION

Early improvement in preoperative motor deficits can be expected in majority of the patients with degenerative cervical PIVD following ACDF.

REFERENCES

1. Bhaganagare AS, Nagesh SA, Shrihari BG, Naik V, Nagarjun MN, Pai BS: Management of cervical monoradiculopathy due to prolapsed intervertebral disc, an institutional experience. *Journal of craniovertebral junction & spine* 8:132, 2017.
2. Boakye M, Mummaneni PV, Garrett M, Rodts G, Haid R: Anterior cervical discectomy and fusion involving a polyetheretherketone spacer and bone morphogenetic protein. *Journal of Neurosurgery: Spine* 2:521-525, 2005.
3. Bohlman HH, Emery SE, Goodfellow DB, Jones PK: Robinson anterior cervical discectomy and arthrodesis for cervical radiculopathy. Long-term follow-up of one hundred and twenty-two patients. *JBJS* 75:1298-1307, 1993.
4. Chiles III BW, Leonard MA, Choudhri HF, Cooper PR: Cervical spondylotic myelopathy: patterns of neurological deficit and recovery after anterior cervical decompression. *Neurosurgery* 44:762-769, 1999.
5. Chung JY, Park J-B, Seo H-Y, Kim SK: Adjacent Segment Pathology after Anterior Cervical Fusion. *Asian Spine J* 10:582-592, 2016.
6. Dardis RM, Saxena A, Shad A, Chitnavis B, Gullan R (2012) *Disc Replacement Technologies in the Cervical and Lumbar Spine*. Elsevier, pp 1777-1788.
7. Davis RA: A long-term outcome study of 170 surgically treated patients with compressive cervical radiculopathy. *Surgical neurology* 46:523-533, 1996.
8. Emery SE, Bohlman HH, Bolesta MJ, Jones PK: Anterior cervical decompression and arthrodesis for the treatment of cervical spondylotic myelopathy. Two to seventeen-year follow-up. *JBJS* 80:941-951, 1998.
9. Fountas KN, Kapsalaki EZ, Nikolakakos LG, Smisson HF, Johnston KW, Grigorian AA, Lee GP, Robinson Jr JS: Anterior cervical discectomy and fusion associated complications. *Spine* 32:2310-2317, 2007.
10. Gore DR, Sepic SB: Anterior Discectomy and Fusion for Painful Cervical Disc Disease: A Report of 50 Patients With an Average Follow-up of 21 Years. *Spine* 23:2047-2051, 1998.
11. Hessler C, Boysen K, Westphal M, Regelsberger J: Functional and radiological outcome after ACDF in 67 cases. *Zeitschrift fur Orthopadie und Unfallchirurgie* 149:683-687, 2011.
12. Jacobs WC, Anderson PG, Limbeek J, Willems PC, Pavlov P: Single or double-level anterior interbody fusion techniques for cervical degenerative disc disease. *Cochrane Database Syst Rev* 4:CD, 2004.
13. Jagannathan J, Shaffrey CI, Oskouian RJ, Dumont AS, Herrold C, Sansur CA, Jane Sr JA: Radiographic and clinical outcomes following single-level anterior cervical discectomy and allograft fusion without plate placement or cervical collar. 2008.
14. Kim W-K (2011) *Role of Minimally Invasive Cervical Spine Surgery in the Aging Spine*. Elsevier, pp 198-203.
15. Lebl DR, Hughes A, Cammisa FP, O'Leary PF: Cervical spondylotic myelopathy: pathophysiology, clinical presentation, and treatment. *HSS journal* 7:170-178, 2011.
16. Lehmann CL, Buchowski JM, Stoker GE, Riew KD: Neurologic Recovery after Anterior Cervical Discectomy and Fusion. *Global Spine J* 4:41-46, 2014.
17. Levinthal R: Anterior cervical discectomy with and

- without fusion. Results, complications, and long-term follow-up. *Spine* 19:2343-2347, 1994.
18. Murrey D, Janssen M, Delamarter R, Goldstein J, Zigler J, Tay B, Darden B: Results of the prospective, randomized, controlled multicenter Food and Drug Administration investigational device exemption study of the ProDisc-C total disc replacement versus anterior discectomy and fusion for the treatment of 1-level symptomatic cervical disc disease. *The Spine Journal* 9:275-286, 2009.
 19. Nurjck S: The pathogenesis of the spinal cord disorder associated with cervical spondylosis. *Brain* 95:87-100, 1972.
 20. Orr RD, Zdeblick TA: Cervical Spondylotic Myelopathy: Approaches to Surgical Treatment. *Clinical orthopaedics and related research* 359:58-66, 1999.
 21. Saldia NS, Okafor C, Harris EB, Vaccaro AR (2011) The Role of Spinal Fusion and the Aging Spine: Stenosis without Deformity. Elsevier, pp 329-335.
 22. Truumees E, Herkowitz HN: Cervical spondylotic myelopathy and radiculopathy. *Instr Course Lect* 49:339-360, 2000.
 23. Wu T-k, Wang B-y, Deng M-d, Hong Y, Rong X, Chen H, Meng Y, Liu H: A comparison of anterior cervical discectomy and fusion combined with cervical disc arthroplasty and cervical disc arthroplasty for the treatment of skip-level cervical degenerative disc disease: A retrospective study. *Medicine* 96:2017.
 24. Yolas C, Ozdemir NG, Okay HO, Kanat A, Senol M, Atci IB, Yilmaz H, Coban MK, Yuksel MO, Kahraman U: Cervical disc hernia operations through posterior laminoforaminotomy. *Journal of Craniovertebral Junction and Spine* 7:91, 2016.
 25. Yu J, Ha Y, Shin JJ, Oh JK, Lee CK, Kim KN, Yoon DH: Influence of plate fixation on cervical height and alignment after one-or two-level anterior cervical discectomy and fusion. *British journal of neurosurgery*:1-8, 2017.