

DOLABRANE-TYPE DITERPENOID AND LIGNAN CONSTITUENTS FROM THE STEM BARKS OF *CERIOPS DECANDRA* (GRIFF.) W. THEOB.

Kieu Thi Phuong Linh¹, Nguyen Van Chien², Nguyen Quoc Trung³,
Vu Huy Thong⁴, Nguyen Van Tuyen⁵, Nguyen Phuong Thao^{1,*}

¹Institute of Marine Biochemistry, Vietnam Academy of Science and Technology (VAST),
18 Hoang Quoc Viet, Cau Giay, Ha Noi, Viet Nam

²Institute for Tropical Technology, VAST, 18 Hoang Quoc Viet, Cau Giay, Ha Noi, Viet Nam

³Institute of Materials Science, VAST, 18 Hoang Quoc Viet, Cau Giay, Ha Noi, Viet Nam

⁴University of Fire Fighting and Prevention, 243 Khat Duy Tien, Ha Noi, Viet Nam

⁵Institute of Chemistry, VAST, 18 Hoang Quoc Viet, Cau Giay, Ha Noi, Viet Nam

*Email: thaonp@imbc.vast.vn

Received: 19 December 2019; Accepted for publication: 16 May 2020

Abstract. Three dolabrane-type diterpenoids (**1–3**) and a lignan (**4**) were isolated from a methanolic extract of *C. decandra* (Griff.) W. Theob. stem barks using various chromatographic separations. Their structures were elucidated to be tagalsine X (**1**), tagalsin P (**2**), *ent*-5 α ,2-oxodolabr-3-ene-3,15,16-triol (**3**), and pinoresinol (**4**) by detailed analysis via spectroscopic techniques (1D, 2D NMR, and ESI-MS data) as well as comparison with those reported.

Keywords: *Ceriops decandra*, Rhizophoraceae, diterpenoid, lignan.

Classification numbers: 1.1.1, 1.1.6.

1. INTRODUCTION

Plants of the Rhizophoraceae family contain approximately 24 species in 4 genera including *Bruguiera* (7 species), *Ceriops* (5 species), *Kandelia* (2 species), and *Rhizophora* (10 species) [1-3]. The species have a wide distribution through both tropical and sub-tropical intertidal estuarine regions worldwide [1, 4]. Among them, the chemical compositions of the *Ceriops* genus have been investigated. Aurane, abietane, beyrane, dolabrane, and podocarpane-type diterpenoids are among the most frequently found secondary metabolites in this genus, which possess diverse structures due to many substituted moieties [2, 3]. Interestingly, dolabrane-type diterpenoids were found only in the *Ceriops* genus of the Rhizophoraceae family, making it a significant chemotaxonomic marker of that specific genus.




Figure 1. Structures of compounds 1–4 isolated from *C. decandra*.

Ceriops decandra (Griff.) W. Theob. is a mangrove species of ethnomedicinal significance having potent activity against a wide range of diseases like angina, boils, diabetes, diarrhea, dysentery, hepatitis, ulcers, and wounds [2, 3, 5 - 10]. Although the chemical constituents and pharmacological effects of the leaves and roots of *C. decandra* have been previously studied [2, 3, 5-8], however, the isolated metabolites (as abietane and podocarpane-type diterpenoids) from the stem barks of *C. decandra* have been limited [9, 10]. To date, a non-systematic phytochemical study that does not contain dolabrane-type diterpenoids has been described as the isolation from *C. decandra* distribution in Viet Nam. The current paper deals with detailed structure elucidation of four compounds (1–4, Figure 1) from this plant.

2. EXPERIMENTAL

2.1. General experimental procedures

The procedure and instruments used correspondingly to isolate compounds, measure optical rotation, and record IR, NMR, ESI-MS data collection, TLC and MPLC are similar to those described in a previous paper [11].

2.2. Plant material

The stem barks of *C. decandra* were collected at Ca Mau National Park, Ca Mau province, Viet Nam in May 2018, and taxonomically identified by Dr. Nguyen The Cuong (Institute of Ecology and Biological Resources, VAST). A voucher specimen (TĐPCCC-2018.01) was deposited at the Herbarium of Institute of Marine Biochemistry and Institute of Ecology and Biological Resources, VAST.

2.3. Extraction and isolation

The dried stem barks of *C. decandra* (2.0 kg) were cut into pieces and extracted with 95 % aqueous MeOH by percolation at room temperature to obtain 210 g of extract. The concentrated

95 % MeOH extract was suspended in H₂O and defatted with *n*-hexane and then was partitioned into an ethyl acetate-soluble fraction.

The EtOAc-soluble fraction (E, 20.5 g) was separated on silica gel MPLC (column: Biotage SNAP Cartridge, KP-SIL, 100 g) using a mobile phase of CH₂Cl₂-EtOAc (0 - 5 min 50 % EtOAc, 6-65 min 50 - 75 % EtOAc, 66 - 75 min 100 % EtOAc, 76 - 90 min 100 % MeOH, 15 mL/min, 90 min) to give twelve fractions (E-1 to E-12). Fractions E-7 (3.2 g) was further separated on a silica gel column, using CH₂Cl₂-acetone (40:1, v/v) as the mobile phase, to give five subfractions (E-7.1 to E-7.5). Subfraction E-7.5 (0.42 g) was chromatographed over an open ODS column eluted with acetone-H₂O (3:2, v/v) to yield three subfractions (E-7.5a to 7.5c). Subfraction E-7.5a was purified using preparative TLC with CH₂Cl₂-acetone (15:1, v/v), to give compound **2** (1.5 mg). Similarly, subfraction E-7.5b was chromatographed over a silica gel column with *n*-hexane-acetone (50:50 → 0:100) and *n*-hexane-EtOAc-MeOH (50:50:0.1 → 0:100:0) mixtures, and the resulting fraction was separated by a Sephadex LH-20 column using acetone-H₂O (1:1, v/v) to afford compound **1** (2.1 mg). Fraction E-11 (0.54 g) was subjected to a Sephadex LH-20 column eluted with MeOH to obtain three subfractions (E-11.1 to E-11.3). Compound **3** (2.5 mg) was purified from subfraction E-11.1 using silica gel CC eluting with *n*-hexane-CH₂Cl₂-acetone (2.5:1:1, v/v). Compound **4** (3.6 mg) was obtained by purifying subfraction E-12 on the YMC*GEL column and followed by separation on a Sephadex LH-20 column using a mixture of acetone-H₂O (1:1).

Tagalsine X (1): Pale yellow, amorphous powder; mp. 40 - 42 °C; $[\alpha]_D^{24} +63.4$ (c 0.2, MeOH); ESI-MS m/z 299 [M + Na]⁺; ¹H NMR (500 MHz, CDCl₃) and ¹³C NMR (125 MHz, CDCl₃) spectroscopic data, see Table 1.

Tagalsin P (2): Colorless crystals; mp. 175 - 177 °C; $[\alpha]_D^{24} +50.6$ (c 0.2, MeOH); UV (MeOH) λ_{max} (log ϵ) 288 (4.06) nm; ESI-MS m/z 315 [M + Na]⁺; ¹H NMR (500 MHz, CDCl₃) and ¹³C NMR (125 MHz, CDCl₃) spectroscopic data, see Table 1.

Ent-5 α ,2-oxodolabr-3-ene-3,15,16-triol (3): White, needle-like solid; mp. 126 - 128 °C; $[\alpha]_D^{24} +28.9$ (c 0.15, MeOH); ESI-MS m/z 337 [M + H]⁺; ¹H NMR (500 MHz, CDCl₃) and ¹³C NMR (125 MHz, CDCl₃) spectroscopic data, see Table 1.

Pinoresinol (4): White, amorphous powder; mp. 120 - 121 °C; $[\alpha]_D^{24} -16.6$ (c 0.05, MeOH); UV λ_{max}^{MeOH} (ϵ) 231 (14000), 280 (5800) nm; IR (KBr) ν_{max} : 3448, 1608, 1520, 1511, 1420, 1380, 1101, and 899 cm⁻¹; EI-MS m/z 341 [M - H₂O + H]⁺; ¹H NMR (500 MHz, CD₃OD): δ_H 6.96 (2H, d, $J = 1.5$ Hz, H-2, H-2'), 6.79 (2H, d, $J = 8.0$ Hz, H-5, H-5'), 6.83 (2H, dd, $J = 1.5, 8.0$ Hz, H-6, H-6'), 4.72 (2H, d, $J = 4.0$ Hz, H-7, H-7'), 3.15 (2H, m, H-8, H-8'), 3.86 (2H, dd, $J = 3.5, 9.0$ Hz, H-9a, H-9'a), 4.24 (2H, dd, $J = 7.5, 9.0$ Hz, H-9b, H-9'b), 3.87 (6H, s, 3,3'-OCH₃); ¹³C NMR (125 MHz, CD₃OD): δ_C 133.8 (C-1, C-1'), 111.0 (C-2, C-2'), 149.1 (C-3, C-3'), 147.3 (C-4, C-4'), 116.1 (C-5, C-5'), 120.1 (C-6, C-6'), 87.5 (C-7, C-7'), 55.4 (C-8, C-8'), 72.6 (C-9, C-9'), 56.5 (3,3'-OCH₃).

3. RESULTS AND DISCUSSION

Compound **1** was obtained as a pale yellow, amorphous powder. Its molecular formula was found to be C₁₈H₂₈O₂ via the ¹³C NMR spectroscopic data and a positive ESI-MS ion at m/z 299 [M + Na]⁺. From this formula and its NMR data **1** was determined to have five degrees of unsaturation, two of which were due to a double bond and a ketone group. The ¹H NMR

spectroscopic data displayed resonances for three tertiary methyls [δ_{H} 1.31 (3H, s, H₃-17), 0.87 (3H, s, H₃-19), and 0.92 (3H, s, H₃-20)], a secondary methyl [δ_{H} 1.03 (3H, d, $J = 6.5$ Hz, H₃-18)], a pair of olefinic protons [δ_{H} 6.84 (1H, dd, $J = 6.0, 10.5$ Hz, H-1) and 6.12 (1H, d, $J = 10.5$ Hz, H-2)], a methine [δ_{H} 2.82 (1H, q, $J = 6.5, 13.5$ Hz, H-4)], along with a series of aliphatic protons [δ_{H} 1.26-1.96 ppm] (Table 1).

Table 1. ^1H and ^{13}C NMR spectroscopic data for **1–3** (in CDCl_3).

Pos.	1		2		3	
	$\delta_{\text{C}}^{\text{a}}$	$\delta_{\text{H}}^{\text{b}}$ (mult., J in Hz)	$\delta_{\text{C}}^{\text{a}}$	$\delta_{\text{H}}^{\text{b}}$ (mult., J in Hz)	$\delta_{\text{C}}^{\text{a}}$	$\delta_{\text{H}}^{\text{b}}$ (mult., J in Hz)
1	147.9	6.84 (dd, 6.0, 10.5)	33.5	2.72 (br d, 18.5) 2.85 (dd, 6.5, 18.5)	33.2	2.72 (br d, 18.5) 2.83 (dd, 6.5, 18.5)
2	130.2	6.12 (d, 10.5)	192.9	-	193.1	-
3	202.6	-	144.6	-	144.6	-
4	45.0	2.82 (q, 6.5, 13.5)	135.4	-	135.5	-
5	39.0	-	39.0	-	39.0	-
6	37.5	1.22 (m) 1.96 (m)	38.0	1.25 (m) 2.17 (dd, 3.0, 14.0)	38.0	1.26 (m) 2.16 (m)
7	25.3	1.21 (m)/1.33 (m)	26.6	1.19*	26.8	1.13 (m)/1.27 (m)
8	44.5	1.31 (m)	44.3	1.19 (m)	41.2	1.36 (m)
9	39.3	-	37.9	-	37.9	-
10	57.5	1.87 (br d, 6.0)	54.4	1.60 (br d, 6.0)	54.4	1.63 (dd, 2.0, 6.0)
11	37.5	1.26 (m) 1.72 (m)	36.4	0.97 (dt, 4.5, 13.5) 1.76 (ddd, 3.5, 4.5, 13.5)	33.8	1.06 (ddd, 3.5, 4.5, 13.5) 1.66 (m)
12	35.6	1.57 (m)/1.68 (m)	35.6	1.52 (m)/1.63 (m)	28.4	1.32 (m)/1.52 (m)
13	71.1	-	71.1	-	36.4	-
14	43.0	1.41 (d, 13.0) 1.52 (d, 13.0)	42.6	1.32 (m) 1.49 (m)	36.3	0.88 (m) 1.37 (m)
15	-	-	-	-	81.0	3.31 (br d, 9.0)
16	-	-	-	-	62.5	3.73 (br d, 10.0) 3.51 (dd, 9.0, 10.0)
17	27.0	1.31 (s)	26.9	1.26 (s)	19.1	0.93 (s)
18	7.9	1.03 (d, 6.5)	11.6	1.87 (s)	11.6	1.87 (s)
19	26.3	0.87 (s)	31.7	1.23 (s)	31.7	1.23 (s)
20	13.5	0.92 (s)	13.9	0.68 (s)	11.5	0.60 (s)
-OH						6.10 (s)

^a125 MHz, ^b500MHz. *Overlapped signals assigned by HSQC and HMBC spectra without designating multiplicity.




Figure 2. Key HMBC correlations of **1**.

Consistent with these observations, its ^{13}C NMR and HMQC spectrum denoted the presence of 18 resonances for the presence of four methyls, five sp^3 methylenes, five sp^3 methines, and five sp^2 quaternary carbons, of which the olefinic carbons [δ_{C} 147.9 (C-1) and 130.2 (C-2)] were attributed to a disubstituted double-bond and a carbon signal [δ_{C} 202.6 (C-3)] was assigned to a ketone group. Apart from a double-bond and a ketone group, the remaining elements of unsaturation were suggested to a tricyclic skeleton in the molecule of **1**. These spectroscopic data indicated that **1** was a dinordolabrane-type diterpenoid [12, 13]. On the other hand, the 6/6/6 tricyclic skeleton of the diterpenoid with 4,5,9,13-tetramet--hyl and 3- α,β -unsaturated ketone substitutions in the A-ring was established by 2D NMR experiments. This assignment was confirmed by the observation of HMBC correlations between δ_{H} 1.03 (H₃-18) to C-3 (δ_{C} 202.6), C-4 (δ_{C} 45.0), and C-5 (δ_{C} 39.0); between δ_{H} 0.87 (H₃-19) to C-4 (δ_{C} 45.0), C-5 (δ_{C} 39.0), C-6 (δ_{C} 37.5), and C-10 (δ_{C} 57.5); between δ_{H} 0.92 (H₃-20) to C-8 (δ_{C} 44.5) C-9 (δ_{C} 39.3), C-10 (δ_{C} 57.5) and C-11 (δ_{C} 37.5); between δ_{H} 1.31 (H₃-17) to C-12 (δ_{C} 35.6), C-13 (δ_{C} 71.1), and C-14 (δ_{C} 43.0). A detailed 2D NMR spectral analysis, including HMQC and HMBC experiments, resulted in a gross structure of **1** (Figures 1-2).

The comparison of NMR spectroscopic data of **1** with reported literature found that they were similar suggesting that **1** was (4*S**,5*S**,8*S**,9*S**,10*R**)-13*S**-hydroxy-15,16-dinordolabr-1(2)-en-3-one and named tagalsine X (Table 1) [12]. From the above evidence, the structure of **1** was determined as tagalsine X. This compound was previously obtained from the leaves of *C. tagal* and had no cytotoxicity against CNE-2, A549, HepG2, and HCT-116 cell lines (IC₅₀ > 50 μM), even with the concentration of 50 μM [12].

Compound **2** was isolated as a colorless crystal. Its molecular formula was determined to be C₁₈H₂₈O₃ based on a sodium adduct molecular ion peak at m/z 315 [M + Na]⁺, consistent with five degrees of unsaturation. Analysis of the ^1H , ^{13}C NMR, and HSQC spectroscopic data of **2** displayed signals for all 18 carbons and 28 protons, including four tertiary methyls, six methylenes, two methines, and six non-protonated carbons. A detailed comparison of 1D and 2D NMR spectroscopic data showed that the structures of **2** and **1** [13] share the same B- and C-ring substitution patterns, with differences observed for the A-ring (Table 1). Further comparison of the ^1H and ^{13}C NMR data of **2** with those of **1** showed that both compounds exhibit closely comparable data, except for the replacement of a disubstituted double-bond at C-2 in **1** by a methylene group [δ_{H} 2.72 (1H, br d, $J = 18.5$ Hz, H-1a), 2.85 (1H, dd, $J = 6.5, 18.5$ Hz, H-1b); δ_{C} 33.5 (C-1)] and a conjugated ketone group [δ_{C} 192.9 (C-2)] in **2**. In addition, α,β -unsaturated ketone group in **2** was a tetrasubstituted double-bond [δ_{C} 144.6 (C-3) and 135.4 (C-4)] with a hydroxygroup [δ_{H} 6.10 (1H, s), suggesting a 15,16-dinor-dolabrane containing a 3-hydroxy-4-methyl-2-enone cyclohexane moiety in its A-ring [13]. Furthermore, this was confirmed by the key HMBC correlations from δ_{H} 1.87 (H₃-18) to C-3 (δ_{C} 144.6), C-4 (δ_{C} 135.4), and C-5 (δ_{C} 39.0), as well as from δ_{H} 1.60 (H-10) to C-1 (δ_{C} 33.5), C-2 (δ_{C} 192.9), C-5 (δ_{C} 39.0), and C-9 (δ_{C} 37.9), respectively. NMR spectroscopic data of **2** were identical to those of tagalsin P [13]. Thus, the structure of compound **2** was determined as tagalsin P, named (5*S**,8*S**,9*S**,10*R**)-3,13*S**-dihydroxy-15,16-dinordolabr-3-en-2-one (Figure 1).

Compound **3** was obtained as a white, needle-like solid. The ESI-MS showed a protonated molecular ion peak at m/z 337 [M + H]⁺, corresponding to a molecular formula of C₂₀H₃₂O₄, which is two carbons, four hydrogens, and one oxygen atom more than in **2**. The 1D NMR spectroscopic data of **3** were very similar to those of **2**, except for the presence of an additional dihydroxyethyl group [δ_{H} 3.31 (1H, br d, $J = 9.0$ Hz, H-15); δ_{C} 81.0 (C-15) and δ_{H} 3.51 (1H, dd, $J = 9.0, 10.0$ Hz, H-16a), 3.73 (1H, br d, $J = 10.0$ Hz, H-16b); δ_{C} 62.6 (C-16)] (Table 1). Furthermore, the location of the attached dihydroxyethyl group at C-13 was supported by a

downfield chemical shift of δ_C 36.4 (C-13) in the ^{13}C NMR spectra and a key HMBC correlation from δ_H 0.93 (s, H₃-17) to δ_C 81.0 (CH, C-15) of the dihydroxyethyl group. However, to date, the relative configuration at C-15 in **3** has been not yet reported. Comparisons of the NMR data of **3** with those of *ent*-5 α ,2-oxodolabr-3-ene-3,15,16-triol [14], as well as detailed analysis of HSQC and HMBC experiments led to identification of **3** as *ent*-5 α ,2-oxodolabr-3-ene-3,15,16-triol (Figure 1). Previously, compound **3** was obtained from the barks of *Endospermum diadenum* [14].

The remaining compound **4** was identified as pinoresinol based on our spectroscopic data and by comparison with those of reported data given in CDCl_3 [15]. This compound is widely distributed throughout the plants in Viet Nam, e.g, *Silybum marianum* [16], *Mallotus macrostachyus* [17], *Rhizophora stylosa* [18], *Trichosanthes kirilowii* [19], *Knema pachycarpa* [20], and *Balanophora laxiflora* [21].

4. CONCLUSIONS

In summary, we report here the isolation and structure elucidation of three dolabrane-type diterpenoids, agalsine X (**1**), tagalsin P (**2**), *ent*-5 α ,2-oxodolabr-3-ene-3,15,16-triol (**3**), and a lignan compound, pinoresinol (**4**) from a methanolic extract of *C. decandra* stem barks, using various chromatographic separations. The structures of these isolates were accomplished using comprehensive spectroscopic methods and comparison with those reported. The present work reports for the first time dolabrane-type diterpenoids study of this species distribution in Viet Nam. This work presents the discovery of dolabrane-type diterpenoid and lignan constituents and provides additional evidence to support mangrove plants as a promising source of chemical diversity.

Acknowledgments. This research is funded by a grant from the Vietnam Academy of Science and Technology (code: TĐPCCC.04/18-20). The authors are grateful to MSc. Dang Vu Luong (Institute of Chemistry, VAST) for measurement of the NMR spectra and Dr. Nguyen The Cuong (Institute of Ecology and Biological Resources, VAST) for taxonomic classification of plant.

REFERENCES

1. Wu J., Xiao Q., Xu J., Li M.Y., Pana J.Y., Yang M. H. - Natural products from true mangrove flora: source, chemistry and bioactivities, *Natural Product Reports* **25** (2008) 955-981.
2. Nebula M., Harisankar H. S., Chandramohanakumar. N. - Metabolites and bioactivities of Rhizophoraceae mangroves, *Natural Products and Bioprospecting* **3** (2013) 207-232.
3. Ethnomedicinal, Mahmud I., Shahria N., Yeasmin S., Iqbal A., Mukul E. H., Gain S., Shilpi J. A., Islam M. K. - Ethnomedicinal, phytochemical and pharmacological profile of amangrove plant *Ceriops decandra* Griff Din Hou, *Journal of Complementary and Integrative Medicine* **16** (2019) 1-22.
4. Chen Y., Hou Y., Guo Z., Wenqing Wang, Zhong C., Zhou R., Shi S. - Applications of multiple nuclear genes to the molecular phylogeny, population genetics and hybrid identification in the mangrove genus *Rhizophora*, *Plos One* (2015) e0145058.
5. Anjaneyulu A. S. R., Rao V. - Ceriopsins F and G, diterpenoids from *Ceriops decandra*, *Phytochemistry* **62** (2003) 1207-1211.

6. Anjaneyulu A. S. R., Rao V. - Ceriopsins A-D, diterpenoids from *Ceriops decandra*, *Phytochemistry* **60** (2002) 777-782.
7. Anjaneyulu A. S. R., Rao V. L., Lobkovsky E., Clardy J. - Ceriopsin E, a new epoxy *ent*-kaurene diterpenoid from *Ceriops decandra*, *Journal of Natural Products* **65** (2002) 592-594.
8. Ponglimanont C., Thongdeeying P. - Lupane-triterpene esters from the leaves of *Ceriops decandra* (Griff.) Ding Hou, *Australian Journal of Chemistry* **58** (2005) 615-618.
9. Jiang Z. P., Tian L. W., Shen L., Wu J. - *Ent*-abietanes from the Godavari mangrove, *Ceriops decandra*: absolute configuration and NF- κ B inhibitory activity, *Fitoterapia* **130** (2018) 272-280.
10. Wang H., Li M. Y., Satyanandamurty T., Wu J. - New diterpenes from a godavari mangrove, *Ceriops decandra*, *Planta Medica* **79** (2013) 666-672.
11. Luyen N. T., Binh P. T., Tham P. T., Hung T. M., Dang N. H., Dat N. T., Thao N. P. - Wedtrilosides A and B, two new diterpenoid glycosides from the leaves of *Wedelia trilobata* (L.) Hitchc. with α -amylase and α -glucosidase inhibitory activities, *Bioorganic Chemistry* **85** (2019) 319-324.
12. Wu X., Liao H., Lu H., Zhang C. - A new dolabrane dinorditerpene from *Ceriops tagal*, *Open Access Library Journal* **3** (2016) e2957.
13. Hu W. M., Li M. Y., Li J., Xiao Q., Feng G., Wu J. - Dolabranes from the Chinese mangrove, *Ceriops tagal*, *Journal of Natural Products* **73** (2010) 1701-1705.
14. Kijjoa A., Pinto M. M. M., Anantachoke C., Gedris T. E., Herz W. - Dolabranes from *Endospermum diadenum*, *Phytochemistry* **40** (1995) 191-193.
15. Xie L. H., Akao T., Hamasaki K., Deyama T., Hattori M. - Biotransformation of pinoresinol diglucoside to mammalian lignans by human intestinal microflora, and isolation of *Enterococcus faecalis* strain PDG-1 responsible for the transformation of (+)-pinoresinol to (+)-lariciresinol, *Chemical and Pharmaceutical Bulletin* **51** (2003) 508-515.
16. Diep T. T., Kiem P. V., Dong N. T., Tung N. H., Bang B. T., Minh C. V., Braca A. - Pinoresinol and 3,4',5,7-tetrahydroxy-3'-methoxyflavanone from the fruits of *Silybum marianum* (L.) Gaertn, *Journal of Chemistry* **45** (2007) 219-222.
17. Nam N. H., Kiem P. V., Ban N. K., Thao N. P., Nhiem N. X., Cuong N. X., Tistaert C., Dejaegher B., Heyden Y. V., Joëlle Quetin-Leclercq, Thao D. T., Minh C. V. - Chemical constituents of *Mallotus macrostachyus* growing in Vietnam and cytotoxic activity of some cycloartane derivatives, *Phytochemistry Letters* **4** (2011) 348-352.
18. Huong P. T. T., Thanh N. V., Diep C. N., Cuong N. T., Cuong N. X., Nam N. H., Thai T. H., Kiem P. V., Minh C. V. - Five lignans from the mangrove *Rhizophora stylosa* Griff, *Vietnam J. of Chem.* **53** (2015) 42-47.
19. Minh C. V., Nhiem N. X., Yen H. T., Kiem P. V., Tai B. H., Anh H. L. T., Hien T. T. T., Park S., Kim N., Kim S. H. - Chemical constituents of *Trichosanthes kirilowii* and their cytotoxic activities, *Archives of Pharmacal Research* **38** (2015) 1443-1448.
20. Tung T. H., Hue C. T., Giap T. H., Thoa H. T., Dung N. A., Hang N. T. M., Hung N. V., Thanh L. N. - Lignans isolated from the ethyl acetate extract of *Knema pachycarpa* fruit, *Vietnam J. of Chem.* **55** (2017) 406-410.
21. Dai T. D., Thuy T. T., Tuan N. N., An N. Q., Nga T. T. T., Cuong H. D., Tien N. Q. - Chemical constituents of *Balanophora laxiflora* Hemsley collected in Tuyen Quang, *Vietnam J. of Chem.* **56** (2018) 254-259.