

Georgia Southern University

Digital Commons@Georgia Southern

Physics and Astronomy Faculty Publications

Physics and Astronomy, Department of

1-20-2006

Evolution of Stellar Bars in Live Axisymmetric Halos: Recurrent Buckling and Secular Growth

Inma Martinez-Valpuesta

University of Kentucky, martinez@pa.uky.edu

Isaac Shlosman

University of Kentucky, isaac.shlosman@uky.edu

Clayton Heller

Georgia Southern University, cheller@georgiasouthern.edu

Follow this and additional works at: <https://digitalcommons.georgiasouthern.edu/physics-facpubs>



Part of the [Astrophysics and Astronomy Commons](#), and the [Physics Commons](#)

Recommended Citation

Martinez-Valpuesta, Inma, Isaac Shlosman, Clayton Heller. 2006. "Evolution of Stellar Bars in Live Axisymmetric Halos: Recurrent Buckling and Secular Growth." *Astrophysical Journal*, 637 (1): 214-226: The American Astronomical Society. doi: 10.1086/498338 source: <https://iopscience.iop.org/article/10.1086/498338>
<https://digitalcommons.georgiasouthern.edu/physics-facpubs/140>

This article is brought to you for free and open access by the Physics and Astronomy, Department of at Digital Commons@Georgia Southern. It has been accepted for inclusion in Physics and Astronomy Faculty Publications by an authorized administrator of Digital Commons@Georgia Southern. For more information, please contact digitalcommons@georgiasouthern.edu.

EVOLUTION OF STELLAR BARS IN LIVE AXISYMMETRIC HALOS: RECURRENT BUCKLING AND SECULAR GROWTH

INMA MARTINEZ-VALPUESTA,^{1,2} ISAAC SHLOSMAN,² AND CLAYTON HELLER³

Received 2005 July 4; accepted 2005 September 27

ABSTRACT

Evolution of stellar bars in disk galaxies is accompanied by dynamical instabilities and secular changes. Following the vertical buckling instability, the bars are known to weaken dramatically and develop a pronounced boxy/peanut shape when observed edge-on. Using high-resolution N -body simulations of stellar disks embedded in live axisymmetric dark matter halos, we have investigated the long-term changes in the bar morphology, specifically the evolution of the bar size, its vertical structure, and the exchange of angular momentum. We find that following the initial buckling, the bar resumes its growth from deep inside the corotation radius and follows the ultraharmonic resonance thereafter. We also find that this secular bar growth triggers a spectacular *secondary* vertical buckling instability that leads to the appearance of characteristic boxy/peanut/X-shaped bulges. The secular bar growth is crucial for the recurrent buckling to develop. Furthermore, the secondary buckling is milder, persists over a substantial period of time, ~ 3 Gyr, and can have observational counterparts. Overall, the stellar bars show recurrent behavior in their properties and evolve by increasing their linear and vertical extents, both dynamically and secularly. We also demonstrate explicitly that the prolonged growth of the bar is mediated by continuous angular momentum transfer from the disk to the surrounding halo and that this angular momentum redistribution is resonant in nature: a large number of lower resonances trap disk and halo particles, and this trapping is robust, in broad agreement with the earlier results in the literature.

Subject headings: galaxies: bulges — galaxies: evolution — galaxies: formation — galaxies: halos — galaxies: kinematics and dynamics — galaxies: spiral — stellar dynamics

Online material: color figures, mpeg animation

1. INTRODUCTION

Observations and numerical modeling of galactic stellar bars have been frequently accompanied with basic controversies about their origin and evolution. While modern understanding of bar growth in a live and responsive environment is rooted in the angular momentum redistribution between the inner and outer disks, bulges, and dark matter halos (Athanassoula 2003), the efficiency of this process is hardly known and its details are still to be investigated. Recent efforts include but are not limited to the issues related to the bar lifetime cycles, gas-star interactions, bar amplitudes and sizes, and bar slowdown (e.g., Bournaud & Combes 2002; Valenzuela & Klypin 2003; Shen & Sellwood 2004; Weinberg 1985; Hernquist & Weinberg 1992; Debattista & Sellwood 1998, 2000). The bigger issue, of course, is how the observational and theoretical aspects of bar evolution fit within the emerging understanding of cosmological galaxy evolution (e.g., Jogee et al. 2004; Elmegreen et al. 2004), specifically, the bar evolution in triaxial halos (El-Zant & Shlosman 2002; Berentzen et al. 2006).

On the one hand, early self-consistent models of numerical stellar bars have relied heavily on the Ostriker & Peebles (1973) result, which emphasized the Maclaurin sequence parameter $T/|W|$, the ratio of bulk kinetic to gravitational energy, as the threshold of bar instability. On the other hand, they revealed robustness of the bars: once formed, the bars persisted (Athanassoula 1984 and references therein). The subsequent increase in the

particle number above $N \sim 10^5$, switching from two-dimensional to three-dimensional models with responsive spheroidal components, and the introduction of nonlinear physics tools in the orbital analysis have shown a much more complex bar evolution and morphology than was anticipated originally (e.g., review by Athanassoula 2002b). This refers especially to the numerical confirmation that live halos can indeed drive the bar instability rather than damp it (Athanassoula & Misiriotis 2002; Athanassoula 2003). Finally, it is still unclear to what extent and how closely the numerical bars correspond to their observed counterparts. This issue exacerbates galaxy studies because overall, both theoretically and observationally, the bars appear to be among the most important drivers of galactic evolution across a wide range of spatial scales.

In this paper we have revisited some aspects of a self-consistent evolution of stellar bars originating in *live* stellar disks embedded in *live* dark matter halos by focusing on dynamical and secular changes⁴ in these systems. The evolution of *numerical* collisionless bars has been characterized so far in the literature by three distinct phases: the initial growth, the rapid vertical buckling, and the prolonged quasi-steady regime, i.e., when the bars preserve their basic parameters (e.g., Sellwood & Wilkinson 1993). However, some indication that bars can grow even in the last phase has been noticed already in low-resolution three-dimensional models with live halos (e.g., Sellwood 1980). This ability of the bars to grow over an extended period of time due to the momentum exchange with the outer disk and especially with the halo has been confirmed recently and analyzed in self-consistent three-dimensional simulations (Athanassoula 2003).

¹ Department of Physics, Astronomy, and Mathematics, University of Hertfordshire, Hatfield, Hertfordshire AL10 9AB, UK.

² Department of Physics and Astronomy, University of Kentucky, Lexington, KY 40506-0055.

³ Department of Physics, Georgia Southern University, P.O. Box 8031, Statesboro, GA 30460.

⁴ We refer to a dynamical evolution when changes develop on the timescale of about one disk rotation and to a secular evolution when they develop on a much longer timescale of $\sim 10\text{--}100$ rotations.

Here we attempt to quantify this secular growth in terms of the bar size and its ellipticity, of the angular momentum exchange, and of the ratio of vertical to radial dispersion velocities. Moreover, we look into the corollaries of such recurrent growth and find that it leads to additional and substantial two-dimensional and three-dimensional structural changes in the bar. We therefore discuss the observational consequences of this evolution.

Early in their growth stage, numerical stellar bars experience a dynamical instability: the vertical buckling. The bars thicken profoundly, become more centrally concentrated, and acquire a characteristic peanut/boxy shape when seen edge-on (Combes et al. 1990; Pfenniger & Friedli 1991; Raha et al. 1991; Berentzen et al. 1998; Patsis et al. 2002b), while nearly dissolving the outer half of the bar, beyond the vertical inner Lindblad resonance (ILR; Martinez-Valpuesta & Shlosman 2004). This happens in live models with both axisymmetric and mildly triaxial halos (Berentzen et al. 2006). These peanut/boxy shapes are similar to bulge shapes observed in edge-on galaxies (e.g., Jarvis 1986; Shaw 1987; Bureau & Freeman 1999; Merrifield & Kuijken 1999), which can be found in nearly half of all edge-on disk galaxies (Lüticke et al. 2000). Although they were observed in numerical simulations a long time ago (Combes & Sanders 1981), the origin of boxy/peanut bulge shapes still has two alternative explanations, the well-known fire-hose instability (e.g., Toomre 1966; Raha et al. 1991; Merritt & Sellwood 1994) and resonance heating (e.g., Combes et al. 1990; Pfenniger & Friedli 1991; Patsis et al. 2002a). These two views can be reconciled if buckling is responsible for shortening the secular timescale of particle diffusion out of the disk plane and for accelerating the buildup of boxy/peanut-shaped bulges, which proceeds on a much shorter dynamical timescale instead (Martinez-Valpuesta & Shlosman 2004).

However, is the buckling really necessary for a buildup of these boxy/peanut-shaped bulges? After all, even *imposing* vertical symmetry did not eliminate this effect, albeit the buildup proceeded on a much longer timescale (Friedli & Pfenniger 1990). Where are the observational counterparts of these asymmetric buckled bars? Due to a particular importance of the buckling instability for the evolution of numerical bars and its plausible connection to the buildup of the pronounced three-dimensional structure there, we have analyzed the bar behavior during and following this instability. Specifically, we find that the bars in a live environment are capable of recurrent growth, that the buckling instability is a recurrent event, and that the buildup of the three-dimensional shape is not necessarily a dynamic phenomenon.

Much of the analysis of bar evolution and the accompanying instabilities is implemented here by means of nonlinear orbit analysis, because for such strong departures from axial symmetry the epicyclic approximation cannot be relied upon: being wrong quantitatively, it frequently leads to qualitative errors. We try to avoid the specific jargon associated with this technique where possible. The angular momentum redistribution in the model is quantified using orbital spectral analysis.

In § 2 we provide the details of numerical modeling and analysis. Section 3 describes the overall results of secular bar evolution. The three-dimensional bar orbital structure and the inferred vertical structure in the bar are analyzed in § 4, and the resonant interaction between the disk and the halo is analyzed in § 5. Discussion and conclusions are given in §§ 6 and 7.

2. NUMERICAL TOOLS AND MODELING

2.1. *N*-Body Simulations

To simulate the stellar disk embedded in a live dark matter halo, we have used version FTM-4.4 of the *N*-body code




FIG. 1.—Initial circular rotation velocities for the disk (dashed line) and halo (dotted line) components. The total is given by the solid line.

(Heller & Shlosman 1994; Heller 1995) with $N = 10^5 - 1.1 \times 10^6$. The gravitational forces are computed using Denhen's (2002) falcON force solver, a tree code with mutual cell-cell interactions and complexity $O(N)$. It conserves momentum exactly and is about 10 times faster than an optimally coded Barnes & Hut (1986) tree code.

The initial density distribution is derived from the Fall & Efstathiou (1980) disk-halo analytical model. The system is not in exact virial equilibrium and therefore must be relaxed iteratively. The halo-to-disk mass ratio within 10 kpc is fixed to unity. The halo has a flat density core of 2 kpc to avoid excessive stochastic behavior associated with the central cusps (El-Zant & Shlosman 2002). The disk is exponential, and its radial and vertical scale lengths are taken as 2.85 and 0.5 kpc, respectively. The disk and halo cutoff radii are 25 and 30 kpc, and the initial circular velocity curves for disk and halo components and their sum are given in Figure 1. The gravitational softening used is 160 pc, and Toomre's parameter $Q = 1.5$. The adopted units are those of $G = 1$, mass $M = 10^{11} M_\odot$, and distance $r = 10$ kpc. This leads to the time unit of $\tau_{\text{dyn}} = 4.7 \times 10^7$ yr and a velocity unit of 208 km s^{-1} . The energy and angular momentum in the system are conserved to within approximately 1% and 0.05% accuracy, respectively. Above $N \sim 10^5$, our results appear to be reasonably independent of N . The model evolution presented here has $N = 1.1 \times 10^6$, with 8×10^5 particles in the disk.

2.2. Orbital and Spectral Analysis

The self-consistent evolution of stellar bars can be only understood by studying their three-dimensional structure (e.g., Pfenniger & Friedli 1991; Skokos et al. 2002a, 2002b). For such in-depth investigation we use the updated algorithm described in Heller & Shlosman (1996), which is based on a comprehensive search for periodic orbits in arbitrary gravitational potentials, and display them in characteristic diagrams (§ 4). These orbits close in the bar frame and provide the backbone for any meaningful analysis of the bar properties. Periodic orbits characterize the overall orbital structure of the bar phase space, because each of them traps a region of phase space around it. These trapped orbits have shapes similar to the shapes of the parent periodic orbit. The algorithm is run using the package Parallel Virtual Machine (PVM), which distributes the search for orbits among different processors and computes the orbits using adaptive step size, with a relative accuracy of 10^{-7} . We also calculate the stability of these periodic orbits to estimate which orbits can be populated. We track both the two-dimensional orbital families and the three-dimensional families bifurcating from planar (i.e., equatorial) orbits at the (vertically) unstable gaps. The shapes of these three-dimensional orbital families will contribute to the shape of the simulated bar, when populated, including the evolving shape of the growing boxy/peanut-shaped bulge.

To quantify the disk-halo interaction and the angular momentum redistribution in the system, we have developed a package that is based on the orbital analysis algorithm and that uses the fast Fourier transform (FFT) to find the main orbital frequencies: the angular frequency, Ω , and the radial and vertical epicyclic frequencies, κ and ν , in the disk and the halo. The particle distribution with the frequency ratio is then determined to find the population of resonant orbits. Finally, the change in the angular momentum is computed for each of these particles (see § 5).

3. RESULTS

We first describe the overall bar evolution during the simulation period of $\tau \sim 14$ Gyr and analyze some of its more important behaviors. The bar develops in an otherwise axisymmetric model during the first two to three disk rotations. Initially it has an axial ratio (flatness) of $c/a \sim 0.1$, the same thickness as the disk, but this ratio increases with time by a factor of ~ 2 . The bar starts to brake against the outer disk and the halo as seen in Figure 2b, reaching maximum strength at $\tau \sim 1.4$ Gyr. At around $\tau \sim 1.8$ Gyr the bar experiences a vertical buckling instability that affects its two-dimensional and three-dimensional appearance. The outer part of the bar nearly dissolves, while overall the bar is weakened dramatically, as shown by the $m = 2$ amplitudes, A_2 , in Figure 2a. Immediately following this buckling, the bar resumes its growth, which saturates again at $\tau \sim 6$ Gyr. At this time A_2 in the outer part weakens again. The growth is resumed after $\tau \sim 7.5$ –8 Gyr. A close inspection of the edge-on bar frames during time intervals of 1.8–2.8 and 6–7.5 Gyr and the analysis described in the next sections reveals sufficient similarities between these two events; both represent the vertical buckling instability in the bar, i.e., the breaking of symmetry of the bar (Fig. 3). Such a recurrent buckling of bars has never been reported in the literature.

3.1. Secular Growth of Stellar Bars

The stellar bar evolution presented here is characterized by substantial changes in the bar size and strength and by changes in its three-dimensional shape (see Fig. 3 and accompanying animation). Determination of the live bar size in numerical simulations is not trivial. For example, Athanassoula & Misiriotis (2002) and O'Neill & Dubinski (2003) used a variety of methods and found that some of them give erroneous and unreliable results. The position of the $A_2(r)$ maximum does not provide any meaningful estimate for the bar size because the contribution of higher harmonics, such as $m = 4$ and 8 , is neglected. To quantify the bar size changes, we have used two alternative methods, an ellipse fitting to the isodensity curves and the characteristic orbital diagrams. The former method has been previously used to detect and to characterize “observational” bars (e.g., Knapen et al. 2000; Laine et al. 2002; Hunt & Malkan 2004). Its main deficiency when applied to numerical bars is an excessive noise for low to moderate values of N and the resulting distribution of the bar ellipticities, $\epsilon(r)$, being flat for young unbuckled bars (Martinez-Valpuesta & Shlosman 2004). However, at later times, with the growth of the central mass concentration, this method becomes more reliable. We find that a consistently reliable estimate of the bar size, r_{bar} , is the radius at which $\epsilon(r)$ declines $\sim 15\%$ from its maximal value. The alternative, and a new method used by us here, relates the bar extent to the size of the maximal stable orbit of the main orbital family supporting the bar (more in § 4). Both methods produce consistent results with each other.




FIG. 2.—(a) Evolution of the bar $m = 2$ amplitude, A_2 , for the inner and outer bar parts, $r = 0$ –11 kpc (thick solid line) and $r = 7$ –11 kpc (thin solid line). (b) Bar pattern speed Ω_b . [See the electronic edition of the Journal for a color version of this figure.]

The bar size evolution is shown in Figure 4a for both methods. It grows initially to 11 kpc, then buckles and shortens to 6 kpc. It then grows again to 13 kpc, where the growth stagnates due to the secondary buckling, for about 3 Gyr. After this the bar resumes its growth to about 16 kpc. The size evolution in the xy -plane is accompanied by the vertical thickening of the bar. It does not stop after the first buckling instability, but continues gradually due to the vertical resonance scattering, amplified by the recurrent buckling instability of the bar. The accompanied boxy/peanut-shaped bulge also grows with time (Fig. 3).

With the bar length, r_{bar} , and its corotation radius, r_{CR} , we can quantify some of the dynamical characteristics of an evolving bar. The ratio $r_{\text{CR}}/r_{\text{bar}}$ is shown in Figure 4b. This ratio determines the shape of the offset dust lanes in barred galaxies that delineate shocks in the gas flow (Athanassoula 1992). The observed shapes constrain the ratio $r_{\text{CR}}/r_{\text{bar}}$ to 1.2 ± 0.2 . The modeled ratio typically falls within the required limits except during the first buckling, when it is higher, ~ 1.5 , as noted already by Martinez-Valpuesta & Shlosman (2004). We discuss this issue in more detail in § 6.

3.2. Recurrent Buckling of Stellar Bars

The buckling is a three-dimensional phenomenon and is most visible in the vertical xz -plane (Fig. 3). To quantify the bar asymmetry, we have calculated the vertical $m = 1$ mode amplitude A_{1z} in the xz -plane and follow the maximal distortion of the bar during its buckling periods. Two maxima are apparent at ~ 2.4 and ~ 7 Gyr in Figure 5. Note that the vertical asymmetry given by this figure builds up slower and is weaker for the second buckling; i.e., from $\tau \sim 5$ Gyr to $\tau \sim 7$ Gyr, the amplitude grows to $A_{1z} \sim 0.03$, compared to $A_{1z} \sim 0.08$ at $\tau \sim 2.4$ Gyr. Figure 3 shows this evolution in a more graphical way: while the first buckling affects mostly the central few kiloparsecs, especially the bar’s midplane, the second buckling is most prominent in the outer bar range of 5–10 kpc and affects the midplane much less visibly.




FIG. 3.—Evolution of the vertical structure in the bar: edge-on view along the bar’s minor axis. The length is given in kpc and the values of the projected isodensity contours are kept unchanged in all frames. The time in Gyr is given in the upper right corners. The maximal vertical asymmetries correspond to two recurrent bucklings at $\tau \sim 2.4$ Gyr and at ~ 7 Gyr. Note the bar flip-flop between $\tau = 2.3$ and 2.4 Gyr, the persistent vertical asymmetry in the bar at $\tau = 5.2$ –7.5 Gyr, and the development of narrow features in the bar midplane outside its core of ~ 8 kpc after $\tau \sim 9.4$ Gyr. Those correspond to ansae (“handles”) in the face-on bar (see Fig. 10) and are observed in early-type disk galaxies. [See the electronic edition of the Journal for a color version of this figure. This figure is also available as an mpeg animation in the electronic edition of the Journal.]

The recurrent buckling can be detected in a number of ways, e.g., from A_2 in the xy -plane (Fig. 2a), as mentioned in § 3.1 and above, and from A_{1z} in the xz -plane, which quantifies the breaking of vertical symmetry in the bar (Fig. 5). Each of the coefficients emphasizes a different property of this instability. We are basically looking at the same phenomena at different times: the first buckling extends over ~ 1 Gyr and the second one over ~ 3 Gyr. The changes in the orbital structure of the bar during the bucklings will be analyzed in § 4.

We have commented in § 1 that the buckling instability is a collective breaking of a vertical symmetry in the bar. Toomre (1966) has shown that the coupling between the vertical and radial degrees of motion is the prime driver of this instability and that the main outcome of it is the equalizing of the velocity dispersion in the xy -plane with the vertical velocity dispersion.⁵

This evolution is characterized by changes in the vertical-to-radial velocity dispersion ratio σ_z^2/σ_r^2 . Toomre estimated the critical value for this ratio to lie at ~ 0.1 for a nonrotating plane-parallel slab, Raha et al. (1991) put it at 0.06–0.3 for a three-dimensional stellar disk, and Sotnikova & Rodionov (2005) estimated it at ~ 0.6 for the central regions embedded in the hot halos. Sellwood (1996) have shown that Toomre’s limit is violated for many of his stellar models, some remaining unstable at ratios up to 0.4. We plot this ratio for two areas of the bar at two different times. For the first buckling (Fig. 6), the velocity dispersions are calculated in the central kiloparsec, where the maximum effect is expected (e.g., Fig. 3). During this buckling, we observe first an increase in σ_r (the initial growth of the bar) and after ~ 1 Gyr a decrease in σ_r with a corresponding increase in σ_z . When σ_z^2/σ_r^2 drops below ~ 0.4 , the bar buckles and weakens. The buckling ends when σ_z^2/σ_r^2 increases to unity. The bar thickens and grows and the vertical ILR moves gradually out. We expect and observe the maximal vertical asymmetry in the outer part of the bar during the second buckling and therefore

⁵ Normally, the kinetic energy of oscillations about the equatorial plane, i.e., along the z -axis, is an adiabatic invariant.




FIG. 4.—(a) Evolution of the bar size (semimajor axis). The solid line represents the bar size obtained from the ellipse fitting to the isodensities. The filled circles correspond to the bar sizes obtained from the semimajor axis of the last stable x_1 periodic orbit that supports the bar obtained from Fig. 7, which is a new method introduced here (see § 4 for details). (b) Evolution of the bar corotation-to-size ratio. (c) Evolution of the maximal ellipticity of the bar, $\epsilon = 1 - b/a$, from the ellipse fitting. [See the electronic edition of the Journal for a color version of this figure.]




FIG. 5.—Evolution of the vertical buckling amplitude in the bar, A_{1z} , i.e., the vertical $m = 1$ mode in the xz -plane, integrated over the intervals $r \sim 0$ –12.5 kpc, $-2 \text{ kpc} < y < 2 \text{ kpc}$, and $-\infty < z < \infty$.




FIG. 6.—Evolution of (a) the vertical-to-radial dispersion velocity ratio in the disk, σ_z^2/σ_r^2 , within the central kiloparsec (left) and within a cylindrical shell of radius 7 ± 1 kpc (right), and (b) separately, the radial, σ_r (dashed line), and the vertical, σ_z (dotted line), dispersion velocities within the same regions as in (a). [See the electronic edition of the Journal for a color version of this figure.]

calculate σ_z^2/σ_r^2 at around 7 kpc. Again, the gradual increase in σ_z and decrease in σ_r drives the ratio up from 0.4 to about unity, similar to the first buckling. For both bucklings we observe a very similar evolution in terms of the velocity dispersions and their ratios.

So far we have shown that the bar buckles twice during the simulations: the first time abruptly, fast, and in the central part, and the second time slower, less pronounced, and in a different part of the bar. In both bucklings we observe the loss of symmetry in the vertical plane, the drop in the value of A_2 and the equalizing of the vertical and radial velocity dispersions. After each buckling the bar becomes more symmetric, with the difference that after the first buckling the asymmetry starts to build anew, and after the second buckling the asymmetry is completely washed out.

To summarize, three main factors characterize the end of the buckling instability: A_2 starts to decrease, the asymmetry in the rz -plane decreases (given by A_{1z}), and $\sigma_z^2/\sigma_r^2 \rightarrow 1$. When we apply these conditions to our model, the end of the first buckling appears at ~ 2.4 – 2.8 Gyr and the end of the second buckling at ~ 8 Gyr.

Since the buckling leads to a sudden vertical thickening of the bar and is therefore characterized by particle injection above the disk plane, we can monitor this instability by following the particle distribution. When viewed along the bar minor axis, during the first buckling the bar bends and develops a boxy/peanut-shaped bulge, while during the second buckling the bar acquires an asymmetric shape that leads to the appearance of an X-shaped bulge (Fig. 3). These shapes have a direct relationship to the population of orbits trapped by the main family of periodic orbits in the bar (Pfenniger & Friedli 1991; § 4 below).

4. ANALYSIS: BAR ORBITAL STRUCTURE EVOLUTION

Stellar bars are three-dimensional objects that exhibit dynamical and secular evolutionary trends both in their morphology and in their internal structure. In this section, we study the evolution of the orbital structure in the bar by searching for the main two-dimensional and three-dimensional families of orbits at various snapshots, with a particular emphasis on the recurrent buckling periods. A comprehensive search for the orbits is beyond the scope of this work. We analyze the main parameters of detected orbits and calculate their stability. This section uses a




Fig. 7.—Three snapshots of the bar orbital structure evolution: *left*, immediately after the first buckling; *middle*, during the second buckling; *right*, well after the second buckling. The $\tau = 7.1$ Gyr frame is nonsymmetrized (vertically), while the others are vertically symmetrized. (a) Characteristic diagrams in the xy -plane showing y -intersections of regular orbits with the $x = 0$ axis as a function of Jacobi energy of the orbits. The bar is oriented along the x -axis. Solid lines represent stable orbits, and dotted lines represent unstable ones. The dashed lines show the zero-velocity curves (ZVCs). (b) Characteristic diagrams in the xz -plane showing z -intersections of regular orbits. Since the 7.1 Gyr snapshot is nonsymmetrized and the bar buckles in this plane, the distribution of the BAN orbits is also asymmetric, and most of the stable BAN orbits are concave down (see text). They have a larger curvature and extend further into the $z < 0$ region, leading to the bar asymmetry in the edge-on view (Fig. 3). (c) Characteristic diagrams in the xz -plane showing the \dot{z} families. The 7.1 Gyr snapshot is omitted because all the ABAN orbits are unstable at this time and, therefore, are difficult to calculate in unsymmetrized potential.

specific terminology developed for nonlinear orbital dynamics (e.g., Binney & Tremaine 1987; Sellwood & Wilkinson 1993). Readers unfamiliar with this terminology can skip it and go directly to § 5. We discuss the results of the orbital analysis in the context of the bar evolution in § 6.

The initial vertical axial ratio of the bar that develops in our models is $c/a \sim 0.1$ (nearly a two-dimensional object) and grows dynamically and secularly to ~ 0.2 over the Hubble time. We first analyze the bar's orbital structure in the equatorial plane, then add the three-dimensional effects accounting for the finite thickness and shape of the bar. In doing so, we limit our analysis to the periodic orbits within r_{CR} , which are largely responsible for the bar shape, and to specific times of bar evolution: $\tau = 1.4$ Gyr (just prior to the first buckling), 2.8 Gyr (after the first buckling), 7.1 Gyr (during the second buckling), and 11.8 Gyr (after the second buckling). The orbits are searched in the potential symmetrized horizontally with respect to the four quadrants. Figure 5 confirms that the vertical symmetry is violated during the buckling periods. The first buckling is fast and there is little meaning, therefore, to calculating the orbits without some frame averaging; the stars see only a time-averaged potential. Hence, we resort to a vertically symmetrized potential at 2.8 Gyr. On the other hand, the second buckling is much more gradual, and we calculate the orbits in the actual "raw" potential at $\tau = 7.1$ Gyr to capture the persisting asymmetry. The third snapshot, in which the bar is symmetric again, is treated similarly to the first snapshot for simplicity.

The extent of the orbital families and their stability are displayed by means of characteristic diagrams. The y , z , and \dot{z} intercept values with the $x = 0$ plane are plotted in Figure 7 with

respect to their Jacobi integral, E_J (e.g., Binney & Tremaine 1987). The Jacobi (energy) integral of motion is conserved along any given orbit in the rotating bar frame. The orbits form curves or families in the characteristic diagrams. The actual trajectories do not coincide exactly with these periodic orbits but may be "trapped" in their vicinity. The properties of periodic orbits and their temporal changes, therefore, reflect the bar structural evolution.

The most important single-periodic orbits in the bar midplane are those aligned parallel or normal to the bar's major axis, the so-called x_1 and x_2 orbits (e.g., Contopoulos & Papayannopoulos 1980; Binney & Tremaine 1987). The x_1 family constitutes the main family of orbits that support the bar figure. While this family exists always within r_{CR} , the x_2 family appears only if the planar ILRs are present. Typically, numerical bars form with a pattern speed sufficiently high to avoid the ILRs, at least in the beginning, and hence the x_1 family completely dominates the bar midplane, short of the corotation region.

The vertical shape of the bar is determined by the projection of the populated three-dimensional orbits onto the corresponding planes. We search for the vertically unstable gaps in the x_1 family. It is at these gaps where the three-dimensional families bifurcate through z and \dot{z} bifurcations (e.g., Pfenniger 1984; Pfenniger & Friedli 1991; Skokos et al. 2002a, 2002b; Patsis et al. 2002a). These gaps coincide with the vertical ILR and other vertical resonances. When an orbital family goes from being stable to unstable, in the mentioned gaps, we get the three-dimensional prograde families, i.e., orbits with initial conditions $(y, z, \dot{y}, \dot{z}) = (a, b, 0, 0)$, with a and $b \neq 0$. When it goes from being unstable to stable, we get the retrograde families,

$(y, z, \dot{y}, \dot{z}) = (c, 0, 0, d)$, with c and $d \neq 0$, where x, y , and z are oriented along the bar's major, minor, and vertical axes.

While the vertical ILR is associated with the boxy/peanut bulge shapes, which develop as a result of the resonance heating of midplane orbits, Patsis et al. (2002a) have found that this effect is not limited to a particular resonance and not to the barred galaxies per se, but operates equally well in nearly axisymmetric and/or ovally distorted disks. Furthermore, the buckling itself is not a necessary condition for these shapes to form, but rather accelerates the process from a secular to a dynamical timescale.

The main families of three-dimensional orbits that are responsible for the appearance of the boxy/peanut-shaped bulges are the BAN (prograde to the bar rotation) and ABAN (retrograde) families found by Pfenniger & Friedli (1991). They are called x_1v_1 and x_1v_2 , respectively, and appear as the three-dimensional generalizations of the x_1 and x_2 families in the nomenclature of Skokos et al. (2002a). In general terms, the BAN orbits can be described as concave down ($z > 0$) and concave up ($z < 0$), and ABAN as “ ∞ ,” when projected onto the xz -plane. Both families are $m:n:l = 2:2:1$ (i.e., two radial oscillations for two vertical oscillations for one turn) in the notation of Sellwood & Wilkinson (1993). Their projections onto the xy -plane have the x_1 orbit shapes. We trace the planar x_1 and x_2 families and the three-dimensional BAN and ABAN families from the bar initial growth period. For simplicity, we have divided the simulation into a number of characteristic time intervals and discuss them separately.

Before and during the first buckling: $\tau \simeq 0\text{--}2.5$ Gyr.—During this period the bar develops and buckles (Figs. 2–6). It is geometrically thin and is not centrally concentrated, so the vertical ILR is not present. Neither the planar ILRs nor consequently the x_2 orbits are present. The stable three-dimensional orbits lie close to the midplane. As we stated above, there is little meaning in calculating the orbits in the rapidly varying potential without the proper time-averaging. Because the midplane of the bar is bent in the xz -plane (Fig. 3), so are the x_1 orbits.

After the first buckling: $\tau \simeq 2.5\text{--}3.5$ Gyr.—The bar has increased its central mass concentration and acquired its boxy/peanut-shaped bulge (see Fig. 3). As we stated above, there is little meaning in calculating the orbits in the vertically unsymmetrized potential at the time of the first buckling because of the rapidly varying potential. Moreover, the bar midplane is bent in xz at this time, and so are the x_1 orbits (Fig. 3). Hence, we have vertically symmetrized the potential at this time frame (Fig. 7b, left). In the midplane, the x_1 orbits dominate and no x_2 family exists (Fig. 6a). Most of the x_1 orbits are stable, except for a small gap, where the vertical ILR is located. At $\tau = 2.8$ Gyr the gap is at $E_J \sim -2.66$, where the BAN and ABAN orbits bifurcate (Fig. 7a). The BAN and ABAN orbits are traced (Figs. 7b and 7c): their origin in the $z = 0$ plane moves gradually out with time toward higher Jacobi energies. By the end of the first buckling it stabilizes and subsequently creeps toward higher values of E_J as the bar brakes against the halo. BAN orbits appear stable from 300 pc above the midplane, with a broad unstable gap at $z \sim 1.1\text{--}1.3$ kpc and a narrow stability island (Fig. 7b). Initially, the stable part of these families is limited to a very small height above the bar midplane, compared to the bar vertical thickness. Gradually the stable extent of the BAN family increases both in z and in x (Fig. 8). The ABAN family is unstable when BAN is stable and vice versa (Fig. 7c), as first noted by Pfenniger & Friedli (1991).

Before and during the second buckling: $\tau \simeq 3.5\text{--}8$ Gyr.—The bar is growing both in size and amplitude and exhibits a pronounced boxy profile when viewed along its minor axis. It ap-




FIG. 8.—Projection of three-dimensional orbits onto the xz -plane in three different snapshots: after the first buckling, during the second buckling, and after the second buckling. Only *stable* orbits of the main families described in the text and in Fig. 7 are plotted. The horizontal extension of the orbits increases with time. The general shape evolves from a peanut/boxy shape to a vertically asymmetric shape and then to a peanut/X-shape.

pears symmetric with respect to the bar midplane up to $\tau \sim 5$ Gyr (Fig. 3), although A_{1z} shows that some small residual asymmetry remains between the bucklings (Fig. 5). After $\tau \sim 5$ Gyr, the vertical symmetry of the bar is broken for an extended period of time, ~ 3 Gyr (Figs. 3 and 5). The maximal distortion happens at ~ 7.1 Gyr. No x_2 orbits exist during this time period, and the unstable gap in x_1 moved to somewhat higher energies, approximately $-2.4 \lesssim E_J \lesssim -1.7$, due to the bar braking (Fig. 7a). An additional unstable gap has appeared at $-2.68 \lesssim E_J \lesssim -2.65$, where the 3:2:1 orbit family bifurcates.

Note that the analysis shown in the $\tau = 7.1$ Gyr frame of Figure 7 is performed in the actual gravitational potential of the N -body simulation, i.e., without symmetrization with respect to the $z = 0$ plane. Therefore, we should not expect to find all the families existing in the symmetrized potentials. But we should be able at least to identify the reason for the vertical asymmetry of the particle distribution in the projection of the three-dimensional orbits onto the xz -plane from the orbital shapes (Fig. 8b). We find the BAN family and also the x_1v_9 family, in the nomenclature of Patsis et al. (2002b).

The BAN family appears stable everywhere above the midplane, except around $z \sim 2$ kpc (Fig. 7b), while the ABAN family is unstable everywhere and so is not shown here. For $z \lesssim 0$, the BAN family is detached from the midplane and starts at $z \sim -0.5$ kpc. It forms a kind of loop in the characteristic diagram. We do not find any stable orbit below $z \sim -2$ kpc. The x_1v_9 family is stable only for a low range in energies and only below the midplane, which makes it important for contributing to the vertical asymmetry in the bar (Fig. 8b).

After the second buckling: $\tau > 8$ Gyr.—The bar slowly regains its vertical symmetry and becomes symmetric after ~ 8 Gyr. As a result of this instability, the bar has developed a pronounced X-shape, which differs from the “usual” peanut shape because of the extended concave region between the spikes, i.e., peanuts (Fig. 8c). Orbit that appear at the same z -value in the characteristic diagram but at different time frames differ in their projections. After the second buckling the orbits are longer and extend higher; i.e., the convexity and concavity have significantly changed. The vertical ILR has moved to $E_J \sim -2.36$ by 11.8 Gyr. The BAN orbits are stable, while the ABAN orbits are unstable. Overall, the boxy/peanut-shaped bulge has increased in size from $\sim 1\text{--}2$ kpc (after the first buckling) to ~ 8 kpc (after the second buckling).

The x_2 orbits and consequently two ILRs in the xy -plane appear only after the second buckling, $\gtrsim 7.2$ Gyr. Initially occupying a small range in energies, their extent grows with time. The x_1 family is mostly stable up to the (mostly) unstable “shoulder” in the characteristic diagram (Fig. 7a, right), which has continued to move out to higher Jacobi energies, along with the last stable orbit supporting the bar.

We also use the characteristic diagrams to get an independent estimate for the bar’s physical size. Unlike the case of analytical bars, where this issue is resolved trivially, to estimate the length of the *live* numerical and “observed” bars can be more difficult. For example, if the bar potential or bar nonaxisymmetric force is used for this purpose, one can get an erroneous result that the bar extends beyond its corotation radius. If one relies on the density distribution, one can overestimate the size as well, because the bar can drive a pair of open spirals. Instead, we use the calculated orbital structure of the bar and rely on the properties of the x_1 family, which is generally stable except for narrow gaps (in Jacobi energy) and in the region around the corotation. Specifically, close to the corotation, the x_1 curve bends upward (e.g., Fig. 7a; Binney & Tremaine 1987). We use the x -axis extent of the last stable x_1 orbit that lies on this upward branch. It is the slow drift of unstable gaps and of the elbow of the x_1 -curve toward higher E_J that leads to the trapping of orbits and the secular increase in the bar size. A large number of models here and in Berentzen et al. (2006) have been used for the comparison with the isodensities fitted to the bar (§ 3).

The bar amplitude A_2 shows that the outer half of the bar practically dissolves in the first buckling and weakens substantially in the recurrent buckling (Fig. 2). The ellipse fit to the bar reflects this by a dramatic decrease in the bar length from ~ 11 to ~ 6 kpc in radius. Unfortunately, it is more tricky to estimate the bar size from the orbital analysis at this time: the underlying potential is time-dependent, while the analysis is performed in the frozen potential. Nevertheless, both methods agree before and after the first buckling and capture the change in the growth rate of the bar length after the second buckling. To summarize, the bar size decreases during the first buckling and levels off during the second one.

The simple explanation for the success of the orbital analysis method is that it is the most self-consistent method known to us and relies on the orbit trapping by the stable periodic family whose existence basically defines the bar. We note that the middle frame of Figure 7a does show a large unstable gap just below the elbow of the x_1 curve. However, we note once more that the potential in this frame has not been vertically symmetrized and that the weakening of the bar during this time of buckling is in fact a direct consequence of this gap.

In summary, the characteristic diagrams and their respective orbital families reflect both the dynamical and secular evolution

of the bar shown in Figure 3 and the accompanying animation. The unstable gaps, where the vertical families (BAN and ABAN) bifurcate, move toward higher Jacobi energies, while the orbits become more extended along the bar axes. The orbital families evolve not just quantitatively but also qualitatively; this shows up in the changing shape of the bar. Although we do not measure the population of the different families, the bar shapes appear to be governed mainly by the BAN family, which evolves by changing the concavity of its orbits, thus giving the bar the pronounced peanut/boxy/X-shapes when viewed along the minor axis. The change in the orbital structure of the bar supports our observation, based on independent arguments, that there is a recurrent buckling. At the second buckling time, the vertical families differ above and below the plane. The former are stable orbits extending more along the x -axis and giving it more of a boxy shape. The latter ones are stable orbits that give the bar more of a peanut shape. The appearance of the x_2 family confirms that the ILRs form only after the second buckling of the bar.

5. DISK-HALO ANGULAR MOMENTUM EXCHANGE: THE ROLE OF THE RESONANCES

The disk region that gives rise to a bar is known to lose its angular momentum (J). In principle, a number of components in a galaxy can acquire this momentum, namely, the outer disk, the bulge, and the halo. While the former two components are able to store J , the responsive (i.e., live) halo can serve as a particularly large angular momentum sink due to its large mass and low J . Athanassoula (2002a, 2003) has shown that the disk-halo interaction is mediated by lower resonances. In order to understand more fully the complex behavior exhibited by the model, such as the initial bar growth, its recurrent bucklings, and subsequent secular growth, we find it instructive to determine the balance of the angular momentum and its evolution. We pay particular attention to the resonant interactions between the model components.

We first determine the overall J balance in the different regions of the disk and the halo. Figure 9 shows the total angular momentum evolution for three different regions in the disk and halo: the inner (0–7 kpc), intermediate (7–15 kpc), and outer (15–35 kpc or 15–50 kpc) regions. The inner and intermediate regions are those hosting the bar for various times during its evolution. The time of the bar formation is characterized by a substantial loss of J in the inner disk. This angular momentum is redistributed to the intermediate region and especially to the outer disk (Fig. 8). During the first buckling, there is some indication that the inner disk gains some angular momentum, and we return to this issue in § 6.2. The subsequent growth of the bar is accompanied by a slow J growth in the outer disk and the halo. Interestingly, after the second buckling J stops growing in the outer disk, while the outer halo picks up J at an increasing rate. There is a general loss of angular momentum from the disk and there is a general gain in the halo. The loss of J in the disk correlates with the slowdown of the bar, which is known to anticorrelate with the growth of the bar (e.g., Athanassoula 2003).

Next we calculate the contribution of the resonances between disk and halo particles with the bar’s Ω_b to the angular momentum exchange between the disk and the halo. We generally follow the procedure described in Athanassoula (2002a). The principal frequencies, Ω and κ , of azimuthal motion and of radial oscillations of the three-dimensional orbits, respectively, have been determined by means of spectral analysis (Binney & Spergel 1982). We freeze the gravitational potential in the model at times $\tau = 2.8$ and 5.2 Gyr but allow the bar to tumble with its




FIG. 9.—Evolution of angular momentum in the various regions of the disk (top) and the halo (bottom). The specified radii are given in cylindrical geometry. [See the electronic edition of the Journal for a color version of this figure.]

original Ω_b and integrate the orbits of 1.5×10^5 randomly picked particles in the disk and the same amount in the halo for an additional 9.4 Gyr, about 20 bar tumblings. Results are plotted in Figure 10 as a function of the frequency ratio $\eta \equiv (\Omega - \Omega_b)/\kappa$, where N_R particles are binned in the intervals of $\Delta\eta = 0.005$ (Fig. 10, top). The difference ΔJ in angular momentum between the two times is given as a function of η at $\tau = 5.2$ Gyr (Fig. 10, bottom). The major resonances are located at $\eta = \pm 0.5$ (ILR and OLR), $\pm 1:3$, $\pm 1:4$, etc., with the positive η indicating the resonances inside the corotation radius and $\eta < 0$ outside the corotation. The corotation resonance corresponds to $\eta = 0$.

The resonant and near-resonant particle distributions N_R at the time $\tau = 2.8$ Gyr are clearly nonuniform and are permeated by numerous resonances. The trapping of orbits by the resonances is also evident. The dominant resonance in the disk appears to be the ILR, which traps the largest number of particles and therefore facilitates the loss of the angular momentum from the disk (Fig. 10, bottom). Note that we have cut N_R for this resonance to keep the same scales for disk and halo diagrams: it extends to 39×10^3 . The x_1 orbits within the CR and, later on, the x_2 (if populated) orbits, as well as the BAN and ABAN orbits, are trapped within this resonance. The broad asymmetric peak (Fig. 10, top left) in the range of $\eta \sim \pm 0.35$ – 0.45 is made out of near-

resonant orbits in the (rotating) boxy/peanut-shaped bulge, many of them near-resonant BAN orbits. Those have been injected during the first buckling. They appear on both sides of the CR because J along these orbits oscillates wildly, $\Delta J \sim J$, and it is sometimes difficult to disentangle the prograde from retrograde ones among them. For comparison, we repeated this procedure at $\tau = 11.8$ Gyr and confirmed that more than half of the particles in the above broad peaks have been trapped by the nearby ILR and the outer Lindblad resonance (OLR) by that time. The resonances for $\eta \lesssim 0$, with CR and OLR as the next strongest ones, absorb the angular momentum, and their values of ΔJ are all positive.

The halo is dominated by the corotation resonance, which absorbs J from the disk. Overall, the disk resonances emit and the halo resonances absorb J . This behavior was demonstrated by Athanassoula (2002a, 2003), where similar plots to Figure 10 were given, and there is a very good agreement between our results. The only difference we find is that the halo ILR is actually losing J in our model, albeit a small amount, unlike in Athanassoula's models. The modeled halo extends well beyond the disk, and its outer part appears to be actively storing J , up to $\eta \sim -3$. This effect is confirmed by the overall J evolution in the halo (Fig. 9). We find that the trapping of orbits by the resonances is robust: almost all the particles trapped at $\tau = 2.8$ Gyr remain trapped at 5.2 Gyr. The ability of the barred disk to transfer its angular momentum to the halo while J saturates in the outer disk explains why the bar is able to grow during this time interval as exhibited by the A_2 amplitude and the size of the bar (e.g., Figs. 2–5). It is this growth of the bar that is ultimately responsible for the recurrent buckling phenomenon analyzed here.

6. DISCUSSION

We have modeled the evolution of collisionless (stellar) bars embedded in responsive (live) axisymmetric dark matter halos using high-resolution N -body simulations. We find that bars experience secular growth over the simulation (\sim Hubble) time, except during well-defined time periods when the bars encounter spontaneous breaks of vertical symmetry, so-called buckling (or fire-hose) instability (e.g., Fig. 3 and accompanying animation). We detect such a recurrent buckling during which the bar, and especially its outer half, weakens substantially, but the growth is resumed subsequently. Two different techniques have been used to measure the bar size, namely, the nonlinear orbital analysis and the isodensity ellipse fitting. The bar strength has been measured using the amplitude of the $m = 2$ mode, and the bar vertical asymmetry has been measured using the vertical $m = 1$ mode. Moreover, we have analyzed the bar evolution and its buckling periods by means of the nonlinear orbit analysis and the ratio of vertical to planar velocity dispersions. Finally, we have examined the bar development in terms of the angular momentum redistribution between various components in the disk-halo system mediated by the resonant interactions using orbital spectral analysis.

Low-resolution N -body simulations exhibit fast (e.g., during one disk rotation) stellar bar growth, followed by a vertical buckling and secular weakening. High-resolution simulations with $N \gtrsim 10^5$ – 10^6 show a more complex evolution. They allow for modeling the disks embedded in live halos and account for the resonant interactions between the halo and the disk, adding a new and crucial element to the simulations in the form of angular momentum transfer between these components. The main difference between the previous low-resolution models and that presented in this work is the ability of the stellar bar to strengthen again after its original weakening following the buckling, a




FIG. 10.—Resonance interaction in the live disk-halo system. *Top:* Histogram of particle distribution N_R with the principal frequency ratio $\eta \equiv (\Omega - \Omega_b)/\kappa$ for the disk and (rotating) boxy/peanut-shaped bulge (*left*) and for the halo (*right*), with about 1.5×10^5 randomly picked particles each, at $\tau = 2.8$ Gyr. The cusp at the ILR ($\eta = 0.5$) for the disk is shown only partially; it extends to $\sim 39 \times 10^3$. The broad peaks near $\eta \sim \pm 0.4$ are made mostly of near-resonant bulge particles injected during the first buckling. *Bottom:* Angular momentum difference, ΔJ , between the particles at $\tau = 5.2$ and 2.8 Gyr in the disk (*left*) and halo (*right*) as a function of η at a time of 5.2 Gyr. The lower resonances, such as the ILR ($\eta = 0.5$), the corotation ($\eta = 0$), the OLR ($\eta = -0.5$), $1:3$ ($\eta = \pm 0.33$), $1:4$ ($\eta = 0.25$), etc., are clearly visible. The ILR dominates the loss of the angular momentum in the disk, and the corotation dominates the absorption of the momentum by the halo (see more in the text). The frequency ratio is binned into intervals of $\Delta\eta = 0.005$. Note that the η scales are different for the disk and the halo.

process that leads to the bar growth and consequently to its recurrent buckling.

6.1. Disk-Halo Resonant Interactions

The bar growth has been associated with the existence of “sinks” of angular momentum located elsewhere. Athanassoula (2002a, 2003) has shown that a galactic halo can play such a role and absorb large quantities of angular momentum from the disk/bar region. This effect appears to completely invert the original suggestion by Ostriker & Peebles (1973) about the stabilizing function of dark matter halos in disk galaxies against the bar formation instability. Angular momentum redistribution, at least in principle, can have contributions both from the nonresonant and resonant interactions between the bar and orbits in the halo. To capture the latter requires large N to stabilize the population of resonant particles (e.g., Weinberg & Katz 2002).

It is therefore important to demonstrate that resonant particles are indeed responsible for the angular momentum transfer in the first place. We start with the relative contributions to the J transfer between the disk and the halo and between the inner (i.e., bar unstable, ~ 15 kpc) and outer disks (Fig. 9). Although the efficiency of this redistribution is of course model dependent, we nevertheless quote the numbers, assuming that to a certain degree, they are representative. We find that $\sim 39\%$ of the angular momentum in the disk is lost to the halo during the evolution. The bar unstable region, $\lesssim 15$ kpc, has lost about 71% of its original J , of which about 38% went to the outer disk and the

rest was absorbed by the halo. The most intensive flow of J to the outer disk happens in the early stage, which ends with the first buckling. The halo particles have much larger dispersion velocities than the disk, and until the bar fully develops, their resonant interaction with the $m = 2$ asymmetry in the disk is virtually nonexistent.

After the first buckling, it is the halo that absorbs most of the angular momentum lost by the bar region, and from Figure 9 (*bottom*) it is clear that the lion’s share of this exchange is resonant and mediated by the CR and the OLR in the halo and by the ILR and the CR in the disk, with some contribution from the numerous minor resonances. This results in the increase of the bar’s size and strength and leads to the gradual decrease in the ratio of the velocity dispersion, σ_z^2/σ_r^2 , in a close analogy with the disk evolution preceding the first buckling. The second buckling of the bar, therefore, can be directly traced to its robust growth in our model.

The issue of a bar dissolution versus growth has become controversial recently. While it is beyond the scope of this work, we do indirectly touch on it. First, we confirm the results of Martinez-Valpuesta & Shlosman (2004) that vertical bucklings do not destroy the bar, unlike those suggested by Raha et al. (1991; see also Sellwood & Wilkinson 1993). Next, in various axisymmetric models, here and in Berentzen et al. (2006), a robust growth in the bar is observed after the first buckling, supplemented by the angular momentum exchange between the disk and the halo. Athanassoula (2003) has analyzed the effect of

initial conditions on this J redistribution, namely, of the initial disk-to-halo mass ratio and of the Toomre Q parameter: the angular momentum exchange varied from $\sim 2\%$ to $\sim 35\%$, with a clear correlation between the amount of the exchanged momentum and the size of the bar. In comparison, our models have a halo-to-disk mass ratio of unity within the central 10 kpc, $Q \sim 1.5$, and a halo core of ~ 2 kpc. They lie between the models MH1, M γ 3–4, and MQ4 of Athanassoula.

Interestingly, Valenzuela & Klypin (2003) argued that it is the mass and force resolutions of numerical models that dictate the efficiency of the J transfer to the halo and hence play an important role in the evolution of the bar. Insufficient resolution results in an excessive growth of numerical bars and an excessive decrease in the bar pattern speed. However, it seems rather that the physical conditions mentioned above take priority: their high-resolution models have extended and hot halos. High dispersion velocities in the halo affect the particle trapping by the resonances and so are expected to lower substantially the J transfer.

6.2. Bar Size Evolution

If one neglects the dissipative component in a modeled galaxy, the bar pattern speed decreases secularly, except during brief time intervals of internal instabilities in the bar itself. At least in numerical models of stellar bars, their overall slowdown is accompanied by an increase in the bar length, so the bar roughly extends to its corotation radius, which increases with time (Athanassoula 1992). Alternative theoretical models of bars terminating at the ILR exist (Lynden-Bell 1979), but difficulties remain in actually reproducing them in numerical simulations (but see, e.g., Polyachenko & Polyachenko 1994 for the so-called slow bars). A sole exception consists of a system of nested bars, where the inner (nuclear) bars form within the ILR (e.g., Shlosman et al. 1989; Friedli & Martinet 1993; Englmaier & Shlosman 2004). However, a dissipative component is required to be present for self-consistency in this case. How exactly the bar traps the disk orbits in order to increase its length is not known at present.

We have demonstrated that while the size of the bar, r_{bar} , drops during the first buckling (Fig. 2a), the ratio of $r_{\text{CR}}/r_{\text{bar}}$ (Fig. 4b) does *not* increase dramatically above the range of 1.2 ± 0.2 determined by Athanassoula (1992) to fit the observed shapes of the offset dust lanes in barred galaxies. Amazingly, the reason for this is that r_{CR} drops as well due to the sudden increase in the bar pattern speed in the same time interval. This is not unexpected due to a clear trend between Ω_b and A_2 (Athanassoula 2003) and a long-known fact that the bar lengthens with its slowdown in numerical simulations. What is new here is that we find that Ω_b increases sharply during the first buckling, and this increase apparently correlates with the sudden decrease in the bar length. Hence, the relation between Ω_b , A_2 , and r_{bar} appears to be more fundamental than anticipated and holds for both decelerating and accelerating bars.

We associate the continued bar growth with the ability of the halo to absorb angular momentum from the disk region lying within the corotation radius, a region that itself expands with time. This process leads to a stronger bar and can be seen as a counterbalance to a number of other processes that have been discussed in the literature within the framework of bar dissolution. Recent observational results from the Galaxy Evolution from Morphology and SEDs (GEMS; Rix et al. 2004) survey have shown that the bar size and axial ratio distributions at intermediate redshifts of $z \sim 0.2$ –1 are compatible with those in the local universe (Jogee et al. 2004; Elmegreen et al. 2004;

Sheth et al. 2003). Our models, which exhibit a slow secular bar growth over the Hubble time, are in general agreement with these results. They do lack the gaseous component that can dramatically shorten the bar life cycle (e.g., Bournaud & Combes 2002). However, it is difficult to understand how the short-lived (e.g., less than 1.5 Gyr; Bournaud & Combes 2002; Combes 2004) bars can form at the same rate and with the same sizes and axial ratios at which they are being destroyed by the gas. Yet, taken at face value, in order to agree with the GEMS results, the bars must preserve their size and strength distributions nearly unchanged over the last 8 Gyr. Clearly, the precise role of gas in the bar evolution must still be determined.

6.3. Bar Shape Evolution

Next we focus on some aspects of the bar's three-dimensional shape evolution in the model, specifically on the axial ratio in the bar midplane and on the bar symmetry in the vertical plane. The midplane axial ratio, or alternatively the bar ellipticity ϵ , is measured from the ellipse fitting (Fig. 4c) and agrees with the evolution of the bar amplitude A_2 . At the same time, the variations in ϵ appear less dramatic than those in A_2 , although recurrent bucklings are clearly visible as a decrease or saturation.

The ratio $r_{\text{CR}}/r_{\text{bar}}$ has also dynamical implications for the bar. For example, it determines the shape of the offset dust lanes in barred galaxies, which in turn delineate shocks in the gas flow (Athanassoula 1992). The strength of the underlying shocks determines the gas inflow toward the central kiloparsec. A bar that is considerably weakened will slow down the radial gas inflow. The observed shapes constrain the ratio $r_{\text{CR}}/r_{\text{bar}}$ to 1.2 ± 0.2 . The modeled ratio (Fig. 4b) typically falls within the required limits, except during the first buckling when it is higher, ~ 1.5 .

After the second buckling, the bar shows the so-called ansae (handles) on both its ends. We observe them as characteristic density enhancements in the face-on (Fig. 11) or edge-on (Fig. 3) disks. Athanassoula (2001) related the appearance of ansae to initial conditions in the models (e.g., the halo-to-disk mass ratio). It remains unknown why they appear at this particular evolutionary stage, $\tau > 8$ Gyr. The ansae can be seen in some early-type barred galaxies, e.g., NGC 4262, NGC 2859, and NGC 2950 (Sandage 1961), NGC 4151 (Mundell & Shone 1999), and ESO 509-98 (Buta et al. 1998).

Vertically, the bar evolves from a geometrically thin configuration, similar to the disk hosting it. The vertical bar buckling, when viewed along the bar's minor axis, shows a rapidly evolving bending that relaxes to a boxy bulge, i.e., a bulge with flat or mildly convex isodensities (Fig. 8a). Subsequently, the bulge acquires a peanut shape. During the second buckling, for ~ 3 Gyr, the vertical asymmetry persists with one-sided boxy and peanut symmetries that derive from the asymmetric BAN family (Fig. 8b). With the asymmetry washed out, the boxy/peanut-shaped bulge/bar has a pronounced X-shape: two pointed spikes with a large concave region in between (Fig. 8c). The X-shapes have been seen before in numerical simulations (e.g., Athanassoula & Misiriotis 2002) and have been observed as well: e.g., NGC 4845, NGC 1381, and IC 4767 (Whitmore & Bell 1988), IC 3370 (Jarvis 1986), and AM 1025–401 (Arp & Madore 1987; Bureau & Freeman 1999; Patsis et al. 2002b). Mihos et al. (1995) proposed a merging scenario for the formation of the X-shaped bulges, e.g., a minor merger for the Hickson 87a galaxy. This merger triggers the bar instability in the disk, followed by the buckling and the X-shaped bulge, when viewed from a specific aspect angle. In principle, there may be more than one cause for the bar buckling and for the formation of peanut/boxy/X-shaped bulges. But when taken together, a high




FIG. 11.—Ansae at time $\tau = 9.9$ Gyr. A face-on view of the modeled barred disk. The gray-scale contours represent the surface density and are spaced logarithmically. The box size is $18 \text{ kpc} \times 18 \text{ kpc}$. [See the electronic edition of the *Journal for a color version of this figure.*]

observed frequency of these bulges (Lüticke et al. 2000) and the need for only mild asymmetry in the disk for their appearance (Patsis et al. 2002a) may hint at their intrinsic origin.

6.4. Vertical Asymmetry of the Bar

The vertical asymmetry of the modeled bar has been detected during its recurrent bucklings for about 1 and ~ 3 Gyr, respectively (Figs. 3 and 5). Its characteristic shape, in principle, is detectable by observations of edge-on galaxies (e.g., Fig. 3 and accompanying animation), especially during the second buckling due to its prolonged period. Observationally, the bucklings differ when viewed edge-on along the bar's minor axis. The main difference is the location of the maximal asymmetry: it is close to the rotation axis during the first buckling and around the middle region of the bar during the second buckling.

Lüticke et al. (2000) present first statistics of edge-on galaxies with the boxy/peanut-shaped bulges and show some isophote fits (their Fig. 2). They distinguish between peanut-shaped and boxy bulges. A total of 27% of the sample of 734 galaxies with boxy/peanut-shaped bulges have bulges that are either close to boxy or, due to low resolution, could not be distinguished between boxy and peanut-shaped. Some of these bulges show a mild vertical asymmetry similar to that found during the second buckling; e.g., NGC 4289 is a good example of a possible observed secondary buckling. Of course, to corroborate the numerical simulations one needs a large, statistically significant sample with high resolution. The main difficulty lies in the unambiguous determination of the inclination angle of a galactic disk: it must be edge-on within $\pm 5^\circ$ – 7° .

From a theoretical point of view, it is unclear how widespread is the recurrent buckling in stellar bars: how sensitive it is to disk and especially halo parameters, such as mass distribution and dispersion velocities. While the condition for a second buckling seem to be directly related to the ability of the bar to strengthen

after the initial weakening, this process can depend on a number of additional parameters. For example, Athanassoula (2003) has found that the angular momentum transfer between the disk and the halo depends on the mass distribution in the halo and weakens substantially for hotter halos. Berentzen et al. (2006) have detected a recurrent buckling in their LS2 model with a 2 kpc flat core (live) halo with a logarithmic potential. At the time of the second buckling, the bar in this model appears somewhat stronger than in the first buckling. Moreover, the secondary buckling can be apparently seen in Figure 4 of O'Neill & Dubinski (2003) and also in Figure 11 of Valenzuela & Klypin (2003), the latter based on the evolution of the bar pattern speed, strength, and the angular momentum rate change. Secondary buckling is also present in some models of L. Athanassoula (2005, private communication). In all cases they went unnoticed. Finally, we comment on the addition of the dissipative component to the stellar disk (Berentzen et al. 1998). The effect of the clumpy gas component is to weaken the buckling instability, but without quantifying the degree of clumpiness within the central kiloparsec, it is not clear whether the isothermal equation of state used has led in fact to overdamping of this instability.

7. CONCLUSIONS

To summarize this work, we have studied the long-term stellar bar evolution in a high-resolution self-consistent model of a disk and a responsive halo. We find that a developing bar goes through the vertical buckling instability, which weakens it and dissolves its outer half. Subsequently the bar experiences a renewed growth that leads to a recurrent buckling. This evolution is driven by the resonant interaction between the barred disk and the surrounding halo; we quantify this effect by means of the spectral analysis of individual orbits in the disk and the halo and show that the halo particles are trapped by numerous lower resonances with the bar and that this trapping is robust. During these periods of recurrent instability, and especially during the slower second buckling, the bar remains vertically asymmetric for a prolonged ~ 3 Gyr time interval, which in principle can be detected observationally. However, two issues can potentially complicate this detection. First, it is not clear how widespread are the conditions favorable for the recurrent bar growth, although we have detected it in a number of models with different initial conditions. Second, while it was shown that a clumpy gaseous component with an isothermal equation of state in the disk will weaken this instability, the effect of a realistic ISM was never estimated. A statistically significant sample of (nearly) edge-on galaxies is required to test the prediction of a prolonged vertical asymmetry.

We also find that the secular bar growth and the triggered buckling instabilities lead to pronounced changes in the bulge shape: it grows both radially and vertically, acquiring a peanut-shaped, a boxy, and finally the X-shaped appearance. While the bar size approximately follows the 4:1 (ultraharmonic) resonance in the disk, the boxy/peanut-shaped bulge size appears to be guided by the vertical ILR. Concurrently, the bar is going through a structural evolution: new families of three-dimensional periodic orbits appear (or become more pronounced) after the bucklings.

Finally, we find that the bar strength correlates with its pattern speed in both strengthening and weakening bars. While it was known already that a stellar bar becomes stronger as it slows down, we detect the reverse trend as well: bars that weaken during the buckling speed up their tumbling. Moreover, the bar size appears to be sensitive to these changes: bars slowing down become longer, while bars speeding up shorten.

We gratefully acknowledge insightful discussions with Lia Athanassoula, Ingo Berentzen, Amr El-Zant, Johan Knapen, and Alar Toomre. This work has been partially supported by grants from NASA LTSA5-13063, NASA ATP NAG 5-10823, *HSTAR*-

09546 and 10284 (I. S.), and by NSF grant AST-0206251 (C. H. and I. S.). I. M. acknowledges support from PPARC. Simulations and orbital analysis have been performed on a dedicated Linux cluster, and we thank Brian Doyle for technical support.

REFERENCES

- Arp, H. C., & Madore, B. F. 1987, A Catalogue of Southern Peculiar Galaxies and Associations (Cambridge: Cambridge Univ. Press)
- Athanassoula, E. 1984, Phys. Rep., 114, 321
- . 1992, MNRAS, 259, 345
- . 2001, in New Horizons of Computational Science, ed. T. Ebisuzaki & J. Makino (Dordrecht: Kluwer), 69
- . 2002a, ApJ, 569, L83
- . 2002b, in ASP Conf. Ser. 275, Disks of Galaxies: Kinematics, Dynamics, and Interactions, ed. E. Athanassoula, A. Bosma, & R. Mujica (San Francisco: ASP), 141
- . 2003, MNRAS, 341, 1179
- Athanassoula, E., & Misiriotis, A. 2002, MNRAS, 330, 35
- Barnes, J., & Hut, P. 1986, Nature, 324, 446
- Berentzen, I., Heller, C. H., Shlosman, I., & Fricke, K. 1998, MNRAS, 300, 49
- Berentzen, I., Shlosman, I., & Jogee, S. 2006, ApJ, in press (astro-ph/0501352)
- Binney, J., & Spergel, D. 1982, ApJ, 252, 308
- Binney, J., & Tremaine, S. 1987, Galactic Dynamics (Princeton: Princeton Univ. Press)
- Bournaud, F., & Combes, F. 2002, A&A, 392, 83
- Bureau, M., & Freeman, K. C. 1999, AJ, 118, 126
- Buta, R., Alpert, A. J., Cobb, M. L., Crocker, D. A., & Purcell, G. B. 1998, AJ, 116, 1142
- Combes, F. 2004, in IAU Symp. 222, The Interplay Among Black Holes, Stars and ISM in Galactic Nuclei, ed. T. Storchi-Bergmann, L. C. Ho, & H. R. Schmitt (Cambridge: Cambridge Univ. Press), 383
- Combes, F., Debbasch, F., Friedli, D., & Pfenniger, D. 1990, A&A, 233, 82
- Combes, F., & Sanders, R. H. 1981, A&A, 96, 164
- Contopoulos, G., & Papayannopoulos, T. 1980, A&A, 92, 33
- Debattista, V. P., & Sellwood, J. A. 1998, ApJ, 493, L5
- . 2000, ApJ, 543, 704
- Dehnen, W. 2002, J. Comp. Phys., 179, 27
- Elmegreen, B. G., Elmegreen, D. M., & Hirst, A. C. 2004, ApJ, 612, 191
- El-Zant, A., & Shlosman, I. 2002, ApJ, 577, 626
- Englmaier, P., & Shlosman, I. 2004, ApJ, 617, L115
- Fall, S. M., & Efstatiou, G. 1980, MNRAS, 193, 189
- Friedli, D., & Martinet, L. 1993, A&A, 277, 27
- Friedli, D., & Pfenniger, D. 1990, in ESO/CTIO Workshop on Bulges of Galaxies, ed. B. Jarvis & D. M. Terndrup (Garching: ESO), 265
- Heller, C. H. 1995, ApJ, 455, 252
- Heller, C. H., & Shlosman, I. 1994, ApJ, 424, 84
- . 1996, ApJ, 471, 143
- Hernquist, L., & Weinberg, M. D. 1992, ApJ, 400, 80
- Hunt, L. K., & Malkan, M. A. 2004, ApJ, 616, 707
- Jarvis, B. J. 1986, AJ, 91, 65
- Jogee, S., et al. 2004, ApJ, 615, L105
- Knapen, J. H., Shlosman, I., & Peletier, R. F. 2000, ApJ, 529, 93
- Laine, S., Shlosman, I., Knapen, J. H., & Peletier, R. F. 2002, ApJ, 567, 97
- Lüticke, R., Dettmar, R.-J., & Pohlen, M. 2000, A&AS, 145, 405
- Lynden-Bell, D. 1979, MNRAS, 187, 101
- Martinez-Valpuesta, I., & Shlosman, I. 2004, ApJ, 613, L29
- Merrifield, M. R., & Kuijken, K. 1999, A&A, 345, L47
- Merritt, D., & Sellwood, J. A. 1994, ApJ, 425, 551
- Mihos, J. C., Walker, I. R., Hernquist, L., Mendes de Oliveira, C., & Bolte, M. 1995, ApJ, 447, L87
- Mundell, C. G., & Shone, D. L. 1999, MNRAS, 304, 475
- O'Neill, J. K., & Dubinski, J. 2003, MNRAS, 346, 251
- Ostriker, J. P., & Peebles, P. J. E. 1973, ApJ, 186, 467
- Patsis, P. A., Athanassoula, E., Grosbøl, P., & Skokos, Ch. 2002a, MNRAS, 335, 1049
- Patsis, P. A., Skokos, Ch., & Athanassoula, E. 2002b, MNRAS, 337, 578
- Pfenniger, D. 1984, A&A, 134, 373
- Pfenniger, D., & Friedli, D. 1991, A&A, 252, 75
- Polyachenko, V. L., & Polyachenko, E. V. 1994, Astron. Lett., 20, 416
- Raha, N., Sellwood, J. A., James, R. A., & Kahn, F. D. 1991, Nature, 352, 411
- Rix, H.-W., et al. 2004, ApJS, 152, 163
- Sandage, A. 1961, The Hubble Atlas of Galaxies (Washington, DC: Carnegie Inst.)
- Sellwood, J. A. 1980, A&A, 89, 296
- . 1996, in ASP Conf. Ser. 91, Barred Galaxies, ed. R. Buta, D. A. Crocker, & B. G. Elmegreen (San Francisco: ASP), 259
- Sellwood, J. A., & Wilkinson, A. 1993, Rep. Prog. Phys., 56, 173
- Shaw, M. A. 1987, MNRAS, 229, 691
- Shen, J., & Sellwood, J. A. 2004, ApJ, 604, 614
- Sheth, K., Regan, M. W., Scoville, N. Z., & Strubbe, L. E. 2003, ApJ, 592, L13
- Shlosman, I., Frank, J., & Begelman, M. C. 1989, Nature, 338, 45
- Skokos, Ch., Patsis, P. A., & Athanassoula, E. 2002a, MNRAS, 333, 847
- . 2002b, MNRAS, 333, 861
- Sotnikova, N. Y., & Rodionov, S. A. 2005, Astron. Lett., 31, 15
- Toomre, A. 1966, in Notes on the 1966 Summer Study Program in Geophysical Fluid Dynamics at the Woods Hole Oceanographic Institution (Woods Hole: Woods Hole Oceanographic Inst.), 111
- Valenzuela, O., & Klypin, A. 2003, MNRAS, 345, 406
- Weinberg, M. D. 1985, MNRAS, 213, 451
- Weinberg, M. D., & Katz, N. 2002, ApJ, 580, 627
- Whitmore, B. C., & Bell, M. 1988, ApJ, 324, 741