

EFFECT OF THE HERBICIDE PARAQUAT GRAMOXONE (1,1 DIMETIL 4,4 BIPIRIDILUM DICHLORIDE ON THE EMBRYONIC DEVELOPMENT OF RATS

GONÇALVES, N.M.F.M.; MACHADO, C.¹; NAGANO, C.E.¹; GIROLDO, F.²;
STROPARO, G.²; TAFAREL, J.R.²; DITZEL, L.F.²; FUJITA, J.I.²; SCHULZ, R.T.²;
CHAMPION, L.²; SILVA, M.X.²; CARARO, M.²; NIED, R.².; BACILA, M.⁴

¹Departamento de Medicina, PUCPR;

²Graduandos do Curso de Farmácia Bioquímica;

³Graduando do Curso de medicina, PUCPR;

⁴Pontifícia Universidade Católica do Paraná.

The main aim of the present research was to study the possible teratogenic effects caused by the herbicide Paraquat-Gramoxone (1,1'dimetil-4,4'-bypiridilium dicloride) in nestlings female Wistar rats (*Rattus norvegicus* var. *albinus*), during the period of 7° to 9° days of embryonic development. Macroscopic analysis of the nestling showed lesions mainly in the areas of nose, eyelids and shoulders. Histological alterations were found also in the liver and kidney preparations.

Key Words: herbicide, teratogenic effects, Paraquat, wistar rats.