

ARTIGO DE REVISÃO

REVISÃO/REVIEW

IMUNOSENESCÊNCIA E SEU IMPACTO NA PRÁTICA MÉDICA. UM NOVO OLHAR PARA UM VELHO PROBLEMA.

IMMUNOSENESCENCE AND ITS IMPACT ON MEDICAL PRACTICE. A NEW LOOK AT AN OLD PROBLEM.

DOI: 10.5380/rmu.v1i4.40693

Pedro G. Lorencetti^{1,2}, Kornelis S.M. van der Geest¹, Qi Wang¹, Elisabeth Brouwer¹, Annemieke M.H. Boots¹

RESUMO

Introdução: O envelhecimento é um processo complexo que influencia todos os sistemas do corpo humano e seu impacto no sistema imune é chamado Imunosenescência. Essa condição é resultado de várias modulações imunológicas causadas por interações entre fatores genéticos e ambientais e é responsável por importantes condições clínicas em indivíduos idosos como alta prevalência de doenças infecciosas e autoimunes, neoplasias e menor eficácia de vacinas. **Objetivo:** Nesta revisão serão discutidos os problemas clínicos mais comuns na população idosa relacionados a Imunosenescência, e os novos achados da ciência básica relevantes a este tópico. **Conclusão:** A melhor compreensão da Imunosenescência é importante para prevenção de doenças comuns relacionadas à idade e para a promoção de um envelhecimento saudável.

Palavras chaves: Imunologia, Envelhecimento, Envelhecimento Celular.

ABSTRACT

Background: Ageing is a very complex process that modulates all the organ systems of the human body, and its impact on the immune system is called Immunosenescence. This condition is the result of several immune modulations due to genetic and environmental interactions and is responsible for important clinical conditions in elderly subjects, such as a higher incidence of infectious and autoimmune diseases, neoplasias and decreased vaccine efficacy. **Objective:** In this current review we will discuss the most common clinical problems in the elderly population related to Immunosenescence and new findings in basic science relevant to this topic. **Conclusion:** A better understanding of Immunosenescence is important to prevention of common age related diseases and for the promotion of healthy ageing.

Key words: Immunology, Aging, Cell Aging.

1 – Department of Rheumatology and Clinical Immunology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

2- Universidade Federal do Paraná – UFPR, Curitiba, Brasil

Contato do Autor / Mail to:

Pedro Gabriel Lorencetti – plorencetti@gmail.com

General Carneiro, 112, Apartamento 201, Alto da Glória, Curitiba, Paraná, CEP: 80060-150.

INTRODUCTION

One way to measure the quality and success of a society is the life expectancy or in other words, life and health span. The recent social, economic and technological development is causing a fast ageing of the world population, especially in developed countries. Brazil being a developing country, is facing a rapid increase of its elderly population. Currently, 11% of the Brazilian population is composed of subjects older than 60¹. This demographical transition implies a great change in social structures, such as the health care system. The elderly population demands a different and specialized care with focus on treatment of chronic and degenerative diseases, prevention of disability, frailty and infectious diseases.

From a biological point of view, ageing is a very complex process that modulates all the organ systems of the human body, and its impact on the immune system is called Immunosenescence. Franceschi et.al² proposed a simple equation to better understand the interaction between ageing and the immune system : *Stress resistance = Ability to survive*. Adopting an evolutionary point of view, the capacity to deal with environmental and physiological challenges prolongs survivability and as a result expose organisms once more to new stressors, promoting an endless selection process. In other words, the evolution of the immune system resulted in lifespan extension and ageing in its own way modulates the defense mechanisms based on the new environment and physiologic demands. The human innate and adaptive immune responses thus pass through several changes dictated by different challenges during ageing.

The immune system ageing is characterized by a decrease of hematopoietic stem cells and thymus involution, leading to a decline in the replenishment of naïve T and B cells. Interaction with environmental triggers leads to a shift from naïve to memory cells, an expansion of oligoclonal memory cells related to cytomegalovirus (CMV), and a low grade and persistent inflammatory state²⁻⁵. The ageing phenotype is a result of the sum of all these immune modulations and has been correlated with several clinical conditions such as increased incidence of inflammatory, neurodegenerative and infectious diseases, frailty, decreased vaccine efficacy and occurrence of some age-related neoplasms⁶⁻¹¹.

In this current review we will discuss the most common clinical problems in the elderly population related to Immunosenescence and new findings in basic science relevant to this topic.

TÍTULO??

Is there a maker of immune ageing? - The immune risk profile.

Humans have always pursued the secret of longevity. Despite intense investigation, only recently some studies identified certain immune characteristics able to predict mortality in elderly populations. The sum of these immune factors is called "Immune Risk Profile" (IRP), and were identified for the first time in the large *Swedish OCTO/NONA*^{12,13} cohort study. It was observed that elderly subjects with a CD4/CD8 ratio < 1, CMV IgG serum positivity, and expansion of CD8⁺ CD28⁻ cell population exhibited higher mortality irrespective of health status. An extension of this study has found that CD4/CD8 ratio <1 was in fact the best independent predictor of mortality and ratios >1 were correlated with successful ageing^{14,15}. Despite the robust evidence, further studies did not confirm the same IRP variables in different elderly populations around the world, suggesting that immune ageing can be a heterogeneous process.

An interesting note covered in the original IRP was the identification of a viral infection playing a role in the ageing process. Further studies documented a close relationship between CMV chronic infection and modulations of T cell subsets. The CMV IgG serum positivity has been correlated with accumulation of highly differentiated effector T cells¹⁶. This cell subset is characterized by an impaired capacity to migrate to lymph nodes, decreased ability to being stimulated by APC and higher production of pro inflammatory cytokines.

In summary, there is not a robust marker of immune ageing. The decline of CD8 naïve T cells and the occurrence of CD28⁻ T cells are the most common and reproducible findings in different elderly populations, but its impact in healthy ageing and disease development is not clear yet.

Sometimes more is less. - Inflammaging a silent killer.

The inflammatory response is the crucial physiological response to the constant immunological challenges that we are exposed to in daily life. Several studies have observed a progressive low-grade inflammation that follows the ageing process termed Inflammaging. Inflammaging has been characterized as an unbalance between pro and anti-inflammatory factors resulting in increased serum levels of IL-6, TNF- α and acute phase proteins such as CRP². This condition has been correlated with the development of important clinical conditions such as frailty, cancer and cardiovascular diseases¹⁷⁻¹⁹. Although several transversal studies correlated CMV infection with Inflammaging, a recent longitudinal study did not find any association between this viral infection and an increase of inflammatory cytokines²⁰.

In healthy centenarians it has been observed that higher production of anti-inflammatory factors is protective and contributes to longer survival. Lio et.al²¹.found that healthy centenarians have higher serum levels of TGF- β and carry a higher frequency of an IL-10 gene polymorphism that promotes higher IL-10 production when compared to controls. The Leiden 85+ cohort study²² also revealed that individuals with higher IL-10 serum levels presented lower cardiovascular mortality when compared with subjects with lower levels of IL-10.

The Groningen Research Initiative on Healthy Ageing and Immune Longevity (GRAIL) group is conducting several research projects related to Inflammaging. Based on the evidence that B cells can present pro and anti-inflammatory roles through their cytokine production profile, our group explored the impact of ageing on cytokine producing B cells. The population of effector B cells is characterized by production of TNF- α and IL-6, promoting inflammation, and disturbances in this subset have been reported in diseases such as rheumatoid arthritis, giant cell arteritis and polymyalgia rheumatica^{23,24}. On the other hand, B cells can block the inflammatory response inhibiting TNF- α and IFN- γ production by T cells in an IL-10 dependent way. IL-10 producing B cells are termed B regulatory cells (Breg). Decreased numbers of Breg have been correlated with several inflammatory diseases such as systemic lupus erythematosus, multiple sclerosis and rheumatoid arthritis²⁵⁻²⁷. In a

cross sectional study we have found that regulatory B cells numbers tends to decrease with ageing whereas we did not find any modulation of the effector B cell subset. Thus, ratio's of Breg/Breg increase with ageing thereby favoring pro-inflammation. Our findings suggest a new players in Inflammaging: the B effectors and the Bregs (unpublished data).

Preventing severe complications of common diseases -Vaccination response in elderly.

Vaccination should be regarded as a milestone in the quest for extension of life expectancy for mankind, making prevention of several life threatening infectious diseases possible. A particular health issue is the vaccination of the elderly population since infectious diseases incidence increases with age⁶. Several clinical and epidemiological studies have shown a decreased vaccine response in older subjects. Both the innate and the adaptive immune system are involved in this process. The aged innate response displays diminished migration and phagocytic activity of neutrophils and macrophages summed to decreased numbers of APC cells and reduced expression of MHC I and II as well^{28,29}. On the other hand, the aged adaptive response is impaired due to decreased numbers of naïve T and B cells, accumulation of highly differentiated effector T cells, decreased expression of CD40-ligand by CD4⁺ cells, defects in isotype switching and somatic mutation in B cells³⁰⁻³⁵. This highly disturbed immune response contributes to increased susceptibility to infectious diseases and is a challenge to be overtaken by the development of new vaccines.

Changing old for new – The challenge of transplantation medicine in elderly subjects.

The improvement of chronic diseases care is leading to an increase in the prevalence of end stage organ failures. Since regenerative medicine and stem cells therapy are not feasible treatments yet, organ transplantation is the best alternative for these conditions. Age is one of the most important factors in transplant outcome and Immunosenescence plays an important role in this clinical scenario. Several epidemiological studies have reported lower incidences of acute graft rejection in elderly recipients³⁶⁻³⁸, however the mortality and chronic rejection are still higher in this population^{39,40}. Most of the authors attributes

this different clinical outcome to the same immune changes that also determine/impact the impaired vaccine response.

Autoimmune diseases: Age marked pathologies.

The correlation between immunosenescence and autoimmune diseases is based on several epidemiological and laboratory data. It is important to note that autoimmune disorders are more prevalent in older subjects particularly with rheumatoid arthritis, giant cell arteritis and polymyalgia rheumatica⁶. This fact can be explained by a breach of tolerance that follows ageing. The low naïve T cell output due to thymus involution may induce an intense proliferation of peripheral naïve T cells, resulting in impaired selection and replication of naïve and memory self-reactive T cells. Insufficiency of telomerase activity, deficiency of DNA repair mechanisms and accumulation of CD28⁻ cells may also contribute to the pathogenesis, promoting inflammation and impaired regulatory capacity⁴¹⁻⁴³. Noteworthy, several of these characteristics can be observed in RA patients, despite disease duration and treatment, suggesting that these subjects present with a premature ageing of the immune system.

CONCLUSIONS

Immunosenescence is a complex phenomenon that affects all the components of the immune system and depends on genetic and environmental factors. Accumulation of oligoclonal highly differentiated T cells and increasing serum levels of pro inflammatory cytokines might be main events responsible for the immune impairment in elderly subjects. Healthy centenarians provide living proof that effective immune responses can be maintained for decades, even after thymus involution. More immunogerontology studies are required to understand immune competence and immune dysfunction with ageing. A better understanding may hold promise for future clinical studies for the prevention of common age related diseases and promotion of healthy ageing.

ACKNOWLEDGMENTS

The main author received a scholarship from Science Without Borders program to one year exchange in Netherlands.

I would like to acknowledge Kornelis S.M. van der Geest, Qi Wang, Elisabeth Brouwer, Annemieke M.H. Boots for all the support, the friendship and the great opportunity to join the GRAIL group and work in several research projects related to Immunosenescence.

REFERÊNCIAS BIBLIOGRÁFICAS

- 1-United Nations, Department of Economic and Social Affairs, Population Division (2013). World Population Ageing 2013. ST/ESA/SER.A/348.
- 2-Franceschi C, Bonafe M, Valensin S, Olivieri F, De Luca M, Ottaviani E, et.al. Inflamm-aging. An evolutionary perspective on immunosenescence. 2000 Ann. N. Y. Acad. Sci. 908, 244–254.
- 3- Boehm T, Swann JB. Thymus involution and regeneration: two sides of the same coin? Nat Rev Immunol. 2013 Nov;13(11):831-8
- 4- Wang C, Liu Y, Xu LT, Jackson KJ, Roskin KM, Pham TD, et.al. Effects of aging, cytomegalovirus infection, and EBV infection on human B cell repertoires. J Immunol. 2014 Jan 15;192(2):603-11.
- 5-Khan N, Shariff N, Cobbold M, Bruton R, Ainsworth JA, Sinclair AJ, Cytomegalovirus seropositivity drives the CD8 T cell repertoire toward greater clonality in healthy elderly individuals. J Immunol. 2002 Aug 15;169(4):1984-92.
- 6- Yoshikawa TT. Epidemiology and unique aspects of aging and infectious diseases. Clin Infect Dis. 2000 Jun;30(6):931-3.
- 7-Crowson CS, Matteson EL, Myasoedova E, Michet CJ, Ernste FC, Warrington KJ, et.al. The lifetime risk of adult onset rheumatoid arthritis and other inflammatory autoimmune rheumatic diseases. Arthritis Rheum. 2011 Mar;63(3):633-9

- 8-Grubeck-Loebenstien B, Wick G. The aging of the immune system. *Adv Immunol* 2002; 80:243-84.
- 9-DePinho RA: The age of cancer. *Nature* 2000, 408(6809):248-254.
- 10- Baylis D, Bartlett DB, Syddall HE, Ntani G, Gale CR, Cooper C, et. al. Immune-endocrine biomarkers as predictors of frailty and mortality: a ten year longitudinal study in community dwelling older people. *Age (Dordr)*. 2013 Jun;35(3):963-71. -8.
- 11-Bruunsgaard H, Andersen-Ranberg K, Jeune B, Pedersen AN, Skinhøj P, Pedersen BK. A high plasma concentration of TNF-alpha is associated with dementia in centenarians. *J Gerontol A Biol Sci Med Sci*. 1999 Jul;54(7):M357-64.
- 12-Ferguson FG, Wikby A, Maxson P, Olsson J, Johansson B. Immune parameters in a longitudinal study of a very old population of Swedish people: a comparison between survivors and nonsurvivors *J Gerontol A Biol Sci Med Sci*. 1995 Nov;50(6):B378-82.
- 13- Wikby, A.,Johansson, B.,Ferguson, F. The OCTO and NONA immune longitudinal studies: a review of 11 years studies of Swedish very old humans.2003. *Adv. Cell Aging Gerontol*. 13, 1-16.
- 14- Wikby A, Maxson P, Olsson J, Johansson B, Ferguson FG. Changes in CD8 and CD4 lymphocyte subsets, T cell proliferation responses and non-survival in the very old: the Swedish longitudinal OCTO-immune study *Mech Ageing Dev*. 1998 May 15;102(2 3):187-98.
- 15-Strindhall J, Nilsson BO, Löfgren S, Erneurudh J,Pawelec G, Johansson B, et.al. No Immune Risk Profile among individuals who reach 100 years of age: findings from the Swedish NONAimmune longitudinal study. *Exp Gerontol*. 2007 Aug;42(8):753-61.
- 16- Wikby A, Johansson B, Olsson J, Löfgren S, Nilsson BO, Ferguson F. Expansions of peripheral blood CD8 T- lymphocyte subpopulations and an association with cytomegalovirus seropositivity in the elderly: the Swedish NONA immune study. *Exp Gerontol*. 2002 Jan-Mar;37(2-3):445-53.
- 17- Baylis D, Bartlett DB, Syddall HE, Ntani G, Gale CR, Cooper C, et.al. Immune-endocrine biomarkers as predictors of frailty and mortality: a 10-year longitudinal study in community-dwelling older people. *Age (Dordr)*. 2013 Jun;35(3):963-71.
- 18- Olivieri F, Rippo MR, Monsurrò V, Salvioli S, Capri M, Procopio AD, et.al. MicroRNAs linking inflamm-aging, cellular senescence and cancer. *Ageing Res Rev*. 2013 Sep;12(4):1056-68.
- 19- Libby P, Okamoto Y, Rocha VZ, Folco E. Inflammation in atherosclerosis: transition from theory to practice. *Circ J*. 2010 Feb;74(2):213-20.
- 20- Bartlett DB, Firth CM, Phillips AC, Moss P, Baylis D, Syddall H, et.al. The age-related increase in low grade systemic inflammation (Inflammaging) is not driven by cytomegalovirus infection. *Aging Cell*. 2012 Oct;11(5):912-5.
- 21- Lio D, Scola L, Crivello A, Colonna-Romano G, Candore G, Bonafè M. et.al. Gender-specific association between -1082 IL 10 promoter polymorphism and longevity. *Genes Immun*. 2002 Feb;3(1):30-3.
- 22- van Exel E, Gussekloo J, de Craen AJ, Bootsma-van der Wiel A, Frölich M, Westendorp RG. Inflammation and stroke: the Leiden 85-Plus Study. *Stroke*. 2002 Apr;33(4):1135-8.
- 23- Frasca D, Diaz A, Romero M, Landin AM, Blomberg BB.High TNF- α levels in resting B cells negatively correlate with their response. *Exp Gerontol*. 2014 Jun;54:116-22.
- 24- van der Geest KS, Abdulahad WH, Chalan P, Rutgers A, Horst G, Huitema MG, et.al. Disturbed B cell homeostasis in newly diagnosed giant cell arteritis and polymyalgia rheumatica. *Arthritis Rheumatol*. 2014 Jul;66(7):1927-38.

- 25- Blair PA, Norena LY, Flores-Borja F, Rawlings DJ, Isenberg DA, Ehrenstein MR, et al. CD19⁺CD24^{hi}CD38^{hi} B cells exhibit regulatory capacity in healthy individuals but are functionally impaired in systemic lupus erythematosus patients. *Immunity* 2010;32:129–40.
- 26-Duddy M, Niino M, Adatia F, Hebert S, Freedman M, Atkins H. Distinct effector cytokine profiles of memory and naïve human B cell subsets and implication in multiplesclerosis. *J Immunol.* 2007 May 15;178(10):6092-9.
- 27- Daien CI, Gailhac S, Mura T, Audo R, Combe B, Hahne M, Morel J. Regulatory B10 cells are decreased in patients with rheumatoid arthritis and are inversely correlated with diseaseactivity. *Arthritis Rheumatol.* 2014 Aug;66(8):2037-46.
- 28- Gomez CR, Boehmer ED, Kovacs EJ. The aging innate immune system. *Curr Opin Immunol* 2005; 17:457–62.
- 29- Grewe M. Chronological ageing and photoageing of dendritic cells. *Clin Exp Dermatol* 2001; 26:608–12.
- 30-Fagnoni FF, Vescovini R, Passeri G, et al. Shortage of circulating naïve CD8(+) T cells provides new insights on immunodeficiency in aging. *Blood* 2000; 95:2860–8.
31. Lazuardi L, Jenewein B, Wolf AM, Pfister G, Tzankov A, Grubeck-Loebenstein B. Age-related loss of naive T cells and dysregulation of T-cell/B-cell interactions in human lymph nodes. *Immunology* 2005; 114:37–43.
- 32- Allman D, Miller JP. B cell development and receptor diversity during aging. *Curr Opin Immunol* 2005; 17:463–7.
- 33- Appay V, Dunbar PR, Callan M, Klenerman P, Gillespie GM, Papagno L, et al. Memory CD8⁺ T cells vary in differentiation phenotype in different persistent virus infections. *Nat Med* 2002; 8:379–85
- 34- Haynes L, Eaton SM. The effect of age on the cognate function of CD4⁺ T cells. *Immunol Rev* 2005; 205:220–8.
- 35- Frasca D, Riley RL, Blomberg BB. Humoral immune response and Bcell functions including immunoglobulin class switch are downregulated in aged mice and humans. *Semin Immunol* 2005; 17:378–84.
- 36- Tullius SG, Milford E. Kidney allocation and the aging immune response. *N Engl J Med.* 2011; 364(14):1369–70.
- 37-Vail A, Gore SM, Bradley BA, Easty DL, Rogers CA, Armitage WJ. Conclusions of the corneal transplant follow up study. *Br J Ophthalmol.* 1997; 81(8):631–6.
- 38- Zetterman RK, Belle SH, Hoofnagle JH, Lawlor S, Wei Y, Everhart J, et al. Age and liver transplantation: a report of the Liver Transplantation Database. *Transplantation.* 1998; 66(4):500–6.
- 39- Meier-Kriesche HU, Ojo AO, Cibrik DM, Hanson JA, Leichtman AB, Magee JC, et al. Relationship of recipient age and development of chronic allograft failure. *Transplantation.* 2000; 70(2):306–10.
- 40- Tullius SG, Reutzel-Selke A, Bachmann U, Jurisch A, Nieminen-Kelha M, Pratschke J, et al. Influence of recipient and donor age on long-term renal allograft function in an experimental model. *Transplant Proc.* 2001; 33(7-8):3345–6.
- 41- Schmidt, D., Martens, P. B., Weyand, C. M. & Goronzy, J. J. The repertoire of CD4⁺ CD28⁻ T cells in rheumatoid arthritis. *Mol Med.* 1996 Sep;2(5):608-18.
- 42- Koetz K, Bryl E, Spickschen K, O'Fallon WM, Goronzy JJ, Weyand CM. T cell homeostasis in patients with rheumatoid arthritis. *Proc Natl Acad Sci U S A.* 2000 Aug 1;97(16):9203-8.
- 43- Boots AM, Maier AB, Stinissen P, Masson P, Lories RJ, De Keyser F. The influence of ageing on the development and management of rheumatoid arthritis. *Nat Rev Rheumatol.* 2013 Oct;9(10):604-13.