

2016

Nucleophile Promoted Gold Redox Catalysis with diazonium: CBr, C-S and C-P Bond Formation through Catalytic Sandmeyer Coupling

Haihui Peng
University of South Florida

Rong Cai
West Virginia University

Chang Xu
Ohio University - Main Campus

Hao Chen
Ohio University - Main Campus

Xiaodong Shi
University of South Florida

Follow this and additional works at: https://researchrepository.wvu.edu/faculty_publications

 Part of the [Chemistry Commons](#)


Digital Commons Citation

Peng, Haihui; Cai, Rong; Xu, Chang; Chen, Hao; and Shi, Xiaodong, "Nucleophile Promoted Gold Redox Catalysis with diazonium: CBr, C-S and C-P Bond Formation through Catalytic Sandmeyer Coupling" (2016). *Faculty & Staff Scholarship*. 2364.
https://researchrepository.wvu.edu/faculty_publications/2364

This Article is brought to you for free and open access by The Research Repository @ WVU. It has been accepted for inclusion in Faculty & Staff Scholarship by an authorized administrator of The Research Repository @ WVU. For more information, please contact ian.harmon@mail.wvu.edu.

Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

COMMUNICATION

Nucleophile Promoted Gold Redox Catalysis with diazonium: C-Br, C-S and C-P Bond Formation through Catalytic Sandmeyer Coupling

Received 00th January 20xx,
Accepted 00th January 20xx

Haihui Peng,^a Rong Cai,^b Chang Xu,^c Hao Chen^{*,c} and Xiaodong Shi^{*,a}


DOI: 10.1039/x0xx00000x

www.rsc.org/

Gold-catalyzed C-heteroatom (C-X) coupling reactions are evaluated without using sacrificial oxidants. Vital to the success of this methodology is the nucleophile-assisted activation of aryl diazonium salts, which could be an effective oxidant to convert Au(I) to Au(III) even without the addition of assisting ligand or photocatalyst. By accelerating reaction kinetics to outcompete C-C homo-coupling or diazonium dediazonation, gold-catalyzed Sandmeyer reactions were achieved with different nucleophiles, forming C-Br, C-S and C-P bonds in high yields and selectivity.

Homogeneous gold catalysis has been well developed for the activation of C-C multiple bond in the past two decades.¹ However, compared with Pd(0), a d¹⁰ isoelectronic counterpart, traditional redox chemistry with Au(I) are relatively rare due to the higher oxidation potential between Au(I) and Au(III).² To maximize the potential of gold catalysis, extensive effort has been put into the development of this new branch of gold chemistry.³ Typically, strong external oxidants, such as Selectfluor and hypervalent iodine, are usually required to access catalytically active Au(III) intermediates. The need for strong oxidants made gold redox chemistry less attractive, especially for the synthesis of complex molecules. One of the most significant improvements of gold redox chemistry is the dual photoredox and gold catalysis, first reported by Glorius' and Toste's groups (Scheme 1).⁴ In their studies, photocatalyst was used to promote the gold redox chemistry under mild conditions. More recently, Hashmi and coworkers further extended the chemistry into photosensitizer-free conditions, achieving alkyne 1,2-difunctionalization with only gold catalyst under visible-light.⁵ In this study, a gold(III) intermediate was successfully isolated, which supported a gold redox catalytic

mechanism under photo-initiated conditions. Herein, we report the investigations on the nucleophile promoted diazonium activation in promoting gold(I) oxidation. Through the mechanistic investigation using NMR and Electrospray ionization mass spectrometry (ESI-MS), the nucleophile was identified as a critical factor in promoting this gold redox chemistry. In addition, through suppressing the undesired C-C homocoupling (via trans-metallation and reductive elimination), catalytic Sandmeyer coupling was achieved and C-X bonds (X = Br, S, P) were prepared with good to excellent yields.⁶ Under this new condition, no strong oxidants or photocatalysts are required to promote gold oxidation, which will potentially open new avenues for future development of gold redox chemistry.



Scheme 1. Redox gold catalysis

The high-oxidation potential between Au(I) and Au(III) has been a major concern that hindered the development of gold redox catalysis for a long time. Thus, achieving gold oxidation under mild conditions is crucial. Our group recently reported the gold catalyzed C-C coupling between alkyne and aryl diazonium salts.⁷ Based on that study, diazonium activation could be achieved with the help of 2,2'-bipyridine (bpy) ligand even without light. Although visible-light condition is extremely mild and readily accessible, understanding the function of bpy ligand will certainly help the elucidation of reaction mechanism, which will beneficially add to further development of gold redox chemistry under mild conditions.

Notably, Shin and coworkers have reported the detection of Au(III) intermediate (using XPS) through mixing PPh₃AuCl and aryl diazonium in MeOH/CH₃CN (20:1) at 60 °C.⁸ To explore the role of bpy ligands, we monitored the reaction of diazonium **1a** (*p*-F-C₆H₄N₂BF₄) and PPh₃AuCl using ³¹P NMR. Interestingly, when mixing **1a** and PPh₃AuCl in CH₃CN, no reaction was observed, even under long exposure to light at 50 °C (Figure 1a). In contrast, with MeOH/CH₃CN (9:1) as the solvent, phosphonium salt **2a** was

^a H. Peng and Prof. Dr. X. Shi, Department of Chemistry
University of South Florida, Tampa, FL 33620, USA
E-mail: xmshi@usf.edu Address here.

^b R. Cai, C. Eugene Bennett Department of Chemistry
West Virginia University, Morgantown, WV 26506, USA.

^c C. Xu and Prof. Dr. H. Chen, Center for Intelligent Chemical Instrumentation,
Department of Chemistry and Biochemistry, Edison Biotechnology Institute,
Ohio University, Athens, OH 45701, USA.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

detected (22.5 ppm, Figure 1b), though in low yield (23% based on NMR). Interestingly, with the addition of 1.0 equiv bpy, **2a** was formed under a much faster rate and PPh₃AuCl was totally consumed within an hour (Figure 1c/d).⁹





Figure 1. Monitoring the reaction of **1a** and PPh₃AuCl with ³¹P NMR

The formation of phosphonium **2a** strongly suggested that Au(III) intermediate should be included during the reaction of PPh₃AuCl and diazonium salts with the assistance from bpy. Thus, it is likely that the combination of bpy and diazonium is the actual oxidant for the oxidation of PPh₃AuCl. Notably, it has been reported in the literature that pyridine could promote diazonium activation through nucleophilic addition.¹⁰ Thus, similar function of bpy is expected as nucleophile in assisting diazonium activation, which accounts for the observed gold oxidation even without photo initiation. ESI-MS was used to explore the reaction intermediates. As expected, treating PPh₃AuCl/ArN₂⁺/bpy (*m/z* = 745.12) gave the clear formation of [PPh₃Au(Ar)bpy]⁺ cation which was also supported by further collision induced dissociation (CID) studies (MS/MS, see details in SI). This result confirmed the gold oxidation by diazonium salts with the assistance of bpy ligand. Encouraged by this discovery of nucleophilic ligand assisted diazonium activation, we wondered whether it is possible to further extend this gold redox chemistry into challenging C-X bond coupling. Our hypothesis was to explore proper anionic nucleophile to achieve both diazonium activation (for gold activation) and coupling (through reductive elimination) under this mild gold redox condition with no need of additional photosensitizers (Scheme 2).¹¹



Scheme 2. Proposed Ar-Nu coupling with gold-redox catalysis

It is well known that conversion of ArN₂⁺ to ArCl or ArBr can be achieved through standard Sandmeyer conditions using stoichiometric amount of CuX.¹² Successful examples of catalytic Sandmeyer reactions are rare. More importantly, CuX could not promote effective C-S and C-P bond formation through coupling mechanism. Compared with C-C bond coupling, formation of a C-X bond from coupling reaction is thermodynamically less favorable. Thus, there were only few successful examples reported to achieve

this important transformation catalytically.¹³ Therefore, the proposed gold-catalyzed coupling is attractive not only due to the mechanistic novelty (no need for strong oxidant or photo-activation), but also because of its potential synthetic applications (formation of challenging C-X bonds under catalytic conditions).

In the NMR studies shown in Figure 1, only a trace amount of aryl chloride was observed, although stoichiometric amount of PPh₃AuCl was used. One possibility is that the reductive elimination of Ar-Cl from Au(III) is unfavorable under gold redox conditions. In fact, Toste group recently confirmed the reductive elimination rate as I > Br > Cl through careful evaluation on different Au(III)-X bond dissociation energies.¹⁴ To explore the proposed catalytic C-X bond formation using gold redox chemistry, we started our investigation from the C-Br bond. To our great satisfaction, excellent yield of aryl bromide **5a** was achieved with the gold catalyst under mild conditions (3% PPh₃AuCl, 81% in 5 h). Results from some alternative conditions are shown in Table 1.

Table 1. Exploring reaction conditions^{a, b}

Entry	Variations from above conditions	Time	Conv. (%)	5a (%)	3a (%)	4a (%)
1	none	5 h	100	83	7	<5
2	blue LED, NO Ph ₃ PAuCl	12 h	50	<10	trace	33
3	LiBr instead of NaBr	12 h	100	78	8	<5
4	Acetone instead of ACN	5 h	100	11	37	<5
5	Ph ₃ PAuNTf ₂ instead of Ph ₃ PAuCl	5 h	100	68	10	7
6	Ph ₃ PAuNTf ₂ and 20 mol% bpy	12 h	100	63	8	15
7	3 mol% Ph ₃ PAuCl	5 h	100	81	7	<5
8	1 mol% Ph ₃ PAuCl	5 h	100	63	13	9
9	No Light (dark)	5 h	100	76	8	<5


^a Reaction conditions: **1** (0.1 mmol), NaBr (0.4 mmol), *cat.* Au (5 mol%) in acetonitrile (ACN), 50 °C; ^b ¹⁹F NMR yield with benzotrifluoride as the internal standard.

First, bpy ligand is not required in this reaction, which suggests that Br⁻ could act as an activation factor for diazonium. In fact, reacting diazonium with I⁻ gave the formation of aryl iodide even without any catalyst.¹⁵ Less than 10% Ar-Br was observed without gold catalyst (entry 2). Switching solvent to acetone caused a significantly increased yield of homo-coupling product **3a**, which suggested either different reductive elimination reaction rate (relative to transmetalation) or alternative radical reaction path. Lowering catalyst loading to 1% led to reduced yield of **5a** (entry 8, 63%) due to the increased aryl homo-coupling and diazonium decomposition (formation of ArH **4a**). Cationic gold(I) catalyst Ph₃PAuNTf₂ also promoted the reaction, though with lower yield (entries 5, 6), which is similar to the performance of Ph₃PAuBr catalyst. Importantly, similar reaction rate was observed while conducting the reaction under dark condition (entry 9), confirming the reaction as nucleophile-promoted activation rather than light-promoted diazonium decomposition. Overall, to the best of our knowledge, this is the first example of catalytic Sandmeyer reaction using only gold catalyst (no photo-activation). With this new optimal condition, various substrates are tested. The reaction substrate scope is shown in Table 2.

Excellent substrate compatibility was found. Diazonium salts with EWGs and EDGs all furnished the products in good yields (**5a-5g**). Notably, aryl iodide is also compatible in this catalytic system (**5e**), highlighting the orthogonal reactivity of Au catalyst over Pd, Cu, and Ni (for which oxidative addition could occur). Carbonyl groups (**5h**, **5i**, **5j**), benzodioxo (**5k**), azobenzene (**5q**) were well tolerated in this reaction. Hetero-aromatic diazonium salts, such as

pyridines (**5m**, **5n**) and indole (**5p**) also worked well in this reaction. Moreover, this reaction proceeded with high efficiency and selectivity with α , β -unsaturated ester (**5s**) and *p*-acetylide aryl diazonium (**5r**) to give the corresponding products. To further evaluate the synthetic utility and generality of this reaction, we tested coumarin derivative (**5l**) and estrone derivative (**5t**) to the reaction conditions. The desired products were achieved with good yields, highlighting the good potential of this catalytic system for complex molecular synthesis.

Table 2. Catalytic Ar-Br cross coupling reaction scope ^{a, b}



^a Reaction conditions: **1** (0.2 mmol), NaBr (0.8 mmol), PPh₃AuCl (3 mol%) in acetonitrile (ACN), 50 °C, 5 h; ^b Isolated yield; ^c determined by ¹⁹F NMR with benzotrifluoride as the internal standard; ^d determined by GC-Mass using decane as the internal standard; ^e LiBr (1.0 mmol) instead of NaBr.

ESI-MS studies were performed to explore the reaction mechanism. As shown in Figure 2, bisbromide-aryl-gold(III) intermediate was observed in MS under the standard reaction conditions. Through collision induced dissociation (CID) studies (MS/MS), the composition of this intermediate was confirmed (see details in SI). This result provided strong evidence for the formation of Au(III) intermediate as proposed.




Figure 2. Evidence of Au(III) intermediate from ESI-MS.

Encouraged by the success of gold catalyzed C-Br bond formation, we turned our attention to the synthesis of more challenging C-S and C-P bonds. Unlike C-Br bond, which can be alternatively prepared using stoichiometric amount of CuBr, sulfur and phosphine are not valid nucleophiles under Sandmeyer conditions due to the strong coordination of sulfur or phosphine with Cu cation (completely quenched metal reactivity).

Thiols (RSH) are good nucleophiles in general and can react with arenediazonium through S_NAr mechanism with the assistance of a base, especially for acidic thiophenols.¹⁶ However, as demonstrated above, one major side reaction of diazonium decomposition is the dediazonation (formation of Ar-H). This side reaction was more prevalent when using proton-containing nucleophiles (NuH). For example, as shown in Table 3, reaction of cysteine derivative **6a** with diazonium **1a** gave only dediazonation product **4a** in 23% yield. Addition of base (2 equiv Na₂CO₃) did help the formation of the desired thioether **7a** (37% yield), however, significant amount of dediazonation byproduct **4a** was obtained (55%). Application of stoichiometric amount of Cu(OAc)₂ did not help the reaction at all.

Table 3. Gold catalyzed C-S bond formation ^{a, b}

$$\text{HS-CH}_2\text{-CH}_2\text{-COOMe} + [\text{ArN}_2]^+[\text{BF}_4]^- \xrightarrow[\text{CH}_3\text{CN}, \text{rt}]{\text{conditions}} \text{C}_6\text{H}_4\text{-P-F-S-CH}_2\text{-CH}_2\text{-COOMe} + \text{Ar-H}$$


$$\text{6a} \qquad \qquad \qquad \text{7a} \qquad \qquad \qquad \text{4a}$$

Catalyst (mol%)	Base (equiv)	Time	Conv.(%)	7a (%)	4a (%)
None	None	24 h	30	0	23
None	Na ₂ CO ₃ (2)	10 h	100	37	55
Cu(OAc) ₂ (100)	Na ₂ CO ₃ (2)	10 h	100	31	65
PPh ₃ AuCl (5)	None	10 h	55	49	38
PPh ₃ AuCl (5)	Na ₂ CO ₃ (2)	3 h	100	87	8
PPh ₃ AuCl (3)	Na ₂ CO ₃ (2)	3 h	100	86	7
PPh ₃ AuCl (1)	Na ₂ CO ₃ (2)	7 h	100	53	30

^a Reaction conditions: **1a** (0.2 mmol), **6a** (0.1 mmol), cat. (5 mol%) and Na₂CO₃ (0.2 mmol) in acetonitrile (ACN), 50 °C; ^b ¹⁹F NMR yield with benzotrifluoride as internal standard.

Interestingly, with PPh₃AuCl as the catalyst, the desired thioether **7a** was obtained even without base (49% yield). These results suggested that with the help of thiol nucleophile, PPh₃AuCl can be an effective catalyst for diazonium decomposition, forming Au(III) even at room temperature. With the aid from base, this challenging C-S coupling was achieved in 86% yield with only 3 mol% gold catalyst loading. Based on the reaction kinetics, the C-S bond formation was dramatically improved with gold catalyst.¹⁷ The reaction scope is shown in Table 4.

Table 4. C-S cross-coupling reaction scope ^{a, b}



^a Reaction conditions, C-S formation: **1** (0.4 mmol), **6** (0.2 mmol), PPh₃AuCl (3 mol%) and Na₂CO₃ (0.4 mmol) in acetonitrile (ACN), rt, 3 h; ^b isolated yield; ^c determined by ¹H NMR using 1,3,5-trimethoxybenzene as internal standard with E/Z selectivity 1:1.

Various cysteine derivatives were successfully prepared in good yields. Both electron-rich (**7b**, **7c**, **7j**) and electron-deficient (**7a**, **7e-7i**) diazonium salts were suitable for this transformation with excellent yields. Diazonium with iodide substituent is also tolerated in this reaction (**7f**), which could be a potential synthetic handle for further functionalization. Heterocycles, including various substituted pyridines (**7k-7m**) and benzothiazole (**7n**), gave the

desired products efficiently with this catalytic system. Acetylide (**7o**) and α , β -unsaturated ester (**7p**) also reacted with good yields. Notably, sulfur containing molecules, as an important class of compounds for both chemical and biological research, are challenging to construct through traditional cross-coupling strategies because of the potential coordination between sulfur and transition metal catalysts.¹⁸ This new catalytic system thus provided an efficient strategy to achieve bioactive amino acids.

Our last attempt is to explore the possibility of C-P bond using this gold redox catalysis. Compared with C-S bond, the C-P bond formation is more challenging as H-phosphonate is much less nucleophilic and it could also be a potential reductant for diazonium.¹⁹ Thus, the C-P bond formation with diazonium cannot be achieved through either S_NAr or Cu-promoted Sandmeyer reactions. Recently, Toste and coworkers reported the application of photocatalyst in gold-catalyzed oxidative coupling to achieve this C-P bond formation.²⁰ Based on the results discussed above, we were wondering whether this nucleophile-promoted gold redox catalysis can be used to achieve this C-P bond formation.

Table 5. Ligand-assisted gold-catalyzed C-P bond formation^{a, b}

$\text{HP(O)(OEt)}_2 + [\text{ArN}_2]^+[\text{BF}_4]^- \xrightarrow[\text{CH}_3\text{CN}, 50^\circ\text{C}]{\text{conditions}}$		$\text{Ar}-\text{P(O)(OEt)}_2$	Ar-H	$\text{Ar}-\text{N}=\text{N}-\text{P(O)(OEt)}_2$		
1a		8a	4a	9a		
Catalyst (mol%)	Additives (equiv)	Time	conv. (%)	8a (%)	4a (%)	9a (%)
None	None	10 h	50	0	31	0
None	Na ₂ CO ₃ (2)	10 h	100	0	70	0
Cu(OAc) ₂ (100)	Na ₂ CO ₃ (2)	10 h	100	0	75	0
PPh ₃ AuCl (5)	None	10 h	50	25	13	0
PPh ₃ AuCl (5)	Na ₂ CO ₃ (2)	10 h	100	11	38	0
PPh ₃ AuNTf ₂ (5)	bpy (0.2), Na ₂ CO ₃ (2)	10 h	100	<5	53	11
PPh ₃ AuCl (5) ^c	3-Cl-py (2)	3 h	100	83	7	0
PPh ₃ AuNTf ₂ (5) ^c	3-Cl-py (2)	3 h	100	70	15	0
none ^d	3-Cl-py (2)	10 h	69	0	5	44
none	3-Cl-py (2)	10 h	>90	0	25	4


^a Reaction conditions: **1a** (0.2 mmol), HP(O)(OEt)₂ (0.1 mmol), cat. (5 mol%) and base (0.2 mmol) in acetonitrile (ACN), 50 °C; ^b ¹⁹F NMR yield with benzotrifluoride as internal standard; ^c ACN : EtOH = 6:1; ^d room temperature.²¹

As shown in Table 5, no desired arylphosphonate (**8a**) was obtained using base and/or copper acetate. Impressively, **8a** was formed even with solely PPh₃AuCl, though in low yield (25%). Addition of Na₂CO₃ did not improve the cross-coupling but promoted the Ar-H formation. Combination of PPh₃AuNTf₂ and bpy in the presence of Na₂CO₃ (previously reported C-C bond coupling conditions) also failed to increase the yield of the desired C-P coupling product. Considering that nucleophilic ligand is crucial in this gold redox catalysis, we put our attention on other pyridine derivatives. Through a comprehensive screening, 3-Cl-pyridine was identified as the optimal nucleophile (see detailed screenings in SI), giving the desired C-P bond coupling product **8a** in 83% isolated yield. Using PPh₃AuNTf₂ as catalyst led to a lower yield of **8a** due to the increasing rate of side reactions. Notably, without gold catalyst, diaza compound **9a** was formed as the major product at room temperature whereas no desired coupling product **8a** was detected.²² At 50 °C, the reaction was very messy and **8a** was not detected at all, which suggested that **8a** was not formed from decomposition of diaza compound **9a**. The reaction substrate scope is shown in Table 6.

Similar to the C-Br and C-S coupling, broad substrate scope is received in the C-P bond formation reactions. Aryl phosphonates with electron rich (**8b**, **8c**) and electron deficient (**8a**, **8d-8h**) substituents could all be generated with good yields. Halogen substitute groups (**8a**, **8e**, **8f**) were all tolerated. Alkyne (**8i**), α , β -

unsaturated ester (**8j**) and azobenzene (**8k**) also gave good results, suggesting great synthetic potential of this methodology.

Table 6. Catalytic C-P cross-coupling reaction scope^{a, b}



^a Reaction conditions: **1** (0.4 mmol), HP(O)(OEt)₂ (0.2 mmol), PPh₃AuCl (5 mol%) and 3-Cl-Py (0.4 mmol) in acetonitrile (ACN), 50 °C, 5 h; ^b isolated yield; ^c determined by ¹H NMR using 1,3,5-trimethoxybenzene as internal standard.

Conclusions

In summary, herein we reported the C-Br, C-S, and C-P bond formation through gold redox catalysis. We demonstrated that nucleophiles played a crucial role in the Au(I) promoted diazonium decomposition. With this strategy, various C-X coupling could be achieved with excellent yields and broad substrate scope simply using LAuCl (no need for external oxidant). These results not only provide a new practical strategy to achieve challenging C-X bond coupling, but also, more importantly, revealed some new mechanistic insight regarding gold redox catalysis, which will likely further enrich the pedigree of gold catalysis.

Acknowledgement

We are grateful for financial support from the NSF (CHE-1619590) and NSFC (21228204). CX and HC thank support from NSF career Award (CHE-1149367) and NSF IDBR (CHE-1455554).

Notes and references

- Selected recent reviews on Au catalysis: (a) A. S. K. Hashmi, *Chem. Rev.* 2007, **107**, 3180; (b) A. Fürstner and P. W. Davies, *Angew. Chem., Int. Ed.* 2007, **46**, 3410; (c) D. Gorin and F. D. Toste, *Nature*, 2007, **446**, 395; (d) A. Arcadi, *Chem. Rev.*, 2008, **108**, 3266; (e) Z. Li, C. Brouwer and C. He, *Chem. Rev.*, 2008, **108**, 3239; (f) D. J. Gorin, B. D. Sherry and F. D. Toste, *Chem. Rev.*, 2008, **108**, 3351; (g) R. A. Widenhoefer, *Chem. Eur. J.*, 2008, **14**, 5382; (h) A. Fürstner, *Chem. Soc. Rev.*, 2009, **38**, 3208; (i) N. Krause and C. Winter, *Chem. Rev.*, 2011, **111**, 1994; (j) M. Bandani, *Chem. Soc. Rev.*, 2011, **40**, 1358; (k) M. N. Hopkinson, A. D. Gee and V. Gouverneur, *Chem. Eur. J.*, 2011, **17**, 8248; (l) A. Corma, A. Leyva-Pérez, and M. Sabater, *Chem. Rev.*, 2011, **111**, 1657; (m) T. de Haro and C. Nevado, *Synthesis*, 2011, 2530; (n) L. Liu and G. B. Hammond, *Chem. Soc. Rev.*, 2012, **41**, 3129; (o) M. Rudolph and A. S. K. Hashmi, *Chem. Soc. Rev.*, 2012, **41**, 2448; (p) L. Zhang, *Acc. Chem. Res.*, 2014, **47**, 877; (q) R. Dorel and A. M. Echavarren, *Chem. Rev.*, 2015, **115**, 9028; (r) D. Pflästerer, A. S. K. Hashmi, *Chem. Soc. Rev.* 2016, **45**, 1331.
- E^0 (Au^I/Au^{III}) = +1.40 V. See: *CRC Handbook of Chemistry and Physics*, 84th ed.; D. R. Lide, CRC Press, Boca Raton, FL, 2004. For a stable example, including an X-ray crystal structure analysis, of a dicoordinated gold(I) center not undergoing oxidative addition with an aryl iodide that could intramolecularly react by a six-membered transition state, see: A. S. K. Hashmi, C. Lothschütz, R. Döpp, M. Ackermann, J. D. B. Becker, M. Rudolph, C. Scholz and F. Rominger, *Adv. Synth. Catal.*, 2012, **354**, 133.

3 For selected examples, see: (a) A. S. K. Hashmi, T. D. Ramamurthi and F. Rominger, *J. Organomet. Chem.*, 2009, **694**, 592; (b) G. Zhang, Y. Peng, L. Cui and L. Zhang, *Angew. Chem., Int. Ed.*, 2009, **48**, 3112; (c) M. N. Hopkinson, J. E. Ross, G. T. Giuffredi, A. D. Gee and V. Gouverneur, *Org. Lett.*, 2010, **12**, 4904; (d) G. Zhang, L. Cui, Y. Wang and L. Zhang, *J. Am. Chem. Soc.*, 2010, **132**, 1474; (e) A. D. Melhado, W. E. Brenzovich, A. D. Lackner and F. D. Toste, *J. Am. Chem. Soc.*, 2010, **132**, 8885; (f) T. de Haro and C. Nevado, *J. Am. Chem. Soc.*, 2010, **132**, 1512; (g) A. Leyva-Pórez, A. Domónech, S. I. Al-Resayes and A. Corma, *ACS Catal.*, 2012, **2**, 121; (h) S. Shi, T. Wang, W. Yang, M. Rudolph and A. S. K. Hashmi, *Chem. Eur. J.*, 2013, **19**, 6576; (i) L. T. Ball, G. C. Lloyd-Jones and C. A. Russell, *Science*, 2012, **337**, 1644; (j) W. J. Wolf, M. S. Winston and F. D. Toste, *Nat. Chem.*, 2014, **6**, 159; (k) L. T. Ball, G. C. Lloyd-Jones and C. A. Russell, *J. Am. Chem. Soc.*, 2014, **136**, 254; (l) M. Joost, A. Zeineddine, L. Estóvez, S. Mallet-Ladeira, K. Miqueu, A. Amgoune and D. Bourissou, *J. Am. Chem. Soc.*, 2014, **136**, 14654; (m) Q. Wu, C. Du, Y. Huang, X. Liu, Z. Long, F. Song and J. You, *Chem. Sci.*, 2015, **6**, 288; (n) M. Joost, L. Estóvez, K. Miqueu, A. Amgoune and D. Bourissou, *Angew. Chem., Int. Ed.*, 2015, **54**, 5236; (o) M. D. Levin, S. Kim and F. D. Toste, *ACS Cent. Sci.*, 2016, **2**, 293.

4 (a) B. Sahoo, M. N. Hopkinson and F. Glorius, *J. Am. Chem. Soc.*, 2013, **135**, 5505; (b) X.-Z. Shu, M. Zhang, Y. He, H. Frei and F. D. Toste, *J. Am. Chem. Soc.*, 2014, **136**, 5844; (c) M. Wieteck, Y. Tokimizu, M. Rudolph, F. Rominger, H. Ohno, N. Fujii and A. S. K. Hashmi, *Chem. Eur. J.*, 2014, **20**, 16331; (d) J. Xie, S. Shi, T. Zhang, N. Mehrkens, M. Rudolph and A. S. K. Hashmi, *Angew. Chem., Int. Ed.*, 2015, **54**, 6046; (e) A. Tlahuext-Aca, M. N. Hopkinson, B. Sahoo and F. Glorius, *Chem. Sci.*, 2016, **7**, 89; (f) S. Kim, J. Rojas-Martin and F. D. Toste, *Chem. Sci.*, 2016, **7**, 85; (g) J. Xie, T. Zhang, F. Chen, N. Mehrkens, F. Rominger, M. Rudolph and A. S. K. Hashmi, *Angew. Chem., Int. Ed.*, 2016, **55**, 2934; (h) S. Hosseyni, L. Wojtas, M. Li, X. Shi, *J. Am. Chem. Soc.*, 2016, **138**, 3994.

5 (a) L. Huang, M. Rudolph, F. Rominger, A. S. K. Hashmi, *Angew. Chem., Int. Ed.*, 2016, **55**, 4808; (b) L. Huang, F. Rominger, M. Rudolph, A. S. K. Hashmi, *Chem. Commun.*, 2016, **52**, 6435.


6 The monitored quantitative rate constant k_{obs} for sp^2 C-C coupling is $(1.5 \pm 0.1) \times 10^{-4} \text{ s}^{-1}$ ($-52 \text{ }^\circ\text{C}$), see reference [3h]; the homo-coupling product was generated from transmetalation between Au(III) and Au(I) intermediate followed by reductive elimination, see reference [3f].

7 R. Cai, M. Lu, E. Y. Aguilera, Y. Xi, N. G. Akhmedov, J. L. Peterson, H. Chen and X. Shi, *Angew. Chem., Int. Ed.*, 2015, **54**, 8772.

8 D. V. Patil, H. Yun and S. Shin, *Adv. Synth. Catal.*, 2015, **357**, 2622.

9 The structure with 27.7 ppm in ^{31}P NMR is not clear right now. According to literature, ^{31}P NMR signal of Au(III) intermediate is around 27 ppm (ref. 4b, 5).

10 A pyridine promoted diazonium decomposition through the formation of aryl radical was proposed: T. Sakakura, M. Hara and M. Tanaka, *J. Chem. Soc., Chem. Commun.*, 1985, 1545.



11 Another possible mechanism is that the nucleophiles are reacting with LAuCl to generate more electron-rich LAuNu, which would be more susceptible to oxidation by diazonium salts.

12 For reviews on diazonium salts in cross-couplings, see: (a) S. Bräse, *Acc. Chem. Res.*, 2004, **37**, 805; (b) A. Roglans, A. Pla-Quintana and M. Moreno-Mañas, *Chem. Rev.*, 2006, **106**, 4622; (c) F. Mo, G. Dong, Y. Zhang and J. Wang, *Org. Biomol. Chem.*, 2013, **11**, 1582. For selected examples using diazonium, see: (a) C. Molinaro, J. Mowat, F. Gosselin, P. D. O'Shea, J.-F. Marcoux, R. Angelaud and I. W. Davies, *J. Org. Chem.*, 2007, **72**, 1856; (b) I. P. Beletskaya, A. S. Sigeev, A. S. Peregudov and P. V. Petrovskii, *Synthesis*, 2007, 2534; (c) G. Danoun, B. Bayarmagnai, M. F. Grünberg and L. J. Gooßen, *Angew. Chem., Int. Ed.*, 2013, **52**, 7972; (d) J. Dai, C. Fang, B. Xiao, J. Yi, J. Xu, Z. Liu, X. Lu, L. Liu and Y. Fu, *J. Am. Chem. Soc.*, 2013, **135**, 8436; (e) X. Wang,

Y. Xu, F. Mo, G. Ji, D. Qiu, J. Feng, Y. Ye, S. Zhang, Y. Zhang and J. Wang, *J. Am. Chem. Soc.*, 2013, **135**, 10330; (f) C. Matheis, V. Wagner and L. J. Gooßen, *Chem. Eur. J.*, 2016, **22**, 79.

13 For selected examples, see: (a) X. Shen, A. M. Hyde and S. L. Buchwald, *J. Am. Chem. Soc.*, 2010, **132**, 14076; (b) J. Pan, X. Wang, Y. Zhang and S. L. Buchwald, *Org. Lett.*, 2011, **13**, 4974; (c) Y. Imazaki, E. Shirakawa, R. Ueno and T. Hayashi, *J. Am. Chem. Soc.*, 2012, **134**, 14760; (d) M. Feller, Y. Diskin-Posner, G. Leitens, L. J. W. Shimon and D. Milstein, *J. Am. Chem. Soc.*, 2013, **135**, 11040; (e) C. M. Le, P. J. C. Menzies, D. A. Petrone and M. Lautens, *Angew. Chem., Int. Ed.*, 2015, **54**, 254; (f) C. Chen, L. Hou, M. Cheng, J. Su and X. Tong, *Angew. Chem., Int. Ed.*, 2015, **54**, 3092.

14 M. S. Winston, W. J. Wolf and F. D. Toste, *J. Am. Chem. Soc.*, 2015, **137**, 7921.

15 See recent example: A. Hubbard, T. Okazaki, K. K. Laali, *J. Org. Chem.*, 2008, **73**, 316.

16 P. C. B. Page, R. D. Wilkes and D. Reynolds, in *Comprehensive Organic Functional Group Transformations*, ed. A. R. Katritzky, O. Meth-Cohn and C. W. Rees, Elsevier, Oxford, 1995, p. 113; For selected recent examples, see: (a) M. Majek and A. J. V. Wangelin, *Chem. Commun.*, 2013, **49**, 5507; (b) M. Barbero, I. Degani, N. Diulgheroff, S. Dughera, R. Fochi and M. Migliaccio, *J. Org. Chem.*, 2000, **65**, 5600; (c) X. Wang, G. D. Cuny and T. Noël, *Angew. Chem., Int. Ed.*, 2013, **52**, 7860.

17 Without catalyst, C-S coupling product **7a** has nearly same kinetics with byproduct **4a**; with the gold catalyst, the formation of **7a** is accelerated with less than 10% **4a** formed. See detailed kinetic study in SI.

18 For selected examples, see: (a) P. Johannesson, G. Lindeberg, A. Johannesson, G. V. Nikiforovich, A. Gogoli, B. Synergren, M. Le Greves, F. Nyberg, A. Karlen and A. Hallberg, *J. Med. Chem.*, 2002, **45**, 1767; (b) L. Llauger, H. Z. He, J. Kim, J. Aguirre, N. Rosen, U. Peters, P. Davies and G. Chiosio, *J. Med. Chem.*, 2005, **48**, 2892; (c) A. Gangjee, Y. B. Zheng, T. Talreja, J. J. Mc Guire, R. L. Kisliuk and S. F. Queener, *J. Med. Chem.*, 2007, **50**, 2046.

19 For recent examples on C-P bond formation, see: (a) J. Xuan, T. Zeng, J. Chen, L. Lu and W. Xiao, *Chem. Eur. J.*, 2015, **21**, 4962; (b) G. Hu, W. Chen, T. Fu, Z. Peng, H. Qiao, Y. Gao and Y. Zhao, *Org. Lett.*, 2013, **15**, 5362; (c) O. Berger, C. Petit, E. L. Deal and J. L. Montchamp, *Adv. Synth. Catal.*, 2013, **355**, 1361; (d) K. Xu, H. Hu, F. Yang and Y. Wu, *Eur. J. Org. Chem.*, 2013, **2013**, 319; (e) A. J. Bloomfield and S. B. Herzon, *Org. Lett.*, 2012, **14**, 4370; (f) S. M. Rummelt, M. Ranocchiari and J. A. van Bokhoven, *Org. Lett.*, 2012, **14**, 2188; (g) C. R. Shen, G. Q. Yang and W. B. Zhang, *Org. Biomol. Chem.*, 2012, **10**, 3500; (h) E. L. Deal, C. Petit and J. L. Montchamp, *Org. Lett.*, 2011, **13**, 3270.

20 Y. He, H. Wu and F. D. Toste, *Chem. Sci.*, 2015, **6**, 1194.

21 Reaction of L-AuCl with pyridine did not give any [LAu(pyridine)]⁺ based on NMR, which confirmed that [LAuNu]⁺ was unlikely the actual catalyst for this reaction.

22 In palladium catalyzed C-P coupling with diazonium in the presence of CuI, **9a** was also isolated as the major byproduct. See: R. Berrino, S. Cacchi, G. Fabrizi, A. Goggiani and P. Stabile, *Org. Biomol. Chem.*, 2010, **8**, 4518.