Available online on 15.06.2020 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Research Article

Comparative Biological Activity Profile of *Nava* (Freshly Collected) and *Purana* (Old) *Dhanyaka* (*Coriandrum sativum* Linn.) through Gas Chromatography and *In-Vitro* Anti-Microbial Study

Chaithra¹, Mallya Suma V^{2*}, Prabhu Suchitra³¹ PG Scholar, Sri Dharmasthala Manjunatheshwara College of Ayurveda, Kuthpady, Udipi, Karnataka 574 118, India² Associate Professor, Sri Dharmasthala Manjunatheshwara College of Ayurveda, Kuthpady, Udipi, Karnataka 574 118, India³ Research Officer, Sri Dharmasthala Manjunatheshwara College of Ayurveda, Kuthpady, Udipi, Karnataka 574 118, India

ABSTRACT

Introduction: *Dhanyaka* (*Coriandrum sativum* Linn.) belonging to family *Apiaceae* is one such common aromatic edible herb, the fruits of which are used as *carminative, antimicrobial and cooling*. Classical texts of *Ayurveda* advocate these fruits to be used in old condition. With this background, a study was done to compare biological activity profile of *Nava* and *Purana Dhanyaka* (*Coriandrum sativum* Linn.) through, Gas chromatography and *in-vitro* anti-microbial activity. **Materials and methods:** Fresh fruits of *Dhanyaka* (*Coriandrum sativum* Linn.) were collected from Rona, Gadag district and authenticated. The fresh fruits named as **S₁** with which essential oil extracted using clevengers apparatus. Few fruits were kept in an air tight container and stored for one year, and at the end of a year named as sample (**S₂**), essential oil extracted and stored. Both samples were subjected for GC and *in-vitro* antimicrobial activity. **Results and Discussion:** The Essential oil of *Nava Dhanyaka* (**S₁**) included 17 identified compounds which was dominated by monoterpenes and sesquiterpene; whereas **S₂** showed 6 identified compounds in which 3 constituents *isoborneol, citronellyl acetate* and *geraniol* were new findings. *In-vitro* anti-microbial study of essential oil extract showed that *Purana Dhanyaka* (**S₂**) had better efficacy with respect to the standard against the strain *Escherichia coli* and *Candida albicans*; whereas that of *Nava Dhanyaka* (**S₁**) against only one bacterial strain i.e., *Streptococcus pyogenes*. **Conclusion:** Preservation of *Dhanyaka* (*Coriandrum sativum* Linn.) showed biological changes in terms of antimicrobial activity compared to fresh sample.

Key words: *Dhanyaka, Coriandrum sativum* Linn., Gas chromatography, *In-vitro* antimicrobial study

Article Info: Received 21 March 2020; Review Completed 28 April 2020; Accepted 11 May 2020; Available online 15 June 2020



Cite this article as:

Chaithra, Mallya SV, Prabhu S, Comparative Biological Activity Profile of *Nava* (Freshly Collected) and *Purana* (Old) *Dhanyaka* (*Coriandrum sativum* Linn.) through Gas Chromatography and *In-Vitro* Anti-Microbial Study, Journal of Drug Delivery and Therapeutics. 2020; 10(3-s):12-16 <http://dx.doi.org/10.22270/jddt.v10i3-s.4060>

*Address for Correspondence:

Mallya Suma V, Associate professor, Sri Dharmasthala Manjunatheshwara College of Ayurveda, Kuthpady, Udipi, Karnataka 574 118, India

INTRODUCTION:

Traditional knowledge on herbs, their potential therapeutic effect was well known among people, also documented in classical text books of *Ayurveda*. Few they used directly from plant, few parts advised to use after drying, other after special processing, the main aim to get target therapeutic effect through hidden metabolites of these natural products¹. *Dhanyaka* (*Coriandrum sativum* Linn.) belonging to family *Apiaceae* is a common aromatic edible herb, the fruits of which are used as *Ahara* and *Oushadha* in many therapeutic conditions². The whole plant, as well as the fruits are used in the preparation of many formulations having therapeutic efficacy in *trishna, Krimiroga, Daha, Mutravikara* etc³. The fruits are to be preserved for a period of 1 year and to be used for medicine. *Adamalla* explains that keeping certain

drugs for period of one year can be taken as *purana dravya* and it increases the potency of the drug.⁴

Biological activity shown by different herbal drug depend on its chemical constituents⁵. Essential oils extracts are the products from the secondary metabolism of plants that have applications in *ethno-medicine, food flavouring, food preservation, cosmetics and pharmaceutical industries*.⁶ *Dhanyaka* (*Coriandrum sativum* Linn.) has been known to have abundant essential oil which has antimicrobial properties⁷.

Hence with above all mentioned facts, a research plan has been designed to investigate chemical constituents of fresh and old sample of *Dhanyaka* (*Coriandrum sativum* Linn.) through Gas chromatography, and to evaluate the anti-

microbial activity of Nava And Purana Dhanyaka (*Coriandrum sativum* Linn.) through In-Vitro method.

MATERIALS AND METHOD:

Sample collection:

Matured whole fruits of Dhanyaka(*Coriandrum sativum* Linn.) were collected from the Rona, Gadag district, authenticated and voucher specimen deposited at department of Pharmaceutical Chemistry and Pharmacognosy, SDM Centre for research in Ayurveda and Allied sciences, Udupi(V/No.18060103).The fresh fruits named as S₁ with which essential oil extracted using clevengers apparatus⁸. Few fruits were kept in an air tight container and stored for one year, and at the end of a year named as sample(S₂), essential oil extracted and stored. Both samples were subjected for GC and in-vitro antimicrobial activity.

Gas chromatography

Methodology:

Volatile oil in the drug was estimated by distilling the 500g of coarsely powdered fruits of *Coriandrum sativum* Linn. with a mixture of 800ml of water, taken in a round bottom flask and connected to the Clevenger's apparatus and distillation was carried out. Volume of the essential oil was measured and sent for the Gas Chromatography study⁹. Methylated test sample was introduced into a stream of helium. The results were tabulated as per the retention time.

In-vitro antimicrobial study

Methodology:

The In-vitro Antimicrobial study is performed using well diffusion method¹⁰. Agar media was inoculated with loop of

desired microorganism (Table 2). Test was conducted for different concentrations of extract at 10, 25, 50 and 80µl and kept for incubation. The presence of zone of inhibition was observed on petridish with respect to standard drug.

Table 1: Microorganism strains used

| Microorganism strain | Code | Standard used |
|-------------------------------|-----------|---------------|
| <i>Staphylococcus aureus</i> | MTCC 3160 | Ampicillin |
| <i>Streptococcus pyogenes</i> | MTCC-86 | Ampicillin |
| <i>Escherichia coli</i> | MTCC-42 | Ampicillin |
| <i>Candida albicans</i> | MTCC- 183 | Clotrimazole |

OBSERVATION AND RESULT

Gas chromatography

The fruits of Nava Dhanyaka (*Coriandrum sativum* Linn.) S₁ yielded about 0.7ml of yellow coloured essential oil with pleasant aroma. The GC-MS analysis of the fruits showed presence of totally 40 compounds in which 17 compounds were identified by comparing their Mass Spectra with MS library. The fruits of Purana Dhanyaka (*Coriandrum sativum* Linn.) S₂ yielded about 0.5ml of colourless essential oil with pleasant aroma. The GC-MS analysis of the fruits showed presence of totally 18 compounds in which 6 compounds were identified by comparing their Mass Spectra with MS library. Comparative essential oil composition of Nava (S₁) and Purana Dhanyaka(S₂) (*Coriandrum sativum* Linn.) are given in Table 1, Figure 1.




Figure 1: Gas chromatography of Essential oil Nava Dhanyaka(*Coriandrum sativum* Linn.) (S₁)




Figure 2: Gas chromatography of Essential oil of Purana Dhanyaka(*Coriandrum sativum* Linn.) (S₂)

Table 2: Essential oil composition of Nava (S₁) and Purana(S₂) Dhanyaka(Coriandrum sativum Linn.)

| Phytochemical constituents | S ₁ | S ₂ |
|---|----------------|----------------|
| Bicyclo[2.2.1]heptan-2-ol, 1,7,7-trimethyl-, (1S-endo)- | - | + |
| alpha.-Terpineol | + | - |
| Decanal | + | - |
| Bicyclo[3.1.1]hept-3-en-2-one, 4,6,6-trimethyl | + | - |
| Geraniol | - | + |
| Citronellol | + | + |
| 2,6-Octadien-1-ol, 3,7-dimethyl-(Z) | + | - |
| 1-Decanol | + | - |
| Undecanal | + | - |
| trans-Geranic acid methyl ester | + | - |
| Myrtenyl acetate | + | - |
| 6-Octen-1-ol, 3,7-dimethyl-, acetate | - | + |
| 3,7-Dimethyloct-6-enyl ethyl carbonate | + | - |
| 2,6-Octadien-1-ol, 3,7-dimethyl-, acetate, (Z)- | + | + |
| Dodecanal | + | - |
| Bicyclo[5.2.0]nonane,2-methylene-4,8,8-trimethyl-4-vinyl- | + | + |
| (E)-Hexadec-2-enal | + | - |
| Caryophyllene oxide | + | - |
| 2-Hexadecenal(Z) | + | - |
| 2-Pentadecanone, 6,10,14-trimethyl- | + | - |

In-vitro antimicrobial study

Zone of inhibition of essential oil extract of Nava (S₁) and Purana (S₂) Dhanyaka (Coriandrum sativum Linn.) against *Escherichia coli*, *Staphylococcus aureus*, *Streptococcus pyogenes* and *Candida albicans* displayed in Table 3,4,5 and 6 respectively.

Table 3: Zone of inhibition of Nava (S₁) and Purana (S₂) Dhanyaka (Coriandrum sativum Linn.) against Escherichia coli

| Sample | Volumes | S ₁ - Zone of inhibition(Radius in mm) | S ₂ - Zone of inhibition (Radius in mm) |
|--------------------------------|---------|---|--|
| Essential oil extract | 10 µl | 0 | 0 |
| | 20 µl | 0 | 0 |
| | 50 µl | 5 | 8 |
| | 80 µl | 6 | 9 |
| Control (DW) | 50 µl | 0 | 0 |
| Standard (<i>Ampicillin</i>) | 30 µg | 10 | 11 |

Table 4: Zone of inhibition of Nava (S₁) and Purana (S₂) Dhanyaka (Coriandrum sativum Linn.) against Staphylococcus aureus

| Sample | Volume | S ₁ - Zone of inhibition(Radius in mm) | S ₂ - Zone of inhibition(Radius in mm) |
|--------------------------------|--------|---|---|
| Essential oil extract | 10 µl | 5 | 0 |
| | 20 µl | 6 | 0 |
| | 50 µl | 8 | 0 |
| | 80 µl | 9 | 6 |
| Control (DW) | 50 µl | 0 | 0 |
| Standard (<i>Ampicillin</i>) | 30 µg | 8 | 8 |

Table 5: Zone of inhibition of *Nava* (S₁) and *Purana* (S₂) *Dhanyaka* (*Coriandrum sativum* Linn.) against *Streptococcus pyogenes*

| Sample | Volumes | S ₁ - Zone of inhibition(Radius in mm) | S ₂ - Zone of inhibition(Radius in mm) |
|--------------------------------|---------|---|---|
| Essential oil extract | 10 µl | 5 | 0 |
| | 20 µl | 6 | 0 |
| | 50 µl | 7 | 0 |
| | 80 µl | 8 | 6 |
| Control (DW) | 50 µl | 0 | 0 |
| Standard (<i>Ampicillin</i>) | 30 µg | 15 | 20 |

Table 6: Zone of inhibition of *Nava* (S₁) and *Purana* (S₂) *Dhanyaka* (*Coriandrum sativum* Linn.) Against *Candida albicans*

| Sample | Volumes | S ₁ - Zone of inhibition(Radius in mm) | S ₂ - Zone of inhibition(Radius in mm) |
|----------------------------------|---------|---|---|
| Essential oil extract | 10 µl | 12 | 0 |
| | 20 µl | 13 | 11 |
| | 50 µl | 15 | 12 |
| | 80 µl | 15 | 13 |
| Control (DW) | 50 µl | 0 | 0 |
| Standard (<i>Clotrimazole</i>) | 30 µg | 10 | 8 |

DISCUSSION:

Herbal drugs are seat of various active principle, proper collection time, preservation, processing decide its biological activity¹¹. *Dhanyaka* (*Coriandrum sativum* Linn.) the dried fruits of which are used in medicine, which are said to be antimicrobial, appetizers, cooling. Fruits said to contain various phytochemical constituents with rich essential oil content. Classical texts of Ayurveda, advices them to use after a year⁴. Gas chromatographic studies and anti-microbial study of *Nava* (fresh) and *Purana* (after 1 year) *Dhanyaka* (*Coriandrum sativum* Linn.) will help us to evaluate biological changes in a drug after preserving for a year.

The Essential oil of *Nava Dhanyaka* included 17 identified compounds which was dominated by monoterpenes like *Terpineol*, *Verbenone*, *citronellol*, acyclic monoterpenes like *Methyl geranate*, *Myrtenyl acetate*, sesquiterpene like *Caryophyllene oxide*. *Terpineol* is a common ingredient in perfumes, cosmetics and is abundantly used as a flavouring agent¹². *α-terpineol* has application in therapeutics due to its anti-oxidant, anti-ulcer and insecticidal properties. *Verbenone*, *Methyl geranate*, *Myrtenyl acetate* give a pleasant characteristic aroma. It also has anti-microbial properties¹³. *Citronellol* is also used as an anti-infective agent in many conditions¹⁴. The major aldehydes identified were *Decanal*, *Undecanal*, *1-Decanol* and *Dodecanal*. *Decanal* is used in foods as a flavouring agent and also has anti-fungal property¹⁵.

The essential oil of *Purana(old)Dhanyaka*(*C. Sativum*) showed 6 identified compounds in which 3 constituents *isoborneol*, *citronellyl acetate* and *geraniol* were specifically identified in *Purana Dhanyaka*.

Isoborneol is extensively used as a food additive. It has camphor like aroma and is also used in insect repellants. It is used as a potential anti-viral agent and has anti-microbial, anti-tumour properties. *Citronellyl acetate* has potential larvicidal, bactericidal and fungicidal properties. *Geraniol* is one of the major constituent of *Coriandrum sativum* Linn. responsible for characteristic aroma. Along with being used as a flavouring agent, it is used as antibacterial, antifungal¹⁷.

In-vitro anti-microbial study of essential oil extract showed that *Purana Dhanyaka*(S₂) had better efficacy with respect to the standard against the strain *Escherichia coli* and *Candida albicans*. The essential oil extract of *Nava Dhanyaka*(S₁) had better efficacy with respect to the standard against only one bacterial strain ie., *Streptococcus pyogens*.

CONCLUSION:

Preserving certain crude drugs will make changes among biological activity by the influence of time, environmental factors, climatic changes etc and also brings about changes in the arrangement, structure and phyto-constituents of the crude drug. *Dhanyaka*(*Coriandrum sativum* Linn.) the fruits which advised to use in medicine after an year were evaluated for thier phyto-constituents and anti-bacterial activity after preserving for a period of an year and has shown significant results. This study gives us a better understanding of rationality behind the effect of collection and preservation of a crude drug and its impact on phyto-chemical constituents, pharmacotherapeutic actions.

REFERENCES:

1. Thangavel A, Duraisamy S, Blakrishnan S; Phytochemical screening, Gas chromatography- Mass spectrometry(GC-MS)analysis of phytochemical constituents and antibacterial activity of *Coriandrum sativum*(L.) SE; International journal of pharmacy and pharmaceutical sciences, 2015; 7(9):153-59.
2. Nadakarni KM., Indian Material Medica Vol-I, Dehradun; International book Distributor: 2006, p395
3. Anonymous. The wealth of India, A dictionary of Indian raw materials & industrial products Vol 2, Newdelhi: Council of scientific & industrial research: 2009 P.270- 275
4. Murthy Srikantha KR, Sarangadhara Samhita, Chaukambha Orientalia, 1997, Varanasi, p7-8
5. Sharma Rakesh K, Arora Rajesh; Herbal Drugs–A twenty first century perspective; New Delhi: Jaypee Brothers Medical Publishers, 2006p488-89
6. Evans WC., Trease and Evan's Pharmacognosy, 15th ed. London: WB Saunders Ltd: 2002; p45- 150
7. Reneta nurzynska- Wierdak; Essentail oil composition of the coriander (*Coriandrum sativum* L.) herb depending on the development stage; Acta agrobotanica; 2013; 66(1):53-60.
8. Nicholes J. Sadgrove, Graham Lloyd Jones; A contemporary introduction to essential oil; Chemistry, bioactivity and prospects for Australian agriculture; Agriculture 2015; 5:48-102.
9. Kanthal LK, Dey A, Bhojarju P; GC-MS analysis of bio-active compounds in methanolic extract of *Lacuca runcinate* DC; Pharmacognosy Research; 2014; 6(1):58-61.
10. Mallya SV, Nesari T; Antibacterial activity profile and quality standards of *Cymbopogon citratus* Stapf- an aromatic grass used in Indian system of medicine; Journal of Ayurvedic and Herbal Medicine 2016; 2(3):63-66
11. Mukharjee PK, Quality Control of Herbal Drugs; New Delhi: Business Horizons; 2002; p68
12. Khaleel C, Tabanca N, Buchbauer G; Alpha Terpineol, a natural monoterpene; A review of its biological properties; open chemistry; 2018;16:349-361
13. Lindgren BS, Miller DR; Effect of Verbenone on fie species of Bark Beetles(Coleoptera; Scolytidae) in Lodgepole Pine Forests; Environmental Entomology, 2002 ; 31(5):759-65. <https://pubchem.ncbi.nlm.nih.gov>
14. <https://en.m.wikipedia.org>wiki>
15. Chen W, Viljoen AM; Geraniol- A review of a commercially important fragrance material; South African Journal of Botany; 2010 ; 76(4):643-51
16. Leite Maria Clerya Alvino, Bezerra André Parente de Brito, de Sousa Janiere Pereira, de Oliveira Lima Edeltrudes. Investigating the antifungal activity and mechanism(s) of geraniol against *Candida albicans* strains. Medical Mycology; 2015; 53(3):275-284.

Journal of Drug Delivery & Therapeutics



JDDT