



Sacred Heart University
DigitalCommons@SHU

Athletic Training Faculty Publications

Physical Therapy & Human Movement Science

7-2015

Predictors of Orthopaedic Surgery in NCAA Athletes

Dean Wang

University of California, Los Angeles

Caitlin M. Rugg

University of California, Los Angeles

Erik Mayer

University of California, Los Angeles

Pamela Sulzicki

Sacred Heart University

Jeremy Vail

University of California, Los Angeles

See next page for additional authors

Follow this and additional works at: https://digitalcommons.sacredheart.edu/pthms_atfac



Part of the [Biomechanics Commons](#), [Orthopedics Commons](#), and the [Sports Sciences Commons](#)

Recommended Citation

Wang, D., Rugg, C. M., Mayer, E., Sulzicki, P., Vail, J., & Hame, S. L. (2015). Predictors of orthopaedic surgery in NCAA athletes. *Orthopaedic Journal of Sports Medicine*, 3(7 suppl2), 2325967115S00096. <https://doi.org/10.1177/2325967115S00096>

This Peer-Reviewed Article is brought to you for free and open access by the Physical Therapy & Human Movement Science at DigitalCommons@SHU. It has been accepted for inclusion in Athletic Training Faculty Publications by an authorized administrator of DigitalCommons@SHU. For more information, please contact ferribyp@sacredheart.edu, lysobeyb@sacredheart.edu.

Authors

Dean Wang, Caitlin M. Rugg, Erik Mayer, Pamela Sulzicki, Jeremy Vail, and Sharon L. Hame

Predictors of Orthopaedic Surgery in NCAA Athletes

Dean Wang, MD¹, Caitlin Marie Rugg¹, Erik Mayer¹, Pamela Sulzicki, MS, ATC², Jeremy Vail, MPT, OCS, MTC, ATC¹, Sharon L. Hame, MD¹

¹UCLA, Los Angeles, CA, USA, ²Sacred Heart University, Fairfield, CT, USA.

Objectives: Orthopaedic injury and surgery is relatively common in National Collegiate Athletic Association (NCAA) athletes and can have devastating career consequences. However, there is a paucity of data regarding predictors of orthopaedic surgery in collegiate athletes. The purpose of this study was to analyze player-related predictors of orthopaedic surgery, including that of the shoulder, hip, and knee, in NCAA athletes.

Methods: All NCAA Division I collegiate athletes at a single institution who began participation from the 2003-2004 through 2008-2009 seasons were retrospectively identified. Player-related factors, including gender, sport, and any pre-college upper or lower extremity orthopaedic surgery, were elicited through pre-participation evaluations (PPEs). Athletes who underwent an orthopaedic surgery in college were identified through the Sports Injury Monitoring System and medical records. All patient-related independent variables were included in a multiple Cox regression model. Exposure time was calculated from the date of PPE to the date of surgery (event) or to the end of the collegiate athletic career (censored). Hazard ratios (HRs) with 95% confidence intervals (CIs) were calculated. Significance was set at $P < 0.05$.

Results: In total, 1,142 athletes in 12 sports (baseball/softball, basketball, football, golf, gymnastics, rowing, swimming & diving, soccer, tennis, track & field/cross country, volleyball, water polo) were identified. There were 262 documented orthopaedic surgeries, including those involving the shoulder ($n = 34$), hip ($n = 25$), and knee ($n = 72$), in 182 athletes. Using the multiple Cox regression model, pre-college lower extremity surgery was an independent predictor of orthopaedic ($P = 0.004$, HR = 1.88) and knee ($P < 0.001$, HR = 3.91) surgery, and type of sport was an independent predictor of orthopaedic ($P < 0.001$), shoulder ($P = 0.002$), and knee surgery ($P < 0.001$) (Table 1). Participation in gymnastics, basketball, and volleyball were the strongest predictors of orthopaedic surgery. Similarly, participation in volleyball, gymnastics, and baseball/softball were the strongest predictors of shoulder surgery, and participation in basketball, football, and volleyball were the strongest predictors of knee surgery. Lastly, athletes with a history of a pre-college orthopaedic surgery were more susceptible to another surgery in the same extremity during college ($P = 0.046$, HR = 2.18). Gender was not a significant predictor of any surgery. No independent predictors of hip surgery were identified.

Conclusion: NCAA athletes who underwent a pre-college lower extremity surgery were more likely to undergo orthopaedic and knee surgery during their collegiate careers. Those in overhead sports (e.g., volleyball, baseball/softball) were more likely to undergo shoulder surgery, and those in cutting and jumping sports (e.g., basketball, football) were more likely to undergo knee surgery. Furthermore, athletes with a history of a pre-college orthopaedic surgery were more likely to undergo another surgery in the same extremity during college, suggesting inadequate rehabilitation or less than full return of function after surgery. The time lost from athletic participation due to an orthopaedic surgery and its potential career impact underscores the need for injury prevention and improved surgical outcomes in the amateur athlete.

Table 1. Multiple Cox Regression Analysis of Player-Related Predictors of Orthopaedic Surgery in NCAA Athletes

Variable	HR	95% CI	P value
Gender (males/females)	0.95	0.65-1.40	0.807
Pre-college upper extremity surgery ¹	1.15	0.65-1.88	0.608
Pre-college lower extremity surgery ²	1.88	1.24-2.78	0.004
Sport			
Baseball / Softball	3.82	0.74-69.89	0.123
Basketball	14.84	3.05-267.38	<0.001
Football	10.33	2.23-183.56	0.001
Golf ³	1.00		
Gymnastics	15.38	3.03-280.58	<0.001
Rowing	3.37	0.59-63.45	0.184
Swimming & Diving	1.90	0.24-38.63	0.560
Soccer	8.29	1.75-148.30	0.003
Tennis	3.21	0.47-62.91	0.239
Track & Field / Cross Country	3.58	0.75-64.25	0.125
Volleyball	10.69	2.23-191.75	0.001
Water Polo	3.79	0.72-69.69	0.130
Surgery on previously operated extremity ⁴	2.18	1.01-4.79	0.046

HR, hazard ratio; CI, confidence interval.

¹Reference group = no pre-college upper extremity surgery

²Reference group = no pre-college lower extremity surgery

³Reference group in analysis

⁴Reference group = surgery on other extremities

The Orthopaedic Journal of Sports Medicine, 3(7)(suppl 2)

DOI: 10.1177/2325967115S00096

©The Author(s) 2015