

Research Article/Artículo de investigación

EFFECT OF DIFFERENT CONCENTRATIONS OF GLYCEROL IN CRYOPRESERVATION OF GADDI GOAT SEMEN

EFFECTO DE DIFERENTES CONCENTRACIONES DE GLICEROL EN LA CRIOPRESERVACIÓN DE SEMEN DE MACHOS CABRÍOS GADDI

AMIT SHARMA¹, PANKAJ SOOD¹, JITENDRA KUMAR CHAUDHARY²

¹CSK Himachal Pradesh Krishi Vishvavidyalya, Palampur-Himachal Pradesh 176062, India.

²Central Agricultural University, Aizwal, Mizorum, India.

Correspondence should be addressed to (La correspondencia debe dirigirse a): A. Sharma, vet05amy@gmail.com

ABSTRACT

Goats have greater importance under small familiar systems in India. The unabated decline in Gaddi males of superior genetic merit due to extensive culling and unplanned breeding (inbreeding) accentuates the importance of conserving superior males and their use for artificial insemination. The available literature is unequivocal about the importance of glycerol during sperm cryopreservation. However, the optimal level seems to vary between species and breeds. The present study aimed to evaluate the effect of different concentrations of glycerol (6, 7, and 8%) on the quality of thawed semen from Gaddi bucks. A total of 180 ejaculates from six Gaddi buck were frozen in Tris citrate extender containing 10% of Egg Yolk with 6, 7, or 8% of glycerol. Sperm quality parameters in thawed semen (morphological abnormalities, sperm viability, progressive motility, and HOST response) were compared; in addition, the percentage of change in thawed semen with respect to fresh was determined. No effect of concentration of glycerol was observed on progressive motility and HOST response. Sperm viability was better in semen frozen at 6 and 8% ($45.26 \pm 1.32\%$ and $45.10 \pm 2.81\%$ respectively) in comparison with 7% of glycerol ($34.81 \pm 2.78\%$, $P < 0.05$). Lower sperm morphological abnormalities were observed in semen frozen at 6% of glycerol (7.93 ± 0.28 , $P < 0.05$) in comparison with 7% ($9.18 \pm 0.69\%$) and 8% ($9.90 \pm 0.55\%$). A fertility rate of 41.25% was achieved following AI with semen frozen containing 6% of glycerol. In conclusions, 6% of glycerol was a valid option to cryopreservation of semen from Gaddi buck, resulting in better viability and lower abnormalities. In addition, a good fertility response was observed.

Keywords: Gaddi goats, glycerol, extender, semen, cryopreservation, fertility.

JOURNAL OF VETERINARY ANDROLOGY (2020) 5(1):01-06

RESUMEN

Las cabras están teniendo mayor importancia en los pequeños sistemas familiares en India. Sin embargo, la disminución deliberada de machos Gaddi de alto merito genético debido a la eliminación excesiva y a los cruzamientos no planificados (consanguinidad) acentúan la importancia de conservar machos superiores y su uso para inseminación artificial. La literatura disponible ha confirmado la importancia del glicerol durante la criopreservación espermática. Sin embargo, el nivel óptimo parece variar entre especies y razas. El propósito de este estudio fue evaluar el efecto de diferentes concentraciones de glicerol (6, 7 y 8%) sobre la calidad del semen descongelado de machos de raza Gaddi. Un total de 180 eyaculados de seis machos fueron congelado en diluyente a base de Tris citrato conteniendo 10% de yema de huevo con 6, 7 u 8% de glicerol. Los parámetros de calidad espermática en el semen descongelado (motilidad progresiva, viabilidad, anomalías morfológicas y respuesta a HOST) fueron comparadas, además el porcentaje de cambio de cada parámetro con respecto al semen fresco fue determinado. No se observó un efecto del nivel de glicerol sobre la motilidad progresiva y la respuesta al HOST. La viabilidad espermática fue mejor en el semen congelado con 6 y 8% de glicerol ($45,26 \pm 1,32\%$ y $45,10 \pm 2,81\%$ respectivamente) en comparación con el semen congelado con 7% de glicerol ($34,81 \pm 2,78\%$, $P < 0,05$). Menos morfoanomalías fueron observadas en el semen congelado con 6% de glicerol ($7,93 \pm 0,28$, $P < 0,05$) en comparación con el congelado con 7% ($9,18 \pm 0,69\%$) y 8% ($9,90 \pm 0,55\%$). Una tasa de 41,25% de fertilidad fue alcanzada luego de la inseminación con semen congelado con 6% de glicerol. En conclusión, 6% de glicerol es una opción válida para la criopreservación de semen de machos cabríos Gaddi, resultando en una mayor viabilidad y menos morfoanomalías espermáticas. Además, una buena fertilidad fue observada.

Palabras clave: cabras Gaddi, glicerol, diluyente, semen, criopreservación, fertilidad.

JOURNAL OF VETERINARY ANDROLOGY (2020) 5(1):01-06

Received/Recibido: 01/05/2019; Accepted/Aceptado: 19/01/2020

Copyright: © 2020 Sharma et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited / Este es un artículo de acceso abierto distribuido bajo los términos de la [Licencia Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/), la cual permite el uso irrestricto, la distribución y reproducción en cualquier medio, dando el crédito correspondiente al autor y la fuente original.



INTRODUCTION

Goats are assuming greater importance under the small household system in India. Accordingly, the Gaddi breed of goats is economically important for a large portion of nomadic and hilly tribes of Himachal Pradesh, India (Dogra & Thakur, 2010). A progressively declining cattle population accentuates goat rearing under a small household system. Sperm sensitivity to cryopreservation protocols varies among different breeds as well as the animals within the same breed and has been incriminated to variation in components of sperm plasma membranes (Bailey et al., 2000). In general, the sperms of small ruminants are more cryosensitive than other species (Purdy, 2006).

To determine ideal extender, different membrane-permeable cryoprotectants (Glycerol, Dimethyl Sulfoxide, Ethylene Glycol, and Propylene Glycol) and their combinations have been tested at different concentrations with buck semen (Leboeuf et al., 2000; Purdy, 2006; Gangwar et al., 2016; Rasad et al., 2017). Glycerol, however, remains the most frequently used penetrating cryoprotectant. Limited literature is available using variable glycerol concentration for semen cryopreservation in goats. Due to variability in reports regarding best glycerol concentration in extender, a varying number of ejaculates in the present study were utilized to establish the most promising glycerol concentration using 6, 7, or 8% concentration *per se* for Gaddi goat semen cryopreservation and its further usage for artificial insemination.

MATERIALS AND METHODS

Selection of animals and screening for diseases

The study was carried out on apparently healthy Gaddi bucks ($n=11$; aged 2.16 ± 0.36 years, weighing 39.1 ± 2.82 kg). These bucks were selected based on breeding history, breeding soundness evaluation, and testicular diameters. All the bucks were maintained under identical conditions and were screened for diseases, Brucellosis (RBPT, OIE guidelines, 2008), Chlamydiosis (AGPT, Chahota et al., 2015) to eliminate the possible transmission of infection.

Location of animals and period of investigation

The study was conducted at University Livestock Farm of CSK Himachal Pradesh Krishi Vishvavidyalaya, Palampur (32.6°N , 76.3°E , altitude 1290.8m) from September-December 2016, January and March 2017 and September-December 2017, respectively. Average light:dark hours (h) and temperature ($^{\circ}\text{C}$) during the period of the study comprised the average of the values considered on the day of semen collection and the preceding days. The average light:dark hours (h) and temperature ($^{\circ}\text{C}$) ranged from 7.07 ± 0.41 : 16.93 ± 0.41 to 12.29 ± 0.13 : 11.71 ± 0.13 and $9.88\pm 0.51^{\circ}\text{C}$ to $21.94\pm 0.18^{\circ}\text{C}$, respectively.

Animal housing and feeding

All bucks were subjected to grazing for five hours and remained under confinement in a shed where they were fed as per the standards of the Indian Council of Agricultural Research (ICAR, 2013). All males had rounded the clock access to the clean drinking water.

Semen collection and evaluation

A total of 180 ejaculates from eleven healthy adult bucks were collected using an estrus doe. The collections were made twice weekly with an artificial vagina (AV) maintained at $42\text{--}43^{\circ}\text{C}$. The fresh ejaculates were evaluated for any gross abnormality, including color, as well as volume and concentration (Caprine Photometer, IMV 1409[®]) followed by microscopic examination for mass motility. The suitability of ejaculates for further processing was determined according to Rather et al., 2016. Absence of any gross abnormality, milky to creamy semen color, mass motility of ≥ 3 , and initial progressive motility of $\geq 70\%$ were the criteria for selection of semen. The post-thaw cryopreserved semen samples were evaluated for progressive motility (Hafez and Hafez, 2000), viability (Hannock, 1951), morphological abnormalities (Bloom, 1977), and HOST reactivity (Pant et al., 2002). Percent change in the parameters described above between fresh and post-thaw semen was also calculated to determine the effect of cryopreservation.

Seminal plasma removal and extension

The seminal plasma was removed as described previously (Nutti, 2007; Sharma et al., 2018). The semen pellet thus obtained was extended with two equal fractions of tris citrate egg yolk extender, TEY (TRIS 1.21 gr, Citric acid 0.685 gr, D-Fructose 0.5 gr, Benzyl Penicillin 1000 IU/ml, Streptomycin Sulphate 1mg/ml), EY (10%) with variable glycerol concentrations (6, 7 or 8%). The TEY was added to semen at a gap of 2-3 minutes to yield a final concentration of 150×10^6 spermatozoa/straw. The pH of the buffer was adjusted to 6.7 to 6.9.

Filling and sealing of straws

Extended semen was filled in 0.25 ml French mini straws (IMV Technologies, L'Aigle, Cedex, France), by aspiration using a micropipette (Minitube, Germany) and subsequently sealed at free end with the help of polyvinyl alcohol (PVA) (IMV Technologies, L'Aigle, Cedex, France) powder. All the steps of semen handling were undertaken at a controlled air temperature of 30°C.

Equilibration, vapor exposure, freezing, and thawing

The straws were laid on a stainless-steel rack and placed in cooling cabinet 4°C (Macro Scientific Works Pvt. Ltd. India) for four hours and after that exposed to liquid nitrogen vapors for 7 minutes. After that, the straws were plunged into liquid nitrogen for storage. The inventory of semen storage was also maintained. Thawing of semen straws was done at 37°C for 30 s (Sariozkan et al., 2010) in a water bath.

Artificial Insemination

It was decided to perform AI using semen found to be the best among the investigated glycerol concentrations. A total of 80 Gaddi goats were synchronized in two lots using Cloprostenol sodium @187.5µg (Pragma®; Intas Pharmaceuticals Ltd., India) IM. Sixty hours later, intra-cervical insemination was carried out using two 0.25 ml straws, each containing 150×10^6 spermatozoa.

Ethical regulations

Proper ethical considerations related to animal handling and semen collections were observed and ensuring not to cause any injury during sampling.

Statistical analysis

The data obtained were analyzed using package R version 3.4.3. A paired sample t-test was used to determine significant differences in the ejaculates processed in varying glycerol concentrations. Pregnancy rates for different bucks were analyzed using the Chi-square test. A difference of $P < 0.05$ or less was significant.

RESULTS

A perusal of the results (Table 1) revealed significantly lower morphological abnormalities ($7.93 \pm 0.28\%$) along with minimal change (20.15) in the said parameter for the semen processed in 6% glycerol. Also, the sperm viability for 6% ($45.26 \pm 1.32\%$) and 8% of glycerol ($45.10 \pm 2.81\%$) were significantly higher than the samples processed in 7% of glycerol ($34.81 \pm 2.78\%$). The numerical values for average progressive motility were also higher in the samples for 6% of glycerol ($35.18 \pm 0.87\%$). There was no difference in HOST reactive sperms across different glycerol concentrations.

In the present study, 1.12 kids per doe (thirty-seven kids: males, 40.54%; females, 59.46%) were delivered. Buck no 3509 yielded best conception rate (53.84%), whereas buck no 3076 gave the lower (30.76%), with an overall conception rate of 41.25% (Table 2) following AI with frozen-thawed semen utilizing TEY extender containing 10% of egg yolk and 6% of glycerol.

DISCUSSION

Breed specificity to varying glycerol concentrations in goats, and accordingly, the quality of cryopreserved semen is on record (Purdy, 2006; Kulaksiz et al., 2013; Gangwar et al., 2016). Our findings using different glycerol concentrations, with 6% yielding the best results, corroborates to a recent observation (Rasad et al., 2017) in which 6% glycerol was better than 5, 7, 8 and 9% for goat (Eta wah crossbred) semen cryopreservation. Similarly, 6% of glycerol was better than 4 and 8% of glycerol in Sirohi bucks (Sikarwar et al., 2015), as revealed by significantly higher motility and viability. Biswas et al. (2002) used 5, 7, and 10% of glycerol and found better motility and viability of frozen-thawed sperm in 7% than other glycerol concentrations. Similarly, Farshad et al. (2009) recorded better post-thaw semen quality with 5 and 7% glycerol than other glycerol concentrations (1 and 3 %) in Markhoz goat semen.

Table 1: Average (Mean±SEM) of post-thaw semen quality parameters extended with different concentration of Glycerol in Gaddi bucks.

Parameters (%)	Percentage of Glycerol		
	6% (n=106) (% change due to processing)	7% (n=37) (% change due to processing)	8% (n=37) (% change due to processing)
Progressive motility	35.18±0.87 ^A (52.03)	29.00±1.79 ^A (58.99)	34.70±3.03 ^A (51.63)
Viability	45.26±1.32 ^A (40.12)	34.81±2.78 ^B (53.30)	45.10±2.81 ^A (37.96)
Morphological abnormalities	7.93±0.28 ^B (20.15)	9.18±0.69 ^A (21.75)	9.90±0.55 ^A (30.26)
HOST reactive	52.48±1.43 ^A (26.26)	53.90±1.47 ^A (28.47)	53.40±1.50 ^A (30.01)

^{A-B} Values with different superscripts within same row differs ($P < 0.05$).

n=Number of ejaculates.

Table 2: Individual buck fertility.

Buck No	Does inseminated (Number)	Does pregnant (Number)	Fertility rate (%)
3067	14	5	35.71
3076	13	4	30.76
1201	15	7	46.60
3509	13	7	53.84
3002	12	5	41.60
3075	13	5	38.46
Overall	80	33	41.25

Chi-square value = 1.842; P value = 0.87

Kulaksiz et al. (2013) used variable concentrations of glycerol (5-10%) in different breeds of goats and concluded that 5% of glycerol is best for Angora, 5-9% for Kilis and 7% for Saanen breeds, respectively. Contrastingly, Deka & Rao (1986) showed that the different concentrations (i.e., 4, 6.4, and 9%) of glycerol did not differ in post-thaw sperm motility and morphological abnormalities in goat semen. Similar observations were recorded by Biswas et al. (2002) in Black Bengal goats.

Sperm membrane lipid composition might be a significant factor responsible for variability in susceptibility of spermatozoa to damage produced during the cryopreservation in different breeds of goats (Holt, 2000). Membrane destabilization occurs when the sperm plasma membrane undergoes a phase transition from the liquid crystalline phase to the gel phase due to a decrease in temperature, which could affect the semen during processing (Barrea-Compean et al., 2005). Glycerol remains to be the most effective cryoprotective compound for freezing goat semen, and no enhancement was shown by the addition of other compounds (Farshad et al., 2009). Ranjan et al. (2015) demonstrated the usage of a combination of egg yolk (10%) and glycerol (6%) to significantly ($P < 0.05$) improve the percentage of intact acrosome in Jamunapari buck semen.

Rizal et al. (2013) reported the toxic effects of a higher concentration of glycerol. Similar observations about the harmful effects of higher concentrations of glycerol (9%; Etawah goats, Rasad et al., 2017) and (7%; Sonmez & Demira, 2004) in ram semen. Glycerol in higher concentrations can cause osmotic damage to spermatozoa because glycerol passes through the sperm membrane much slower than other cryoprotectants (Guthrie et al., 2002). The toxicity of glycerol in semen extender results in a reduction in sperm motility and alterations of the acrosome integrity by interfering with the permeability of the sperm membrane, which intern results in lower fertility rates when such semen is used for artificial insemination (Salmon & Maxwell, 2000).

Higher kids per doe have been observed by Gacitua and Arav, 2005 (Saanen goats; 1.78), Kharche et al., 2013 (Jamunapari goats; 1.64). Variable fertility rates (38 to 65 %) with the use of frozen-thawed semen, have been earlier observed in Indian and exotic breeds (Gacitua & Arav, 2005; Kharche et al., 2013). The reasons per se could be due to differences in freezability, a result of chilling injury (Drobnis et al., 1993) and fertilizing capacity of semen.

CONCLUSIONS

Extender containing 6% of glycerol was best concerning least morphological abnormalities and better progressive motility, viability for Gaddi goat semen cryopreservation. These results, together with the fertility of 41.25% achieved, indicate that glycerol at 6% represents an excellent alternative to preserve the quality of frozen-thawed semen from Gaddi bucks.

REFERENCES

- Bailey JL., Bilaodeau JF., Cormier N. 2000. Semen cryopreservation in domestic animals: a damaging capacitating phenomenon. **Journal of Andrology** 20: 1-7.
- Barrea-Compean MH., Purdy PH., Dzakurna JM., Newton GR., Nuti LC. 2005. Cholesterol-loaded cyclodextrin improves post-thaw goat sperm motility. **Journal of Animal Sciences** 83(1): 153-153.
- Biswas D., Bari FY., Shamsuddin M., Rahman MM. 2002. Determination of glycerol percentages for preserving the Black Bengal Buck (*Capra hircus*) spermatozoa for long time. Pak. **Journal of Biological Sciences** 5: 715-718.
- Bloom E. 1977. Sperm morphology with reference to bull infertility. In: Some papers contributed to the 1st All India Symposium on Animal Reproduction, Punjab Agricultural University, Ludhiana, India, pp. 66-8.
- Chahota R., Gupta S., Bhardwaj B., Malik P., Verma S., Sharma M. 2015. Seroprevalence studies on animal Chlamydia amongst ruminants I five states of India. **Veterinary World** 8(1): 72-75.
- Deka BC., Rao AR. 1986. Effect of Glycerol Level In Tris based extender and equilibration period on quality of frozen goat semen. **Theriogenology** 2: 231-237.
- Dogra PK., Thakur YP. 2010. Livestock biodiversity of Himachal Pradesh with special reference to pashmina goats. In: Proceeding of National training on "Production, processing and utilization of pashmina fibre" 1-7 Dec Palampur, H.P, India pp 1-12.
- Drobnis ES., Crowe LM., Berger T., Anchodoguy TJ., Overstreet JW., Crowe J.H. 1993. Cold shock damage is due to lipid phase transition in cell membranes: a demonstration using sperm as a model. **Journal of Experimental Zoology** 265(4): 432-437.
- Farshad A., Khalili B., Fazeli P. 2009. The effect of different concentrations of Glycerol and DMSO on the viability of Markhoz Goat spermatozoa during different freezing temperatures steps. **Pakistan Journal of Biological Sciences** 12: 239-245.
- Gacitua H., Arav A. 2005. Successful pregnancies with directional freezing of large volume buck semen. **Theriogenology** 63: 931-938.
- Gangwar C., Kharche SD., Kumar S., Jindal SK. 2016. Cryopreservation of goat semen: status and prospects. **Indian Journal of Small Ruminants** 22: 1-10.
- Guthrie HD., Liu J., Critser JK. 2002. Osmotic tolerance limits and effects of cryoprotectants on motility of bovine spermatozoa. **Biology of Reproduction** 67: 1811-1816.
- Hafez B., Hafez ESE. 2000. **Reproduction in Farm Animals**. 7th ed. Lippincott Williams and Wilkens, New York. pp. 509.
- Hannock JL. 1951. A staining technique for study of temperature shock in semen. **Nature** 167, 323-324.
- Holt WV. 2000. Basic aspects of frozen storage of semen. **Animal Reproduction Science** 62: 3-22.
- ICAR. 2013. **Nutrient Requirements of Sheep, Goats and Rabbits**. 3rd ed. ICAR, New Delhi.
- Kharche SD., Jindal SK., Priyadarshini R., Kumar S., Goel AK., Ramachandran N., Rout PK. 2013. Fertility following frozen semen artificial insemination in Jamunapari goats. **Indian Journal of Animal Sciences** 83(10): 1071-1073.
- Kulaksiz R., Ari UÇ., Daskin A., Üner AG. 2013. The effect of different Glycerol concentrations on freezability of semen from Angora, Kilis and Saanen goats. **Slovakian Journal of Animal Sciences** 46: 39-44.
- Leboeuf B., Restall B., Salamon S. 2000. Production and storage of goat semen for artificial insemination. **Animal Reproduction Science** 62:113-141.
- Nuti I. 2007. Techniques of artificial insemination in goats. In: Youngquist, R.S. and Threlfall, W.R (eds) **Current Therapy in Large Animal Theriogenology**. 2nd ed. Saunders-Elsevier, St. Louis, MO. pp. 529-534.
- OIE 2008. **Quality Standards and Guidelines for Veterinary Laboratories**, 2nd ed., OIE, Paris.
- Pant HC., Barot LR., Kasiraj R., Misra AK., Prabhakar JH. 2002. Acrosomal integrity of buffalo spermatozoa and its relation to fertility. **Indian Journal of Animal Reproduction** 23: 110-112.

Purdy PH. 2006. A review on goat sperm cryopreservation. **Small Ruminant Research** 63: 215-225.

R Core Team 2013. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Ranjan R., Goel AK., Ramchandran N., Kharche SD., Jindal SK. 2015. Effect of egg yolk levels and equilibration periods on freezability of Jamunapari buck semen. **Indian Journal of Small Ruminant** 21(1): 32-36.

Rasad SD., Soeparna, Nurcholiah, Setiawan R., Widyastuti R. 2017. Effect of Glycerol level in two different extenders on post thawed sperm quality of crossbred Etawah goat. **Journal of Animal & Veterinary Advances** 16(8-12): 87-91.

Rather HA., Islam R., Malik AA., Lone FA. 2016. Addition of antioxidants improves quality of ram spermatozoa during preservation at 4°C. **Small Ruminant Research** 141: 24-28.

Rizal M., Toeliher, MR., Yusuf TL., Purwantara B., Situmorang P. 2013. Quality of Garut frozen semen in various concentrations of Glycerol. **Indonesian Journal of Animal & Veterinary Sciences** 7: 194-199.

Salamon S., Maxwell WM. 2000. Storage of ram semen. **Animal Reproduction Science** 62: 77-111.

Sariözkan S., Bucak MN., Tuncer PB., Taşdemir U., Kinet H., Ulutaş PA. 2010. Effects of different extenders and centrifugation/washing on post thaw microscopic-oxidative stress parameters and fertilizing ability of Angora buck sperm. **Theriogenology** 73, 316-323.

Sharma A., Sood P., Dogra P. 2018. Seminal plasma removal improves cryopreserved semen quality in Gaddi bucks. **Indian Journal of Animal Reproduction** 39(2): 25-28.

Sikarwar AKS., Ramachandra N., Ranjan R., Gangwar C., Aggarwal JK., 2015. Effect of different levels of glycerol and MSO on freezability of buck semen. **Livestock Research International** 3: 71-73.

Sonmez M., Demira E. 2004. The effect of ascorbic acid on the freezability of ram semen diluted with extenders containing different proportions of Glycerol. **Turkish Journal of Veterinary & Animal Sciences** 28: 893- 899.