



タイトル Title	Identification of novel serum markers for the progression of coronary atherosclerosis in WHHLMI rabbits, an animal model of familial hypercholesterolemia
著者 Author(s)	Shiomi, Masashi / Takeda, Hiroaki / Irino, Yasuhiro / Kimura, Norie / Yamada, Satoshi / Kuniyoshi, Nobue / Kikumori, Akio / Koike, Yu / Koike, Tomonari / Yoshida, Masaru / Izumi, Yoshihiro / Shinohara, Masakazu / Bamba, Takeshi / Ishida, Tatsuro
掲載誌・巻号・ページ Citation	Atherosclerosis,284:18-23
刊行日 Issue date	2019-05
資源タイプ Resource Type	Journal Article / 学術雑誌論文
版区分 Resource Version	author
権利 Rights	© 2019 Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
DOI	10.1016/j.atherosclerosis.2019.02.020
JaLDOI	
URL	http://www.lib.kobe-u.ac.jp/handle_kernel/90005706

1 Identification of novel serum markers for the progression of coronary atherosclerosis in
2 WHHLMI rabbits, an animal model of familial hypercholesterolemia

3

4 Masashi Shiomi^{a,b}, Hiroaki Takeda^c, Yasuhiro Irino^d, Norie Kimura^b, Satoshi Yamada^a,
5 Nobue Kuniyoshi^a, Akio Kikumori^a, Yu Ying^e, Tomonari Koike^e, Masaru Yoshida^f, Yoshihiro
6 Izumi^c, Masakazu Shinohara^g, Takeshi Bamba^c, Tatsuro Ishida^h

7

8 ^a Institute for Experimental Animals, Kobe University Graduate School of Medicine, Kobe,
9 Japan

10 ^b Division of Comparative Pathophysiology, Department of Physiology and Cell Biology, Kobe
11 University Graduate School of Medicine, Kobe, Japan

12 ^c Division of Metabolomics, Research Center for Transomics Medicine, Medical Institute of
13 Bioregulation, Kyushu University, Fukuoka, Japan

14 ^d Division of Evidence-based Laboratory Medicine, Department of Internal Related, Kobe
15 University Graduate School of Medicine, Kobe, Japan

16 ^e University of Michigan Medical Center, Ann Arbor, MI, USA

17 ^f Division of Metabolomics Research, Department of Internal Medicine, Kobe University
18 Graduate School of Medicine, Kobe, Japan

19 ^g Division of Epidemiology, Department of Community Medicine and Social Healthcare
20 Science, Kobe University Graduate School of Medicine, Kobe, Japan

21 ^h Cardiovascular Medicine, Department of Internal Medicine, Kobe University Graduate
22 School of Medicine, Kobe, Japan

23

24 * Corresponding author: Masashi Shiomi, Institute for Experimental Animals, Kobe
25 University Graduate School of Medicine, 7-5-1 Kusunoki-cho, Chuo-ku, Kobe 650-0017,
26 Japan. (ieakusm@med.kobe-u.ac.jp)

1 **ABSTRACT**

2 *Background and aims:* The development of serum markers specific for coronary lesions is
3 important to prevent coronary events. However, analyses of serum markers in humans are
4 affected by environmental factors and non-target diseases. Using an appropriate model
5 animal can reduce these effects. To identify specific markers for coronary atherosclerosis, we
6 comprehensively analyzed the serum of WHHLMi rabbits, which spontaneously develop
7 coronary atherosclerosis.

8 *Methods:* Female WHHLMi rabbits were fed standard chow. Serum and plasma were
9 collected under fasting at intervals of 4 months from 4 months old, and a total of 313 lipid
10 molecules, 59 metabolites, lipoprotein lipid levels, and various plasma biochemical
11 parameters were analyzed. The severity of coronary lesions was evaluated with cross-
12 sectional narrowing (CSN) corrected with a frequency of 75% - 89% CSN and CSN > 90%.

13 *Results:* There was a large variation in the severity of coronary lesions in WHHLMi rabbits
14 despite almost no differences in plasma biochemical parameters and aortic lesion area
15 between rabbits with severe and mild coronary lesions. The metabolites and lipid molecules
16 selected as serum markers for coronary atherosclerosis were lysophosphatidylcholine (LPC)
17 22:4 and diacylglycerol 18:0-18:0 at 4 months old, LPC 20:4 (sn-2), ceramide d18:1-18:2, citric
18 acid plus isocitric acid, and pyroglutamic acid at 8 months old, and
19 phosphatidylethanolamine plasminogen 16:1p-22:2 at 16 months old.

20 *Conclusions:* These serum markers were coronary lesion-specific markers independent of
21 cholesterol levels and aortic lesions and may be useful to detect patients who develop
22 cardiovascular disease.

23

24

25 **Key words**

26 coronary atherosclerosis, lipidome analysis, metabolome analysis, serum marker, WHHLMi
27 rabbit

1 1. Introduction

2 The discovery of serum markers specific for coronary lesions is important to prevent
3 coronary events. In epidemiological studies, several marker candidates for coronary artery
4 disease (CAD), such as the levels of serum cholesterol, low-density lipoprotein (LDL)
5 cholesterol, high-density lipoprotein (HDL) cholesterol, the ratio of LDL cholesterol to HDL
6 cholesterol, lipoprotein (a), small dense LDL, and oxidized LDL, have been reported. These
7 conventional factors are not always associated with the onset of CAD [1-3], and metabolome
8 and lipidome analyses have revealed other candidates for serum markers of CAD or
9 atherosclerosis in animal models [1,4-8] and humans [9-14]. Because serum levels of
10 metabolites or lipid molecules are regulated by gene expression and environmental stresses,
11 these metabolites or lipid molecules reflect the development and progression of the disease
12 [15]. In clinical or epidemiological studies, several factors, such as dietary habits,
13 environmental conditions, non-target diseases, social stress, and alcohol, affect analysis of
14 markers related to CAD [16-19]. In studies using animals, these confounding factors can be
15 kept constant, and age-related changes can be investigated in candidate markers. Although
16 animals fed a cholesterol diet or a high fat diet [6-8] or raised in a smoke-filled environment
17 [1,4] develop atherosclerotic lesions, these conditions directly or indirectly affect the serum
18 concentration of metabolites or lipid molecules. Therefore, animal models that develop
19 atherosclerosis spontaneously without the influence of environmental factors that regulate
20 the concentration of serum parameters are better alternative animal models. In previous
21 studies using these animals, candidate marker substances were analyzed for aortic
22 atherosclerosis. These studies were performed under the assumption that there are no
23 differences between aortic lesions and coronary lesions. Because the volume of aortic lesions
24 is extremely large compared with that of coronary lesions, serum concentrations of marker
25 substances for aortic lesions may mask those of coronary lesions. Therefore, markers that are
26 specific for coronary lesions are unknown. Previous studies using WHHLMI rabbits, a
27 spontaneous animal model for hypercholesterolemia from LDL receptor deficiency, coronary
28 atherosclerosis, and myocardial lesions [20], suggest that factors associated with coronary

1 lesions differ from those of aortic lesions [21,22]. In this study, we identified serum markers
2 specific for coronary lesions in WHHLMI rabbits using lipidome and metabolome analyses.

3

4 **2. Materials and Methods**

5 *2.1. Animals*

6 We examined serum markers for coronary lesions using 22 WHHLMI rabbits (12 females
7 and 10 males), and 363 WHHLMI rabbits born between 2010 and 2016 were examined for
8 age-dependent changes in the progression of coronary lesions. In addition, 6 male JW rabbits
9 were used as normal controls. The details are described in the Supplementary Methods.

10

11 *2.2. Fractionation of plasma lipoproteins and measurement of lipid concentration*

12 Lipoproteins were fractionated with an ultracentrifuge to yield very low-density
13 lipoproteins (VLDL, $d < 1.006$ g/ml), LDL ($1.006 < d < 1.063$ g/ml), and HDL ($d > 1.063$ g/ml). The
14 details are described in the Supplementary Methods.

15

16 *2.3. Metabolome analyses of serum*

17 Metabolomic profiling using gas chromatography/mass spectrometry (GC/MS) was
18 performed as reported previously [23]. Fifty-nine metabolites were recorded. The details are
19 described in the Supplementary Methods.

20

21 *2.4. Lipidome analyses of serum*

22 Lipids were quantified using supercritical fluid chromatography triple quadrupole mass
23 spectrometry (SFC/MS/MS) in the multiple reaction monitoring mode [24]. There were 313
24 lipid molecules detected in the serum. The relative concentration (RC) of lipid molecules was
25 corrected with serum total cholesterol concentrations using the following equation:

26
$$[\text{Corrected relative concentration (CRC)}] = [\text{relative concentration}] / \text{serum total}$$

27 cholesterol concentration x 1,000.

28 The details are described in the Supplementary Methods.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

2.5. Evaluation of atherosclerotic lesions

Aortic lesions were evaluated as the ratio of the surface area of the lesions to the surface area of the entire lumen. The severity of coronary lesions was evaluated using a coronary severity score based on the following equation:

$$[\text{Coronary severity score}] = [\text{average cross-sectional narrowing (CSN)}] \times (1 + [\text{frequency of sections with 75\%-89\% CSN}] \times 10 + [\text{frequency of sections with >90\% CSN}] \times 20)$$

The details are described in the Supplementary Methods.

2.6. Age-dependent progression of coronary lesions

To examine age-dependent progression of coronary lesions in WHHLMI rabbits, we examined the maximum CSN from one month old to 30 months old using 363 dead rabbits or rabbits euthanized in other experiments between 2010 and 2016. Coronary lesions were examined in the manner described above. The details are described in the Supplementary Methods.

2.7. Other assays

At the start of the experiment (4 months old) and at the end of experiment (20 months old or at death), we assayed the plasma levels of glucose, insulin, NO₂, NO₃, and oxidized LDL, and superoxide dismutase (SOD) activity. The details are described in the Supplementary Methods.

2.8. Data and statistical analyses

We used data only from females because of sex differences in coronary lesions and serum levels of metabolites and lipid molecules. The serum marker candidates for coronary atherosclerosis were determined in two steps at 4 and 8 months old, and in 3 steps at over 8 months old, and at 20 months old or death. The details are shown in the supplementary

1 information.

2

3 **3. Results**

4 *3.1. Atherosclerotic lesions of WHHLMI rabbits*

5 Table 1 shows that the parameters of coronary lesions were significantly higher in the
6 severe group than in the mild group, but there were no differences in aortic surface lesion
7 area ($96\pm 2\%$ vs $94\pm 1\%$). These results suggest that the factors related to the progression of
8 atherosclerotic lesions were different between aorta and coronary arteries.

9

10 *3.2. Plasma biochemical parameters in WHHLMI rabbits*

11 There were no significant differences in lipoprotein lipid levels and plasma biochemical
12 parameters between the severe and mild groups (Supplementary Table 5 and 6). These
13 results indicate that factors other than lipoprotein lipids and other plasma parameters were
14 involved in the progression of coronary lesions.

15

16 *3.3. Examination of serum metabolites as markers specific for coronary lesions*

17 In analyses of serum metabolites, the most abundant metabolites were glycine, lactic
18 acid, and alanine (data not shown). In volcano plot analyses of differences in serum
19 metabolites between the severe and mild groups (Figure 1-A) for the first selection, 9
20 metabolites were selected at various ages. However, in the second selection, the Spearman's
21 rank-correlation coefficient at 8 months old was 0.655 ($p = 0.029$) in pyroglutamic acid and
22 0.609 ($p = 0.047$) in citric acid plus isocitric acid (Table 2). The RC of these metabolites did
23 not show a sex difference (data not shown). We selected these metabolites as markers for the
24 rapid progression of coronary atherosclerosis at an early stage. In the sparse partial least
25 square discriminant analysis (SPLS-DA) of each metabolite, the distribution of rabbits was
26 distinctly separated between the two groups, except at 4 months old (Figure 1-B). Comparing
27 normal and WHHLMI rabbits, the RC of 26 of 59 serum metabolites was significantly
28 different (Supplementary Figure 4-A). The RCs of metabolites selected as markers of

1 coronary atherosclerosis (citric acid plus isocitric acid and pyroglutamic acid) were high in
2 the WHHLMi rabbit severe group compared to those in normal rabbits, but the difference
3 was not significant (Supplementary Figure 5-A).

5 *3.4. Examination of serum lipid molecules as markers specific for coronary lesions*

6 In analyses of serum lipid molecules, the most abundant lipids were cholesterol ester
7 (CE), triacylglycerol (TAG), and phosphatidylcholine (PC) (data not shown). In volcano plot
8 analyses of differences in the CRC of serum lipid molecules between the severe and mild
9 groups for the first selection (Figure 2-A), 27 of 293 lipid molecules were selected (5 lipids at
10 4 months old, 4 lipids at 8 months old, 3 lipids at 12 months old, 12 lipids at 16 months old,
11 and 3 lipids at 20 months old or death). In the second selection by the Spearman's rank-
12 correlation coefficient (Table 3), 2 of 5 lipid molecules were selected at 4 months old [LPC
13 22:4 ($r = 0.700$, $p = 0.016$) and DAG 18:0-18:0 ($r = -0.636$, $p = 0.035$)] as markers for the
14 generation of coronary lesions. At 8 months old, 2 of 4 lipid molecules were selected [LPC
15 20:4 (sn-2) ($r = 0.773$, $p = 0.005$) and Cer d18:1-18:2 ($r = 0.673$, $p = 0.023$)] as markers for the
16 rapid progression of coronary lesions. In the third analyses for occluded coronary lesions or
17 myocardial ischemia at over 8 months old (Figure 3-A), only PE 16:1p-22:2 was selected at 16
18 months old. In SPLS-DA of serum lipid molecules (Figure 2-B), the distribution of the rabbits
19 was distinctly separated between the two groups. For sex differences, the CRC of Cer d18:1-
20 18:2 was lower in females than males, but other selected lipid molecules were not different
21 between males and females (data not shown). In comparison to normal JW rabbits, the
22 relative concentrations of many serum lipid molecules in WHHLMi rabbits were extremely
23 high (Supplementary Figure 4-B). In lipid molecules selected for markers of coronary
24 atherosclerosis, the RCs of LPC 22:4, LPC 20:4 (sn-2) and Cer d18:1-18:2 were significantly
25 high in WHHLMi rabbits compared to those in normal JW rabbits (Supplementary Figure 5-
26 B), but others were below the detection limitation in normal JW rabbits.

28 **4. Discussion**

1 We identified serum markers specific for the generation, rapid progression, and occlusion
2 of coronary lesions, which may be different from markers of aortic surface lesion area in
3 female WHHLMi rabbits. Because many factors, such as the environment, lifestyle habits,
4 feeding, social stress, non-target diseases, sex, cholesterol levels, and degree of aortic lesions,
5 that influence serum marker analysis in humans [16-19] and other experimental animals [6-
6 8] were controlled in our study, many of these previously described metabolites and lipid
7 molecules were not selected. Therefore, the serum marker candidates found in the present
8 study may reflect the development and progression of coronary lesions and myocardial
9 ischemia independent of environmental factors.

10 At 4 months old, which corresponded to when coronary lesions were generated
11 (Supplementary Figure 6), the lipids LPC22:4 and DAG18:0-18:0 were selected as markers.
12 LPC is involved in monocyte recruitment, macrophage proliferation, increased expression of
13 endothelial adhesion molecules, and endothelial dysfunction [25]. In patients with coronary
14 artery disease, the serum levels of four LPCs containing unsaturated fatty acids were higher
15 than in normal subjects, whereas the serum levels of two LPCs containing saturated fatty
16 acids were lower than in normal subjects [12]. LPC22:4 is an LPC that contains arachidonic acid,
17 which enhances inflammation [26]. Therefore, differences in the side chain of LPC may be
18 important in the generation of coronary lesions. DAG18:0-18:0 was negatively associated
19 with coronary lesions. The odds ratio of stable CAD and normal subjects was high in
20 unsaturated fatty acid-containing DAG but low in saturated fatty acid-containing DAG [27].
21 This observation indicates that the atherogenicity of DAG differs depending on side chain
22 fatty acids and suggests that DAG18:0-18:0 is not atherogenic at an early stage.

23 At 8 months old, which corresponded to when coronary lesions rapidly progressed
24 (Supplementary Figure 6), citric acid plus isocitric acid, pyroglutamic acid, LPC20:4 (sn-2),
25 and Cer d18:1-18:2 were selected as markers for the rapid progression of coronary lesions at
26 an early stage. For citric acid, Yamashita et al. [28] reported that in macrophage-rich lesions
27 in the iliac artery of fat-fed rabbits, intermediates in the TCA cycle including citric acid was
28 high compared to that of normal control arteries. Intermediates in the citric acid cycle are G-

1 protein-coupled receptor ligands and are involved in atherosclerosis [29]. Pyroglutamic acid
2 is a cyclic derivative of glutamine and is independently associated with heart failure in
3 humans [30]. In vitro studies demonstrated that pyroglutamate is associated with monocyte
4 chemoattractant protein-1 activity and expression of HUVEC adhesion factor [31]. An
5 increase in LPC20:4 (sn-2) is also observed in patients with angina pectoris and myocardial
6 infarction [12] and carotid plaques [32]. We found that LPC22:4 increased when lesions were
7 generated, and LPC20:4 (sn-2) increased when lesions rapidly progressed. These results
8 suggest that there are differences in LPC molecules that are associated with coronary lesions
9 during lesion development. Several studies also showed that several ceramide molecules
10 were related to cardiovascular disease [12,33-36]. Ceramides have several functions related
11 to atherosclerosis involving cell survival or death, the aggregation of lipoproteins,
12 transcytosis of oxidized LDL across endothelial cells, regulation of monocyte adhesion to
13 vascular walls, and promotion of LDL uptake [37]. Furthermore, knockout of sphingomyelin
14 synthase 2, an enzyme for ceramide synthesis, decreases atherosclerosis in mice [38].
15 Therefore, reduction of ceramide by medication may be effective in suppressing
16 arteriosclerosis. However, ceramide levels were lower in females (data not shown) than in
17 males similar to those in mice [39]. Analysis of the relationship between ceramide and
18 coronary artery lesions using males is needed in the future.

19 At 16 months old, which corresponds to when the average of maximum CSN was over
20 90% (Supplementary Figure 6) and the development of myocardial ischemia, the CRC of
21 PE16:1p-22:2, a phosphatidylethanolamine plasminogen species (PE-pln), was selected as a
22 marker for occluded coronary lesions or myocardial ischemia. PE-pln has anti-oxidative and
23 anti-inflammatory functions [40]. However, it is unclear how PE16:1p-22:2 is associated with
24 occluded coronary lesions or myocardial ischemia. The serum levels of several PE-pln species
25 are significantly higher in patients with angina pectoris than in normal subjects [12].
26 Because the frequency of sections with more than 90% CSN was $45 \pm 13\%$ (82%, 32%, and
27 18% of sections in three rabbits), which suggests ischemic damage of the myocardium in the
28 severe group, occlusion of coronary lesions may cause an increase in serum PE16:1p-22:2

1 similar to that of patients with angina pectoris.

2 There were several limitations in the present study. Many serum metabolites and lipid
3 molecules showed sex differences, and variations in coronary lesion severity were large in
4 females, although 7 of 9 males had severe coronary lesions. Therefore, we used only females
5 in the present study. To compensate for the decrease in the number of rabbits by limiting our
6 analysis to female rabbits, we used a rigorous selection criteria for marker candidates.
7 However, we need to analyze serum markers using males in the future. It is unclear whether
8 the serum markers we found can be used in non-FH patients. Because there is no difference
9 in the serum total cholesterol level between the mild group and the severe group, it is
10 considered that the selected markers were not affected by hypercholesterolemia. Therefore,
11 these markers may be useful for patients with cardiovascular disease without
12 hypercholesterolemia.

13 In conclusion, we found serum markers specific for the generation of coronary lesions, for
14 the rapid progression of coronary lesions at an early stage, and for occluded coronary lesions.
15 These serum markers may be useful to detect patients who develop cardiovascular disease.

16

17 **Conflict of interests**

18 The authors do not have anything to disclose or conflicts of interest with respect to this
19 manuscript.

20

21 **Financial support**

22 This study was partly supported by a Grant-in-Aid for Scientific Research from the
23 Ministry of Education, Culture, Sports and Technology, Japan (23300157 to Masashi Shiomi)
24 and the AMED-CREST Program from the Japan Agency for Medical Research and
25 Development (JPMJCR1395 and JP18gm0910013 to Yoshihiro Izumi and Takeshi Bamba).

26

27 **Author contributions**

28 The study was conceived and designed by MShio; the experiments were performed by

1 MShio, HT, YIr, NKi, SY, NKu, AK, YY, TK, and YIz; the data were analyzed by MShio, YIz,
2 MShin, TB, and TI; the manuscript was prepared by MShio, MShin, TB, and TI.

3

4 **References**

- 5 [1] S. Boué, K. Tarasov, M. Jänis, et al., Modulation of atherogenic lipidome by cigarette
6 smoke in apolipoprotein E-deficient mice, *Atherosclerosis* 225 (2012) 328-334.
- 7 [2] W.P. Castell, Lipids, risk factors and ischaemic heart disease, *Atherosclerosis* 124 (1996)
8 S1-S9.
- 9 [3] A. Sachdeva, C.P. Cannon, P.C. Deedwania, et al., Lipid levels in patients hospitalized
10 with coronary artery disease: an analysis of 136 905 hospitalizations in Get With The
11 Guidelines, *Am. Heart J.* 157 (2009) 111–117.
- 12 [4] V.T. Dang, A. Huang, H. Lexy, et al., Comprehensive plasma metabolomic analyses of
13 atherosclerotic progression reveal alterations in glycerophospholipid and sphingolipid
14 metabolism in apolipoprotein E-deficient mice, *Sci. Rep.* 6 (2016) 35037.
- 15 [5] O. Roma, C. Grajeda-Iglesias, M. Najjara, et al., Atherogenicity of amino acids in the
16 lipid-laden macrophage model system in vitro and in atherosclerotic mice: a key role for
17 triglyceride metabolism, *J. Nutr. Biochem.* 45 (2017) 24–38.
- 18 [6] A. Lazar, L. Bojic, D.G. McLaren, et al., Lipidome of atherosclerotic plaques from
19 hypercholesterolemic rabbits, *Int. J. Mol. Sci.* 15(2014) 23283-23293.
- 20 [7] M. Jové, V. Ayala, O. Ramírez-Núñez, et al., Lipidomic and metabolomic analyses reveal
21 potential plasma biomarkers of early atheromatous plaque formation in hamsters,
22 *Cardiovasc. Res.* 97 (2013) 642–652.
- 23 [8] T. Kasumov, L Li, M. Li M, et al., Ceramide as a mediator of non-alcoholic fatty liver
24 disease and associated atherosclerosis, *PlosOne* 10 (2015) e0126910.
- 25 [9] K. Tarasov, K. Ekroos, M. Suoniemi, et al., Molecular lipids identify cardiovascular risk
26 and are efficiently lowered by simvastatin and PCSK9 deficiency, *J. Clin. Endocrinol.*
27 *Metab.* 99 (2014) E45-E52.
- 28 [10] C. Stegemann, I. Drozdov, J. Shalhoub, et al., Comparative lipidomics profiling of human

- 1 atherosclerotic plaque, *Circ. Cardiovasc. Genet.* 4 (2011) 232-242.
- 2 [11] I Sutter, S. Velagapudi, A. Othman, et al., Plasmalogens of high-density lipoproteins
3 (HDL) are associated with coronary artery disease and anti-apoptotic activity of HDL,
4 *Atherosclerosis* 241(2015) 539-546.
- 5 [12] J.Y. Park, S.H. Lee, M.J. Shin, et al., Alteration in metabolic signature and lipid
6 metabolism in patients with angina pectoris and myocardial infarction, *PlosOne*
7 10(2015) e0135228.
- 8 [13] J.M. Cheng, M.Suoniemi, I. Kardys, et al., Plasma concentrations of molecular lipid
9 species in relation to coronary plaque characteristics and cardiovascular outcome:
10 Results of the ATHEROREMO-IVUS study, *Atherosclerosis* 243 (2015) 560-566.
- 11 [14] J. Yu, W. Pan, R. Shi, et al., Ceramide is upregulated and associated with mortality in
12 patients with chronic heart failure, *Can. J. Cardiol.* 31 (2015) 357-363.
- 13 [15] G.J. Patti, O. Yanes, G. Siuzdak, Metabolomics: the apogee of the omics trilogy, *Nat. Rev.*
14 *Mol. Cell Biol.* 13 (2012) 263–269.
- 15 [16] M.S. Grase, P.C. Dempsey, P. Sethi, et al., Breaking up prolonged sitting alters the
16 postprandial plasma lipidomic profile of adults with type 2 diabetes, *J. Clin. Endocrinol.*
17 *Metab.* 102 (2017) 1991-1999.
- 18 [17] G. Marin-Royo, E. Martinez-Martinez, B. Gutierrez, et al. The impact of obesity in the
19 cardiac lipidome and its consequence in the cardiac damage observed in obese rat, *Clin.*
20 *Invest. Arterioscler.* 30 (2018) 10-20.
- 21 [18] B.C. Bergman, J.T. Brozinick, A. Strauss, et al., Serum sphingolipids: relationships to
22 insulin sensitivity and changes with exercise in humans, *Am. J. Physiol. Endocrinol.*
23 *Metab.* 309 (2015) E398–E408
- 24 [19] F. Rosqvist, D. Iggman, J. Kullberg, et al., Overfeeding polyunsaturated and saturated
25 fat causes distinct effects on liver and visceral fat accumulation in humans, *Diabetes* 63
26 (2014) 2356–2368.
- 27 [20] M. Shiomi, T. Ito T, S. Yamada, et al., Development of an animal model for
28 spontaneous myocardial infarction (WHHLMI rabbit), *Arterioscler. Thromb. Vasc. Biol.*

- 1 23 (2003) 1239-1244.
- 2 [21] R. Nagasaka, T. Koike, N. Tsukada, et al., The coronary artery running pattern is one of
3 the causes of individual differences in the progression of coronary atherosclerosis in
4 WHHLMI rabbits, an animal model for coronary atherosclerosis, *J. Atheroscler. Thromb.*
5 25 (2018) 393-404.
- 6 [22] S. Yamada S, T. Koike, T. Nakagawa, et al., Morphological features of coronary plaques
7 in WHHLMI rabbits (*Oryctolagus cuniculus*), an animal model for familial
8 hypercholesterolemia, *Exp. Anim.* 66 (2017) 145-157.
- 9 [23] S. Nishiumi, T. Kobayashi, A. Ikeda, et al., A novel serum metabolomics-based diagnostic
10 approach for colorectal cancer, *PLoS One* 7 (2012) e40459.
- 11 [24] H. Takeda, Y. Izumi, M. Takahashi, et al., Widely-targeted quantitative lipidomics
12 method by supercritical fluid chromatography triple quadrupole mass spectrometry, *J.*
13 *Lipid Res.* 59 (2018) 1283-1293.
- 14 [25] X. Li., P. Fang, Y. Li, et al., Mitochondrial reactive oxygen species mediate
15 lysophosphatidylcholine-induced endothelial cell activation, *Arterioscler. Thromb. Vasc.*
16 *Biol.* 36 (2016) 1090-1100.
- 17 [26] S.H.H. Nababan, S. Nishiumi, Y. Kawano, et al., Adrenic acid as an inflammation
18 enhancer in non-alcoholic fatty liver disease, *Arch. Biochem. Biophys.* 623-624 (2017) 64-
19 75.
- 20 [27] P.J. Meikle, G. Wong, D. Tsorotes, et al., Plasma lipidomic analysis of stable and
21 unstable coronary artery disease, *Arterioscler. Thromb. Vasc. Biol.* 31 (2011) 2723-2732.
- 22 [28] A. Yamashita, Y. Zhao, Y. Matsuura, et al., Increased metabolite levels of glycolysis and
23 pentose phosphate pathway in rabbit atherosclerotic arteries and Hypoxic Macrophage,
24 *PLoS One* 9 (2014): e86426.
- 25 [29] He W, Miao FJ, Lin DC, et al., Citric acid cycle intermediates as ligands for orphan G-
26 protein-coupled receptors, *Nature.* 429 (2004):188-193.
- 27 [30] Y. Zhen, D. Alexander, T.A. Manolio, et al., Associations between metabolome compounds
28 and incident heart failure among African Americans: The ARIC study, *Am. J. Epidemiol.*

- 1 178 (2013): 534-542.
- 2 [31] A. Kehlen, M. Haegele, L. Bohme, et al., N-terminal pyroglutamate formation in
3 CX3CL1 is essential for its full biologic activity, *Biosci. Rep.* 37 (2017): BSR20170712.
- 4 [32] L Menegaut, D. Masson, N. Abello, et al., Specific enrichment of 2-arachidonoyl-
5 lysophosphatidylcholine in carotid atheroma plaque from type 3 diabetic patients,
6 *Atherosclerosis* 251 (2016):339-347.
- 7 [33] L.P. de Carvalho, S.H. Tan, G.S. Ow, et al., Plasma ceramides as prognostic biomarkers
8 and their arterial and myocardial tissue correlates in acute myocardial infarction, *JACC*
9 *Basic, Transl. Sci.* 3 (2018):163-175.
- 10 [34] P.A. Mundra, C.K. Barlow, P.J. Nestel, Large-scale plasma lipidomic profiling identifies
11 lipids that predict cardiovascular events in secondary prevention, *JCI Insight.* 3
12 (2018):e121326.
- 13 [35] J.M. Cheng, M. Suoniemi, I. Kardys, et al., Plasma concentration of molecular lipid
14 species in relation to coronary plaque characteristics and cardiovascular outcome:
15 Results of the ATHEROREMO-IVUS study, *Atherosclerosis* 243 (2015): 560-566.
- 16 [36] B. Chaurasia, SA. Summers. Ceramides - Lipotoxic inducers of metabolic disorders,
17 *Trends Endocrinol. Metabolism* 26 (2015): 538-550.
- 18 [37] G.M. Deevska, M. Sunkara, A.J. Morris, et al., Characterization of secretory
19 sphingomyelinase activity, lipoprotein sphingolipid content and LDL aggregation in *ldlr*-
20 *-/-* mice fed on a high-fat diet, *Biosci. Rep.* 31 (2012) 479-490.
- 21 [38] J. Liu, C. Huan, M. Chakraborty, et al., Macrophage sphingomyelin synthase 2
22 deficiency decreases atherosclerosis in mice, *Circ Res.* 105 (2009):295-303.
- 23 [39] F. Norheim, T. Bjellaas, S.T. Hui et al., Genetic, dietary, and sex-specific regulation of
24 hepatic ceramides and the relationship between hepatic ceramides and IR, *J. Lipid. Res.*
25 59 (2018):1164-1174.
- 26 [40] S. Wallner, G. Schimtz, Plasmalogens the neglected regulatory and scavenging lipid
27 species, *Chem. Phys. Lipids* 164 (2011): 573-589.

1 **Figure legends**



2 Fig. 1. Profiles of serum metabolites in female WHHLMI rabbits with severe and mild
3 coronary atherosclerosis. (A) Upper panels indicate the results of the volcano plot. The cutoff
4 value was set at a p-value of 0.05 and a 30% change in the relative concentration of plasma
5 metabolites between the severe and mild groups. Dotted lines indicate the cutoff line. *P*
6 values were calculated by a Student's t-test or Welch's t-test. (B) Lower panels indicate the
7 results of a direct comparison of the SPLS-DA score plots of 59 metabolites.

8



9 Fig. 2. Profiles of serum lipid molecules in female WHHLMI rabbits with severe and mild
10 coronary atherosclerosis. (A) Upper panels indicate the results of the volcano plot. The cutoff
11 value was set at a p-value of 0.05 and a 30% change in relative concentration of plasma
12 metabolites between the severe and mild groups. Dotted lines indicate the cutoff line. *P*
13 values were calculated by a Student's t-test or Welch's t-test. (B) Lower panels indicate the
14 results of a direct comparison of the SPLS-DA score plots of 310 lipid molecules.

15

16 Fig. 3. The corrected relative concentration (CRC) of serum PE16:1p-22:2. (A) Distribution of
17 the CRC of PE16:1p-22:2 at 16 months old for 11 rabbits. (B) Changes in the CRC of
18 PE16:1p-22:2 with aging. Data are mean \pm SEM. FC in panel B indicates the fold changes
19 between the CRC of PE16:1p-22:2 of the severe group at 16 months old and that of the mild
20 group at the indicated age. Statistical analyses were performed by a Student's t-test in panel
21 A and by a Dunnett test in panel B. A correlation coefficient was calculated by Spearman's
22 rank-correlation coefficient. Diamonds indicate the severe group (n=3), circles indicate the
23 mild group (n=6), and squares indicate the intermediate group (n=2).

Figure 1


Sparse partial least squares discriminant analysis (SPLS-DA) of relative concentration of plasma metabolites in severe and mild groups

Figure 2


Sparse partial least squares discriminant analysis (SPLS-DA) of corrected relative concentration of plasma lipid molecules in severe and mild groups

Figure 3

(A) Distribution of CRC of PE 16:1p-22:2



(B) Changes in CRC of PE16:1p-22:2 with aging



1 Table 1. Development of atherosclerotic lesions in female WHHLMI rabbits

2	<hr/>			
3		Severe group	Mild group	P-value
4	<hr/>			
	Animals	3	6	
5	Coronary lesions			
6	Examined sections	22 ± 0	20 ± 1	0.013
7	Sections with lesions	17 ± 1	10 ± 3	0.010
8	Frequency of sections with lesions (%)	76 ± 15	49 ± 16	0.059
9	Maximum CSN (%)	98 ± 2	49 ± 12	<0.001
10	Average CSN (%)	62 ± 15	26 ± 15	0.005
11	Sections with >90% CSN	10 ± 4	0 ± 0	<0.001
12	Frequency of sections with >90% CSN (%)	45 ± 13	0 ± 0	<0.001
13	Sections with >75% CSN (%)	12 ± 4	0 ± 0	0.001
14	Frequency of sections with >75% CSN (%)	56 ± 20	0 ± 0	0.001
15	Coronary severity score	18 ± 11	0.2 ± 0.1	<0.001
16	Aortic lesions			
17	Percentage of surface lesion area (%)	96 ± 2	94 ± 1	0.157
18	<hr/>			

1 Data are the mean \pm SEM. Statistical analyses were performed with the Mann-Whitney
2 U-test. CSN (cross-sectional narrowing) was calculated by dividing the lesion area by
3 the area of the lumen and lesions. The percentage of the surface lesion area was
4 calculated by dividing the surface area of the lesions by the surface area of the aortic
5 lumen.

1 Table 2. Selected serum metabolites as markers for progression of coronary lesions.

2	Severe group	Mild group	Fold		Correlation coefficient		
3	(n = 3)	(n = 6)	Change	p-value	r	p-value	
4	8 months old						
5	Citric acid + Isocitric acid	1.03±0.16	0.67±0.07	1.552	0.037	0.609	0.047
6	Pyroglutamic acid	0.082±0.012	0.054±0.005	1.513	0.032	0.655	0.029

7 Data indicate the relative concentration of each metabolite and are mean ± SEM. A difference was calculated by dividing the relative
 8 concentration of plasma metabolites in the severe group by that in the mild group. Statistical analyses of the difference were performed with a
 9 Student t-test or Welch's t-test. Correlation coefficients were calculated with Spearman's rank-correlation coefficient using all 11 females in the
 10 intermediate group.

1 Table 3. Selected serum lipid molecules at 4 and 8 months old as markers for the progression
 2 of coronary lesions.

	Severe group (n=3)	Mild group (n=6)	Fold Change	p-value	Correlation coefficient	
					r	p-value
4 months old						
LPC22:4	0.47 ± 0.05	0.34 ± 0.03	1.404	0.030	0.700	0.016
DAG18:0-18:0	0.11 ± 0.01	0.21 ± 0.02	0.510	0.005	-0.636	0.035
8 months old						
LPC20:4 (sn-2)	0.81 ± 0.03	0.60 ± 0.03	1.359	0.005	0.773	0.005
Cer d18:1-18:2	0.08 ± 0.01	0.06 ± 0.00	1.377	0.041	0.673	0.023

11 Data indicate the corrected relative concentration (CRC) of each lipid molecule and are mean
 12 ± SEM. A difference was calculated by dividing the CRC of serum metabolites in the severe
 13 group by that in the mild group. Statistical analyses of the difference were performed with a
 14 Student t-test or Welch's t-test. Correlation coefficients were calculated with Spearman's
 15 rank-correlation coefficient using all 11 females in the intermediate group.

Supplementary Material for online publication only

[Click here to download Supplementary Material for online publication only: Supplemental Data-2019_2_5.pdf](#)

1 **Supplementary information**

2 **Materials and Methods**

3 *2.1. Animals*

4 We examined serum markers for coronary lesions using 22 WHHLMI rabbits (12
5 females and 10 males), and 363 WHHLMI rabbits born between 2010 and 2016 were
6 examined for age-dependent changes in the progression of coronary lesions. WHHLMI
7 rabbits were bred at the Institute for Experimental Animals, Kobe University
8 Graduate School of Medicine (Kobe, Japan). In addition, 6 male JW rabbits (Kitayama
9 Labes Co., Ltd., Ina, Japan) were used as a normal control. Rabbits were fed a
10 standard rabbit chow (LPRC4, Oriental Yeast Co., Ltd., Tokyo, Japan). Rabbits were
11 housed individually in metal cages (550 mm x 600 mm x 450 mm in width, depth, and
12 height, respectively) with a flat metal floor. Animals were maintained under specific
13 pathogen-free conditions with a constant temperature (22 ± 2 °C), relative humidity
14 (50–60%), ventilation rate (15 cycles/h), air supply (through a HEPA filter), and
15 lighting cycle (12 h light/dark). This study was approved by the Kobe University
16 Animal Care and Use Committee (approval number: P110511 and P110511-R1), and all
17 animal experiments were conducted in accordance with the Regulations for Animal
18 Experimentation of Kobe University, the Act on Welfare and Management of Animals

1 (Law No. 105, 1973, revised in 2006), Standards Relating to the Care and Management
2 of Laboratory Animals and Relief of Pain (Notification No. 88, 2006), and Fundamental
3 Guidelines for the Proper Conduct of Animal Experiments and Related Activities in
4 Academic Research Institutions under the Jurisdiction of the Ministry of Education,
5 Culture, Sports, Science and Technology (Notice No. 71, 2006). Blood samples were
6 collected at 4-month intervals from 4 months old to 20 months old. At the end of the
7 experiments (20 months old), rabbits were euthanized with an intravenous injection of
8 pentobarbital, and the heart and aorta were excised.

9

10 *2.2. Fractionation of plasma lipoproteins and measurement of lipid concentration*

11 After overnight fasting, samples were collected every 4 months from the marginal
12 ear vein, and lipoproteins were fractionated with an ultracentrifuge to yield very low-
13 density lipoproteins (VLDL, $d < 1.006$ g/mL), LDL ($1.006 < d < 1.063$ g/mL), and HDL
14 ($d > 1.063$ g/mL). Lipoprotein lipid concentration was assayed using enzymatic
15 methods.

16

17 *2.3. Metabolome analyses of sera*

18 Metabolomic profiling using gas chromatography/mass spectrometry (GC/MS) was

1 performed as reported previously [S-1]. Low molecular weight metabolites were
2 extracted, and sinapinic acid (WAKO, Tokyo, Japan) was used as an internal standard
3 to correct for the losses of analytes during sample preparation. Metabolites were
4 analyzed with a GCMS-QP2010 Ultra (Shimadzu Co, Kyoto, Japan), which was
5 equipped with a fused silica capillary column (CP-SIL 8 CB low bleed/MS; 30 mm ×
6 0.25 mm inner diameter, 0.25- μ m film thickness; Agilent Co., Palo Alto, CA). Fifty-nine
7 metabolites (Supplementary Table 1) were recorded using the Advanced Scanning
8 Speed Protocol (ASSP, Shimadzu Co, Kyoto, Japan) and were analyzed using MetAlign
9 (Wageningen UR The Netherlands) and AI-output software (PMID 21641865,
10 21542920).

11

12 *2.4. Lipidome analyses of sera*

13 Lipids were extracted from sera using Bligh and Dyer's method [S-2] with minor
14 modifications [S-3]. A dodecanoyl or heptadecanoyl-based synthetic internal standard
15 mixture (Supplementary Table 2) was used as an internal standard. Lipid molecules
16 were quantified using supercritical fluid chromatography triple quadrupole mass
17 spectrometry (SFC/MS/MS) in the multiple reaction monitoring mode [S-3]. The
18 SFC/MS/MS system was composed of an ACQUITY UltraPerformance Convergence

1 Chromatography (UPC²) system (Waters Co., Milford, MA, USA) and Xevo TQ-S micro
2 tandem mass spectrometer with an electrospray ionization ion source (Waters Co.),
3 which were controlled by MassLynx software version 4.1 (Waters Co.). An HPLC 515
4 pump (Waters Co.) was used as a make-up pump to enhance ionization efficiency. There
5 were 313 lipid molecules detected in the serum (Supplementary Table 3). The relative
6 concentrations (RC) of detected lipid molecules were corrected with serum total
7 cholesterol concentrations using the following equation:

8
$$[\text{Corrected relative concentration (CRC)}] = [\text{relative concentration}] / [\text{concentration}$$

9
$$\text{of serum total cholesterol}] \times 1,000.$$

10

11 *2.5. Evaluation of atherosclerotic lesions*

12 We examined atherosclerotic lesions of WHHLMi rabbits that died from
13 myocardial ischemia during the experiment in addition to rabbits euthanized at the
14 end of the experiment (20 months old). Coronary sections were prepared as reported
15 previously [S-4]. We examined the atherosclerotic lesions of the left circumflex arteries
16 because this artery is the major coronary artery, and atherosclerotic lesions frequently
17 develop there in rabbits [S-5]. Sections were stained with Victoria blue-HE staining.
18 The degree of coronary atherosclerosis was evaluated as cross-sectional narrowing

1 (CSN). The severity of coronary lesions was evaluated using a coronary severity score
2 based on the following equation:

$$3 \quad [\text{Coronary severity score}] = [\text{average CSN}] \times (1 + [\text{frequency of sections with 75\%} \\ 4 \quad \quad \quad 89\% \text{ CSN}] \times 10 + [\text{frequency of sections with >90\% CSN}] \times 20)$$

5 Supplementary Figure 1 shows the development of atherosclerotic lesions in the
6 aorta and coronary circumflex artery of WHHLMI rabbits. As a result of the evaluation
7 of the coronary severity score, we classified WHHLMI rabbits into three groups (severe,
8 intermediate, or mild). In the severe group, every rabbit had coronary sections with
9 more than 90% CSN, and no rabbits in the mild group had coronary sections with more
10 than 75% CSN. Aorta were excised from the aortic root to the bifurcation of the femoral
11 artery. Aortic lesions were evaluated as the ratio of the surface area of the lesions to
12 the surface area of the entire lumen. Yu et al. [S-6] demonstrated that in cholesterol-fed
13 rabbits, the percentage of surface lesion area on aortic lumen surface correlated with
14 the intimal lesion area of the aortic histopathological sections.

15

16 *2.6. Age-dependent progression of coronary lesions*

17 To examine age-dependent progression of coronary lesions in WHHLMI rabbits, we
18 examined the maximum CSN from one month old to 30 months old using 363 dead

1 rabbits or rabbits euthanized in other experiments between 2010 and 2016. Coronary
2 lesions were examined in the manner described above.

3

4 *2.7. Other assays*

5 At the start of the experiment (4 months old) and at the end of experiment (20
6 months old or at death), we assayed the plasma levels of glucose, insulin, NO₂, NO₃,
7 and oxidized LDL, and superoxide dismutase (SOD) activity using the Autokit Glucose
8 (FUJIFILM Wako Pure Chemical Corporation, Osaka, Japan), Lbis (R) Insulin-Rat-T
9 (FUJIFILM Wako Pure Chemical Corporation), NO₂/NO₃ Assay Kit-CII (Dojindo
10 Laboratories, Co., Ltd., Kumamoto, Japan), Rabbit Oxidized Low-density Lipoprotein
11 (OxLDL) ELISA kit (MyBioSource Inc., San Diego, USA), and SOD Assay Kit-WST
12 (Dojindo Laboratories, Inc.), respectively.

13

14 *2.8. Data and statistical analyses*

15 Three rabbits died during the experiment. Because the frequency of sections with
16 >90% CSN was 86%, 82%, and 87% in each of these rabbits, the cause of death was
17 likely related to occlusion of the coronary arteries from an atherosclerotic lesion. In
18 analyses of serum marker candidates for coronary lesions, we also used data from these

1 three rabbits in addition to data from the 17 rabbits sacrificed at 20 months old
2 because the purpose was to identify serum marker candidates for the progression of
3 coronary atherosclerosis. Two rabbits that died for renal tumor or lymphoma were
4 excluded from analysis. In analyses of sex differences, the severity of coronary lesions
5 and the RC of serum metabolites and the CRC of serum lipid molecules were
6 significantly different between males and females (Supplementary Table 4,
7 Supplementary Figure 2), despite no differences in aortic surface lesion area and
8 conventional plasma biochemical parameters. In addition, 7 of 9 males and 3 of 11
9 females were in the severe group. Consequently, we performed metabolomic and
10 lipidomic analyses using only females. Serum marker candidates for coronary
11 atherosclerosis were determined by the method of Supplementary Figure 3. In lipidome
12 analyses, because 20 of 313 lipid molecules were below the detection limit in at least
13 one rabbit, 293 lipid molecules were used to select serum marker candidates. In brief,
14 for serum metabolites or lipid molecules at 4 and 8 months old, marker candidates
15 were selected using two statistical analysis steps. The first step was a volcano plot
16 analysis [S-7] using a cutoff p -value of 0.05 and 30% fold-changes in the CRC between
17 the severe and mild groups. The second step was a Spearman's rank-correlation
18 analysis using a cutoff correlation coefficient-value of 0.550. For serum metabolites and

1 lipid molecules at 12, 16, and 20 months old or death, marker candidates were selected
2 after an additional Dunnett analyses between the CRC of selected candidate molecules
3 of the severe group at the selected age and those of the mild group at each age.
4 Metabolites or lipid molecules with significant >50% fold changes from the severe
5 group at the selected age from that in the mild group at each age were selected as
6 markers.

7 Differences in the profiles of serum metabolites or lipid molecules between the
8 severe and mild groups were analyzed by sparse partial least square discriminant
9 analysis (SPLS-DA) using MetaboAnalyst 4.0 (<http://www.metaboanalyst.ca/>) [S-8].

10 Data are the mean \pm SEM. $P < 0.05$ was considered significant. Correlation
11 analyses between the coronary severity score and the relative concentration of plasma
12 metabolites or lipid molecules were performed with Spearman's rank correlation
13 coefficient.

14

15 **References**

- 16 S-1. S. Nishiumi, T. Kobayashi, A. Ikeda, et al., A novel serum metabolomics-based
17 diagnostic approach for colorectal cancer, PLoS One 7 (2012): e40459.
18 S-2. E.G. Bligh, W.J. Dyer, A rapid method of total lipid extraction and purification,

- 1 Can. J. Biochem. Phys. 37 (1959), 911-917.
- 2 S-3 H. Takeda, Y. Izumi, M. Takahashi, et al., Widely-targeted quantitative lipidomics
3 method by supercritical fluid chromatography triple quadrupole mass
4 spectrometry, *J. Lipid Res.* 59 (2018) 1283-1293.
- 5 S-4. M. Shiomi, T. Ito, M. Shiraishi, et al., Inheritability of atherosclerosis and the role
6 of lipoproteins as risk factors in the development of atherosclerosis in WHHL
7 rabbits: risk factors related to coronary atherosclerosis are different from those
8 related to aortic atherosclerosis, *Atherosclerosis* 96 (1992) 43–52.
- 9 S-5. R. Nagasaka, T. Koike, N. Tsukada, et al., The coronary artery running pattern is
10 one of the causes of individual differences in the progression of coronary
11 atherosclerosis in WHHLMi rabbits, an animal model for coronary atherosclerosis,
12 *J. Atheroscler. Thromb.* 25 (2018) 393-404.
- 13 S-6. Q. Yu, Y. Li, A.B. Waqar, et al: Temporal and quantitative analysis of
14 atherosclerotic lesions in diet-induced hypercholesterolemic rabbits. *J. Biomed.*
15 *Biotech.* 2012: Article ID 506159, 7 pages
- 16 S-7. J.J. Chen, S-J. Wang, C-A. et al.: Selection of differentially expressed genes in
17 microarray data analysis. *Pharmacogenomics J.* 7(2007):212-220.
- 18 S-8. K. Chog, O. Soufan, C. Li, et al.: MetaboAnalyst 4.0: towards more transparent

1 and integrative metabolomics analysis. *Nucleic Acids Res.* 46 (2018): W486-W494.

Supplementary Table 1. Fifty nine metabolites examined in metabolome analyses

1,5-Anhydro-D-glucitol	1-Hexadecanol	1-Methyl Histidine	2-Aminoethanol	3-Hydroxy-Butyrate	Alanine	a-Phenylglycine
Arabitol	Asparagine	Aspartic acid	b-Alanine	Benzen-1,3-Dicarboxylic acid	Citric acid + Isocitric acid	Citrulline
Cystamine	Cystathionine	Cysteine+Cystine	Fumaric acid	Galactosamine_2	Glucuronate_1	Glutamic acid
Glutamine	Glyceraldehyde_2	Glycerol	Glycine	Glycolic acid	Histidine	Hypoxanthine
Inositol	Ketoisoleucine_1	Ketovaline_1	Lactic acid	Lauric acid	Lysine	Lyxose_2
Malic acid	Mannose_1	meso-erythritol	Methionine	N-Acetyl-D-Glucosamine_1	N-FormylGlycine	Nonanoic acid
O-Phosphoethanolamine	Ornithine	Phenylalanine	Phosphate	Prolinamide	Proline	Pyroglutamic acid
Pyruvate+Oxalacetic acid	Ribitol	Serine	Threo-b-HydroxyAspartic acid	Threonine	trans-4-Hydroxy-L-proline	Tryptophan
Tyrosine	Uracil	Valine				

Supplementary Table 2. Internal standard substances in lipidome analyses

Lipid class	ISD conc. (μ M)	Final conc. (nM)	Injection (fmol)	Total extraction volume (μ mol)	Serum concentration (nmol/ml)
FFA 17:0	500	2244	2244	4999	250
LPC 17:0	25	112	112	250	12.5
LPE 17:1	500	2,243	2,244	4,999	250
PC 17:0-17:0	500	2,243	2,244	4,999	250
PE 17:0-17:0	200	1,794	1,795	3,999	200
PS 17:0-17:0	2,000	17,948	17,948	399,887	1,999
SM d18:1-17:0	5	22.4	22.4	50.0	2.50
Cer d18:1-17:0	5	22.4	22.4	50.0	2.50
CE 17:0	250	2244	2244	4999	250
MAG 17:0	50	224	224	500	25
DAG 12:0-12:0	50	224	224	500	25
TAG 51:0	50	224	224	500	25

ISD, internal standard; FFA, free fatty acids; LPC, Lysophosphatidylcholine; LPE, Lysophosphatidylethanolamine; PC, phosphatidylcholine; PE, Phosphatidylethanolamine; PS, phosphatidylserine; SM, Sphingomyelin; Cer, Ceramide; CE, cholesterol ester; MAG, monoacylglycerol; DAG, Diacylglycerol; TAG, triacylglycerol

Supplementary Table 3. 322 lipid molecules examined in lipidome analyses**Free fatty acids (FFA): 8 lipids**

FFA 14:0 FFA 16:0 FFA 16:1 FFA 18:0 FFA 18:1 FFA 18:2 FFA 18:3 FFA 20:4

Lysophosphatidylcholine (LPC): 26 lipidsLPC 14:0 LPC 16:0 (sn-1) LPC 16:0 (sn-2) LPC 16:1 (sn-1) LPC 16:1 (sn-2) LPC 18:0 (sn-1) LPC 18:0 (sn-2) LPC 18:1 (sn-1) LPC 18:1 (sn-2) LPC 18:2 (sn-1) LPC 18:2 (sn-2)
LPC 18:3 (sn-1) LPC 18:3 (sn-2) LPC 20:1 (sn-1) LPC 20:1 (sn-2) LPC 20:2 (sn-1) LPC 20:2 (sn-2) LPC 20:3 (sn-1) LPC 20:3 (sn-2) LPC 20:4 (sn-1) LPC 20:4 (sn-2) LPC 20:5
LPC 22:0 LPC 22:4 LPC 22:5 LPC 22:6**Lysophosphatidylethanolamine (LPE): 6 lipids**

LPE 16:0 LPE 18:0 (sn-1) LPE 18:0 (sn-2) LPE 18:1 LPE 18:2 (sn-1) LPE 18:2 (sn-2)

Phosphatidylcholine (PC): 33 lipidsPC 16:0-16:0 PC 16:0-16:1 PC 16:0-18:0 PC 16:0-18:1 PC 16:0-18:2 PC 16:0-18:3 PC 16:0-20:0 PC 16:0-20:2 PC 16:0-20:3 PC 16:0-20:4 PC 16:0-22:4
PC 16:0-22:5 PC 16:1-18:1 PC 16:1-18:2 PC 18:0-18:0 PC 18:0-18:1 PC 18:0-18:2 PC 18:0-18:3 PC 18:0-20:2 PC 18:0-20:3 PC 18:0-20:4 PC 18:0-22:4
PC 18:0-22:5 PC 18:1-18:1 PC 18:1-18:2 PC 18:1-18:3 PC 18:1-20:4 PC 18:2-18:2 PC 18:2-18:3 PC 18:2-20:1 PC 18:2-20:2 PC 18:2-20:3 PC 18:2-20:4**Ether Phosphatidylcholine (EPC): 11 lipids**PC 16:1e-16:0 / PC 16:0p-16:0 PC 16:1e-18:1 / PC 16:0p-18:1 PC 16:1e-18:2 / PC 16:0p-18:2 PC 16:1e-20:4 / PC 16:0p-20:4 PC 16:2e-20:5 / PC 16:1p-20:5
PC 18:1e-18:1 / PC 18:0p-18:1 PC 18:1e-18:2 / PC 18:0p-18:2 PC 18:1e-20:4 / PC 18:0p-20:4 PC 18:2e-18:2 / PC 18:1p-18:2 PC 18:2e-20:4 / PC 18:1p-20:4 PC 18:2p-20:5**Phosphatidylethanolamine (PE): 9 lipids**

PE 16:0-18:1 PE 16:0-18:2 PE 16:0-20:4 PE 18:0-18:1 PE 18:0-18:2 PE 18:0-18:3 PE 18:0-20:4 PE 18:1-18:2 PE 18:2-18:2

Phosphatidylethanolamine plasmalogen (PEpln): 40 lipidsPE 16:0p-16:0 PE 16:0p-18:1 PE 16:0p-18:2 PE 16:0p-18:3 PE 16:0p-20:3 PE 16:0p-20:4 PE 16:0p-20:5 PE 16:0p-22:4 PE 16:0p-22:5 PE 16:0p-22:6 PE 16:1p-20:2
PE 16:1p-22:2 PE 16:1p-22:3 PE 18:0p-16:0 PE 18:0p-16:1 PE 18:0p-18:0 PE 18:0p-18:1 PE 18:0p-18:2 PE 18:0p-18:3 PE 18:0p-20:3 PE 18:0p-20:4 PE 18:0p-20:5
PE 18:0p-22:4 PE 18:0p-22:5 PE 18:0p-22:6 PE 18:1p-16:0 PE 18:1p-18:1 PE 18:1p-18:2 PE 18:1p-18:3 PE 18:1p-20:2 PE 18:1p-20:3 PE 18:1p-20:4 PE 18:1p-22:4
PE 18:1p-22:5 PE 18:1p-22:6 PE 18:2p-18:1 PE 18:2p-18:2 PE 18:2p-20:2 PE 18:2p-20:4 PE 18:2p-22:2**Phosphatidylinositol (PI): 10 lipids**

PI 16:0-16:0 PI 16:0-18:1 PI 16:0-18:2 PI 18:0-18:1 PI 18:0-18:2 PI 18:0-20:2 PI 18:0-20:3 PI 18:0-20:4 PI 18:1-18:1 PI 18:1-18:2

Sphingomyelin (SM): 14 lipidsSM d18:1-14:0 SM d18:1-14:1 SM d18:1-16:0 SM d18:1-16:1 SM d18:1-18:0 SM d18:1-18:1 SM d18:1-18:2 SM d18:1-20:0 SM d18:1-20:1 SM d18:1-20:2 SM d18:1-22:0
SM d18:1-22:1 SM d18:1-22:2 SM d18:1-22:3**Ceramide (Cer): 14 lipids**Cer d18:1-16:0 Cer d18:1-18:0 Cer d18:1-18:2 Cer d18:1-18:4 Cer d18:1-20:0 Cer d18:1-20:1 Cer d18:1-20:2 Cer d18:1-20:4 Cer d18:1-22:0 Cer d18:1-22:1 Cer d18:1-22:2
Cer d18:1-22:3 Cer d18:1-22:4 Cer d18:1-22:5

(Supplementary table 3—continued)

Cholesterol ester (CE): 20 lipids

CE 14:0	CE 14:1	CE 16:0	CE 16:1	CE 18:0	CE 18:1	CE 18:2	CE 18:3	CE 18:4	CE 20:1
CE 20:2	CE 20:3	CE 20:4	CE 20:5	CE 22:1	CE 22:2	CE 22:3	CE 22:4	CE 22:5	CE 22:6

Monoacylglycerol (MAG): 2 lipids


MAG 16:0	MAG 18:0
----------	----------

Diacylglycerol (DAG): 51 lipids

DAG 14:0-16:0	DAG 14:0-16:1	DAG 14:0-18:0	DAG 14:0-18:1	DAG 14:0-18:2	DAG 14:0-18:3	DAG 14:0-20:0	DAG 14:1-18:1	DAG 14:1-18:2	DAG 14:1-22:2
DAG 16:0-16:0	DAG 16:0-16:1	DAG 16:0-18:0	DAG 16:0-18:1	DAG 16:0-18:2	DAG 16:0-18:3	DAG 16:0-20:2	DAG 16:0-20:3	DAG 16:0-20:4	DAG 16:0-22:4
DAG 16:0-22:5	DAG 16:1-16:1	DAG 16:1-18:0	DAG 16:1-18:1	DAG 16:1-18:2	DAG 16:1-18:3	DAG 16:1-20:2	DAG 16:1-20:3	DAG 16:1-20:4	DAG 18:0-18:0
DAG 18:0-18:1	DAG 18:0-18:2	DAG 18:0-18:3	DAG 18:0-20:4	DAG 18:1-18:1	DAG 18:1-18:2	DAG 18:1-18:3	DAG 18:1-18:4	DAG 18:1-20:2	DAG 18:1-20:3
DAG 18:1-20:4	DAG 18:1-22:4	DAG 18:1-22:5	DAG 18:2-18:2	DAG 18:2-18:3	DAG 18:2-20:2	DAG 18:2-20:3	DAG 18:2-20:4	DAG 18:2-22:4	DAG 18:3-18:3
DAG 18:3-20:4									

Triacylglycerol (TAG): 69 lipids

TAG 44:0	TAG 44:1	TAG 44:2	TAG 44:4	TAG 44:5	TAG 46:0	TAG 46:1	TAG 46:2	TAG 46:3	TAG 48:0
TAG 48:1	TAG 48:2	TAG 48:3	TAG 48:4	TAG 48:6	TAG 48:7	TAG 48:8	TAG 50:0	TAG 50:1	TAG 50:2
TAG 50:3	TAG 50:4	TAG 50:5	TAG 50:6	TAG 50:7	TAG 50:8	TAG 50:9	TAG 52:0	TAG 52:1	TAG 52:2
TAG 52:3	TAG 52:4	TAG 52:5	TAG 52:6	TAG 52:8	TAG 52:9	TAG 54:0	TAG 54:1	TAG 54:2	TAG 54:3
TAG 54:4	TAG 54:5	TAG 54:6	TAG 54:7	TAG 54:8	TAG 54:9	TAG 54:10	TAG 54:11	TAG 54:12	TAG 56:1
TAG 56:2	TAG 56:3	TAG 56:4	TAG 56:5	TAG 56:6	TAG 56:7	TAG 56:8	TAG 56:9	TAG 56:10	TAG 56:11
TAG 56:12	TAG 56:13	TAG 58:12	TAG 58:13	TAG 60:12	TAG 60:13	TAG 60:15	TAG 62:15	TAG 62:16	





Supplementary Figure 1. Development of atherosclerotic lesions in aorta and coronary arteries of 20 WHHLMI rabbits analyzed in this study. Bars in photomicrographs indicate 500 micrometers.

Coronary severity score = [average CSN] x (1 + [frequency of sections with 75-89% CSN] x 10+ [frequency of sections with >90% CSN] x 20); CSN, cross-sectional narrowing; Solid symbols, female; Open symbols, males


Supplementary Table 4. Sex differences in coronary severity score of WHHLMi rabbits

		Coronary severity score	Examined sections	Sections with lesions	Frequency of sections with lesions	Average CSN	Maximum CSN	Sections with >75% CSN	Frequency of sections with >75% CSN	Sections with >90% CSN	Frequency of sections with >90% CSN	Aortic lesion
Mean	Female (n=11)	5.3	20.3	13.5	65%	39%	66%	5	21%	2.3	10%	94%
	Male (n=9)	15.2	19.4	15.4	82%	63%	94%	9	51%	5.7	33%	92%
SEM	Female	5.4	0.6	2.3	11%	9%	10%	2.1	10%	1.6	7%	2%
	Male	3.5	1.1	1.3	8%	8%	3%	1.8	11%	1.6	11%	1%
	<i>P-value</i>	<i>0.020</i>	<i>0.726</i>	<i>0.909</i>	<i>0.136</i>	<i>0.044</i>	<i>0.030</i>	<i>0.052</i>	<i>0.036</i>	<i>0.035</i>	<i>0.021</i>	<i>0.271</i>

Statistical analyses were performed by Mann-Whitney U-test. CSN, cross-sectional narrowing; average CSN, mean of CSN in all coronary cross-sections; Maximum CSN, The largest CSN in all coronary cross-sections.



Supplementary Figure 2. Sex differences in plasma metabolites (A) and lipid molecules (B). Data were analyzed by volcano plot analyses. Horizontal red broken lines indicate levels of $p=0.05$ in Student's *t*-test or Welch's *t*-test.



Supplementary Figure 3. Method for selecting blood marker candidates relating to progression of coronary lesions.

RC, relative concentration; CRC, corrected relative concentration with plasma total cholesterol concentration

Supplementary Table 5. Plasma lipid levels of WHHLMI rabbits.



	Severe group	Mild group	<i>P-value</i>
Plasma cholesterol (mg/dl)			
At 4 months old	1346 ± 46	1394 ± 35	0.446
At 20 months old or at death	1367 ± 251	1019 ± 79	0.146
Plasma triglyceride (mg/dl)			
At 4 months old	290 ± 71	257 ± 19	0.560
At 20 months old or at death	417 ± 171	303 ± 35	0.576
VLDL cholesterol(mg/dl)			
At 4 months old	149 ± 17	133 ± 11	0.428
At 20 months old or at death	233 ± 110	96 ± 18	0.340
VLDL triglyceride (mg/dl)			
At 4 months old	50 ± 14	39 ± 4	0.348
At 20 months old or at death	132 ± 81	64 ± 10	0.489
LDL cholesterol (mg/dl)			
At 4 months old	1184 ± 38	1248 ± 35	0.290
At 20 months old or at death	1025 ± 204	910 ± 67	0.197
LDL triglyceride (mg/dl)			
At 4 months old	234 ± 56	208 ± 15	0.566
At 20 months old or at death	270 ± 85	216 ± 26	0.454
HDL cholesterol (mg/dl)			
At 4 months old	12.8 ± 2.4	12.6 ± 1.1	0.937
At 20 months old	15.8 ± 2.6	12.1 ± 1.0	0.153
LDL-cholesterol / HDL cholesterol			
At 4 months old	99 ± 17	102 ± 8	0.842
At 20 months old	72 ± 5	78 ± 9	0.640

Data are presented as the mean ± SEM. Statistical analyses were performed with Student's t-test or Welch's t-test.


Supplementary Table 6. Plasma levels of glucose, insulin, oxidized LDL, SOD, NO₂, and NO₃ of WHHLMi rabbits.

	Severe group	Mild group	<i>P</i> -value
Plasma glucose (mg/dl)			
At 4 months old	137 ± 9	133 ± 5	0.712
At 20 months old or at death	115 ± 6	124 ± 4	0.201
Plasma insulin (ng/ml)			
At 4 months old	0.48 ± 0.06	0.54 ± 0.09	0.640
At 20 months old or at death	0.78 ± 0.30	0.74 ± 0.20	0.913
NO₂(μmol/l)			
At 4 months old	16.6 ± 2.0	16.2 ± 2.4	0.913
At 20 months old or at death	21.7 ± 7.4	15.0 ± 2.8	0.329
NO₃ (μmol/l)			
At 4 months old	31.2 ± 14.4	32.2 ± 6.5	0.943
At 20 months old or at death	21.0 ± 2.6	20.6 ± 4.0	0.943
Oxidized LDL (ng/ml)			
At 4 months old	0.55 ± 0.10	0.52 ± 0.06	0.789
At 20 months old or at death	0.90 ± 0.15	0.76 ± 0.20	0.668
Superoxide dismutase activity (U/ml)			
At 4 months old	1580 ± 754	1676 ± 359	0.900
At 20 months old or at death	1227 ± 399	1383 ± 353	0.795


Data are presented as the mean ± SEM. Statistical analyses were performed with Student's t-test or Welch's t-test.



Supplementary Figure 4 Differences in plasma metabolites (A) and lipid molecules (B) between normal JW rabbits and WHHLMI rabbits. Data were analyzed volcano plot analyses. Horizontal red broken lines indicate levels of $p=0.05$ in Student's *t*-test or Welch's *t*-test.



Supplementary Figure 5. Differences in selected serum metabolites (A) and lipid molecules (B) between JW rabbits and WHHLMI rabbits. Error bars indicate SEM. Statistical analyses were performed by the Dunnet method (*, $p=0.05$ for JW vs WHHLMI rabbit severe group, #, $p=0.05$ for JW rabbits vs WHHLMI rabbit mild group). Concentration of other selected lipid molecules were below the detection limit in JW rabbits. RC, relative concentration



Supplementary figure 6. Development and progression of coronary atherosclerosis in 363 WHLMI rabbits that died between 2010 and 2016. Data are represented as the mean \pm SEM. Sex differences were observed in the maximum coronary cross-sectional narrowing at 23-24 months old and 25-26 months old by Welch's t-test.