



Differences in fungal immune recognition by monocytes and macrophages: N-mannan can be a shield or activator of immune recognition



Bhawna Yadav^a, Héctor M. Mora-Montes^{a,1}, Jeanette Wagener^{a,2}, Iain Cunningham^{a,3}, Lara West^{c,4}, Ken Haynes^{c,5}, Alistair J.P. Brown^{a,b}, Neil A.R. Gow^{a,b,*}

^a The Aberdeen Fungal Group, School of Medicine, Medical Sciences & Nutrition, Institute of Medical Sciences, University of Aberdeen, Foresterhill, Aberdeen AB25 2ZD, UK

^b Medical Research Council Centre for Medical Mycology at the University of Exeter, Geoffrey Pope Building, Stocker Road, Exeter EX4 4QD, UK

^c Department of Biosciences, University of Exeter, Geoffrey Pope Building, Stocker Road, Exeter EX4 4QD, UK

ARTICLE INFO

Keywords:

Fungal cell wall
N-linked mannosylation
β-Glucan
Monocytes
Macrophages

ABSTRACT

We designed experiments to assess whether fungal cell wall mannans function as an immune shield or an immune agonist. Fungal cell wall β-(1,3)-glucan normally plays a major and dominant role in immune activation. The outer mannan layer has been variously described as an immune shield, because it has the potential to mask the underlying β-(1,3)-glucan, or an immune activator, as it also has the potential to engage with a wide range of mannose detecting PRRs. To resolve this conundrum we examined species-specific differences in host immune recognition in the *och1Δ* N-mannosylation-deficient mutant background in four species of yeast-like fungi. Irrespective of the fungal species, the cytokine response (TNFα and IL-6) induced by the *och1Δ* mutants in human monocytes was reduced compared to that of the wild type. In contrast, TNFα production induced by *och1Δ* was increased, relative to wild type, due to increased β-glucan exposure, when mouse or human macrophages were used. These observations suggest that N-mannan is not a major PAMP for macrophages and that in these cells mannan does shield the fungus from recognition of the inner cell wall β-glucan. However, N-mannan is a significant inducer of cytokine for monocytes. Therefore the metaphor of the fungal “mannan shield” can only be applied to some, but not all, myeloid cells used in immune profiling experiments of fungal species.

1. Introduction

We set out to understand some unresolved questions about the role of mannose-based outer cell wall components in fungi in either initiating or shielding immune recognition. The innate immune system forms the first line of defence against fungal infections and is essential for protective cell-mediated immunity (da Silva Dantas et al., 2016; Drummond et al., 2014; Netea et al., 2015; Qin et al., 2016; Salazar and Brown, 2018). Innate immune responses involve multiple immune cell types of which monocytes and macrophages are its vital components.

These cells express germline-encoded pattern recognition receptors (PRRs) such as C-Type Lectin Receptors (CLRs), Toll-like-receptors (TLRs) and NOD-like receptors (NLRs) that sense conserved pathogen-specific molecular patterns (PAMPs) that, for fungi, are mainly located in the cell wall. These immune recognition events trigger intracellular signalling pathways leading to a release of pro-inflammatory cytokines and chemokines that orchestrate the recruitment of neutrophils and activate downstream innate and adaptive immune responses (Borriello et al., 2020; Dambuza and Brown, 2015; Erwig and Gow, 2016; Nikolakopoulou et al., 2020; Plato et al., 2015).

* Corresponding author.

E-mail addresses: bhawnayadav1983@yahoo.com (B. Yadav), hmora@ugto.mx (H.M. Mora-Montes), jeanette_wagener@oslerdiagnostics.com (J. Wagener), iaincunningham@abdn.ac.uk (I. Cunningham), lara.edmonstone.west@gmail.com (L. West), a.j.p.brown@exeter.ac.uk (A.J.P. Brown), n.gow@exeter.ac.uk (N.A.R. Gow).

¹ Current address: Departamento de Biología, División de Ciencias Naturales y Exactas, Universidad de Guanajuato, Guanajuato, Gto. 36050, Mexico.

² Current address: Osler Diagnostics Limited, King Charles House, Park End Street, Oxford, OX1 1JD, UK.

³ Current address: Aberdeen Centre for Arthritis and Musculoskeletal Health, Institute of Medical Sciences, University of Aberdeen, Foresterhill, Aberdeen AB25 2ZD, UK.

⁴ Current address: Kempen Capital Management, Octagon Point, 5 Cheapside, London EC2V 6AA, UK.

⁵ This article is submitted in memory of the late Ken Haynes who contributed to this work.

<https://doi.org/10.1016/j.tcs.2020.100042>

Received 13 May 2020; Received in revised form 8 July 2020; Accepted 8 July 2020

Available online 21 July 2020

2468-2330/ © 2020 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

The fungal cell wall is the primary point of contact with the components of the host innate immune system (Erwig and Gow, 2016) and is a highly dynamic organelle essential for cell viability, morphogenesis, and pathogenesis (García-Rubio et al., 2019; Gow et al., 2017; Gow and Yadav, 2017; Vendele et al., 2020). The composition and architecture of the fungal cell wall changes in response to different environmental and stress conditions and varies in different cell morphotypes, helping the fungi to adapt in different growth conditions (Brown et al., 2014; Cottier and Hall, 2019; Hopke et al., 2018). The best studied cell wall of a fungal pathogen is that of *Candida albicans* (Gow et al., 2012; Gow and Hube, 2012). It is a bilayer structure composed of an inner layer of β -(1,3)-glucan and chitin, which is interconnected through β -(1,6)-glucan to the outer layer rich in glycosylphosphatidylinositol-anchored mannoproteins, modified with *N*- and *O*-linked mannans (Hall and Gow, 2013; Klis et al., 2001). The outer cell wall mannan forms a fringe of microfibrils that represent a permeability barrier to high molecular weight molecules, but apparently not to liposomes (Walker et al., 2018). These polysaccharides comprising the cell wall are highly fungal specific and present important pathogen-associated molecular patterns (PAMPs) recognized by PRRs (Gow et al., 2012; Levitz, 2010; Netea et al., 2008; Rappleye and Goldman, 2008; Snarr et al., 2017; van de Veerdonk et al., 2008).

However, during evolution and species divergence, while some of the cell wall components have been conserved, others are species-specific. The inner cell wall polysaccharides have been shown to be largely conserved across the fungal kingdom, and studies have suggested the inner layer to be highly immunogenic and capable of eliciting a strong immune response (Snarr et al., 2017). The β -(1,3)-glucan-dectin-1 interaction is amongst the most extensively studied fungal PAMP–host PRR interactions (Brown and Gordon, 2001; Dambuza and Brown, 2015; Gow et al., 2007; Netea et al., 2006; Rogers et al., 2005; Taylor et al., 2004). β -(1,3)-Glucan is the major component of most fungal cell walls and comprises between 30 and 60% of the dry cell wall weight (Bowman and Free, 2006; Gow et al., 2017; Klis et al., 2001; Latgé, 2007). Although β -(1,3)-glucan is a common element of the cell wall of fungi there are differences in the extent of β -(1,6)-glucan cross linking of this component and some structural differences – for example in the yeast and hyphal β -(1,3)-glucans of *C. albicans* (Lowman et al., 2014).

Outer cell wall mannan comprises about 40% of dry cell wall weight in *C. albicans* (Gow et al., 2017; Klis et al., 2001) and has been shown across several studies to be important for host fungal interactions (Bates et al., 2006; Vendele et al., 2020). While *O*-mannan is mostly a linear structure of α -mannoses, with occasional β -(1–2) side chains, *N*-mannan is a highly branched structure of mannose residues in different linkages (Hall, 2015; Hall and Gow, 2013). It has a highly conserved oligosaccharide core synthesised in the endoplasmic reticulum, and a highly branched outer chain elaborated in the Golgi complex, which consists of a linear α -(1,6)-linked mannose backbone with numerous side chains composed of α -(1,2)-, α -(1,3)-, β -(1,2)-linked mannose residues and phosphomannan moieties (Mora-Montes et al., 2009; Shibata et al., 1992). The *N*-mannan branching patterns and patterns of linkages between mannose residues are fungal species-specific and contribute to the diversity of these structures.

Even though mannan has been considered as a mask for the underlying immunogenic cell wall components to escape immune recognition, numerous PRRs specific for mannans are involved in activating host defence responses, indeed there are more mannan-recognising immune receptors for fungi than for any other class of ligand (Erwig and Gow, 2016; Patin et al., 2019; Salazar and Brown, 2018). For example, it has been demonstrated that the CLR mannose receptor (MR), DC-SIGN, dectin-2 and dectin-3 are involved in the sensing of *N*-linked mannans (Cambi et al., 2008; Netea et al., 2006; Saijo et al., 2010; Vendele et al., 2020; Yamamoto et al., 1997). It has also been established that TLR2 recognises the glycolipid phospholipomannan (Jouault et al., 2003) while TLR4 recognises the *O*-linked mannan (Netea et al., 2006, 2002). Soluble receptors such as mannose-

binding lectin (MBL) and galectin-3 are also involved in the recognition of the fungal mannose structures by the host (Snarr et al., 2017).

Significant progress has been made on understanding fungus-host interactions through studies of *C. albicans*, partly because of the availability of a comprehensive number of genetic mutants lacking, or altered in key immunologically active components of its cell wall. The study of *C. albicans* mutants with defects in the *N*-linked mannosylation pathway has helped to establish the importance of these cell wall structures in the recognition of *C. albicans* by innate immune cells (Netea et al., 2006; Hall, 2015; Hall et al., 2013; Hall and Gow, 2013; Zhang et al., 2016). Mutants of *N*-mannan biogenesis, (for example, mutants with defects in *N*-linked mannan processing enzyme α -glucosidase or mutants with loss of outer chain *N*-mannans), are affected in their ability to induce cytokine production by human peripheral blood mononuclear cells (hPBMCs), dendritic cells, and murine macrophages (Cambi et al., 2008; Hall et al., 2013; McKenzie et al., 2010; Mora-Montes et al., 2010, 2007; Netea et al., 2006; Zhang et al., 2016). They are also affected in their phagocytosis by human dendritic cells and murine macrophages (Bain et al., 2014, 2012; Cambi et al., 2008; Lewis et al., 2012; McKenzie et al., 2010). In contrast to the *C. albicans* glycosylation mutants, *C. glabrata* mannosylation mutants display enhanced virulence in a murine infection model (West et al., 2013).

However, the role of *C. albicans* *N*-linked mannans in the innate immune recognition cannot always be extrapolated to other fungi since there are considerable differences in the chemical structure of the *N*-mannan outer chains (Fabre et al., 2014; Jouault et al., 2006). These differences often define the species- and strain-specific serotype (Fukazawa, 1989; Kozel et al., 2004). Furthermore, the compactness of these structures can also affect the extent of masking or exposure of the inner immunogenic cell wall components and the overall organisation of the cell wall can be significantly affected by yeast-hypha morphogenesis, carbon source and local environmental stresses (Bain et al., 2014; Ballou et al., 2016; Graus et al., 2018; Mukaremera et al., 2017; Pradhan et al., 2018; Vendele et al., 2020).

The outer layer of fungal cell walls displays a large chemical diversity among fungal pathogens (Erwig and Gow, 2016; García-Rubio et al., 2019) and has been said to conceal underlying immunogenic layers. By hiding β -(1,3)-glucan in particular the outer layers have the potential to contribute to immune evasion (Hernández-Chávez et al., 2017; Netea et al., 2008). In this regard α -(1,3)-glucan has been shown to mask immune recognition of underlying β -(1,3)-glucan in *Histoplasma capsulatum* (Garfoot et al., 2016; Rappleye et al., 2007) and outer cell wall mannans have been claimed to do the same for *Candida* species (Graus et al., 2018; Wheeler et al., 2008). However, a striking anomaly in the literature has been the variable assertions as to the immunological role of mannans. Both mannans and α -(1,3)-glucan have been suggested as acting primarily in shielding the underlying pro-inflammatory β -(1,3)-glucan layer from immune surveillance (Rappleye et al., 2007; Rappleye and Goldman, 2008; Wheeler and Fink, 2006). For example, *S. cerevisiae* wild type cells were shown to be poor in stimulating cytokine production, but mutant strains with defects in the *N*-linked mannan elaboration, such as the *och1Δ* null mutant, stimulated the mouse macrophage cell line RAW 264.7 to produce high levels of TNF α (Wheeler and Fink, 2006). These data contrast with the observations made using *C. albicans* *och1Δ* null mutant, where loss of the *N*-linked mannan outer chain led to reduced cytokine production by hPBMCs and dendritic cells (Bates et al., 2006; Cambi et al., 2008; Gow et al., 2007; Hall et al., 2013; Netea et al., 2006).

To further explore the apparent differences between *N*-mannan mediated immune recognition effects of different fungal species we used *och1* mutants from four different fungal species, *C. albicans*, *C. dubliniensis*, *C. glabrata* and *S. cerevisiae* and compared the immune response induced by hPBMCs and mouse macrophages (RAW 264.7). *Och1* is an α -(1,6)-mannosyl transferase catalyzing the addition of first α -(1,6)-mannose to the *N*-glycan core. The *och1Δ* mutants show a loss of the *N*-mannan fibrillar layer leading to the exposure of the inner cell

wall components and also a compensatory increase in β -(1,3)-glucan and chitin. We also compared fungal cells prepared for the immune assays by killing by heat-treatment – which is known to expose inner cell wall ligands (Gow et al., 2007), and chemical fixation cells that better preserves the native cell wall structure. In some experiments we used heat killed fungal cells as the target for immune cell recognition. Heat killed cells have been used to prevent cell wall changes associated with yeast cell growth and yeast-hypha morphogenesis from interfering with cytokine induction assays. In our experiments heat-killing also permeabilises the cell wall, thus allowing PRR engagement with cell wall PAMPs that are not exposed in native cell walls (Gow et al., 2007). We also examined the relative porosity and cell wall composition of the cell walls of these species and their mutants.

We showed that loss of *N*-mannan led to a lower cytokine response (TNF α and IL-6) from hPBMCs, but a higher TNF α induction from RAW 264.7 and human macrophages compared to the controls, and that this was explained by recognition of fungal cells by macrophages being primarily mediated by dectin-1. But, irrespective of the fungal species, fungal cell wall *N*-mannan was important for fungal recognition by human monocytes. This provides an explanation for apparently contradictory results in the literature. Mannan acts only as an immune shield for cells such as macrophages that deploy dectin-1 as the primary immune recognition receptor, whilst other immune cell types depend strongly on recognition of *N*-mannan via a set of mannan detecting PRRs.

2. Methods

2.1. Strains and culture conditions

The strains constructed and used in this work are listed in Table 1. To prepare yeast cells for cytokine stimulation assays, cells were grown overnight at 30 °C in YPD broth with shaking at 200 rpm, transferred to fresh broth with a starting O.D.₆₀₀ of 0.2, and further incubated for 4 h till they reached the mid-log growth. *Candida* strains were maintained at 30 °C in YPD, but *S. cerevisiae* cells were grown in SC medium (0.67% (w/v) yeast nitrogen base with ammonium sulfate without amino acids, 0.2% (w/v) complete supplement mixture minus uracil with 2% (w/v) glucose (SC-glc) or with 2% galactose and 3% (w/v) raffinose (SC-gal-raf). A supplement of 50 μ g/ml uridine was added to media used for the growth of *S. cerevisiae* wild type and all *Ura^ooch1 Δ* strains. *C. glabrata* strains were grown in YPD with and without 20 μ g/ml doxycycline.

Cells were harvested by centrifugation, and washed twice with PBS. The cells were heat-killed at 65 °C for 2 h. To compare the temperature-induced cell-wall changes in wild type strains, 2×10^7 cells/ml cells were killed with 100 mM thimerosal for 1 h, washed thrice with PBS, counted and cell suspensions of 2×10^6 cells/ml were exposed to 30, 37, 45, 50, 55, 65 and 75 °C for 2 h. The loss of cell viability was assessed by colony counting of viable cells on YPD plates.

2.2. Construction of null and conditional mutants

The mini-ura-blaster technique was used for *C. dubliniensis* *OCH1* disruption as follows: the primer pair 5'-ATGCTACAACTAAGGGAACC CAAAATGGTTCATAGACATCTAAACTAGCCATTTTAGGAATAATAGT

Table 1
Strains used in this study.

Species	Strain	Genotype	Source
<i>C. albicans</i>	CAI4	<i>ura3Δ::imm434/ura3Δ::imm434</i>	(Fonzi and Irwin, 1993)
<i>C. albicans</i>	CaWT (NGY152)	As CAI4 but <i>RPS1/rps1Δ::Clp10</i>	(Brand et al., 2004)
<i>C. albicans</i>	<i>Caoch1</i> (NGY357)	As CAI4 but <i>och1Δ::hisG/och1Δ::hisG, RPS1/rps1Δ::Clp10</i>	(Bates et al., 2006)
<i>C. albicans</i>	<i>Caoch1Δ/ CaOCH1</i> (NGY358)	As CAI4 but <i>och1Δ::hisG/och1Δ::hisG, RPS1/rps1Δ::Clp10-CaOCH1</i>	(Bates et al., 2006)
<i>C. albicans</i>	<i>Camnn2Δ⁶</i> (NGY 600)	As CAI4 but <i>mnn2Δ::dpl200/mnn2Δ ::dpl200mnn22Δ::dpl200/mnn2Δ::dpl200mnn23Δ::dpl200/mnn23Δ::dpl200mnn24Δ ::dpl200/mnn24Δ::dpl200mnn26Δ::dpl200/mnn26Δ::dpl200mnn21Δ::dpl200/mnn21Δ ::dpl200 RPS1/rps1Δ::Clp10</i>	(Hall et al., 2013)
<i>C. albicans</i>	<i>Camnn2Δ⁶ + CaMNN2 + CaMNN26</i> (NGY 601)	As CAI4 but <i>mnn2Δ::dpl200/mnn2Δ ::dpl200mnn22Δ::dpl200/mnn2Δ::dpl200mnn23Δ::dpl200/mnn23Δ::dpl200mnn24Δ ::dpl200/mnn24Δ::dpl200mnn26Δ::dpl200/mnn26Δ::dpl200mnn21Δ::dpl200/mnn21Δ ::dpl200 RPS1/rps1Δ::Clp10-NN2/MNN26</i>	(Hall et al., 2013)
<i>C. albicans</i>	<i>pmr1Δ</i> (NGY355)	As CAI4 but <i>pmr1Δ::hisG/pmr1Δ::hisG, RPS1/rps1Δ::Clp10</i>	(Bates et al., 2005)
<i>C. albicans</i>	<i>pmr1Δ + PMR1</i> (NGY356)	As CAI4 but <i>pmr1Δ::hisG/pmr1Δ::hisG, RPS1/rps1Δ::Clp10-PMR1</i>	(Bates et al., 2005)
<i>C. albicans</i>	<i>mnt1-mnt2Δ</i> (NGY337)	As CAI4 but <i>mnt1-mnt2Δ::hisG/mnt1-mnt2Δ::hisG, RPS1/rps1Δ::Clp10</i>	(Munro et al., 2005, p. 1)
<i>C. albicans</i>	<i>mnt1-mnt2Δ + MNT1</i> (NGY335)	As CAI4 but <i>mnt1-mnt2Δ::hisG/mnt1-mnt2Δ::hisG, RPS1/rps1Δ::Clp10-MNT1</i>	(Munro et al., 2005)
<i>C. albicans</i>	<i>mnn4Δ</i> (CDH15)	As CAI4 but <i>mnn4Δ::hisG/mnn4Δ::hisG, RPS1/rps1Δ::Clp10</i>	(Hobson et al., 2004)
<i>C. albicans</i>	<i>mnn4Δ + MNN4</i> (CDH13)	As CAI4 but <i>mnn4Δ::hisG/mnn4Δ::hisG, RPS1/rps1Δ::(Clp10-MNN4)_n</i>	(Hobson et al., 2004)
<i>C. dubliniensis</i>	CdUM4A	As Wü284 but <i>ura3Δ1::FRT/ ura3Δ2::FRT</i>	(Staub et al., 2001)
<i>C. dubliniensis</i>	CdWT(NGY568)	As CdUM4A but <i>RPS1/rps1Δ::Clp10</i>	This study
<i>C. dubliniensis</i>	<i>Cdoch1Δ</i> (NGY563)	As CdUM4A but <i>och1Δ::dp1200/och1Δ::dp1200, RPS1/rps1Δ::Clp10</i>	This study
<i>C. dubliniensis</i>	<i>Cdoch1Δ/CdOCH1</i> (NGY564)	As CdUM4A but <i>och1Δ::dp1200/och1Δ::dp1200, RPS1/rps1Δ::Clp10-CdOCH1</i>	This study
<i>C. glabrata</i>	CgWT(2001)	HETS202 strain	(Ueno et al., 2007)
<i>C. glabrata</i>	<i>P_{TET}-CgOCH1</i>	As CgWT but <i>P_{TET}-OCH1</i>	This study
<i>S. cerevisiae</i>	ScWT(BY4741)	<i>S. cerevisiae</i> MATa his3 Δ 1 leu2 Δ 0 met15 Δ 0 ura3 Δ 0	ATCC
<i>S. cerevisiae</i>	<i>Scoch1Δ</i> (Y04406)	As BY4741 but <i>och1Δ::KanMX4</i>	Euroscarf
<i>S. cerevisiae</i>	<i>Scoch1Δ + P_{GALI1}-ScOCH1</i> (NGS7)	Y04406 transformed with <i>pYES2.1/V5-His-TOPO-ScOCH1</i>	This study

CAGTTTTCCAGTCACGACGTT-3' and 5'-TTACTGCATTTCTGGCATA CATCATCTTTCCAACACTACCAGAAACATATGTTTTGCGTATGCCATTGG ATGTGGAATTGTGAGCGGATA-3' (underline bases correspond to complementary sequences to the 5'- and 3'- regions of ORF Cd36_86020 from the *C. dubliniensis* GeneDB, <http://old.genedb.org/genedb/cdubliniensis/index.jsp>) were used to amplify the disruption cassette from pDDB57 plasmid by PCR (Wilson et al., 2000). The strain CdUM4A, a Ura^r mutant derived from the clinical isolate Wü284 (Staub et al., 2001), was sequentially transformed with the disruption cassette and the *URA3* marker recycled by growing on SC medium supplemented with 1 mg/ml 5-fluoroorotic acid and uridine. The plasmid Clp10 was used to restore *URA3* at the *RPS1* locus as previously described (Murad et al., 2000).

Despite repeated attempts, a null mutant for *OCH1* in *C. glabrata* could not be generated, indicating that this gene may be essential for the viability of this species, at least under conditions of transformation. Therefore a conditional mutant was generated by placing the *OCH1* gene under the control of a tetracycline regulatable promoter 97t (Nakayama et al., 1998). The Tet-P cassette was amplified using plasmid pTK916 as a template, with the primer pairs 5'-TCCTGTTCTG AGACCAAATAGCAAACCGAAGCTGGCTTGATACAGTAAATTCA GTGggccgctgatcagc-3' and 5'-AGAATACCACCAGCACCAGCACCGCCA AGCACTATGTGCCTCTTCTGCCATTTACTATcgtgagctgg-3'. This amplicon was used to transform HETS202 strain (Ueno et al., 2007), thus replacing the wild type promoter of *OCH1* in this strain with the conditional Tet-promoter. The mutants were confirmed by PCR and northern analysis (data not shown). *OCH1* expression was repressed in the conditional mutant using 20 µg/ml doxycycline in the growth medium.

2.3. Complementation of *och1Δ* null mutants.

To generate a re-integrant control strain for *C. dubliniensis och1Δ*, the primer pair 5'-GCGGCCGCAAATGAAAAATATTACC TC and 5'-GCGGCCGCTTGTAGATTAAATTTGGATT (with bases added to create a *NotI* site underlined) were used to amplify by PCR a 2907 bpDNA fragment containing the *C. dubliniensis OCH1* ORF plus 995 bp of its promoter and 731 bp of its terminator regions, and the DNA fragment cloned into pCR[®]2.1-TOPO[®] vector (Invitrogen, Paisley, UK). The insert was released by digesting with *NotI*, and subcloned into the *NotI* site of Clp10, generating plasmid Clp10-Cd*OCH1*. The *StuI*-digested plasmid was integrated at the *RPS1* locus generating strain NGY565.

The *S. cerevisiae* optimized, galactose-inducible protein expression vector pYES2.1/V5-His-TOPO (Invitrogen, Paisley, UK) was used to express *S. cerevisiae OCH1* in the *S. cerevisiae och1Δ* null mutant strain. The *S. cerevisiae OCH1* ORF was amplified by PCR (primer pair 5'-ATGTCTAGGAAGTTGTCCACCTGA and 5'-GATGCTGATAAAAT GCAGTCATAAATAA) and ligated into the pYES2.1/V5-His-TOPO vector according to manufacturers' instructions and the construction used to transform *Escherichia coli* TOP10 cells (Invitrogen, Paisley, UK). The construction of pYES2.1/V5-His-TOPO was confirmed by sequencing and used to transform *S. cerevisiae och1Δ* null mutant.

2.4. Analysis of N-linked mannosylation status and cell wall phosphomannan content

The electrophoretic mobility shift assay of secreted β-N-acetylglucosaminidase and the Alcian Blue binding assay were used to assess the N-linked mannosylation status and the phosphomannan content, respectively, as previously described (Bates et al., 2006).

2.5. Fluorescence microscopy

Mannan, chitin and β-glucan staining in the live fungal cell walls were done as previously described with minor adaptations (Hall et al.,

2013).

For mannan staining, 2.5*10⁶ exponentially growing yeast cells were washed three times with sterile PBS. The cells were then stained with 100 µg/ml ConA-Rh for 45 min in dark and washed three times in PBS. The cells were next stained for chitin using 100 µg/ml freshly prepared WGA-FITC for 45 min in dark. After staining, the cells were washed thrice with PBS, fixed in 4% formaldehyde for 1 h and observed under the Zeiss Axioplan 2 microscope (63X objective). The images were analysed using AxioVision ES64 Rel.4.9.1 software.

For β-glucan staining, 2.5*10⁶ exponentially growing yeast cells were washed thrice with sterile PBS and blocked in Buffer A (0.5% BSA, 5.0 mM EDTA, 2.0 mM NaN₃, 5% heat-inactivated goat serum) for 30 min at room temperature. The cells were next washed thrice with Buffer B (Buffer A without goat serum). An aliquot of 0.5 µg of purified Fc-dectin-1, (kind gift from Prof. Gordon Brown), was added to each of the samples and incubated in ice for 1 h. The cells were again washed three times with Buffer B. A 100.0 µl sample of TRITC-conjugated anti-human IgG Fc goat IgG (from Thermofisher Scientific, catalogue number A18822, diluted 1:200 in Buffer B) was added to each of the samples and incubated in ice for 45 min. The cells were next washed five times with Buffer A, fixed in 4% paraformaldehyde, and observed under the Zeiss Axioplan 2 microscope at 63X magnification. The images were analysed using AxioVision ES64 Rel.4.9.1 software.

2.6. Cell wall composition and porosity

Determination of cell wall mannan, chitin, and β-glucan content was achieved by acid-hydrolysing the polymers, and quantifying mannose, glucosamine, and glucose content, respectively, by high-performance anion-exchange chromatography with pulsed amperometric detection as previously described (Mora-Montes et al., 2007; Plaine et al., 2008).

The cell wall porosity was determined by relative porosity to poly-cations as described (De Nobel et al., 1990). Overnight-grown cells in Sabouraud broth were inoculated into fresh broth, further incubated at 30 °C for 4 h with 200 rpm, and washed twice with PBS. Cell pellets containing 1*10⁸ cells were resuspended in 10 mM Tris-HCl, pH 7.4 (buffer A), buffer A plus 15 µg/ml poly-L-lysine (Mw 30–70 kDa, Sigma Cat. No. P2636) or buffer A plus 30 µg/ml DEAE-dextran (Mw 500 kDa, Sigma Cat. No. D9885), and incubated for 30 min at 30 °C, 200 rpm. Preparations were centrifuged to pellet cells, supernatants were collected, further centrifuged and absorbance at 260 nm measured. The relative cell wall porosity to DEAE-dextran was calculated as described (De Nobel et al., 1990).

2.7. Cytokine stimulation using human peripheral blood mononuclear cells (hPBMCs)

For hPBMC isolation, human blood was collected from healthy volunteers according to the local guidelines and regulations, as approved by the College Ethics Review Board of the University of Aberdeen (CERB/2012/11/676 and CERB/2016/8/1300). The isolation of hPBMCs was essentially performed as previously described (Endres et al., 1988). Samples of 5*10⁵ freshly isolated hPBMCs were placed into round-bottom 96-well Nunclon plates (Nunc, Roskilde, Denmark) and stimulated with heat-killed 2*10⁵ yeast cells for 24 h at 37 °C and 5% (v/v) CO₂. For the preparation of glucan-phosphate-treated human PBMCs, cells were incubated with 10.0 µg of glucan-phosphate (kind gift from Prof. David L Williams) for 2 h at 37 °C and 5% (v/v) CO₂ before challenge with fungal cells. Additionally, for understanding the temperature-induced cell wall changes in wild type strains, the hPBMCs were similarly challenged with thimerosal killed fungal cells exposed to different temperatures, as described in the material and methods section. The supernatants were collected after 24 h of challenge and stored at -20 °C till cytokine analysis was performed. Cytokine measurements were carried out by ELISA using commercially available kits from R&D

systems (Abingdon, UK).

2.8. TNF α stimulation using mouse macrophages from the cell line RAW264.7

The RAW264.7 cell line was obtained from the European Collection of Cell Culture. Cells were grown at 37 °C and 5% (v/v) CO₂ in tissue culture flasks (Nagle Nunc. International, Hereford, UK) containing DMEM medium (Lonza Group Ltd, Braine-l'Alleud, Belgium), added with 1% (w/v) L-glutamine (Invitrogen, Paisley, UK), 10% (v/v) heat inactivated foetal calf serum (Biosera, Ringmer, UK), and 2% (w/v) penicillin/streptomycin antibiotics (Invitrogen Ltd, Paisley, UK). For cytokine stimulation, cells were gently and mechanically detached from bottles, and 2*10⁵ cells/well were seeded in a 96-well flat-bottom plate. The cells were left to adhere overnight at 37 °C and 5% (v/v) CO₂. 100 μ l fresh medium was added to the wells and then stimulated with 2*10⁵ heat-killed yeast cells for 24 h at 37 °C and 5% (v/v) CO₂. For blocking with glucan-phosphate-treated, cells were incubated with 10.0 μ g of the glucan-phosphate for 2 h at 37 °C and 5% (v/v) CO₂ before challenge with fungal cells. To analyze the effect of temperature-induced cell wall changes in wild type strains, the RAW macrophages were similarly challenged with thimerosal killed fungal cells exposed to different temperatures, as described in the methods section. Upon stimulation, supernatants were collected as described above and used to quantify murine TNF α by ELISA using kits from R&D.

2.9. Statistical analysis

The cell wall analysis by HPLC was done using two independent cultures, performed in duplicate. Cytokine stimulations using human PMBCs were performed in duplicate with a total of six healthy donors; whereas assays with RAW264.7 macrophages were conducted four times in duplicate. Student *t*-test or one-way ANOVA (with posthoc correction using Tukey test) were used to establish statistical significance of data, as applicable, with a significance level set at $p < 0.05$.

3. Results

3.1. Cell wall structural differences between the fungal species results in their differential immune recognition

We first examined how cell wall differences between the different fungal species affected immune recognition. We chemically fixed yeast cells of *C. albicans*, *C. dubliniensis*, *C. glabrata* and *S. cerevisiae* with thimerosal before exposing them to different temperatures (from 30 °C to 75 °C) for 2 h. Heat treatment is known to increase exposure of the inner cell wall layers and is commonly used in immune assays to prevent unwanted fungal growth (for example yeast-hypha morphogenesis) during immune assays. These treated cells were exposed to hPBMCs as well as RAW264.7 macrophages and cytokine profiles were measured after 24 h.

Interestingly despite the phylogenetic similarity between *C. albicans* and *C. dubliniensis* there was a marked difference in the temperature profile map of the TNF α response induced by yeast cells (Fig. 1). Induced TNF α levels did not vary significantly when the temperature exposure of the cells was increased. TNF α induced by cells of *C. glabrata* and *S. cerevisiae* displayed a similar pattern but the patterns were different to that for *C. albicans* and *C. dubliniensis*, showing a marked increase in TNF α induction in hPBMCs at higher temperatures. A different trend was observed in RAW264.7 macrophages, suggesting underlying differences in the thermostability of the cell wall of these species (Fig. 1). In the case of hPBMCs, patterns similar to TNF α secretion were observed for another cytokine, IL-6, as well (data not shown). IL-6 levels secreted by RAW macrophages were too low to be detected by our assay method. The microscopic assessment of the mannan, chitin or β -

glucan levels with the increase in temperature did not reveal any striking differences in the relative exposure of β -glucan at different temperatures (data not shown). However, the thermal profiling experiments did suggest important underlying differences in the structure of the cell wall that were differentially affected by heat treatment.

3.2. Generation of *C. dubliniensis* and *C. glabrata och1 Δ* null mutants

In order to investigate the impact of outer cell wall *N*-mannosylation defect in different fungal species on the immune recognition, we generated *och1* mutant and re-integrant strains from the four fungal species (Table 1). The *S. cerevisiae och1 Δ* null mutant strain was complemented with a galactose-inducible expression vector to express *S. cerevisiae OCH1* in the *S. cerevisiae och1 Δ* (ScOCH1). The *C. dubliniensis* CdUM4A, a *ura*⁻ strain derived from the clinical isolate Wü284 (Staub et al., 2001), was used as genetic background for the sequential disruption of both *OCH1* alleles, using the mini-ura-blaster strategy (Wilson et al., 2000). The *Cdoch1 Δ* null mutant (NGY562) was transformed with the *Cip10* plasmid (Murad et al., 2000), introducing *URA3* at the neutral *RPS1* locus (Brand et al., 2004). The re-integrant control strain (NGY564) had the *CdOCH1* allele at the *RPS1* locus as described in Methods section. Attempts to disrupt *CgOCH1* in *C. glabrata* ATCC2001 were unsuccessful hence a conditional mutant using the tetracycline-regulatable system was generated.

Similar to *Caoch1 Δ* and *Scoch1 Δ* null mutants, yeast cells of the *Cdoch1 Δ* null mutant and the *Cgoch1* conditional mutant growing in presence of doxycycline were swollen, formed aggregates, and had small and crenulated colonies (data not shown). The defect in the *N*-linked glycosylation of *Cdoch1 Δ* null mutant was assessed by electrophoretic mobility shift of secreted β -*N*-acetylhexosaminidase (HexNAcase) in native gels using an *in situ* activity assay (Bates et al., 2006). HexNAcase is a highly *N*-linked glycosylated enzyme (Molloy et al., 1994), and changes in its electrophoretic mobility have been used as indicators of *N*-linked glycosylation defects which lead to marked decreases in total molecular mass (Bates et al., 2006; Mora-Montes et al., 2010, 2007). The HexNAcase extracted from the null mutant ran faster than the enzyme obtained from wild type cells, while reintegration of *CdOCH1* restored the enzyme mobility to wild type levels (Fig. S1A). Endoglycosidase H-treated HexNAcase from wild type, *Cdoch1 Δ* null mutant and re-integrant control migrated faster and with the similar mobility (Fig. S1B). No HexNAcase activity was detectable in protein preparations from *C. glabrata* and native zymograms for invertase also failed to generate usable electrophoretic mobility shifts (data not shown). Hence, in this case cell wall phosphomannan levels was assessed by Alcian Blue binding to confirm that the *N*-linked glycosylation status of *C. glabrata* was altered in this mutant (Bates et al., 2006; Herrero et al., 2002; Mora-Montes et al., 2007; Nishikawa et al., 2002). *C. glabrata* wild type cells bound 82.7 \pm 5.6 mg Alcian Blue, whereas the *Cgoch1* conditional mutant bound 74.7 \pm 5.4 (P value 0.15). When the mutant cells were grown in presence of 20 μ g/ml doxycycline there was a significant reduction in the ability to bind Alcian Blue (22.2 \pm 2.9 mg dye bound, P < 0.05). Therefore, these data confirmed that both the *Cdoch1 Δ* null mutant and the *Cgoch1* conditional mutant also had defects in *N*-linked glycosylation.

3.3. *och1* mutants from different fungal species have altered cell wall composition and characteristics

The cell wall organisation and sugar composition of the *och1* mutants for all the fungal species, *C. albicans*, *C. dubliniensis*, *C. glabrata*, and *S. cerevisiae*, were assessed by fluorescence microscopy and HPLC respectively. The *och1 Δ* mutants from all the fungal species displayed higher chitin and β -glucan levels compared to their wild type and re-integrant controls as assessed by microscopy and HPLC (Figs. 2 and 3). Disruption of *OCH1* resulted in gross reduction in the outermost *N*-mannan layer in the fungal cells, and exposure of the inner layers of




Fig. 1. TNFα production from hPBMCs and RAW264.7 macrophages upon stimulation with wild type cells: The wild type cells were thimerosal killed and exposed to different temperatures to expose the inner cell wall components. These fungal cells were then used to challenge hPBMCs and RAW macrophages for 24 h and the TNFα levels were monitored using ELISA. For hPBMCs the experiment was done with six donors, three independent fungal cultures, each time done in duplicates. For RAW macrophages, the experiment was repeated four times in duplicate. Data represent means ± SD.

chitin and β-glucan. As expected, all the mutants displayed reduced levels of mannan compared to the controls.

The walls of *S. cerevisiae* and *C. glabrata* were significantly more porous to DEAE-dextran than walls of *C. albicans* and *C. dubliniensis* (Fig. S2). Wall porosity was increased most in *N*-mannan mutants (*och1Δ* and *pmr1Δ*), intermediate in the *O*-mannan double mutant (*mnt1Δ-mnt2Δ*) and unaffected by loss of cell wall phosphomannan (*mnn4Δ*) (Fig. S2). The *och1* conditional mutant of *C. glabrata* grown under repressing conditions did not show any further reduction in wall porosity – most likely because even wild cells were relatively porous

(Fig. S2).

Thus, disruption of *OCH1* in the four fungal species led to similar qualitative changes in the cell wall: reduced *N*-mannan, increased chitin content, increased internal wall exposure and increased cell wall porosity.

3.4. hPBMC's recognize fungal *N*-mannosylation

We demonstrated previously that hPBMCs produce low cytokine levels when stimulated with *Caoch1Δ* null mutant (e.g. Netea et al.,




Fig. 2. Mannan, chitin and β-glucan levels in live fungal cells: Cell wall structure was assessed in live cells by staining using fluorescence microscopy, as described in the methods section. (A) Mannan was stained using ConA-Rh (red), while cell surface chitin was stained using WGA-FITC (green). (B) β-glucan levels were detected using TRITC-conjugated anti-human IgG Fc goat IgG (red). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)




Fig. 3. Cell wall composition analysis by HPLC: The acid hydrolysed cell walls of the strains were analyzed by HPLC to quantify glucosamine, glucose and mannose indicative of chitin, β -glucan and mannan contents respectively. Independent biological replicates were carried out in duplicate. Data represent means \pm SD. * $P < 0.05$, ** $P < 0.005$.

2006). We therefore assessed the impact of *OCH1* disruption in *C. dubliniensis*, *C. glabrata*, and *S. cerevisiae* on the ability to stimulate cytokine production by human monocytes. TNF α stimulation was significantly lower upon co-incubation of hPBMCs with heat-killed *och1 Δ mutants from all the analyzed yeast species (Fig. 4). The levels of IL-6, were also found to be significantly lower in all species (Fig. S3). We showed above that the *och1 Δ mutants from all the four fungal species resulted in higher β -glucan and chitin exposure (Fig. 2). We therefore tested the effect of the four *och1* mutants on immune recognition via dectin-1. TNF α or IL-6 levels were not affected significantly in the presence or absence of the dectin-1 blocker, glucan phosphate (Fig. 4), and thus dectin-1 did not have a major role in the recognition of fungal cells by hPBMCs. We also tested another *N*-glycosylation mutant from *C. albicans*, *mnn2* Δ ⁶, that also had higher β -glucan exposure (Hall et al., 2013). This mutant also displayed reduced cytokine induction in hPBMCs, confirming the importance of *N*-mannan in immune recognition by human monocytes (Fig. 4 and Fig. S3). Chitin represents another important inner cell wall PAMP which has immunomodulatory effects (Elieh Ali Komi et al., 2018) and has been shown to induce the anti-inflammatory cytokine IL-10 (Wagener et al., 2014). To analyze the impact of exposed chitin in *och1 Δ mutants on IL-10 secretion by hPBMC's, the same samples were also analyzed for IL-10 levels. Even***

though chitin levels were enhanced in *och1 Δ mutants of all the four fungal species, IL-10 levels were found to be reduced upon challenge with *och1 Δ mutants (data not shown), as for TNF α and IL-6, indicating a general reduced cytokine induction response by *och1 Δ mutants in hPBMCs. Therefore, even when the outer 'masking' cell wall layer is absent, and the underlying cell wall layers are exposed, human PBMCs recognition of the cell wall was strongly dependent on *N*-linked mannan.***

3.5. Macrophages primarily recognize fungal β -glucan

Next, we assessed the ability of the *och1 Δ mutants from the four fungal species to stimulate TNF α production in mouse macrophage cell line RAW264.7. We compared the cytokine induction by heat-killed wild type, *och1 Δ mutants and *OCH1* re-integrant controls for these four fungal species. Starkly contrasting with the results obtained with hPBMCs, the heat-killed *och1 Δ null mutants from all four fungal species stimulated significantly more TNF α production in RAW macrophages than did the wild type and *OCH1* re-integrant cells (Fig. 5). Also in contrast with the data using monocytes, TNF α induction in glucan phosphate- treated RAW macrophages was significantly reduced upon challenge with all fungal strains (as compared to the controls without***




Fig. 4. TNF α stimulation by *och1* mutants using the hPBMCs: 2×10^5 heat-killed fungal cells were co-incubated with 5×10^5 hPBMCs and TNF α levels were quantified by ELISA after 24 h of stimulation. For blocking dectin-1, hPBMCs were pre-treated with 10.0 μ g glucan phosphate for 2 h, before challenge with fungal cells. The experiment used monocytes from six donors, with three independent biological replicates, each done in duplicate. Data represent means \pm SD. * $P < 0.05$, ** $P < 0.005$, *** $P < 0.001$.

inhibitor) indicating that recognition was dectin-1 dependent (Fig. 5). Interestingly, TNF α induction triggered by the *och1* mutants in the presence of the dectin-1 blocking agent, although reduced, remained high compared to wild type and re-integrant control strains under the same conditions, suggesting other PRRs were also involved in cytokine stimulation.

These data suggest that the absence of *N*-mannosylation leads to a better recognition of fungal cells by RAW264.7 macrophages, due to unmasking of β -glucan in the inner cell wall layer. A similar increase in TNF α production by *och1* mutants from *C. albicans* and *S. cerevisiae* was observed in human monocyte derived macrophages (Fig. S4A) and in the J774.1 murine macrophage cell line (Fig. S4B). The above data underline significant differences at interaction level of PAMPs with PRRs in macrophages and monocytes.

4. Discussion

The cell wall components play a central role in fungal sensing by the innate immune system (Erwig and Gow, 2016; Netea et al., 2015). The

fungal kingdom displays large heterogeneity and diversity in cell wall structures but most walls consist of identifiable outer and inner cell wall layers (Erwig and Gow, 2016; Garcia-Rubio et al., 2019). β -(1,3)-glucan is a conserved component of most fungal walls and is a structural polymer residing predominately in the inner cell wall layer. β -(1,3)-glucan is a strong immune agonist but it is normally covered by outer wall components and therefore is not immediately accessible to its cognate PRR dectin-1. Therefore components of the outer cell wall have been considered as an impediment to host immune recognition by shielding or masking for the inner cell wall β -(1,3)-glucan. An example is the shielding effect of outer cell wall α -(1,3)-glucan of inner cell wall β -(1,3)-glucan in *H. capsulatum* (Rappleye et al., 2007; Rappleye and Goldman, 2008). In *Candida* species the fibrillar outer cell wall mannan layer that represents 30–40% of the wall mass has also been proposed to have this β -(1,3)-glucan shielding function (Bain et al., 2014; Ballou et al., 2016; Pradhan et al., 2019, 2018; Wheeler et al., 2008; Wheeler and Fink, 2006). Although cell wall mannan undoubtedly covers and obscures the inner cell wall PAMPs, β -(1,3)-glucan and chitin, multiple PRRs are potentially involved in mannan recognition including




Fig. 5. TNF α stimulation by *och1* Δ mutants using the murine macrophage cell line RAW264.7: 2×10^5 RAW macrophages and 2×10^5 heat-killed yeast cells were co-incubated for 24 h at 37 °C, and TNF α concentration was determined by ELISA. For blocking dectin-1, RAW macrophages were pre-treated with 10.0 μ g glucan phosphate for 2 h, before challenge with fungal cells. The experiment had four biological replicates. Data represent means \pm SD. * P < 0.05, ** P < 0.005, *** P < 0.001.

mannose receptor (MR), dectin-2, dectin-3, mincle, DC-SIGN, galectin-3, Fc γ R, CD14, CD23, TLR2, TLR4 and TLR6 (Erwig and Gow, 2016; Vendele et al., 2020) and simultaneous recognition of β -(1,3)-glucan and mannans can result in co-receptor amplification of immune responses (Dennehy et al., 2008). Therefore the superficial mannan layer has the potential to trigger immunity and amplify the inflammatory response as β -(1,3)-glucan recognition also becomes engaged.

N-mannan is the major cell wall protein post-translational modification of outer cell wall proteins and is a significant fraction of the total mass of the *Candida* cell wall. In the elaboration of *N*-mannan, Och1 is a conserved α -(1,6)-mannosyltransferase, that catalyses the addition of first α -(1,6)-mannose residue to the conserved core triantennary *N*-glycan structure (Hall and Gow, 2013). Mutants of Och1 display a severely truncated *N*-glycan that results in exposure of the inner cell wall layers. *C. albicans och1* Δ null mutant only stimulated about 20% of the cytokine levels in hPBMCs compared to the wild type control cells, stressing the importance of *N*-linked mannans for *C. albicans* immune recognition (Netea et al., 2006). In contrast, the *S. cerevisiae och1* Δ null mutant was reported to elicit higher levels of TNF α production in RAW264.7 macrophages than the wild type control

(Wheeler and Fink, 2006). This marked difference in the phenotype of the *och1* mutant in these yeasts could suggest that the mannan structures in *S. cerevisiae* are different from *C. albicans*. However, although there are differences in the cell walls of these two yeasts, we show here that the more likely explanation in this disparity lies in the immune cell deployed in these reports, and not the fungus. We demonstrate that although there are cell wall architectural differences between different yeast species such as wall porosity that could impact the immune response elicited, the apparent discrepancy can be explained by the fact that RAW264.7 macrophages predominantly recognize β -glucan and not *N*-mannan, whereas cytokine induction by hPBMCs is dependent on *N*-linked mannans. This observation was further supported by extending the analysis to two other *Candida* species – *C. glabrata* and *C. dubliniensis* (Butler et al., 2009; Jackson et al., 2009). For all four fungi, TNF α induction was reduced in the *och1* mutant when using human monocytes but was increased when using RAW and other macrophage cell types.

Human PBMCs express most of the well characterised PRRs involved in fungal sensing, including dectin-1, TLR2, TLR4, and mannose receptor (Ferwerda et al., 2008; Netea et al., 2008; van de Veerdonk et al.,

2009). Human PBMCs stimulated with *C. albicans*, *C. dubliniensis*, *C. glabrata* or *S. cerevisiae och1Δ* mutants induced less cytokine production than the wild type controls, indicating that even if the inner cell wall PAMPs β -glucan and chitin were exposed and available for recognition by PRRs, *N*-linked mannan recognition was required for maximum cytokine stimulation. This observation is in line with other reports indicating that *N*-linked mannans are required for fungal recognition by innate immune cells (Cambi et al., 2008; Keppeler-Ross et al., 2010; Mora-Montes et al., 2010, 2007; Netea et al., 2006; van de Veerdonk et al., 2009).

Macrophages have a heterogeneous receptor expression and express most of the PRRs (Brown, 2011). Our results support the view that fungal β -glucan is the primary functional PAMP perceived by this cell line, hence *och1Δ* mutants that have more exposed β -(1,3)-glucan induce higher TNF α levels than the parent wild type. This recognition and TNF α induction was significantly reduced by pre-treatment of the RAW264.7 macrophages with glucan phosphate, a specific blocker of the major β -glucan receptor, dectin-1. In a previous study, macrophage migration had been shown to be enhanced towards *C. albicans* glycosylation mutants with exposed β -glucan, again suggesting better recognition of these mutants by macrophages (Lewis et al., 2012).

Inflammatory monocytes have been shown to have an essential and protective role in the first 48 h post fungal infection (Ngo et al., 2014). While in circulation, monocytes recognize fungi using a combination of TLRs and CTLs. Upon differentiation into macrophages, while expression of TLRs is maintained, there is an upregulation in the expression of CTLs (Netea et al., 2008). Thus, these results highlight the role of monocytes during the early fungal infection stage, when β -glucan is masked by the outer mannan layer (Wheeler et al., 2008). As the infection progresses, β -glucan is increasingly exposed (Hopke et al., 2018) and thus the dectin-1 mediated recognition of fungal cells by macrophages becomes increasingly relevant and important.

Several studies have demonstrated the cross-talk between the CLR and TLRs and have shown that PRRs work co-operatively and synergistically in PAMP recognition. Dectin-1 collaboration with TLR2 and TLR4 upregulates cytokine production upon stimulation (Dennehy et al., 2008; Ferwerda et al., 2008; Gantner et al., 2003). Dectin-1 has also been shown to operate in conjunction with galectin-3 and this interaction has been shown to increase TNF α production by macrophages in response to β -(1,3)-glucan (Esteban et al., 2011).

Cooperative interactions of PRRs have been shown in multiple contexts. For example, cytokine production by human monocytes and macrophages was shown to be dependent on both *N*- and *O*-linked mannans (mediated by MR and TLR4, respectively), and was reduced when the fungal cells had either *N*- or *O*-mannosylation defect (Netea et al., 2006). Dectin-2 forms heterodimers with dectin-3 and these heterodimers were more potent stimulators of cytokine production than the homodimers (Zhu et al., 2013). Galectin-3 association with TLR2 is required for signalling upon recognition of *C. albicans* (Jouault et al., 2006).

A comparison of the expression of different PRRs in various macrophage and monocyte populations using Immunological Genome Project (Heng et al., 2008) showed that macrophage populations display a higher expression of dectin-1 and TLR-4 as compared to the monocytes. Hence a synergistic response from dectin-1 recognition of exposed β -(1,3)-glucan and TLR-4 recognition of *O*-mannan in *och1Δ* mutants could be responsible for higher cytokine production in macrophages. The macrophages also present a higher dectin-2 and dectin-3 expression compared to monocytes (Heng et al., 2008). Dectin-2 recognizes the core mannan structure (Vendele et al., 2020), which is exposed in *och1Δ* mutants, and hence a cross-talk between these two receptors could also contribute to an augmented cytokine response to *och1Δ* mutants.

To conclude, we observe that monocytes recognize fungal cells predominantly through *N*-mannan of the outer *Candida* cell wall, in contrast to RAW264.7 and other primary macrophages that primarily

recognize β -(1,3)-glucan in the inner cell wall layer. These observations reconcile apparent contradictions in the literature about the importance of *N*-mannan which serves both as a shield of β -(1,3)-glucan recognition and an activator of a range of mannan detecting PRRs depending on the type of immune cell being tested. This highlights the importance of the type of immune cells chosen while studying fungal immune responses in general.

CRedit authorship contribution statement

Bhawna Yadav: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Validation, Visualization, Writing - original draft, Writing - review & editing. **Héctor M. Mora-Montes:** Investigation, Methodology, Writing - review & editing. **Jeanette Wagener:** Data curation, Formal analysis, Investigation, Methodology, Writing - review & editing. **Iain Cunningham:** Investigation, Methodology, Writing - review & editing. **Lara West:** Data curation, Formal analysis, Writing - review & editing. **Ken Haynes:** Funding acquisition, Project administration, Conceptualization. **Alistair J.P. Brown:** Conceptualization, Writing - review & editing. **Neil A.R. Gow:** Conceptualization, Funding acquisition, Project administration, Resources, Validation, Visualization, Writing - original draft, Writing - review & editing.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank Professor Gordon Brown for Fc-dectin-1 and Professor David Williams for glucan phosphate. We also thank Kevin MacKenzie, Debbie Wilkinson, Gillian Milne, and Lucy Wright at the University of Aberdeen Core Microscopy & Histology Facility.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.tcs.2020.100042>.

References

- Bain, J.M., Lewis, L.E., Okai, B., Quinn, J., Gow, N.A.R., Erwig, L.-P., 2012. Non-lytic expulsion/exocytosis of *Candida albicans* from macrophages. *Fungal Genet. Biol.* **49**, 677–678. <https://doi.org/10.1016/j.fgb.2012.01.008>.
- Bain, J.M., Louw, J., Lewis, L.E., Okai, B., Walls, C.A., Ballou, E.R., Walker, L.A., Reid, D., Munro, C.A., Brown, A.J.P., Brown, G.D., Gow, N.A.R., Erwig, L.P., 2014. *Candida albicans* hypha formation and mannan masking of β -glucan inhibit macrophage phagosome maturation. *mBio* **5**, e01874. doi: 10.1128/mBio.01874-14.
- Ballou, E.R., Avelar, G.M., Childers, D.S., Mackie, J., Bain, J.M., Wagener, J., Kastora, S.L., Panea, M.D., Hardison, S.E., Walker, L.A., Erwig, L.P., Munro, C.A., Gow, N.A.R., Brown, G.D., MacCallum, D.M., Brown, A.J.P., 2016. Lactate signalling regulates fungal β -glucan masking and immune evasion. *Nat. Microbiol.* **2**, 16238. <https://doi.org/10.1038/nmicrobiol.2016.238>.
- Bates, S., Hughes, H.B., Munro, C.A., Thomas, W.P.H., MacCallum, D.M., Bertram, G., Atrih, A., Ferguson, M.A.J., Brown, A.J.P., Odds, F.C., Gow, N.A.R., 2006. Outer chain N-glycans are required for cell wall integrity and virulence of *Candida albicans*. *J. Biol. Chem.* **281**, 90–98. <https://doi.org/10.1074/jbc.M510360200>.
- Bates, S., MacCallum, D.M., Bertram, G., Munro, C.A., Hughes, H.B., Buurman, E.T., Brown, A.J.P., Odds, F.C., Gow, N.A.R., 2005. *Candida albicans* Pmr1p, a secretory pathway P-type Ca²⁺/Mn²⁺-ATPase, is required for glycosylation and virulence. *J. Biol. Chem.* **280**, 23408–23415. <https://doi.org/10.1074/jbc.M502162200>.
- Borriello, F., Zononi, I., Granucci, F., 2020. Cellular and molecular mechanisms of anti-fungal innate immunity at epithelial barriers: The role of C-type lectin receptors. *Eur. J. Immunol.* **50**, 317–325. <https://doi.org/10.1002/eji.201848054>.
- Bowman, S.M., Free, S.J., 2006. The structure and synthesis of the fungal cell wall. *BioEssays News Rev. Mol. Cell. Dev. Biol.* **28**, 799–808. <https://doi.org/10.1002/bies.20441>.
- Brand, A., MacCallum, D.M., Brown, A.J.P., Gow, N.A.R., Odds, F.C., 2004. Ectopic expression of URA3 can influence the virulence phenotypes and proteome of *Candida*

- albacans but can be overcome by targeted reintegration of URA3 at the RPS10 locus. *Eukaryot. Cell* 3, 900–909. <https://doi.org/10.1128/EC.3.4.900-909.2004>.
- Brown, A.J.P., Brown, G.D., Netea, M.G., Gow, N.A.R., 2014. Metabolism impacts upon Candida immunogenicity and pathogenicity at multiple levels. *Trends Microbiol.* 22, 614–622. <https://doi.org/10.1016/j.tim.2014.07.001>.
- Brown, G.D., 2011. Innate antifungal immunity: the key role of phagocytes. *Annu. Rev. Immunol.* 29, 1–21. <https://doi.org/10.1146/annurev-immunol-030409-101229>.
- Brown, G.D., Gordon, S., 2001. Immune recognition. A new receptor for beta-glucans. *Nature* 413, 36–37. <https://doi.org/10.1038/35092620>.
- Butler, G., Rasmussen, M.D., Lin, M.F., Santos, M.A.S., Sakthikumar, S., Munro, C.A., Rheinbay, E., Grabherr, M., Forche, A., Reedy, J.L., Agrafioti, I., Arnaud, M.B., Bates, S., Brown, A.J.P., Brunke, S., Costanzo, M.C., Fitzpatrick, D.A., de Groot, P.W.J., Harris, D., Hoyer, L.L., Hube, B., Klis, F.M., Kodira, C., Lennard, N., Logue, M.E., Martin, R., Neiman, A.M., Nikolaou, E., Quail, M.A., Quinn, J., Santos, M.C., Schmitzberger, F.F., Sherlock, G., Shah, P., Silverstein, K.A.T., Skrzypek, M.S., Soll, D., Staggs, R., Stansfield, I., Stumpf, M.P.H., Sudbery, P.E., Srikantha, T., Zeng, Q., Berman, J., Berriman, M., Heitman, J., Gow, N.A.R., Lorenz, M.C., Birren, B.W., Kellis, M., Cuomo, C.A., 2009. Evolution of pathogenicity and sexual reproduction in eight *Candida* genomes. *Nature* 459, 657–662. <https://doi.org/10.1038/nature08064>.
- Cambi, A., Netea, M.G., Mora-Montes, H.M., Gow, N.A.R., Hato, S.V., Lowman, D.W., Kullberg, B.-J., Torensma, R., Williams, D.L., Figdor, C.G., 2008. Dendritic cell interaction with *Candida albicans* critically depends on N-linked mannans. *J. Biol. Chem.* 283, 20590–20599. <https://doi.org/10.1074/jbc.M709334200>.
- Cottier, F., Hall, R.A., 2019. Face/off: the interchangeable side of *Candida Albicans*. *Front. Cell. Infect. Microbiol.* 9, 471. <https://doi.org/10.3389/fcimb.2019.00471>.
- da Silva Dantas, A., Lee, K.K., Raziunaite, I., Schaefer, K., Wagener, J., Yadav, B., Gow, N.A., 2016. Cell biology of *Candida albicans*-host interactions. *Curr. Opin. Microbiol.* 34, 111–118. <https://doi.org/10.1016/j.mib.2016.08.006>.
- Dambuzza, I.M., Brown, G.D., 2015. C-type lectins in immunity: recent developments. *Curr. Opin. Immunol.* 32, 21–27. <https://doi.org/10.1016/j.coi.2014.12.002>.
- De Nobel, J.G., Klis, F.M., Munnik, T., Priem, J., van den Ende, H., 1990. An assay of relative cell wall porosity in *Saccharomyces cerevisiae*, *Kluyveromyces lactis* and *Schizosaccharomyces pombe*. *Yeast Chichester Engl.* 6, 483–490. <https://doi.org/10.1002/yea.320060605>.
- Dennehy, K.M., Ferwerda, G., Faro-Trindade, I., Pyz, E., Willment, J.A., Taylor, P.R., Kerrigan, A., Tsoni, S.V., Gordon, S., Meyer-Wentrup, F., Adema, G.J., Kullberg, B.-J., Schweighoffer, E., Tybulewicz, V., Mora-Montes, H.M., Gow, N.A.R., Williams, D.L., Netea, M.G., Brown, G.D., 2008. Syk kinase is required for collaborative cytokine production induced through Dectin-1 and Toll-like receptors. *Eur. J. Immunol.* 38, 500–506. <https://doi.org/10.1002/eji.200737741>.
- Drummond, R.A., Gaffen, S.L., Hise, A.G., Brown, G.D., 2014. Innate defense against fungal pathogens. *Cold Spring Harb. Perspect. Med.* 5. <https://doi.org/10.1101/cshperspect.a019620>.
- Elieh Ali Komi, D., Sharma, L., Dela Cruz, C.S., 2018. Chitin and Its Effects on Inflammatory and Immune Responses. *Clin. Rev. Allergy Immunol.* 54, 213–223. doi: 10.1007/s12016-017-8600-0.
- Endres, S., Ghorbani, R., Lonnemann, G., van der Meer, J.W., Dinarello, C.A., 1988. Measurement of immunoreactive interleukin-1 beta from human mononuclear cells: optimization of recovery, intrasubject consistency, and comparison with interleukin-1 alpha and tumor necrosis factor. *Clin. Immunol. Immunopathol.* 49, 424–438.
- Erwig, L.P., Gow, N.A.R., 2016. Interactions of fungal pathogens with phagocytes. *Nat. Rev. Microbiol.* 14, 163–176. <https://doi.org/10.1038/nrmicro.2015.21>.
- Esteban, A., Popp, M.W., Vyas, V.K., Strijbis, K., Ploegh, H.L., Fink, G.R., 2011. Fungal recognition is mediated by the association of dectin-1 and galectin-3 in macrophages. *Proc. Natl. Acad. Sci. U.S.A.* 108, 14270–14275. <https://doi.org/10.1073/pnas.1111415108>.
- Fabre, E., Hurtaux, T., Fradin, C., 2014. Mannosylation of fungal glycoconjugates in the Golgi apparatus. *Curr. Opin. Microbiol.* 20, 103–110. <https://doi.org/10.1016/j.mib.2014.05.008>.
- Ferwerda, G., Meyer-Wentrup, F., Kullberg, B.-J., Netea, M.G., Adema, G.J., 2008. Dectin-1 synergizes with TLR2 and TLR4 for cytokine production in human primary monocytes and macrophages. *Cell. Microbiol.* 10, 2058–2066. <https://doi.org/10.1111/j.1462-5822.2008.01188.x>.
- Fonzi, W.A., Irwin, M.Y., 1993. Isogenic strain construction and gene mapping in *Candida albicans*. *Genetics* 134, 717–728.
- Fukazawa, Y., 1989. Antigenic structure of *Candida albicans*. *Immunochemical basis of the serologic specificity of the mannans in yeasts. Immunol. Ser.* 47, 37–62.
- Gantner, B.N., Simmons, R.M., Canavera, S.J., Akira, S., Underhill, D.M., 2003. Collaborative induction of inflammatory responses by dectin-1 and Toll-like receptor 2. *J. Exp. Med.* 197, 1107–1117. <https://doi.org/10.1084/jem.20021787>.
- Garcia-Rubio, R., de Oliveira, H.C., Rivera, J., Trevijano-Contador, N., 2019. The fungal cell wall: *Candida*, *Cryptococcus*, and *Aspergillus* species. *Front. Microbiol.* 10, 2993. <https://doi.org/10.3389/fmicb.2019.02993>.
- Garfoot, A.L., Shen, Q., Wüthrich, M., Klein, B.S., Rappleye, C.A., 2016. The Eng1 β -glucanase enhances histoplasma virulence by reducing β -glucan exposure. *mBio* 7, e01388–1315. <https://doi.org/10.1128/mBio.01388-15>.
- Gow, N.A.R., Hube, B., 2012. Importance of the *Candida albicans* cell wall during commensalism and infection. *Curr. Opin. Microbiol.* 15, 406–412. <https://doi.org/10.1016/j.mib.2012.04.005>.
- Gow, N.A.R., Latgé, J.-P., Munro, C.A., 2017. The fungal cell wall: structure, biosynthesis, and function. *Microbiol. Spectr.* 5. <https://doi.org/10.1128/microbiolspec.FUNK-0035-2016>.
- Gow, N.A.R., Netea, M.G., Munro, C.A., Ferwerda, G., Bates, S., Mora-Montes, H.M., Walker, L., Jansen, T., Jacobs, L., Tsoni, V., Brown, G.D., Odds, F.C., Van der Meer, J.W.M., Brown, A.J.P., Kullberg, B.J., 2007. Immune recognition of *Candida albicans* beta-glucan by dectin-1. *J. Infect. Dis.* 196, 1565–1571. <https://doi.org/10.1086/523110>.
- Gow, N.A.R., van de Veerdonk, F.L., Brown, A.J.P., Netea, M.G., 2012. *Candida albicans* morphogenesis and host defence: discriminating invasion from colonization. *Nat. Rev. Microbiol.* 10, 112–122. <https://doi.org/10.1038/nrmicro2711>.
- Gow, N.A.R., Yadav, B., 2017. Microbe Profile: *Candida albicans*: a shape-changing, opportunistic pathogenic fungus of humans. *Microbiol. Read. Engl.* 163, 1145–1147. <https://doi.org/10.1099/mic.0.000499>.
- Graus, M.S., Wester, M.J., Lowman, D.W., Williams, D.L., Kruppa, M.D., Martinez, C.M., Young, J.M., Pappas, H.C., Lidke, K.A., Neumann, A.K., 2018. Mannan molecular substructures control nanoscale glucan exposure in *Candida*. *Cell Rep.* 24, 2432–2442.e5. <https://doi.org/10.1016/j.celrep.2018.07.088>.
- Hall, R.A., 2015. Dressed to impress: impact of environmental adaptation on the *Candida albicans* cell wall. *Mol. Microbiol.* 97, 7–17. <https://doi.org/10.1111/mmi.13020>.
- Hall, R.A., Bates, S., Lenardon, M.D., Maccallum, D.M., Wagener, J., Lowman, D.W., Kruppa, M.D., Williams, D.L., Odds, F.C., Brown, A.J.P., Gow, N.A.R., 2013. The Mnn2 mannosyltransferase family modulates mannoprotein fibril length, immune recognition and virulence of *Candida albicans*. *PLoS Pathog.* 9, e1003276. <https://doi.org/10.1371/journal.ppat.1003276>.
- Hall, R.A., Gow, N.A.R., 2013. Mannosylation in *Candida albicans*: role in cell wall function and immune recognition. *Mol. Microbiol.* 90, 1147–1161. <https://doi.org/10.1111/mmi.12426>.
- Heng, T.S.P., Painter, M.W., Immunological Genome Project Consortium, 2008. The Immunological Genome Project: networks of gene expression in immune cells. *Nat. Immunol.* 9, 1091–1094. doi: 10.1038/ni1008-1091.
- Hernández-Chávez, M.J., Pérez-García, L.A., Niño-Vega, G.A., Mora-Montes, H.M., 2017. Fungal Strategies to Evade the Host Immune Recognition. *J. Fungi Basel Switz.* 3. <https://doi.org/10.3390/jof3040051>.
- Herrero, A.B., Uccelletti, D., Hirschberg, C.B., Dominguez, A., Abejón, C., 2002. The Golgi GDPase of the fungal pathogen *Candida albicans* affects morphogenesis, glycosylation, and cell wall properties. *Eukaryot. Cell* 1, 420–431.
- Hobson, R.P., Munro, C.A., Bates, S., MacCallum, D.M., Cutler, J.E., Heinsbroek, S.E.M., Brown, G.D., Odds, F.C., Gow, N.A.R., 2004. Loss of cell wall mannosylphosphate in *Candida albicans* does not influence macrophage recognition. *J. Biol. Chem.* 279, 39628–39635. <https://doi.org/10.1074/jbc.M405003200>.
- Hopke, A., Brown, A.J.P., Hall, R.A., Wheeler, R.T., 2018. Dynamic fungal cell wall architecture in stress adaptation and immune evasion. *Trends Microbiol.* 26, 284–295. <https://doi.org/10.1016/j.tim.2018.01.007>.
- Jackson, A.P., Gamble, J.A., Yeomans, T., Moran, G.P., Saunders, D., Harris, D., Aslett, M., Barrell, J.F., Butler, G., Citiulo, F., Coleman, D.C., de Groot, P.W.J., Goodwin, T.J., Quail, M.A., McQuillan, J., Munro, C.A., Pain, A., Poulter, R.T., Rajandream, M.-A., Renaud, H., Spiering, M.J., Tivey, A., Gow, N.A.R., Barrell, B., Sullivan, D.J., Berriman, M., 2009. Comparative genomics of the fungal pathogens *Candida dubliniensis* and *Candida albicans*. *Genome Res.* 19, 2231–2244. <https://doi.org/10.1101/gr.097501.109>.
- Jouault, T., El Abed-El Behi, M., Martínez-Esparza, M., Breuilh, L., Trinel, P.-A., Chamailard, M., Trottein, F., Poulain, D., 2006. Specific recognition of *Candida albicans* by macrophages requires galectin-3 to discriminate *Saccharomyces cerevisiae* and needs association with TLR2 for signaling. *J. Immunol. Baltim. Md 1950 (177)*, 4679–4687.
- Jouault, T., Ibata-Ombetta, S., Takeuchi, O., Trinel, P.-A., Sacchetti, P., Lefebvre, P., Akira, S., Poulain, D., 2003. *Candida albicans* phospholipomannan is sensed through toll-like receptors. *J. Infect. Dis.* 188, 165–172. <https://doi.org/10.1086/375784>.
- Kepler-Ross, S., Douglas, L., Konopka, J.B., Dean, N., 2010. Recognition of yeast by murine macrophages requires mannan but not glucan. *Eukaryot. Cell* 9, 1776–1787. <https://doi.org/10.1128/EC.00156-10>.
- Klis, F.M., de Groot, P., Hellingwerf, K., 2001. Molecular organization of the cell wall of *Candida albicans*. *Med. Mycol.* 39 (Suppl. 1), 1–8.
- Kozel, T.R., MacGill, R.S., Percival, A., Zhou, Q., 2004. Biological activities of naturally occurring antibodies reactive with *Candida albicans* mannan. *Infect. Immun.* 72, 209–218. <https://doi.org/10.1128/iai.72.1.209-218.2004>.
- Latgé, J.-P., 2007. The cell wall: a carbohydrate armour for the fungal cell. *Mol. Microbiol.* 66, 279–290. <https://doi.org/10.1111/j.1365-2958.2007.05872.x>.
- Levitz, S.M., 2010. Innate recognition of fungal cell walls. *PLoS Pathog.* 6, e1000758. <https://doi.org/10.1371/journal.ppat.1000758>.
- Lewis, L.E., Bain, J.M., Lowes, C., Gillespie, C., Rudkin, F.M., Gow, N.A.R., Erwig, L.-P., 2012. Stage specific assessment of *Candida albicans* phagocytosis by macrophages identifies cell wall composition and morphogenesis as key determinants. *PLoS Pathog.* 8, e1002578. <https://doi.org/10.1371/journal.ppat.1002578>.
- Lowman, D.W., Greene, R.R., Bearden, D.W., Kruppa, M.D., Pottier, M., Monteiro, M.A., Soldatov, D.V., Ennsley, H.E., Cheng, S.-C., Netea, M.G., Williams, D.L., 2014. Novel structural features in *Candida albicans* hyphal glucan provide a basis for differential innate immune recognition of hyphae versus yeast. *J. Biol. Chem.* 289, 3432–3443. <https://doi.org/10.1074/jbc.M113.529131>.
- McKenzie, C.G.J., Koser, U., Lewis, L.E., Bain, J.M., Mora-Montes, H.M., Barker, R.N., Gow, N. a. R., Erwig, L.P., 2010. Contribution of *Candida albicans* cell wall components to recognition by and escape from murine macrophages. *Infect. Immun.* 78, 1650–1658. doi: 10.1128/IAI.00001-10.
- Molloy, C., Cannon, R.D., Sullivan, P.A., Shepherd, M.G., 1994. Purification and characterization of two forms of N-acetylglucosaminidase from *Candida albicans* showing widely different outer chain glycosylation. *Microbiol. Read. Engl.* 140 (Pt 7), 1543–1553. <https://doi.org/10.1099/13500872-140-7-1543>.
- Mora-Montes, H.M., Bates, S., Netea, M.G., Castillo, L., Brand, A., Buurman, E.T., Díaz-Jiménez, D.F., Jan Kullberg, B., Brown, A.J.P., Odds, F.C., Gow, N.A.R., 2010. A multifunctional mannosyltransferase family in *Candida albicans* determines cell wall mannan structure and host-fungus interactions. *J. Biol. Chem.* 285, 12087–12095.

- <https://doi.org/10.1074/jbc.M109.081513>.
- Mora-Montes, H.M., Bates, S., Netea, M.G., Díaz-Jiménez, D.F., López-Romero, E., Zinker, S., Ponce-Noyola, P., Kullberg, B.J., Brown, A.J.P., Odds, F.C., Flores-Carreón, A., Gow, N.A.R., 2007. Endoplasmic reticulum alpha-glycosidases of *Candida albicans* are required for N glycosylation, cell wall integrity, and normal host-fungus interaction. *Eukaryot. Cell* 6, 2184–2193. <https://doi.org/10.1128/EC.00350-07>.
- Mora-Montes, H.M., Ponce-Noyola, P., Villagómez-Castro, J.C., Gow, N.A., Flores-Carreón, A., López-Romero, E., 2009. Protein glycosylation in *Candida*. *Future Microbiol.* 4, 1167–1183. <https://doi.org/10.2217/fmb.09.88>.
- Mukaremera, L., Lee, K.K., Mora-Montes, H.M., Gow, N.A.R., 2017. *Candida albicans* yeast, pseudohyphal, and hyphal morphogenesis differentially affects immune recognition. *Front. Immunol.* 8, 629. <https://doi.org/10.3389/fimmu.2017.00629>.
- Munro, C.A., Bates, S., Buurman, E.T., Hughes, H.B., Maccallum, D.M., Bertram, G., Atri, A., Ferguson, M.A.J., Bain, J.M., Brand, A., Hamilton, S., Westwater, C., Thomson, L.M., Brown, A.J.P., Odds, F.C., Gow, N.A.R., 2005. Mnt1p and Mnt2p of *Candida albicans* are partially redundant alpha-1,2-mannosyltransferases that participate in O-linked mannosylation and are required for adhesion and virulence. *J. Biol. Chem.* 280, 1051–1060. <https://doi.org/10.1074/jbc.M411413200>.
- Murad, A.M., Lee, P.R., Broadbent, I.D., Barelle, C.J., Brown, A.J., 2000. Clp10, an efficient and convenient integrating vector for *Candida albicans*. *Yeast* 16, 325–327. [https://doi.org/10.1002/1097-0061\(20000315\)16:4<325::AID-YEA538>3.0.CO;2-#](https://doi.org/10.1002/1097-0061(20000315)16:4<325::AID-YEA538>3.0.CO;2-#).
- Nakayama, H., Izuta, M., Nagahashi, S., Sihta, E.Y., Sato, Y., Yamazaki, T., Arisawa, M., Kitada, K., 1998. A controllable gene-expression system for the pathogenic fungus *Candida glabrata*. *Microbiol. Read. Engl.* 144 (Pt 9), 2407–2415. <https://doi.org/10.1099/00221287-144-9-2407>.
- Netea, M.G., Brown, G.D., Kullberg, B.J., Gow, N.A.R., 2008. An integrated model of the recognition of *Candida albicans* by the innate immune system. *Nat. Rev. Microbiol.* 6, 67–78. <https://doi.org/10.1038/nrmicro1815>.
- Netea, M.G., Gow, N.A.R., Munro, C.A., Bates, S., Collins, C., Ferwerda, G., Hobson, R.P., Bertram, G., Hughes, H.B., Jansen, T., Jacobs, L., Buurman, E.T., Gijzen, K., Williams, D.L., Torensma, R., McKinnon, A., MacCallum, D.M., Odds, F.C., Van der Meer, J.W.M., Brown, A.J.P., Kullberg, B.J., 2006. Immune sensing of *Candida albicans* requires cooperative recognition of mannans and glucans by lectin and Toll-like receptors. *J. Clin. Invest.* 116, 1642–1650. <https://doi.org/10.1172/JCI27114>.
- Netea, M.G., Joosten, L.A.B., van der Meer, J.W.M., Kullberg, B.-J., van de Veerdonk, F.L., 2015. Immune defence against *Candida* fungal infections. *Nat. Rev. Immunol.* 15, 630–642. <https://doi.org/10.1038/nri3897>.
- Netea, M.G., Van Der Graaf, C.A.A., Vonk, A.G., Verschuere, I., Van Der Meer, J.W.M., Kullberg, B.J., 2002. The role of toll-like receptor (TLR) 2 and TLR4 in the host defense against disseminated candidiasis. *J. Infect. Dis.* 185, 1483–1489. <https://doi.org/10.1086/340511>.
- Ngo, L.Y., Kasahara, S., Kumasaka, D.K., Knoblauch, S.E., Jhingran, A., Hohl, T.M., 2014. Inflammatory monocytes mediate early and organ-specific innate defense during systemic candidiasis. *J. Infect. Dis.* 209, 109–119. <https://doi.org/10.1093/infdis/jit413>.
- Nikolakopoulou, C., Willment, J.A., Brown, G.D., 2020. C-Type Lectin Receptors in Antifungal Immunity. *Adv. Exp. Med. Biol.* 1204, 1–30. https://doi.org/10.1007/978-981-15-1580-4_1.
- Nishikawa, A., Poster, J.B., Jigami, Y., Dean, N., 2002. Molecular and phenotypic analysis of CaVRG4, encoding an essential Golgi apparatus GDP-mannose transporter. *J. Bacteriol.* 184, 29–42.
- Patin, E.C., Thompson, A., Orr, S.J., 2019. Pattern recognition receptors in fungal immunity. *Semin. Cell Dev. Biol.* 89, 24–33. <https://doi.org/10.1016/j.semcdb.2018.03.003>.
- Plaine, A., Walker, L., Da Costa, G., Mora-Montes, H.M., McKinnon, A., Gow, N.A.R., Gaillardin, C., Munro, C.A., Richard, M.L., 2008. Functional analysis of *Candida albicans* GPI-anchored proteins: roles in cell wall integrity and caspofungin sensitivity. *Fungal Genet. Biol.* 45, 1404–1414. <https://doi.org/10.1016/j.fgb.2008.08.003>.
- Plato, A., Hardison, S.E., Brown, G.D., 2015. Pattern recognition receptors in antifungal immunity. *Semin. Immunopathol.* 37, 97–106. <https://doi.org/10.1007/s00281-014-0462-4>.
- Pradhan, A., Avelar, G.M., Bain, J.M., Childers, D., Pelletier, C., Larcombe, D.E., Shekhova, E., Netea, M.G., Brown, G.D., Erwig, L., Gow, N.A.R., Brown, A.J.P., 2019. Non-canonical signalling mediates changes in fungal cell wall PAMPs that drive immune evasion. *Nat. Commun.* 10, 5315. <https://doi.org/10.1038/s41467-019-13298-9>.
- Pradhan, A., Avelar, G.M., Bain, J.M., Childers, D.S., Larcombe, D.E., Netea, M.G., Shekhova, E., Munro, C.A., Brown, G.D., Erwig, L.P., Gow, N.A.R., Brown, A.J.P., 2018. Hypoxia Promotes Immune Evasion by Triggering β -Glucan Masking on the *Candida albicans* Cell Surface via Mitochondrial and cAMP-Protein Kinase A Signaling. *mBio* 9. doi: 10.1128/mBio.01318-18.
- Qin, Y., Zhang, L., Xu, Z., Zhang, J., Jiang, Y.-Y., Cao, Y., Yan, T., 2016. Innate immune cell response upon *Candida albicans* infection. *Virulence* 7, 512–526. <https://doi.org/10.1080/21505594.2016.1138201>.
- Rappleye, C.A., Eissenberg, L.G., Goldman, W.E., 2007. Histoplasma capsulatum alpha-(1,3)-glucan blocks innate immune recognition by the beta-glucan receptor. *Proc. Natl. Acad. Sci. U.S.A.* 104, 1366–1370. <https://doi.org/10.1073/pnas.0609848104>.
- Rappleye, C.A., Goldman, W.E., 2008. Fungal stealth technology. *Trends Immunol.* 29, 18–24. <https://doi.org/10.1016/j.it.2007.10.001>.
- Rogers, N.C., Slack, E.C., Edwards, A.D., Nolte, M.A., Schulz, O., Schweighoffer, E., Williams, D.L., Gordon, S., Tybulewicz, V.L., Brown, G.D., Reis e Sousa, C., 2005. Syk-dependent cytokine induction by Dectin-1 reveals a novel pattern recognition pathway for C type lectins. *Immunity* 22, 507–517. <https://doi.org/10.1016/j.immuni.2005.03.004>.
- Saijo, S., Ikeda, S., Yamabe, K., Kakuta, S., Ishigame, H., Akitsu, A., Fujikado, N., Kusaka, T., Kubo, S., Chung, S., Komatsu, R., Miura, N., Adachi, Y., Ohno, N., Shibuya, K., Yamamoto, N., Kawakami, K., Yamasaki, S., Saito, T., Akira, S., Iwakura, Y., 2010. Dectin-2 recognition of alpha-mannans and induction of Th17 cell differentiation is essential for host defense against *Candida albicans*. *Immunity* 32, 681–691. <https://doi.org/10.1016/j.immuni.2010.05.001>.
- Salazar, F., Brown, G.D., 2018. Antifungal innate immunity: a perspective from the last 10 years. *J. Innate Immun.* 10, 373–397. <https://doi.org/10.1159/000488539>.
- Shibata, N., Arai, M., Haga, E., Kikuchi, T., Najima, M., Satoh, T., Kobayashi, H., Suzuki, S., 1992. Structural identification of an epitope of antigenic factor 5 in mannans of *Candida albicans* NIH B-792 (serotype B) and J-1012 (serotype A) as beta-1,2-linked oligomannosyl residues. *Infect. Immunol.* 60, 4100–4110.
- Snarr, B.D., Qureshi, S.T., Sheppard, D.C., 2017. Immune recognition of fungal polysaccharides. *J. Fungi Basel Switz.* 3. <https://doi.org/10.3390/jof3030047>.
- Staib, P., Moran, G.P., Sullivan, D.J., Coleman, D.C., Morschhäuser, J., 2001. Isogenic strain construction and gene targeting in *Candida dubliniensis*. *J. Bacteriol.* 183, 2859–2865. <https://doi.org/10.1128/JB.183.9.2859-2865.2001>.
- Taylor, P.R., Brown, G.D., Herre, J., Williams, D.L., Willment, J.A., Gordon, S., 2004. The role of SIGNR1 and the beta-glucan receptor (dectin-1) in the nonopsonic recognition of yeast by specific macrophages. *J. Immunol. Baltim. Md* 1950 (172), 1157–1162.
- Ueno, K., Uno, J., Nakayama, H., Sasamoto, K., Mikami, Y., Chibana, H., 2007. Development of a highly efficient gene targeting system induced by transient repression of YKU80 expression in *Candida glabrata*. *Eukaryot. Cell* 6, 1239–1247. <https://doi.org/10.1128/EC.00414-06>.
- van de Veerdonk, F.L., Kullberg, B.J., van der Meer, J.W.M., Gow, N.A.R., Netea, M.G., 2008. Host-microbe interactions: innate pattern recognition of fungal pathogens. *Curr. Opin. Microbiol.* 11, 305–312. <https://doi.org/10.1016/j.mib.2008.06.002>.
- van de Veerdonk, F.L., Marijnissen, R.J., Kullberg, B.J., Koenen, H.J.P.M., Cheng, S.-C., Joosten, I., van den Berg, W.B., Williams, D.L., van der Meer, J.W.M., Joosten, L.A.B., Netea, M.G., 2009. The macrophage mannose receptor induces IL-17 in response to *Candida albicans*. *Cell Host Microbe* 5, 329–340. <https://doi.org/10.1016/j.chom.2009.02.006>.
- Vendele, I., Willment, J.A., Silva, L.M., Palma, A.S., Chai, W., Liu, Y., Feizi, T., Spyrou, M., Stappers, M.H.T., Brown, G.D., Gow, N.A.R., 2020. Mannan detecting C-type lectin receptor probes recognise immune epitopes with diverse chemical, spatial and phylogenetic heterogeneity in fungal cell walls. *PLoS Pathog.* 16, e1007927. <https://doi.org/10.1371/journal.ppat.1007927>.
- Wagener, J., Malireddi, R.K.S., Lenardon, M.D., Köberle, M., Vautier, S., MacCallum, D.M., Biedermann, T., Schaller, M., Netea, M.G., Kannekanti, T.-D., Brown, G.D., Brown, A.J.P., Gow, N.A.R., 2014. Fungal chitin dampens inflammation through IL-10 induction mediated by NOD2 and TLR9 activation. *PLoS Pathog.* 10, e1004050. <https://doi.org/10.1371/journal.ppat.1004050>.
- Walker, L., Sood, P., Lenardon, M.D., Milne, G., Olson, J., Jensen, G., Wolf, J., Casadevall, A., Adler-Moore, J., Gow, N.A.R., 2018. The Viscoelastic Properties of the Fungal Cell Wall Allow Traffic of AmBisome as Intact Liposome Vesicles. *mBio* 9. doi: 10.1128/mBio.02383-17.
- West, L., Lowman, D.W., Mora-Montes, H.M., Grubb, S., Murdoch, C., Thornhill, M.H., Gow, N.A.R., Williams, D., Haynes, K., 2013. Differential virulence of *Candida glabrata* glycosylation mutants. *J. Biol. Chem.* 288, 22006–22018. <https://doi.org/10.1074/jbc.M113.478743>.
- Wheeler, R.T., Fink, G.R., 2006. A drug-sensitive genetic network masks fungi from the immune system. *PLoS Pathog.* 2, e35. <https://doi.org/10.1371/journal.ppat.0020035>.
- Wheeler, R.T., Kombe, D., Agarwala, S.D., Fink, G.R., 2008. Dynamic, morphotype-specific *Candida albicans* beta-glucan exposure during infection and drug treatment. *PLoS Pathog.* 4, e1000227. <https://doi.org/10.1371/journal.ppat.1000227>.
- Wilson, R.B., Davis, D., Enloe, B.M., Mitchell, A.P., 2000. A recyclable *Candida albicans* URA3 cassette for PCR product-directed gene disruptions. *Yeast* 16, 65–70. [https://doi.org/10.1002/\(SICI\)1097-0061\(20000115\)16:1<65::AID-YEA508>3.0.CO;2-M](https://doi.org/10.1002/(SICI)1097-0061(20000115)16:1<65::AID-YEA508>3.0.CO;2-M).
- Yamamoto, Y., Klein, T.W., Friedman, H., 1997. Involvement of mannose receptor in cytokine interleukin-1beta (IL-1beta), IL-6, and granulocyte-macrophage colony-stimulating factor responses, but not in chemokine macrophage inflammatory protein 1beta (MIP-1beta), MIP-2, and KC responses, caused by attachment of *Candida albicans* to macrophages. *Infect. Immunol.* 65, 1077–1082.
- Zhang, S.Q., Zou, Z., Shen, H., Shen, S.S., Miao, Q., Huang, X., Liu, W., Li, L.P., Chen, S.M., Yan, L., Zhang, J.D., Zhao, J.J., Xu, G.T., An, M.M., Jiang, Y.-Y., 2016. Mnn10 maintains pathogenicity in *Candida albicans* by extending α -1,6-mannose backbone to evade host dectin-1 mediated antifungal immunity. *PLoS Pathog.* 12, e1005617. <https://doi.org/10.1371/journal.ppat.1005617>.
- Zhu, L.-L., Zhao, X.-Q., Jiang, C., You, Y., Chen, X.-P., Jiang, Y.-Y., Jia, X.-M., Lin, X., 2013. C-type lectin receptors Dectin-3 and Dectin-2 form a heterodimeric pattern-recognition receptor for host defense against fungal infection. *Immunity* 39, 324–334. <https://doi.org/10.1016/j.immuni.2013.05.017>.