

**How to cite:** Rodrigo Ramirez-Campillo, Daniel Castillo, Javier Raya-González, Jason Moran, Eduardo Sáez de Villarreal, Rhodri S. Lloyd. Effects of plyometric jump training on jump and sprint performance in young male soccer players: a systematic review and meta-analysis. **Sports Medicine**. In press.

**Short Title:** Plyometric jump training in soccer

Rodrigo Ramirez-Campillo<sup>1</sup>, Daniel Castillo<sup>2</sup>, Javier Raya-González<sup>2</sup>, Jason Moran<sup>3</sup>, Eduardo Sáez de Villarreal<sup>4</sup>, Rhodri S. Lloyd<sup>5,6,7</sup>

<sup>1</sup> Human Performance Laboratory. Quality of Life and Wellness Research Group. Department of Physical Activity Sciences. Universidad de Los Lagos. Osorno, Chile.

<sup>2</sup> Faculty of Health Sciences. Universidad Isabel I, Burgos, Spain.

<sup>3</sup> School of Sport, Rehabilitation and Exercise Sciences, University of Essex, Colchester, UK.

<sup>4</sup> Universidad Pablo de Olavide, Physical Performance & Sports Research Center, Seville, Spain.

<sup>5</sup> Youth Physical Development Centre, Cardiff School of Sport and Health Sciences, Cardiff Metropolitan University, Cardiff, UK.

<sup>6</sup> Sport Performance Research Institute, New Zealand (SPRINZ), AUT University, Auckland, New Zealand.

<sup>7</sup> Centre for Sport Science and Human Performance, Waikato Institute of Technology, Hamilton, New Zealand.

### **Corresponding author**

Rodrigo Ramirez-Campillo, PhD. Human Performance Laboratory. Department of Physical Activity Sciences. Universidad de Los Lagos. Osorno, Chile. Mail: r.ramirez@ulagos.cl

ORCID: Rodrigo Ramirez-Campillo (0000-0003-2035-3279); Daniel Castillo (0000-0002-4159-6475); Javier Raya-González (0000-0002-3570-7159); Jason Moran; Eduardo Sáez de Villarreal; Rhodri S. Lloyd.

### **Key Points**

-Jumping and sprinting are key physical fitness proxies of soccer performance for young players.

-Plyometric jump training may improve both jumping and sprinting performance.

-Plyometric jump training may be particularly effective after interventions >7 weeks.

## Abstract

**Background:** Even from a young age, modern soccer requires high levels of physical fitness development, particularly jumping and sprinting. Plyometric jump training (PJT), combined with young athletes' regular soccer sessions, has the potential to improve jumping and sprinting. However, studies exploring the effects of PJT are generally limited by small sample sizes. This problem of underpowered studies may thus be resolved by pooling study results in a meta-analysis.

**Objective:** The objective of this systematic review with meta-analysis (SRMA) was to assess the effects of plyometric jump training (PJT) on jumping and sprinting among young male soccer players.

**Methods:** The SRMA included peer-reviewed articles that incorporated PJT in healthy players (i.e., <23 years of age), a control group, and a measure of jumping or sprinting. Means and standard deviations of outcomes were converted to Hedges' *g* effect sizes (ES), using the inverse variance random-effects model. Moderator analyses were conducted for PJT duration, frequency, total number of sessions, participants' chronological age, and FIFA age categories (i.e., U-17 vs. U-20 vs. U-23). A multivariate random effects meta-regression was also conducted.

**Results:** Thirty-three studies were included, comprising 1,499 participants. PJT improved vertical jump tests (ES = 0.60-0.98; all  $p < 0.01$ ) and linear sprint performance (ES = 0.60-0.98;  $p < 0.03$ ). Interventions of >7 weeks and >14 PJT sessions induced greater effects compared to PJT with  $\leq 7$  weeks and  $\leq 14$  total sessions on 10-m sprint performance (between-group  $p = 0.038$ ).

**Conclusion.** Therefore, PJT is effective in improving jumping and sprinting performance among young male soccer players. Greater 10-m linear sprinting improvements were noted after interventions >7 weeks duration and >14 sessions, suggesting a greater return from exposure to longer PJT interventions, partially in support for the adoption of a long-term approach to athletic development in young athletes. However, with reference to the findings of the meta-regression, and those from the remaining subgroup and single factors analysis, a robust confirmation regarding the moderator role of participant's age, or PJT configuration effects on young soccer player's fitness qualities needs future confirmation.

## 1 Introduction

Even from a young age, modern soccer requires high levels of physical fitness development [1-3]. Although aerobic physical fitness is important during a soccer game [2], maximal or near-maximal intensity single-bout efforts are key factors for optimal physical performance [1, 4, 5]. Therefore, aside from endurance activity, soccer players must also perform numerous explosive actions [6], including jumping, kicking, accelerating, decelerating and changing of movement direction, with most of these preceding goal-scoring opportunities in competitive leagues [4, 7]. Specifically, the straight sprint (45%) and the vertical jump (16%) have been shown to be the two most frequent actions preceding goal situations in soccer [4]. Moreover, a significant relationship between team average for vertical jump height and the final league standing of teams has been observed [1]. Furthermore, youth elite and sub-elite players were shown to jump higher and run faster than non-elite [8, 9], whereas future international and professional players had superior jump and speed ability at youth level than future amateur players [10]. Therefore, jump and sprint-related actions may not only be important qualities at youth-soccer level [8, 9], but also at a later stage of a player's career [10]. Because of this, jumping and straight sprint qualities should be developed at an early age to help players to cope with the increased competitive demands of modern play [11, 12]. On this basis, the investigation of methods to improve jumping and straight sprint actions in young soccer players is essential to optimize on-field performance.

It has previously been shown that the inclusion of plyometric jump training (PJT), combined with young athletes' regular soccer sessions, has the potential to improve many components of physical fitness [13], and may even reduce the risk of sustaining injuries [14]. A PJT program is characterized by exercises that utilize the stretch-shortening cycle of the musculotendinous unit [15, 16]. Typically, PJT exercises can be conducted with short (<250 ms; fast stretch-shortening cycle) or long duration (>250 ms; long stretch-shortening cycle) ground contact times [17-19]. Regarding PJT's effects on sprinting and jumping in young soccer players, previous research in U-17 male soccer players showed that PJT can substantially improve these physical fitness traits after eight weeks of exercise [20]. Similar benefits have also been shown in U-20 [21] and U-23 [22] male soccer players after six weeks of PJT. However, not all PJT studies corroborate these findings. For example, among U-17 [23] male soccer players, six weeks of PJT did not facilitate a significant improvement in sprinting or jumping performance; while, among U-20 [24] male soccer players, 8 weeks of PJT did not induce a significant improvement in sprinting. Moreover, among U-23 [25] male soccer players, six weeks of PJT did not induce a significant improvement in sprinting nor jumping performance. Indeed, in the last two studies [24, 25], it was noted that a significant reduction in sprinting and jumping performance occurred after PJT. Such contrasting findings may be related to several factors, such as the methodological characteristics of PJT interventions (e.g., duration, intensity), participants' characteristics (e.g., initial fitness level) [26, 27] or the sprint or jump testing protocols [28, 29]. Moreover, a common limitation of PJT interventions, which could limit conclusive recommendations on prescription for sprinting and jumping performance, is the commonly low number of participants included in PJT interventions [26, 27]. Indeed, from 420 articles analyzed in a previous PJT scoping review, an average of 10 participants per group was noted [27]. In this way, studies exploring the effects of PJT are generally limited by small sample sizes, affecting the generalizability of the results. This problem of underpowered studies may thus be resolved by pooling study results in a meta-analysis.

Specifically, by pooling the results of several primary studies, the overall statistical power facilitates the drawing of more robust conclusions on the effectiveness of PJT on sprinting and jumping among young soccer players. However, to our knowledge, only one systematic review with meta-analysis (SRMA) has been conducted regarding the effects of PJT on sprinting and jumping among soccer players [30]. This SRMA [30] included only adult soccer players and no moderator analyses were incorporated (e.g., effects

of PJT according to duration, frequency or total number of PJT sessions), thus limiting knowledge of factors that could affect the main effect. Further, in some of the analysed outcomes (i.e., 15-m sprint) very few studies were included (n=2) [30], precluding a robust conclusion regarding the effects of PJT on sprint performance among soccer players. Furthermore, although a previous systematic review [13] analysed the effects of PJT on the physical fitness of young soccer players, no meta-analysis was conducted. Additionally, both male and female soccer players aged between 10 to 17 years were included in the aforementioned systematic review [13]. As males and females experience different effects from PJT according to sex-specific maturational development [31-37], the pooling of the sexes in this way can be misleading.

Given the increased scientific awareness of the relevance of PJT, evidenced by a 25-fold increase in PJT-related scientific publications between the years of 2000-2020, the contrasting findings among PJT interventions, and the typically small sample sizes used in intervention studies, a contemporary SRMA on the topic is warranted. Therefore, the aim of this SRMA was to assess the effects of PJT on jumping and sprinting among young male soccer players. Considering the beneficial effects of PJT on physical fitness in adult female [38] and male soccer players [30], and in athletes from other team sports similar to soccer in terms of intermittent profile and requirements of power expression, such as handball [39] and volleyball [40], we hypothesised PJT would exert beneficial effects on jumping and sprinting among young male soccer players.

## **2 Methods**

A SRMA was conducted following the guidelines of the Cochrane Collaboration [41]. Findings were reported in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) [42].

### **2.1 Eligibility criteria**

The *a priori* inclusion criteria for this SRMA were the following: i) studies that incorporated a PJT programme of at least two weeks in duration, defined as “lower body unilateral or bilateral bounds, jumps and hops that commonly utilise a pre-stretch or countermovement that incites usage of the stretch–shortening cycle” [26, 43, 44], ii) cohorts of healthy young male soccer players, aged 23 years or less (as per FIFA competitions and tournaments regulations according to player’s age) [45], iii) a control group (including active controls) of young male soccer players, iv) a measure of physical fitness (i.e., jumping, sprinting) that was selected based on a logically defensible rationale [4, 44, 46], usually with a high measurement reliability (ICC >0.75; CV <8%) [47, 48]. Trials that included combined training (e.g. PJT and strength training) were included when the control group included the same training, except for the PJT component. Only peer-reviewed articles were included in this SRMA. Articles were excluded if they were cross-sectional, a review, or a training-related study not focused on the effect of PJT exercises. Description of the study selection process is detailed in the results section of the manuscript. Briefly, we also excluded retrospective studies, prospective studies, studies in which the use of jump exercises was not clearly described, studies for which only the abstract was available, case reports, studies with ambiguous study protocols, non-human investigations, special communications, repeated-bout effect interventions, letters to the editor, invited commentaries, errata, overtraining studies, and detraining studies. In the case of detraining studies, if there was a training period prior to a detraining period, the study was considered for inclusion. Non-English language studies were not explored [26].

## 2.2 Search strategy

The PubMed, MEDLINE, Web of Science, and SCOPUS electronic databases were searched from inception until 9 December 2019. Keywords were collected through experts' opinion, a systematic literature review, and controlled vocabulary (e.g., Medical Subject Headings: MeSH). Boolean search syntax using the operators “AND”, “OR” was applied. The words “ballistic”, “complex”, “explosive”, “force-velocity”, “plyometric”, “stretch-shortening cycle”, “jump”, “training”, “male”, “men”, “football”, and “soccer” were used. An example of a PubMed search is: ("randomized controlled trial"[Publication Type]) OR "controlled clinical trial"[Publication Type] OR "randomized"[Title/Abstract] OR "trial"[Title] OR "clinical trials as topic"[MeSH Major Topic] AND "soccer"[Title/Abstract] AND "training"[Title/Abstract] OR "plyometric"[Title/Abstract]. After an initial search, accounts were created in the respective databases. Through these accounts, the lead investigator received automatically generated emails for updates regarding the search terms used. These updates were received on a daily basis (if available), and studies were eligible for inclusion until June 2020. Following the main systematic searches, additional hand-searches were conducted. This SRMA was approved by \*\*\*blind for review purposes\*\*\*.

## 2.3 Study selection and data collection process

In selecting studies for inclusion, a review of all relevant article titles was conducted before an examination of article abstracts and then full-published articles. Two authors conducted the process independently. Potential discrepancies between the two reviewers about study conditions were resolved by consensus with a third author. Full-text articles excluded, with reasons, were recorded. Data were extracted from gathered articles by two authors independently (JRG, RRC), using a form created in Microsoft Excel (Microsoft Corporation, Redmond, WA, USA).

## 2.4 Data items

For the current SRMA, vertical jumping (i.e., vertical height) and linear sprint (i.e., time to complete different distances) performance were chosen as the main outcomes based on establishing a degree of consistency between analysed studies. We sought to analyse different jumping actions and sprint distances as they are considered as separate indicators of fitness relevant to soccer performance, particularly at youth level where maturational changes are taking place [12, 49, 50]. Extracted data also included the following information: quality of PJT treatment description (e.g., well described versus insufficiently described), type of control, type of randomization, number of participants per group. In addition, sample demographics, including age (years), body mass (kg), height (m), fitness level, and previous experience with PJT were extracted. Regarding PJT characteristics, extracted data also included the frequency of training (days/week), duration of training (weeks), and number of total jumps completed during the intervention. A complete description of the aforementioned PJT characteristics have been previously published [26, 27].

## 2.5 Methodological quality in primary studies

The Physiotherapy Evidence Database (PEDro) scale was used to assess the methodological quality of eligible studies included in the SRMA. This scale evaluates internal study validity on a scale from 0 (high risk of bias) to 10 (low risk of bias). As in previous PJT meta-analyses [51, 52], the study quality assessment was interpreted using the following 10-point scale:  $\leq 3$  points was considered poor quality, 4–5 points as moderate quality, and 6–10 points as high quality. Two independent reviewers (JRG, DC), performed this

process. Disagreements in the rating of the studies between the reviewers were resolved through discussion and consensus with a third author. Agreement between reviewers was assessed using a Kappa correlation for risk of bias. The a priori agreement rate between reviewers was set at  $k \geq 0.8$ . If trials had already been assessed and listed on the PEDro database (or similar sources), these scores were adopted. However, methodological quality was not an inclusion criterion. Moreover, the Cochrane Collaboration has previously discouraged the use of these scales, stating that the practice is not underpinned by empirical evidence and assessment criteria may apply inaccurate study weights [53]. In this sense, the subjectivity of personal opinion may undermine the accuracy of such scales.

## 2.6 Summary measures

For analysis and interpretation of results, meta-analyses were conducted if at least three studies provided baseline and follow-up data for the same parameter [44, 54, 55]. Means and standard deviations for a measure (jumping; sprinting) of pre-post-intervention were converted to Hedges'  $g$  effect size (ES) for between-group comparisons. An example (including equations) for Hedges'  $g$  ES calculation for between-group comparisons is provided in *Electronic Supplementary Material Table S1*. The inverse variance random-effects model for meta-analyses was used because it allocates a proportionate weight to trials based on the size of their individual standard errors [56] and facilitates analysis while accounting for heterogeneity across studies [57]. In this sense, the likelihood approach with random effects was used to better account for the inaccuracy in the estimate of between-study variance [58]. The ESs were presented alongside 95% confidence intervals (CIs). The calculated ES were interpreted using the thresholds outlined for standardised mean difference:  $<0.2$ , trivial;  $0.2-0.6$ , small;  $>0.6-1.2$ , moderate;  $>1.2-2.0$ , large;  $>2.0-4.0$ , very large;  $>4.0$ , extremely large [59]. In some studies in which there was more than one intervention group, the control group was proportionately divided to facilitate comparison across all participants [60]. All meta-analyses were carried out using the Comprehensive Meta-Analysis program (version 2; Biostat, Englewood, NJ, USA). Comparisons between the control and the experimental groups for the mean, median, and inter-quartile range (IQR) relative change in a given outcome were calculated from the studies raw data using a form created in Microsoft Excel (Microsoft Corporation, Redmond, WA, USA).

## 2.7 Synthesis of results

To gauge the degree of heterogeneity amongst the included studies, the percentage of total variation across the studies due to heterogeneity (Cochran's  $Q$ -statistic) [61] was used to calculate the  $I^2$  statistic. This represents the proportion of effects that are due to heterogeneity as opposed to chance [42]. Low, moderate and high levels of heterogeneity correspond to  $I^2$  values of  $<25\%$ ,  $25-75\%$ , and  $>75\%$ , respectively [61, 62]. However, these thresholds are considered tentative [61]. The Chi square test assesses if any observed differences in results are compatible with chance alone. A low  $p$  value, or a large Chi square statistic relative to its degree of freedom, provides evidence of heterogeneity of intervention effects beyond those attributed to chance [56].

## 2.8 Risk of bias across studies

Risk of bias across studies was assessed using the extended Egger's test [63]. Sensitivity analyses were conducted to assess the robustness of the summary estimates in order to determine whether a particular study accounted for the heterogeneity. Thus, in order to examine the effects of each result from each study on the overall findings, results were analysed with each study deleted from the

model once. It is acknowledged that other factors, such as differences in trial quality or true study heterogeneity, could produce asymmetry.

## 2.9 Additional analyses

To assess the potential effects of moderator variables, subgroup analyses were performed. Using a random-effects model, potential sources of heterogeneity likely to influence the effects of training were selected a priori. Using the median split technique [64-66], the moderator variables of programme duration (i.e.,  $\leq 7$  vs.  $> 7$  weeks), training frequency (i.e.,  $\leq 2$  vs.  $> 2$  sessions per week), and total number of training sessions (i.e.,  $\leq 14$  vs.  $> 14$  sessions), were chosen based on the accepted influence of these variables on adaptations to exercise [64, 65, 67], in addition to participants' chronological age (i.e.,  $\leq 13.2$  years vs.  $> 13.2$  years). Additionally, FIFA age categories (i.e., U-17 vs. U-20 vs. U-23) were also considered as potential moderator variables. Meta-analyses stratification by each of these factors was performed, with a  $p$  value of  $< 0.05$  considered as the threshold for statistical significance.

## 2.10 Meta-regression

A multivariate random effects meta-regression was conducted to verify if any of the training variables (i.e. frequency, duration, and total number of sessions) or participant's age predicted the effects of PJT on measures of physical fitness. According to the Cochrane handbook for systematic reviews, computation of meta-regression was performed with at least ten studies per covariate [41].

## 3 Results

Figure 1 provides a graphical schematization of the study selection process. Through database searching, 7,859 records were initially identified, and 33 [20, 22, 68-98] were considered in the meta-analysis. The included studies provided mean and standard deviation pre-post-intervention data for at least one main outcome. The included studies comprised 48 individual experimental groups and 752 participants (747 in the control groups).

\*\*\*Figure 1\*\*\*

### 3.1 Study characteristics

The basic characteristics of the participants and the programming parameters of the PJT interventions from the included studies are displayed in Table 1.

\*\*\*Table 1\*\*\*

### 3.2 Methodological quality of included studies

The 33 included studies achieved 6–7 points (i.e., high quality) (Table 2).

\*\*\*Table 2\*\*\*

### 3.3 Meta-analysis results for vertical jump performance

#### 3.3.1 Countermovement jump

From the included studies, 21 provided data for countermovement jump height performance, involving 30 experimental groups. The relative weight of each study in the analysis varied between 1.8% and 4.6%. Of note, in the sensitivity analysis, the results remained consistent ( $p < 0.05$ ) across all deletions. There was a significant favouring of PJT for increase in countermovement jump height performance (ES = 0.79 [95% CI = 0.56 to 1.02],  $p < 0.001$ ) (*Electronic Supplementary Material Fig. S1*). A moderate heterogeneity ( $I^2 = 68.7\%$ ) was observed, and the Egger's test indicated  $p = 0.002$ . After the trim and fill method the adjusted values indicated a point estimate of ES = 0.88 (95% CI = 0.65 to 1.12). Compared to the control groups, the mean relative improvement in the PJT groups was 8.6% (median = 7.9; IQR = 4.2 – 12.1).

#### 3.3.2 Countermovement jump with arms

From the included studies, 6 provided data for countermovement jump with arms height performance, involving 12 experimental groups. The relative weight of each study in the analysis varied between 5.7% and 12.8%. Of note, in the sensitivity analysis, the results remained consistent ( $p < 0.05$ ) across all deletions. There was a significant favouring of PJT for increase in countermovement jump with arms height performance (ES = 0.48 [95% CI = 0.25 to 0.71],  $p < 0.001$ ) (*Electronic Supplementary Material Fig. S2*). A low heterogeneity ( $I^2 = 9.2\%$ ) was observed, and the Egger's test indicated  $p = 0.869$ . Compared to the control groups, the mean relative improvement in the PJT groups was 7.9% (median = 7.0; IQR = 5.9 – 9.1).

#### 3.3.3 Squat jump

From the included studies, 9 provided data for squat jump height performance, involving 10 experimental groups. The relative weight of each study in the analysis varied between 8.7% and 12.5%. Of note, in the sensitivity analysis, the results remained consistent ( $p < 0.05$ ) across all deletions. There was a significant favouring of PJT for increase in squat jump height performance (ES = 0.73 [95% CI = 0.29 to 1.16],  $p = 0.001$ ) (*Electronic Supplementary Material Fig. S3*). A high heterogeneity ( $I^2 = 77.4\%$ ) was observed, and the Egger's test indicated  $p = 0.241$ . After we removed either the Sedano et al. study [94] or the Vaczi et al. study [97] from the analysis, although the significant effect of PJT remained ( $p < 0.001$  to 0.003), the heterogeneity was reduced (70.3 to 71.8%). Compared to the control groups, the mean relative improvement in the PJT groups was 8.2% (median = 7.6; IQR = 4.9 – 12.9).

### 3.4 Meta-analysis results for linear sprint performance

#### 3.4.1 5-m linear sprint

From the included studies, 6 provided data for 5-m linear sprint performance, involving 6 experimental groups. The relative weight of each study in the analysis varied between 15.4% and 17.3%. Of note, in the sensitivity analysis, the results remained consistent ( $p < 0.05$ ) across all deletions. There was a significant favouring of PJT for increase in 5-m linear sprint performance (ES = 0.98 [95% CI = 1.83 to 0.13],  $p = 0.024$ ) (*Electronic Supplementary Material Fig. S4*). A high heterogeneity ( $I^2 = 82.1\%$ ) was observed, and the



Egger's test indicated  $p = 0.296$ . After we removed one study from the analysis [81], although the significant effect of PJT remained ( $p < 0.001$ ), the heterogeneity was reduced to 31.3%. Compared to the control groups, the mean relative improvement in the PJT groups was 7.5% (median = 11.7; IQR = 6.6 – 12.7).

#### 3.4.2 10-m linear sprint

From the included studies, 10 provided data for 10-m linear sprint performance, involving 12 experimental groups. The relative weight of each study in the analysis varied between 7.1% and 9.8%. Of note, in the sensitivity analysis, the results remained consistent ( $p < 0.05$ ) across all deletions. There was a significant favouring of PJT for increase in 10-m linear sprint performance (ES = 0.60 [95%CI = 1.04 to 0.17],  $p = 0.007$ ) (*Electronic Supplementary Material Fig. S5*). A moderate heterogeneity ( $I^2 = 70.1%$ ) was observed, and the Egger's test indicated  $p = 0.280$ . Compared to the control groups, the mean relative improvement in the PJT groups was 2.8% (median = 1.7; IQR = 0.1 – 3.7).

#### 3.4.3 20-m linear sprint

From the included studies, 14 provided data for 20-m linear sprint performance, involving 21 experimental groups. The relative weight of each study in the analysis varied between 2.9% and 6.2%. Of note, in the sensitivity analysis, the results remained consistent ( $p < 0.05$ ) across all deletions. There was a significant favouring of PJT for increase in 20-m linear sprint performance (ES = 0.62 [95%CI = 0.90 to 0.33],  $p < 0.001$ ) (*Electronic Supplementary Material Fig. S6*). A moderate heterogeneity ( $I^2 = 73.3%$ ) was observed, and the Egger's test indicated  $p = 0.425$ . Compared to the control groups, the mean relative improvement in the PJT groups was 4.8% (median = 4.5; IQR = 2.3 – 6.0).

#### 3.4.4 30-m linear sprint

From the included studies, 10 provided data for 30-m linear sprint performance, involving 16 experimental groups. The relative weight of each study in the analysis varied between 4.5% and 8.8%. Of note, in the sensitivity analysis, the results remained consistent ( $p < 0.05$ ) across all deletions. There was a significant favouring of PJT for increase in 30-m linear sprint performance (ES = 0.64 [95%CI = 0.89 to 0.39],  $p < 0.001$ ) (*Electronic Supplementary Material Fig. S7*). A moderate heterogeneity ( $I^2 = 37.1%$ ) was observed, and the Egger's test indicated  $p = 0.679$ . Compared to the control groups, the mean relative improvement in the PJT groups was 3.6% (median = 4.1; IQR = 1.6 – 5.2).

#### 3.4.5 40-m linear sprint

From the included studies, 4 provided data for 40-m linear sprint performance, involving 4 experimental groups. The relative weight of each study in the analysis varied between 22.0% and 26.7%. Of note, in the sensitivity analysis, the results remained consistent ( $p > 0.05$ ) across all deletions. There was a no significant favouring of PJT for increase in 40-m linear sprint performance (ES = 0.94 [95%CI = 1.95 to -0.08],  $p = 0.070$ ) (*Electronic Supplementary Material Fig. S8*). A high heterogeneity ( $I^2 = 81.8%$ ) was observed, and the Egger's test indicated  $p = 0.162$ . After we removed one study from the analysis [71], although the non-significant effect of PJT remained ( $p = 0.072$ ), the heterogeneity was reduced to 0.0%. Compared to the control groups, the mean relative improvement in the PJT groups was 1.6% (median = 1.8; IQR = 1.3 – 1.9).

### 3.5 Additional meta-analyses for vertical jump and linear sprint performance

Regarding interventions with  $\leq 14$  total PJT sessions, also comprising a duration of  $\leq 7$  weeks (5 study groups; ES = 0.11 [95%CI = 0.65 to -0.42],  $p = 0.677$ ; within-group  $I^2 = 39.7\%$ ) and those with  $>14$  sessions, also comprising a duration of  $>7$  weeks (7 study groups; ES = 0.93 [95%CI = 1.47 to 0.38],  $p = 0.001$ ; within-group  $I^2 = 71.9\%$ ), only the latter induced a significant improvement on 10-m linear sprint performance, with a significant between-group difference ( $p = 0.038$ ) (Figure 2).

#### \*\*\*Figure 2\*\*\*

No other significant between-group difference was noted for the remaining of the additional analyses; including PJT frequency, PJT duration, and total number of PJT sessions, participant's age, and FIFA age categories. A detailed description of all additional analyses is indicated as supplementary material (*Electronic Supplementary Material Appendix S1*).

### 3.6 Results of meta-regression

The meta-regression analysis was computed for the outcomes countermovement jump and 10-m linear sprint performance, and included three different training variables (i.e., training frequency, training duration, and total number of training sessions) and participants' chronological age (Table 3). Irrespective of the training type, none of the training variables predicted the effects of PJT on countermovement jump height or 10-m linear sprint performance ( $p=0.095$  to  $0.713$ ). The coefficient of determination was  $R^2=0.07$  and  $0.0$  for countermovement jump height and 10-m linear sprint performance. The regression was not computed for 20-m and 30-m linear sprint due to a problem with collinearity. For the remaining outcomes less than 10 studies were available, precluding a robust meta-regression.

#### \*\*\*Table 3\*\*\*

### 3.7 Adverse effects

Among the included studies, three [83-85] reported low level of pain experienced by participants. Although prevalence was not reported in the aforementioned studies, authors reported relatively low pain level among participants (all  $<3$ , in a 10-point scale). Moreover, mean pain levels of 0, 1.3, 0.8, 0.3, 0.1 and 0 were observed at times points before, immediately after, 24h, 48h, 72h, and 96h after the first plyometric training session, respectively. Furthermore, compared to the first week of PJT, muscle pain after plyometric training sessions was reduced toward the last week of PJT [83-85]. Another study [98] reported soreness in the lower leg muscle groups (13% of participants), pain in the knees mainly during the last stage of the intervention (8% of participants), and fatigue (13% of participants). However, no intervention-related injuries were reported. The remaining studies reported no soreness, pain, fatigue, injury, damage or adverse effects related to the PJT intervention.

## 4 Discussion

The aim of this SRMA was to assess the effects of PJT on jumping and sprinting among young male soccer players. The data showed that PJT resulted in significant small-moderate improvements in vertical jump performance and linear sprint. Regarding training prescription effects, PJT interventions of longer duration (>7 weeks and >14 PJT sessions) induced significantly greater moderate improvements in linear sprint performance. Findings are explored in more detail throughout the rest of the discussion.

Improvements in jumping and sprinting after PJT can likely be attributed to enhanced neural drive to agonist muscles, alterations to musculotendinous stiffness, improved intermuscular coordination (e.g., enhanced antagonist muscle inhibition), greater excitability of the stretch reflex, changes in muscle fiber mechanics, and changes in muscle size and architecture [99, 100]. In fact, a 6-week intervention, comprised of three sessions per week [101], in young adult female and male team sport players (including soccer), resulted in a significant improvement of 8.5-13.2% in unloaded jumping height performance. This improvement was in line with increases in maximal voluntary force and electromyographic activity of the leg extensor muscles, as well as greater thickness, fascicle length and pennation angle of knee flexor and extensor muscles [101]. However, when compared to the aforementioned study [101], others have found greater (~28%) [80], similar (~9%) [102] or lower (~3%) [103] improvement in vertical jump height. Differences in the participants' characteristics, including age, may help to explain differences in physical fitness changes, including jumping and sprinting, after PJT among young male soccer players [33, 35, 104]. However, differences between PJT programmes (e.g., frequency, duration, total number of PJT sessions) may also help explain the different magnitudes of physical fitness changes among studies. To analyse these possibilities, the effects of potential moderator variables were explored.

Regarding PJT frequency, moderator analyses and meta-regression analyses were available only for countermovement jump height and 10-m linear sprint performance. No between-group differences were noted for the improvements in countermovement jump and 10-m linear sprinting after interventions with  $\leq 2$  sessions per week or  $> 2$  sessions per week. Indeed, previous PJT meta-analyses also observed no significant subgroup differences or correlation for training frequency and vertical jump height [28, 64] or linear sprinting [105] changes. Furthermore, studies in adult futsal and soccer players [106, 107] compared the relative effects of one and two PJT sessions per week, equated for total volume, intensity and jump type, and found similar effects in vertical jump height and linear sprinting. Of note, results of the multivariate random-effect meta-regression revealed that training frequency did not predict PJT effects on countermovement jump and linear sprint performance in young male soccer players. Although two sessions per week seemed more effective than one for the improvement of linear sprinting among young male athletes [105], three sessions per week may have a lower effect than two sessions per week [105]. Indeed, greater training frequencies are associated with higher training volumes and because of this, could increase the risk of injury [108]. However, the lack of a significant difference between  $\leq 2$  sessions/week compared to  $> 2$  sessions/week in our meta-analysis may be related to an imbalance of studies in the respective subgroups. For countermovement jump height, 24 study groups were available for  $\leq 2$  sessions/week, whereas only three studies were available for  $> 2$  sessions/week. Indeed, compared to the moderate heterogeneity for  $\leq 2$  sessions/week, the presence of high heterogeneity after subgroup analysis for  $> 2$  sessions/week suggests that moderators of the main effect may not have been found, meaning other factors (aside from training frequency) could account for training adaptations. This would seem to imply a potential synergy between programming variables and other factors, such as biological maturity, in determining the magnitude of response to PJT in young athletes [105]. To this end, a moderator analysis accounting for player age would clarify this issue.

Although the maturity status of the players would be ideal in order to perform moderator analyses for the effects of PJT, maturity is often not reported. In fact, a recent scoping review [27] from 420 PJT studies, observed that 37% of the included studies involved youth groups, with only a third of these reporting physiological maturity status. This important research gap seems common among resistance training research literature [109]. This limitation is compounded by the utilization of different measures of physiological maturity across studies, making it difficult to compare results [26, 27]. This could be viewed as a critical limitation among PJT interventions performed with youth, especially since physiological maturity seems to affect adaptations to PJT interventions with young males. In the current SRMA, with regard to player age, moderator analyses were available for countermovement jump height and squat jump height performance, and 10-m, 20-m and 30-m linear sprint. From these studies, data indicate that players >13.2 years old experienced a similar moderate (ES = 0.64-0.77) beneficial training effect on jumping performance compared those ≤13.2 years old (ES = 0.81-0.94). For the 10-m, 20-m and 30-m linear sprint, players >13.2 years old experienced a moderate beneficial training effect (ES = 0.83-0.89), compared to a small beneficial effect on those ≤13.2 years old (ES = 0.40-0.53). Results of the multivariate random-effect meta-regression revealed that player's age did not predict PJT effects on jump and linear sprint performance in young male soccer players. Although for the analysed outcomes in the current SRMA no significant between-group differences were noted regarding player age, the moderate beneficial effect of PJT on linear sprint performance among the older young players, compared to only a small beneficial effect among their younger counterpart, is in line with previous meta-analyses. In one meta-analysis [105], greater improvements in sprinting were noted among participants with mean ages of 14.1 years (ES = 1.15) and 16.8 y (ES = 1.39), compared with those with a mean age of 11.2 y (ES = -0.18) after sprinting programmes (also involving high-intensity, stretch-shortening cycle muscle actions). In another meta-analysis [110], improvements in non-linear sprinting (i.e., change of direction speed) were noted among participants with a mean age of 14.5 y (ES = 0.95) and 17 years (ES = 0.99), compared with those with a mean age of 11.5 y (ES = 0.68) after PJT interventions. Moreover, when participants between the mean ages of 10 and 12.9 years, 13 and 15.9 years, as well as 16 and 18 years were exposed to PJT, the greatest magnitude of improvement in countermovement jump height performance was noted among the older group (ES = 1.02) [33]. However, in the aforementioned meta-analysis [33] the magnitude of adaptation to PJT between the mean ages of 13 and 15.9 years was lower (ES = 0.47) compared to the younger group (ES = 0.91). In relation to this finding, complex changes occur in physical performance during growth and maturation and these can affect both jumping [31] and sprinting [111, 112]. During growth and maturation, the natural development of the stretch-shortening cycle is of key relevance for both jump and sprint performance and this occurs due to greater muscular size, increased limb length, changes to musculotendinous tissue, enhanced neural and motor development and better movement quality and coordination [31, 111, 112]. As the timing and tempo of these factors seems highly variable across individuals [32, 33, 113], this can make it difficult for coaches to determine how best to structure training during this highly sensitive period of development. Therefore, soccer coaches involved with youth populations should consider not only the characteristics of the applied training programme, but also the dynamic physiological change that takes place across the adolescent years. Such training principle related to the interaction between training and maturation have been termed “synergistic adaptation” [34, 114, 115] and should be considered of utmost importance when working with young soccer players.

Regarding intervention duration and total PJT sessions, results of the multivariate random-effect meta-regression revealed that none of these training factors predict PJT effects on countermovement jump and 10-m linear sprint performance in young male soccer players. However, analyses for intervention duration and total PJT sessions as single-factor moderators were available for countermovement and squat jump height, and for 10-m, 20-m and 30-m linear sprint performance. From these, interventions with a duration of >7 weeks and >14 total PJT sessions induced a greater beneficial training effect compared to those interventions with ≤7 weeks and ≤14 total PJT sessions on 10-m linear sprint performance. Unsurprisingly, the moderator analysis supported the use of

longer programmes (>7 weeks) and more training sessions per programme (>14) for the enhancement of horizontally-orientated outcomes and skills such as short sprints. It was surprising, however, that longer programmes (>7 weeks, >14 session) were no more effective than shorter programmes in eliciting increases in vertically-orientated outcome measures such as countermovement and squat jump height performance. The reasons for these contrasting findings are unclear but could suggest that increases in vertically-orientated performance are achievable in the short term whilst the attainment of the more sport-specific horizontally-orientated performance [4] could take longer to achieve. This could indicate a differential in the time-course of adaptation of vertically- and horizontally-orientated performance or could also represent a bias towards the selection of vertically-orientated exercises in modern strength and conditioning programmes for young soccer players [12]. Alternatively, current findings may indicate that the longer-term programs were not sufficiently periodized and the players were not exposed to sufficient PJT load, particularly PJT intensity. In general, coaches have traditionally been cautious of higher training intensities, however, this prescription variable is crucial for long-term PJT programmes [35, 116, 117].

Our meta-analyses demonstrated that young soccer players may improve vertical jump (ES = 0.48 to 0.79) and linear sprint performance (ES = 0.60 to 0.98) to a similar extent after PJT. Although this may be considered not in line with the principle of training specificity, most of the included studies in our meta-analyses involved mixed PJT programmes that combined horizontal and vertical drills. Indeed, while vertically-oriented PJT may induce greater improvements in vertical jump performance, horizontal-oriented PJT may induce greater improvements in linear sprint performance [87]. However, a combination of both may be of particular relevance to improve both vertical jump (12.3%; ES = 0.51) and linear sprint performance (5.8-6.0%; ES = 0.63-0.99) among young male soccer players [87]. In addition to the PJT characteristics, similar improvements in vertical jump and linear sprint performance among young soccer players in our meta-analysis are in line with the findings from previous PJT meta-analyses, which have shown an improvement in vertical jump (ES = 0.84) [28] and linear sprint performance (ES = 0.37) [29]. However, such effects were noted for participants with a wide range of sport backgrounds. In the aforementioned meta-analyses soccer players demonstrated vertical jump improvements of ES = 0.51 [28] and linear print ES = 0.69 [29]. The reasons for these findings are unclear but could suggest that the underlying mechanisms responsible for vertical jump and linear sprint performance may be similarly improved after PJT. Indeed, PJT can increase neural drive to agonist muscles, lower-limb stiffness, intermuscular coordination, excitability of the stretch reflex, among others [99, 100]. Such factors are important for both jumping and sprinting [15, 118-123]. The underlying mechanisms (e.g. physiological; biomechanical) responsible for improvements in vertical jump and linear sprint after PJT should be considered in future studies. From a practical point of view, combination of both vertical-oriented and horizontal-oriented PJT drills in the young soccer player's programme seems a sound approach.

Our meta-analyses revealed linear sprint improvements after PJT in a range from ES = 0.60 to 0.98 for distances between 5-m to 40-m. Although linear sprint performance may correlate across different distances [118], the underlying mechanisms (e.g., physiological; biomechanical) responsible for the athlete performance across different linear sprint distances may be differentially affected. Such effects may be related to the distinct characteristics of the PJT interventions across analysed studies (e.g., total programme duration; as previously discussed) and the nature of the plyometric exercises. Indeed, depending on the training approach, one may expect greater improvements in one particular distance over another. For example, at shorter-distances (e.g. 5-m) horizontal force application on the ground is of paramount importance, thus a greater load of horizontal PJT may lead to larger improvements during the early acceleration phase (horizontal GRF; push-off phase) [87, 119, 124]. In addition, exercises with horizontal orientation and longer ground contact times will allow for more time to generate force, thus greater impulse and, therefore, acceleration. In contrast, PJT with a greater emphasis in the vertical direction may induce larger improvements when nearing top speed (vertical GRF) [87, 119,

124], particularly after vertical exercises with faster ground contact times and higher rate of force development. In this meta-analysis, most of the included studies involved mixed PJT programmes that combined horizontal and vertical drills. This may help to explain the improvements noted across different linear sprint distances.

Among the included studies, no intervention-related injuries were reported. The relative safety of PJT programmes have been previously reported [26, 27, 99]. Moreover, when adequately programmed and well coached, PJT interventions may reduce the risk of injury among young soccer players [14, 125]. Although PJT seems safe for young male soccer players, caution is recommended when applying this type of training in poor-conditioned athletes with lower strength levels and an inability to decelerate their body mass during landing tasks. Of note, in a study by Vlachalopoulos et al. [98], participants reduced the volume of jumps in the last 12 weeks due to soreness and some muscle problems (not injuries). It is possible that a volume-based taper in the last stage of a PJT can increase control over inflammation caused by the overload induced by large eccentric loads [126, 127] and, in this way, a taper strategy may facilitate the processes of adaptation of the musculoskeletal system and physical fitness [128, 129]. In addition to taper strategies, low volumes of high intensity work may be more advantageous at the long-term compared to greater volumes [28, 106, 108, 130-132]. In other words, intervention-related injuries may be reduced, and physical fitness improved to a greater extent using sufficiently periodized longer-term programs, taking into account PJT intensity as a key prescription variable for young athletes in long-term athletic development programmes [35, 116, 117].

To our knowledge, this is the first SRMA to examine the effects of PJT on jumping and sprinting performance in young soccer players. In the current SRMA 752 participants (in addition to 747 controls) partook in PJT among the single studies reported in the literature. This large pooled sample size is a strength of the current SRMA, addressing the ongoing problem of underpowered studies due to reduced sample size [133]. However, aside from the aforementioned strengths, some potential limitations should be acknowledged. For some outcomes (i.e. 5-m linear sprint), additional analyses regarding PJT frequency, duration, total PJT sessions or participants' age were not possible as <3 studies were available for at least one of the moderators. Moreover, for some outcomes (i.e., 40-m linear sprint) and/or moderator analysis only three studies were available in total, suggesting that results should be interpreted with caution, and confirmed in the future. Additionally, the dichotomisation of continuous data (e.g.,  $\leq 7$  weeks compared to  $> 7$  weeks) with the median split technique could result in residual confounding and reduced statistical power [134]. Furthermore, the effects of these programming variables were calculated independently, and not interdependently. Univariate analysis must be interpreted with caution because the programming parameters were calculated as single factors, irrespective of between-parameter interactions. However, our meta-analysis also incorporated a meta-regression, revealing that none of the analyzed training factors predicted PJT effects on either jump or linear sprint performance in young male soccer players. Finally, the current SRMA was focused on young male participants. As young males and females clearly experience different effects from PJT according to sex-specific maturational development [31-35, 37], future SRMAs should take a similar approach for female participants. Additionally, although our analyses did not reveal a significant difference between participants aged <13.2 compared to >13.2 (or between FIFA age categories U17, U21, and U23), these were limited only to chronological age. A moderator analysis for biological maturity was limited somewhat with the evidence that is available. Indeed, not many PJT studies report well controlled measures of maturity status [27]. Considering that biological maturity may affect adaptations to strength and conditioning practices in general, and resistance training and PJT in particular [31, 33, 35, 112, 115], future PJT meta-analyses should strive to include youth athletes biological maturity as a moderator in the analyses. Despite these limitations, the current SRMA makes an original and significant contribution to the literature and clearly shows the merits on including PJT as part of a well-rounded athletic development program to enhance jumping and sprinting performance in young soccer players.

## 5 Conclusion

In conclusion, PJT seems safe and was proved to be effective in improving vertical jumping and linear sprinting performance among young male soccer players. Greater 10-m linear sprinting improvements were noted after interventions >7 weeks duration and >14 sessions, suggesting a greater return from exposure to longer PJT interventions, partially in support for the adoption of a long-term approach to athletic development in young athletes. However, with reference to the findings of the meta-regression, and those from the remaining subgroup and single factors analysis, a robust confirmation regarding the moderator role of participant's age, or PJT configuration, including duration, on its effects on young soccer player's fitness qualities needs future confirmation. Practitioners working in youth soccer should take into account the dose-response trends identified in this SRMA to prescribe the appropriate level of training for the young male soccer player. Importantly, rather than an independent entity, PJT should be a component of an integrated approach to youth physical development, which targets multiple physical fitness qualities and aligns with the goals of long-term physical development strategies. Practitioners should seek to periodize PJT for young athletes by manipulating both volume and intensity to ensure ongoing adaptations.

## 6 Data Availability Statement

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

## Compliance with Ethical Standards

### *Funding*

No sources of funding were used to assist in the preparation of this article.

### *Conflicts of Interest*

Rodrigo Ramirez-Campillo, Daniel Castillo, Javier Raya-González, Jason Moran, Eduardo Sáez de Villarreal and Rhodri Lloyd declare that they have no conflicts of interest relevant to the content of this review.

### *Authorship Contributions*

RRC, DC and JRG wrote the first draft of the manuscript. RRC, DC and JRG collected data. RRC and JM analyzed and interpreted the data. ESV and RD revised the original manuscript. All authors read and approved the final manuscript.

## REFERENCES

1. Arnason A, Sigurdsson SB, Gudmundsson A, Holme I, Engebretsen L, Bahr R. Physical fitness, injuries, and team performance in soccer. *Med Sci Sports Exerc.* 2004 Feb;36(2):278-85.
2. Stolen T, Chamari K, Castagna C, Wisloff U. Physiology of soccer: an update. *Sports Med.* 2005;35(6):501-36.
3. Castagna C, D'Ottavio S, Abt G. Activity profile of young soccer players during actual match play. *J Strength Cond Res.* 2003 Nov;17(4):775-80.

4. Faude O, Koch T, Meyer T. Straight sprinting is the most frequent action in goal situations in professional football. *J Sports Sci.* 2012;30(7):625-31.
5. Barnes C, Archer DT, Hogg B, Bush M, Bradley PS. The evolution of physical and technical performance parameters in the English Premier League. *Int J Sports Med.* 2014 Dec;35(13):1095-100.
6. Turner E, Munro AG, Comfort P. Female Soccer: Part 1—A Needs Analysis. *Strength & Conditioning Journal.* 2013;35(1):51-7.
7. Datson N, Hulton A, Andersson H, Lewis T, Weston M, Drust B, et al. Applied physiology of female soccer: an update. *Sports Med.* 2014 Sep;44(9):1225-40.
8. Hansen L, Bangsbo J, Twisk J, Klausen K. Development of muscle strength in relation to training level and testosterone in young male soccer players. *J Appl Physiol.* 1999 Sep;87(3):1141-7.
9. Vaeyens R, Malina RM, Janssens M, Van Renterghem B, Bourgois J, Vrijens J, et al. A multidisciplinary selection model for youth soccer: the Ghent Youth Soccer Project. *Br J Sports Med.* 2006 Nov;40(11):928-34; discussion 34.
10. le Gall F, Carling C, Williams M, Reilly T. Anthropometric and fitness characteristics of international, professional and amateur male graduate soccer players from an elite youth academy. *J Sci Med Sport.* 2010 Jan;13(1):90-5.
11. Reilly T, Gilbourne D. Science and football: a review of applied research in the football codes. *J Sports Sci.* 2003 Sep;21(9):693-705.
12. Meylan C, Cronin J, Oliver J, Hughes M, Manson S. An evidence-based model of power development in youth soccer. *Int J Sports Sci Coach.* 2014;9(5):1241-64.
13. Bedoya AA, Miltenberger MR, Lopez RM. Plyometric training effects on athletic performance in youth soccer athletes: a systematic review. *J Strength Cond Res.* 2015 Aug;29(8):2351-60.
14. Rossler R, Donath L, Verhagen E, Junge A, Schweizer T, Faude O. Exercise-based injury prevention in child and adolescent sport: a systematic review and meta-analysis. *Sports Med.* 2014 Dec;44(12):1733-48.
15. Taube W, Leukel C, Gollhofer A. How neurons make us jump: the neural control of stretch-shortening cycle movements. *Exerc Sport Sci Rev.* 2012 Apr;40(2):106-15.
16. Komi PV, Gollhofer A. Stretch reflex can have an important role in force enhancement during SSC-exercise. *Journal of Applied Biomechanics.* 1997;13:451-9.
17. Duda M. Plyometrics: A Legitimate Form of Power Training? *Phys Sportsmed.* 1988 Mar;16(3):212-8.
18. Faigenbaum A, Chu D. *Plyometric Training for Children and Adolescents.* Indianapolis, IN: American College of Sports Medicine. 2017.
19. Sands WA, Wurth JJ, Hewitt JK. Speed and Agility Training. In: (NSCA) TNSaCAs, editor. *Basics of Strength and Conditioning manual;* 2012.
20. Negra Y, Chaabene H, Fernandez-Fernandez J, Sammoud S, Bouguezzi R, Prieske O, et al. Short-term plyometric jump training improves repeated-sprint ability in prepuberal male soccer players. *J Strength Cond Res.* In press Jun 14.
21. Loturco I, Pereira LA, Kobal R, Maldonado T, Piazza AF, Bottino A, et al. Improving Sprint Performance in Soccer: Effectiveness of Jump Squat and Olympic Push Press Exercises. *PLoS One.* 2016;11(4):e0153958.
22. Ramirez-Campillo R, Vergara-Pedrerros M, Henriquez-Olguin C, Martinez-Salazar C, Alvarez C, Nakamura FY, et al. Effects of plyometric training on maximal-intensity exercise and endurance in male and female soccer players. *J Sports Sci.* 2016 Apr;34(8):687-93.
23. Thomas K, French D, Hayes PR. The effect of two plyometric training techniques on muscular power and agility in youth soccer players. *J Strength Cond Res.* 2009 Jan;23(1):332-5.



24. Kobal R, Loturco I, Barroso R, Gil S, Cuniyochi R, Ugrinowitsch C, et al. Effects of different combinations of strength, power, and plyometric training on the physical performance of elite young soccer players. *J Strength Cond Res.* 2017 Jun;31(6):1468-76.
25. Yanci J, Los Arcos A, Camara J, Castillo D, García A, Castagna C. Effects of horizontal plyometric training volume on soccer players' performance. *Res Sports Med.* 2016;24(4):308-19.
26. Ramirez-Campillo R, Alvarez C, Garcia-Hermoso A, Ramirez-Velez R, Gentil P, Asadi A, et al. Methodological characteristics and future directions for plyometric jump training research: A scoping review. *Sports Med.* 2018 May;48(5):1059-81.
27. Ramirez-Campillo R, Moran J, Chaabene H, Granacher U, Behm DG, Garcia-Hermoso A, et al. Methodological characteristics and future directions for plyometric jump training research: A scoping review update. *Scand J Med Sci Sports.* 2020 Feb 8.
28. de Villarreal ES, Kellis E, Kraemer WJ, Izquierdo M. Determining variables of plyometric training for improving vertical jump height performance: a meta-analysis. *J Strength Cond Res.* 2009 Mar;23(2):495-506.
29. de Villarreal ES, Requena B, Cronin JB. The effects of plyometric training on sprint performance: a meta-analysis. *Journal of Strength and Conditioning Research.* 2012 Feb;26(2):575-84.
30. van de Hoef PA, Brauers JJ, van Smeden M, Backx FJG, Brink MS. The effects of lower-extremity plyometric training on soccer-specific outcomes in adult male soccer players: a systematic review and meta-analysis. *Int J Sports Physiol Perform.* 2019 Dec 4:1-15.
31. Radnor JM, Oliver JL, Waugh CM, Myer GD, Moore IS, Lloyd RS. The influence of growth and maturation on stretch-shortening cycle function in youth. *Sports Medicine.* 2017 September 12.
32. Moran J, Clark CCT, Ramirez-Campillo R, Davies MJ, Drury B. A meta-analysis of plyometric training in female youth: its efficacy and shortcomings in the literature. *J Strength Cond Res.* 2019 Jul;33(7):1996-2008.
33. Moran J, Sandercock G, Ramirez-Campillo R, Meylan C, Collison J, Parry D. Age-related variation in male youth athletes' countermovement jump after plyometric training: A meta-analysis of controlled trials. *Journal of Strength and Conditioning Research.* 2017;31(2):552-65.
34. Lloyd RS, Cronin JB, Faigenbaum AD, Haff GG, Howard R, Kraemer WJ, et al. National Strength and Conditioning Association position statement on long-term athletic development. *J Strength Cond Res.* 2016 Jun;30(6):1491-509.
35. Lloyd RS, Meyers RW, Oliver JL. The natural development and trainability of plyometric ability during childhood. *Strength Cond J.* 2011 Apr;33(2):23-32.
36. Lloyd RS, Oliver JL. The Youth Physical Development Model: A New Approach to Long-Term Athletic Development. *Strength & Conditioning Journal.* 2012;34(3):61-72.
37. Lloyd RS, Oliver JL, Hughes MG, Williams CA. The influence of chronological age on periods of accelerated adaptation of stretch-shortening cycle performance in pre and postpubescent boys. *J Strength Cond Res.* 2011 Jul;25(7):1889-97.
38. Ramirez-Campillo R, Sanchez-Sanchez J, Romero-Moraleda B, Yanci J, Garcia-Hermoso A, Manuel Clemente F. Effects of plyometric jump training in female soccer player's vertical jump height: A systematic review with meta-analysis. *J Sports Sci.* 2020 Apr 7:1-13.
39. Ramirez-Campillo R, Alvarez C, Garcia-Hermoso A, Keogh JW, García-Pinillos F, Pereira LA, et al. Effects of jump training on jumping performance of handball players: A systematic review with meta-analysis of randomised controlled trials. *Inter J Sports Sci Coach.* 2020:1747954120928932.

40. Ramirez-Campillo R, Andrade DC, Nikolaidis PT, Moran J, Clemente FM, Chaabene H, et al. Effects of plyometric jump training on vertical jump height of volleyball players: a systematic review with meta-analysis of randomized-controlled trial. *J Sport Sci Med.* 2020;19:489-99.
41. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al. *Cochrane handbook for systematic reviews of interventions.* Second edition Chichester (UK): John Wiley & Sons; 2019.
42. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ.* 2009;339:b2700.
43. Chu D, Myer G. *Plyometrics.* Champaign: Human Kinetics. 2013.
44. Moran J, Ramirez-Campillo R, Granacher U. Effects of jumping exercise on muscular power in older adults: a meta-analysis. *Sports Med.* 2018 Dec;48(12):2843-57.
45. FIFA. Federation Internationale de Football Association. Zürich, Switzerland. 2020; Accessed at: <https://www.fifa.com>.
46. Turner HM, Bernard RM. Calculating and synthesizing effect sizes. *Contemp Issues Commun Sci Disord.* 2006;33:42-55.
47. Slinde F, Suber C, Suber L, Edwen CE, Svantesson U. Test-retest reliability of three different countermovement jumping tests. *J Strength Cond Res.* 2008 Mar;22(2):640-4.
48. Altmann S, Ringhof S, Neumann R, Woll A, Rumpf MC. Validity and reliability of speed tests used in soccer: A systematic review. *PLoS One.* 2019;14(8):e0220982.
49. Murtagh CF, Brownlee TE, O'Boyle A, Morgans R, Drust B, Erskine RM. Importance of speed and power in elite youth soccer depends on maturation status. *J Strength Cond Res.* 2018 Feb;32(2):297-303.
50. Moran J, Paxton K, Jones B, Granacher U, Sandercock GRH, Hope E, et al. Variable long-term developmental trajectories of short sprint speed and jumping height in English Premier League academy soccer players: an applied case study. *J Sport Sci.* In press; In press.
51. Stojanović E, Ristić V, McMaster DT, Milanović Z. Effect of plyometric training on vertical jump performance in female athletes: A systematic review and meta-analysis. *Sports Medicine.* 2017;47(5):975-86.
52. Ramirez-Campillo R, Sanchez-Sanchez J, Romero-Moraleda B, Yanci J, García-Hermoso A, Manuel Clemente F. Effects of plyometric jump training in female soccer player's vertical jump height: a systematic review with meta-analysis. *J Sports Sci.* 2020 (in press).
53. Higgins JP, Altman DG. Assessing Risk of Bias in Included Studies. *Cochrane Handbook for Systematic Reviews of Interventions;* 2008. p. 187-241.
54. Skrede T, Steene-Johannessen J, Anderssen SA, Resaland GK, Ekelund U. The prospective association between objectively measured sedentary time, moderate-to-vigorous physical activity and cardiometabolic risk factors in youth: a systematic review and meta-analysis. *Obes Rev.* 2019 Jan;20(1):55-74.
55. Garcia-Hermoso A, Ramirez-Campillo R, Izquierdo M. Is Muscular Fitness Associated with Future Health Benefits in Children and Adolescents? A Systematic Review and Meta-Analysis of Longitudinal Studies. *Sports Med.* 2019 Jul;49(7):1079-94.
56. Deeks JJ, Higgins JP, Altman DG. Analysing data and undertaking meta-analyses. In: Higgins JP, Green S, editors. *Cochrane Handbook for Systematic Reviews of Interventions: The Cochrane Collaboration;* 2008. p. 243-96.
57. Kontopantelis E, Springate DA, Reeves D. A re-analysis of the Cochrane Library data: the dangers of unobserved heterogeneity in meta-analyses. *PLoS One.* 2013;8(7):e69930.
58. Hardy RJ, Thompson SG. A likelihood approach to meta-analysis with random effects. *Statistics in medicine.* 1996 Mar 30;15(6):619-29.

59. Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc.* 2009 Jan;41(1):3-13.
60. Higgins JP, Deeks JJ, Altman DG. Special topics in statistics. In: Higgins JP, Green S, editors. *Cochrane handbook for systematic reviews of interventions: The Cochrane Collaboration*; 2008. p. 481-529.
61. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *Bmj.* 2003 Sep 6;327(7414):557-60.
62. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Statistics in medicine.* 2002 Jun 15;21(11):1539-58.
63. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *Bmj.* 1997 Sep 13;315(7109):629-34.
64. Moran J, Clark CCT, Ramirez-Campillo R, Davies MJ, Drury B. A meta-analysis of plyometric training in female youth: its efficacy and shortcomings in the literature. *J Strength Cond Res.* IN PRESS.
65. Moran J, Sandercock G, Ramirez-Campillo R, Clark CCT, Fernandes JFT, Drury B. A meta-analysis of resistance training in female youth: Its effect on muscular strength, and shortcomings in the literature. *Sports Med.* 2018 Apr 6.
66. Moran J, Sandercock GR, Ramirez-Campillo R, Meylan C, Collison J, Parry DA. A meta-analysis of maturation-related variation in adolescent boy athletes' adaptations to short-term resistance training. *J Sports Sci.* 2017 Jun;35(11):1041-51.
67. Pescatello LS, MacDonald HV, Lamberti L, Johnson BT. Exercise for Hypertension: A Prescription Update Integrating Existing Recommendations with Emerging Research. *Current hypertension reports.* 2015;17(11):87-.
68. Asadi A, Ramirez-Campillo R, Arazi H, Saez de Villarreal E. The effects of maturation on jumping ability and sprint adaptations to plyometric training in youth soccer players. *J Sports Sci.* 2018 Nov;36(21):2405-11.
69. Beato M, Bianchi M, Coratella G, Merlini M, Drust B. Effects of plyometric and directional training on speed and jump performance in elite youth soccer players. *J Strength Cond Res.* 2018;32(2):289-96.
70. Brito J, Vasconcellos F, Oliveira J, Krusturup P, Rebelo A. Short-term performance effects of three different low-volume strength-training programmes in college male soccer players. *J Hum Kinet.* 2014 Mar 27;40:121-8.
71. Chelly MS, Ghenem MA, Abid K, Hermassi S, Tabka Z, Shephard RJ. Effects of in-season short-term plyometric training program on leg power, jump- and sprint performance of soccer players. *J Strength Cond Res.* 2010 Oct;24(10):2670-6.
72. Chtara M, Rouissi M, Haddad M, Chtara H, Chaalali A, Owen A, et al. Specific physical trainability in elite young soccer players: efficiency over 6 weeks' in-season training. *Biology of Sport.* 2017;34(2):137-48.
73. Coratella G, Beato M, Milanese C, Longo S, Limonta E, Rampichini S, et al. Specific adaptations in performance and muscle architecture after weighted jumpsquat vs. body mass squat jump training in recreational soccer players. *J Strength Cond Res.* 2018;32(4):921-9.
74. Hammami M, Negra Y, Aouadi R, Shephard RJ, Chelly MS. Effects of an in-season plyometric training program on repeated change of direction and sprint performance in the junior soccer player. *J Strength Cond Res.* 2016 Dec;30(12):3312-20.
75. Hammami M, Gaamouri N, Shephard RJ, Chelly MS. Effects of contrast strength vs. plyometric training on lower-limb explosive performance, ability to change direction and neuromuscular adaptation in soccer players. *J Strength Cond Res.* 2019 Aug;33(8):2094-103.
76. Jlid MC, Coquart J, Maffulli N, Paillard T, Bisciotti GN, Chamari K. Effects of in Season Multi-Directional Plyometric Training on Vertical Jump Performance, Change of Direction Speed and Dynamic Postural Control in U-21 Soccer Players. *Frontiers in physiology.* 2020;11:374.


77. Jlid MC, Racil G, Coquart J, Paillard T, Bisciotti GN, Chamari K. Multidirectional Plyometric Training: Very Efficient Way to Improve Vertical Jump Performance, Change of Direction Performance and Dynamic Postural Control in Young Soccer Players. *Frontiers in physiology*. 2019;10:1462.
78. McKinlay BJ, Wallace P, Dotan R, Long D, Tokuno C, Gabriel DA, et al. Effects of plyometric and resistance training on muscle strength, explosiveness, and neuromuscular function in young adolescent soccer players. *J Strength Cond Res*. 2018;32(11):3039-50.
79. Meylan C, Malatesta D. Effects of in-season plyometric training within soccer practice on explosive actions of young players. *J Strength Cond Res*. 2009 Dec;23(9):2605-13.
80. Michailidis Y, Fatouros IG, Primpa E, Michailidis C, Avloniti A, Chatzinikolaou A, et al. Plyometrics trainability in preadolescent soccer athletes. *J Strength Cond Res*. 2013;27(1):38-49.
81. Nakamura D, Suzuki T, Yasumatsu M, Akimoto T. Moderate running and plyometric training during off-season did not show a significant difference on soccer-related high-intensity performances compared with no-training controls. *J Strength Cond Res*. 2012 Dec;26(12):3392-7.
82. Negra Y, Chaabene H, Stöggl T, Hammami M, Chelly MS, Hachana Y. Effectiveness and time-course adaptation of resistance training vs. plyometric training in prepubertal soccer players. *J Sport Health Sci*. 2016 2016/07/16/.
83. Ramirez-Campillo R, Andrade DC, Alvarez C, Henriquez-Olguin C, Martinez C, Baez-Sanmartin E, et al. The effects of intersession rest on adaptation to 7 weeks of explosive training in young soccer players. *J Sports Sci Med*. 2014 May;13(2):287-96.
84. Ramirez-Campillo R, Meylan C, Alvarez C, Henriquez-Olguin C, Martinez C, Canas-Jamett R, et al. Effects of in-season low-volume high-intensity plyometric training on explosive actions and endurance of young soccer players. *J Strength Cond Res*. 2014 May;28(5):1335-42.
85. Ramirez-Campillo R, Meylan CM, Alvarez-Lepin C, Henriquez-Olguin C, Martinez C, Andrade DC, et al. The effects of interday rest on adaptation to 6 weeks of plyometric training in young soccer players. *J Strength Cond Res*. 2015 Apr;29(4):972-9.
86. Ramirez-Campillo R, Burgos CH, Henriquez-Olguin C, Andrade DC, Martinez C, Alvarez C, et al. Effect of unilateral, bilateral, and combined plyometric training on explosive and endurance performance of young soccer players. *J Strength Cond Res*. 2015 May;29(5):1317-28.
87. Ramirez-Campillo R, Gallardo F, Henriquez-Olguin C, Meylan CM, Martinez C, Alvarez C, et al. Effect of vertical, horizontal, and combined plyometric training on explosive, balance, and endurance performance of young soccer players. *J Strength Cond Res*. 2015 Jul;29(7):1784-95.
88. Ramirez-Campillo R, Henriquez-Olguin C, Burgos C, Andrade DC, Zapata D, Martinez C, et al. Effect of progressive volume-based overload during plyometric training on explosive and endurance performance in young soccer players. *J Strength Cond Res*. 2015 Jul;29(7):1884-93.
89. Ramirez-Campillo R, Alvarez C, García-Pinillos F, Sanchez-Sanchez J, Yanci J, Castillo D, et al. Optimal reactive strength index: is it an accurate variable to optimize plyometric training effects on measures of physical fitness in young soccer players? *J Strength Cond Res*. 2018;32(4):885-93.
90. Ramirez-Campillo R, Alvarez C, Garcia-Pinillos F, Gentil P, Moran J, Pereira LA, et al. Effects of plyometric training on physical performance of young male soccer players: potential effects of different drop jump heights. *Pediatr Exerc Sci*. 2019 Feb 8;31(3):306-13.
91. Ramirez-Campillo R, Alvarez C, Gentil P, Loturco I, Sanchez-Sanchez J, Izquierdo M, et al. Sequencing effects of plyometric training applied before or after regular soccer training on measures of physical fitness in young players. *J Strength Cond Res*. 2018 Mar 22;34(7):1959-66.

92. Ramirez-Campillo R, Alvarez C, Garcia-Pinillos F, Garcia-Ramos A, Loturco I, Chaabene H, et al. Effects of combined surfaces vs. single-surface plyometric training on soccer players' physical fitness. *J Strength Cond Res.* 2019 Jan 17.
93. Rosas F, Ramirez-Campillo R, Diaz D, Abad-Colil F, Martinez-Salazar C, Caniuqueo A, et al. Jump training in youth soccer players: effects of haltere type handheld loading. *Int J Sports Med.* 2016 Dec;37(13):1060-5.
94. Sedano S, Matheu A, Redondo JC, Cuadrado G. Effects of plyometric training on explosive strength, acceleration capacity and kicking speed in young elite soccer players. *J Sports Med Phys Fitness.* 2011 Mar;51(1):50-8.
95. Söhnlein Q, Müller E, Stöggl TL. The effect of 16-week plyometric training on explosive actions in early to mid-puberty elite soccer players. *J Strength Cond Res.* 2014;28(8):2105-14.
96. Spinetti J, Figueiredo T, Bastos De Oliveira V, Assis M, Fernandes De Oliveira L, Miranda H, et al. Comparison between traditional strength training and complex contrast training on repeated sprint ability and muscle architecture in elite soccer players. *J Sports Med Phys Fit.* 2016 Nov;56(11):1269-78.
97. Vácz M, Tollár J, Meszler B, Juhász I, Karsai I. Short-term high intensity plyometric training program improves strength, power and agility in male soccer players. *J Human Kin.* 2013;36(1):17-26.
98. Vlachopoulos D, Barker AR, Ubago-Guisado E, Williams CA, Gracia-Marco L. The effect of a high-impact jumping intervention on bone mass, bone stiffness and fitness parameters in adolescent athletes. *Archives Osteoporosis.* 2018 Nov 17;13(1):128.
99. Markovic G, Mikulic P. Neuro-musculoskeletal and performance adaptations to lower-extremity plyometric training. *Sports Med.* 2010 Oct 1;40(10):859-95.
100. Slimani M, Chamari K, Miarka B, Del Vecchio FB, Chéour F. Effects of Plyometric Training on Physical Fitness in Team Sport Athletes: A Systematic Review. *Journal of Human Kinetics.* 2016;53(1):231-47.
101. Ullrich B, Pelzer T, Pfeiffer M. Neuromuscular effects to 6 weeks of loaded countermovement jumping with traditional and daily undulating periodization. *Journal of strength and conditioning research.* 2018;32(3):660-74.
102. Sáez De Villarreal E, Suarez-Arrones L, Requena B, Haff GG, Ferrete C. Effects of plyometric and sprint training on physical and technical skill performance in adolescent soccer players. *J Strength Cond Res.* 2015;29(7):1894-903.
103. Faude O, Roth R, Giovine DD, Zahner L, Donath L. Combined strength and power training in high-level amateur football during the competitive season: A randomised-controlled trial. *J Sport Sci.* 2013;31(13):1460-7.
104. Ramirez-Campillo R, Alvarez C, Sanchez-Sanchez J, Slimani M, Gentil P, Chelly MS, et al. Effects of plyometric jump training on the physical fitness of young male soccer players: Modulation of response by inter-set recovery interval and maturation status. *J Sport Sci.* 2019:1-8.
105. Moran J, Sandercock G, Rumpf MC, Parry DA. Variation in responses to sprint training in male youth athletes: A meta-analysis. *Int J Sports Med.* 2017 Jan;38(1):1-11.
106. Ramirez-Campillo R, Garcia-Pinillos F, Garcia-Ramos A, Yanci J, Gentil P, Chaabene H, et al. Effects of different plyometric training frequencies on components of physical fitness in amateur female soccer players. *Frontiers in physiology.* 2018;9:934.
107. Yanci J, Castillo D, Iturricastillo A, Ayarra R, Nakamura FY. Effects of Two Different Volume-Equated Weekly Distributed Short-Term Plyometric Training Programs on Futsal Players' Physical Performance. *J Strength Cond Res.* 2017 Jul;31(7):1787-94.
108. Brumitt J, Heiderscheid BC, Manske RC, Niemuth P, Mattocks A, Rauh MJ. The lower-extremity functional test and lower-quadrant injury in ncaa division iii athletes: a descriptive and epidemiologic report. *J Sport Rehabil.* 2016 Aug;25(3):219-26.

109. Granacher U, Lesinski M, Büsch D, Muehlbauer T, Prieske O, Puta C, et al. Effects of resistance training in youth athletes on muscular fitness and athletic performance: A conceptual model for long-term athlete development. *Frontiers in physiology*. 2016;7(MAY).
110. Asadi A, Arazi H, Ramirez-Campillo R, Moran J, Izquierdo M. Influence of maturation stage on agility performance gains after plyometric training: A systematic review and meta-analysis. *J Strength Cond Res*. 2017 Sep;31(9):2609-17.
111. Oliver JL, Rumpf MC. Speed development in youths. In: Lloyd R, Oliver JL, editors. *Strength and conditioning for young athletes: Science and application*. London/New York: Routledge; 2014. p. 80-93.
112. Oliver JL, Lloyd RS, Rumpf MC. Developing speed throughout childhood and adolescence: the role of growth, maturation and training. *Strength & Conditioning Journal*. 2013;35(3):42-8.
113. Ramirez-Campillo R, Alvarez C, Gentil P, Moran J, Garcia-Pinillos F, Alonso-Martinez AM, et al. Inter-individual variability in responses to 7 weeks of plyometric jump training in male youth soccer players. *Frontiers in physiology*. 2018;9:1156.
114. Faigenbaum AD, Lloyd RS, MacDonald J, Myer GD. Citius, Altius, Fortius: beneficial effects of resistance training for young athletes: Narrative review. *Br J Sports Med*. 2016 Jan;50(1):3-7.
115. Lloyd RS, Radnor JM, De Ste Croix MBA, Cronin JB, Oliver JL. Changes in Sprint and Jump Performances After Traditional, Plyometric, and Combined Resistance Training in Male Youth Pre- and Post-Peak Height Velocity. *Journal of Strength and Conditioning Research*. 2016 May;30(5):1239-47.
116. Lesinski M, Prieske O, Granacher U. Effects and dose-response relationships of resistance training on physical performance in youth athletes: a systematic review and meta-analysis. *Br J Sports Med*. 2016 Jul;50(13):781-95.
117. Ramirez-Campillo R, Moran J, Drury B, Williams M, Keogh JW, Chaabene H, et al. Effects of equal volume but different plyometric jump training intensities on components of physical fitness in physically active young males. *J Strength Cond Res*. 2019 Feb 6.
118. Nesser TW, Latin RW, Berg K, Prentice E. Physiological determinants of 40-meter sprint performance in young male athletes. *J Strength Cond Res*. 1996;10(4):263-7.
119. Morin JB, Bourdin M, Edouard P, Peyrot N, Samozino P, Lacour JR. Mechanical determinants of 100-m sprint running performance. *Eur J Appl Physiol*. 2012 Nov;112(11):3921-30.
120. Bishop DJ, Girard O. Determinants of team-sport performance: implications for altitude training by team-sport athletes. *Br J Sports Med*. 2013 Dec;47 Suppl 1:i17-21.
121. Bobbert MF, Gerritsen KG, Litjens MC, Van Soest AJ. Why is countermovement jump height greater than squat jump height? *Med Sci Sports Exerc*. 1996 Nov;28(11):1402-12.
122. Bobbert MF, Huijing PA, van Ingen Schenau GJ. Drop jumping. II. The influence of dropping height on the biomechanics of drop jumping. *Med Sci Sports Exerc*. 1987 Aug;19(4):339-46.
123. Bobbert MF, Huijing PA, van Ingen Schenau GJ. Drop jumping. I. The influence of jumping technique on the biomechanics of jumping. *Med Sci Sports Exerc*. 1987 Aug;19(4):332-8.
124. Dello Iacono A, Martone D, Milic M, Padulo J. Vertical- vs. horizontal-oriented drop jump training: chronic effects on explosive performances of elite handball players. *J Strength Cond Res*. 2017 Apr;31(4):921-31.
125. Rossler R, Donath L, Bizzini M, Faude O. A new injury prevention programme for children's football--FIFA 11+ Kids--can improve motor performance: a cluster-randomised controlled trial. *J Sports Sci*. 2016;34(6):549-56.
126. Choi SJ. Cellular mechanism of eccentric-induced muscle injury and its relationship with sarcomere heterogeneity. *Journal of exercise rehabilitation*. 2014;10(4):200-4.

127. Fransz DP, Huurnink A, Kingma I, de Boode VA, Heyligers IC, van Dieen JH. Performance on a Single-Legged Drop-Jump Landing Test Is Related to Increased Risk of Lateral Ankle Sprains Among Male Elite Soccer Players: A 3-Year Prospective Cohort Study. *Am J Sports Med.* 2018 Dec;46(14):3454-62.
128. Mujika I. Tapering and peaking for optimal performance: *Human Kinetics*; 2009.
129. Ramirez-Campillo R, Pereira LA, Andrade D, Méndez-Rebolledo G, de la Fuente CI, Castro-Sepulveda M, et al. Tapering strategies applied to plyometric jump training: a systematic review with meta-analysis of randomized-controlled trials. *J Sports Med Phys Fitness.* 2020 Jul 16.
130. de Villarreal ESS, González-Badillo JJ, Izquierdo M. Low and moderate plyometric training frequency produces greater jumping and sprinting gains compared with high frequency. *J Strength Cond Res.* 2008;22(3):715-25.
131. de Villarreal ESS, Requena B, Newton RU. Does plyometric training improve strength performance? A meta-analysis. *Journal of Science and Medicine in Sport.* 2010;13(5):513-22.
132. Brumitt J, Wilson V, Ellis N, Petersen J, Zita CJ, Reyes J. Preseason lower extremity functional test scores are not associated with lower quadrant injury - A validation study with normative data on 395 division III athletes. *Int J Sports Phys Ther.* 2018 Jun;13(3):410-21.
133. Abt G, Boreham C, Davison G, Jackson R, Nevill A, Wallace E, et al. Power, precision, and sample size estimation in sport and exercise science research. *J Sport Sci.* 2020:1-3.
134. Altman DG, Royston P. The cost of dichotomising continuous variables. *BMJ.* 2006;332(7549):1080.

Figure 1. PRISMA flow diagram.








Figure 2. Forest plot of increases in 10-m linear sprint performance in young male soccer players participating in plyometric jump training compared to controls, after  $\leq 14$  total PJT sessions (also comprising a duration of  $\leq 7$  weeks) or  $> 14$  total PJT sessions (also comprising a duration of  $> 7$  weeks) of intervention. Values shown are effect sizes with 95% confidence intervals (CI).

Table 1. Basic characteristics of included study participants and plyometric jump training programs.

	Treat	Ran	Age (years)*	Body mass (kg)*	Height (m)*	SPT	Fit	Frequency (days/week)	Duration (weeks)	NTJ
Asadi et al. (2018) pre-PHV (66)	WD	Yes	11.5	31.0	1.38	No	Moderate	2	6	720
Asadi et al. (2018) mid-PHV (66)			14.0	43.5	1.55					
Asadi et al. (2018) post-PHV (66)			16.6	60.6	1.72					
Beato et al. (2018) (67)	ID	Yes	17.0	69.2	1.75	NR	Moderate	1	6	360
Brito et al. (2014) (68)	WD	Yes	19.9	72.2	1.80	No	Normal	2	9	360 + 90 m
Chelly et al. (2010) (69)	WD	Yes	19.1	70.3	1.76	NR	Moderate - High	2	8	860
Chtara et al. (2017) (70)	ID	Yes	13.6	54.1	1.65	No	Moderate - High	2	6	632
Coratella et al. (2018) body mass (71)	WD	Yes	21.0	73.0	1.78	NR	Moderate	2	8	800
Coratella et al. (2018) loaded (71)			21.1							
Hammami et al. (2016) (72)	WD	Yes	15.7	59.0	1.76	No	High	2	8	722
Hammami et al. (2019) (73)	WD	Yes	15.7	58.9	1.75	NR	Moderate	2	8	722
Jlid et al. 2019 (74)	WD	Yes	11.8	36.5	1.43	NR	Moderate	2	8	1,596
Jlid et al. 2020 (75)	WD	Yes	19.0	67.6	1.76	NR	High	2	6	2,112
McKinlay et al. (2018) (76)	WD	No	12.6	47.2	1.58	No	Moderate	3	8	3,438
Meylan and Malatesta (2009) (77)	ID	No	13.2	48.6	1.59	No	Normal - Moderate	2	8	≥768
Michailidis et al. (2013) (78)	ID	Yes	10.7	42.5	1.47	No	Moderate	2	12	≥1,560
Nakamura et al. (2012) (79)	ID	NR	22.7	68.8	1.75	NR	High	2	3	360 + 198 m
Negra et al. 2016 (80)	ID	Yes	12.7	45.9	1.56	NR	Moderate	2	12	≥1,344
Negra et al. (in press) (20)	WD	Yes	12.7	43.7	1.59	NR	Moderate - High	5	8	1,284
Ramirez-Campillo et al. (2014) (81)	WD	Yes	13.2	47.9	1.54	No	Moderate	2	7	840
Ramirez-Campillo et al. (2014) 30s (82)	WD	Yes	10.4	37.0	1.41	No	Moderate	2	7	840
Ramirez-Campillo et al. (2014) 60 s (82)			10.4	37.2	1.41					
Ramirez-Campillo et al. (2014) 120s (82)			10.3	38.0	1.42					
Ramirez-Campillo et al. (2015) 24h (83)	WD	Yes	14.2	50.3	1.58	No	Normal - Moderate	2	6	2,400
Ramirez-Campillo et al. (2015) 48h (83)			14.1	51.8	1.59					
Ramirez-Campillo et al. (2015) no increase (84)	WD	Yes	12.8	53.9	1.60	No	Moderate	2	6	1,440
Ramirez-Campillo et al. (2015) increase (84)			13.0	53.8	1.61					2,160
Ramirez-Campillo et al. (2015) bilateral (85)			WD	Yes	11.0					43.5
Ramirez-Campillo et al. (2015) unilateral (85)	11.6	45.0			1.47					
Ramirez-Campillo et al. (2015) combined (85)	11.6	42.2			1.44					
Ramirez-Campillo et al. (2015) vertical (86)	WD	Yes	11.6	40.0	1.44	No	Moderate	2	6	1,610
Ramirez-Campillo et al. (2015) horizontal (86)			11.4	44.6	1.50					1,610
Ramirez-Campillo et al. (2015) combined (86)			11.2	44.1	1.41					1,440
Ramirez-Campillo et al. (2016) (22)	WD	Yes	20.4	68.4	1.71	No	Moderate	2	6	1,440
Ramirez-Campillo et al. (2018) fixed (87)	WD	Yes	13.9	46.7	1.53	No	Moderate	2	7	906
Ramirez-Campillo et al. (2018) optimal (87)			13.1	47.2	1.53					
Ramirez-Campillo et al. (2019) (88)	WD	Yes	13.2	48.6	1.54	No	Normal - Moderate	2	7	840
Ramirez-Campillo et al. (in press) before (89)	WD	Yes	16.9	64.9	1.72	No	Moderate	2	7	1,944
Ramirez-Campillo et al. (in press) after (89)			17.1	65.4	1.73					
Ramirez-Campillo et al. (in press) s-surface (90)	WD	Yes	12.9	44.4	1.54	No	Moderate	2	8	810
Ramirez-Campillo et al. (in press) com-surface (90)			12.1	45.6	1.59					
Rosas et al. (2016) jump (91)	WD	Yes	12.3	47.3	1.50	No	Moderate	2	6	1,152
Rosas et al. (2016) jump+haltere (91)			12.1	45.0	1.50					
Sedano et al. (2011) (92)	WD	Yes	18.4	70.7	1.74	Yes	Moderate - High	3	10	2,880
Sohnlein et al. (2014) (93)	ID	No	13.0	51.0	1.62	NR	Moderate	2	16	~3,467
Spinetti et al. (2016) (94)	ID	Yes	18.4	70.2	1.80	NR	High	2	8	1,440
Vaczi et al. (2013) (95)	WD	Yes	21.9	75.9	1.80	Yes	Moderate - High	2	6	925
Vlachopoulos et al. (2018) (96)	ID	Yes	13.8	49.3	1.61	NR	Normal - High	3 to 5	36	8800

\*: mean values reported for the experimental groups; **Fit**: physical fitness/playing level. The physical fitness/playing level was categorized following an adaptation of previous recommendations for plyometric jump training reviews and meta-analysis, and after consensus among authors from current review. In this sense, the physical fitness/playing level of experimental groups was classified as: i) **high**, for professional/elite athletes with regular enrollment in national and/or international competitions, highly trained participants with  $\geq 10$  training hours per week or  $\geq 6$  training sessions per week and a regularly scheduled official and friendly competitions, ii) **moderate**, for non-elite/professional athletes, with a regular attendance in regional and/or national competitions, between 5 - 9.9 training hours per week or 3-5 training sessions per week and a regularly scheduled official and friendly competitions, iii) **normal**, for recreational athletes with  $< 5$  training hours per week with sporadic or no competitions' participation, and school-age youths regularly involved in after-school soccer classes. The jump-training load was not considered as part of the regular training load of participants, hence, it was not considered to classify participants' physical fitness/playing level; **Freq**: frequency (sessions per week); **ID**: insufficiently described; **NR**: not clearly reported information; **NTJ**: number of total jumps; **Ran**: randomization; **SPT**: systematic plyometric jump training experience before intervention. If authors stated that participants had previous experience, a dichotomy characterization identifier was used as yes or no, without consideration the extent of the experience; **Treat**: treatment description quality. The plyometric jump training treatments were further categorized as i) **well described**, when treatment description allowed for adequate study replication, including the reporting of duration, frequency, intensity, type of exercises, sets, and repetitions, or ii) **insufficiently described**, when the treatment description omitted any of the six aforementioned descriptors; **WD**: well described.

Table 2. Physiotherapy Evidence Database (PEDro) scale ratings.												
	N° 1*	N° 2	N° 3	N° 4	N° 5	N° 6	N° 7	N° 8	N° 9	N° 10	N° 11	Total**
Asadi et al. (2018) (66)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Beato et al. (2018) (67)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Brito et al. (2014) (68)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Chelly et al. (2010) (69)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Chtara et al. (2017) (70)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Coratella et al.(2018) (71)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Hammami et al. (2016) (72)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Hammami et al. (2019) (73)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Jlid et al. (2019) (74)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Jlid et al. (2020) (75)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
McKinlay et al. (2018) (76)	Yes	No	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	6
Meylan and Malatesta (2009) (77)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Michailidis et al.(2013) (78)	Yes	Yes	Yes	Yes	No	No	No	No	Yes	Yes	Yes	6
Nakamura et al. (2012) (79)	Yes	No	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	6
Negra et al. (2016) (80)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Negra et al. (in press) (20)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Ramirez-Campillo et al. (2014) (81)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Ramirez-Campillo et al. (2014) (30 vs. 60 vs. 120s) (82)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Ramirez-Campillo et al. (2015) (24 vs 48h) (83)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Ramirez-Campillo et al. (2015) (progressive vs- non-progressive) (84)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Ramirez-Campillo et al. (2015) (unilateral vs. bilateral vs. combi) (85)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Ramirez-Campillo et al. (2015) (vertical vs. horizontal vs. combi) (86)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Ramirez-Campillo et al. (2016) (22)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Ramirez-Campillo et al. (2018) (fixed vs. optimal RSI) (87)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Ramirez-Campillo et al. (2019) (88)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Ramirez-Campillo et al. 2020 (before vs. after) (89)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Ramirez-Campillo et al. (in press) (surface vs. comb surfaces) (90)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Rosas et al. (2016) (91)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Sedano et al. (2011) (92)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Sohnlein et al. (2014) (93)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Spineti et al. (2016) (94)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Vaczi et al. (2013) (95)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
Vlachopoulos et al. (2018) (96)	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7

\*: PEDro scale items number. A detailed explanation for each PEDro scale item can be accessed at <https://www.pedro.org.au/english/downloads/pedro-scale>  
(Access for this review: March 1, 2020); \*\*: the total number of points from a possible maximal of 10.

Table 3. Results of the multivariate random-effect meta-regression for training variables to predict PJT effects on vertical jump and linear sprint performance in young male soccer players.

Covariate	Coefficient	95% CI, lower	95% CI, upper	Z value	P value
<b>CMJ (N = 27)</b>					
Intercept	-0.7915	-4.7029	3.1199	-0.40	0.6917
Frequency	0.4507	-0.9436	1.8450	0.63	0.5264
Training duration	0.2979	-0.0519	0.6477	1.67	0.0951
Total sessions	-0.0786	-0.1710	0.0138	-1.67	0.0954
Participant's chronological age	-0.0211	-0.1332	0.0911	-0.37	0.7128
<b>10-m linear sprint (N = 12)</b>					
Intercept	4.2645	-6.8485	15.3775	0.75	0.4520
Frequency	-1.1923	-6.4320	4.0475	-0.45	0.6556
Training duration	-0.5130	-1.8177	0.7917	-0.77	0.4409
Total sessions	0.1909	-0.4495	0.8313	0.58	0.5591
Participant's chronological age	-0.0924	-0.2346	0.0498	-1.27	0.2029
N: number of study groups; CI: Confidence interval; PJT: plyometric jump training; CMJ: countermovement jump height.					

### **Electronic Supplementary Material Table S1**

**Article title:** Effects of plyometric jump training on jump and sprint performance in young male soccer players: a systematic review and meta-analysis.

**Journal name:** Sports Medicine.

**Author names:** Rodrigo Ramirez-Campillo, Daniel Castillo, Javier Raya-González Jason Moran, Eduardo Sáez de Villarreal, Rhodri S. Lloyd.

**Affiliation and e-mail of the corresponding author:** Rodrigo Ramirez-Campillo, PhD. Human Performance Laboratory. Department of Physical Activity Sciences. Universidad de Los Lagos. Osorno, Chile. Mail: [r.ramirez@ulagos.cl](mailto:r.ramirez@ulagos.cl)

Table S1. Example for 5-m linear sprint Hedges' g effect size (ES) calculation for between-group comparisons.										
	Plyometric group					Control group				
	Pre		Post		n	Pre		Post		n
	Mean	SD	Mean	SD		Mean	SD	Mean	SD	
Sohnlein et al. (2014)	1.06*	0.04	1.02	0.04	12	1.1	0.05	1.08	0.03	10

\*: data in seconds. SD: standard deviation; ES: effect size. Pre and post: before and after intervention.

All analyses were carried out using the Comprehensive Meta-Analysis program (version 2; Biostat, Englewood, NJ, USA).

### Standardized difference in means (standardize by change SD)

MeanChange(1) = Group 1 mean difference

MeanChange(2) = Group 2 mean difference

RawDiff = MeanChange(1) - MeanChange(2)

$SDChange(1) = \sqrt{SDPre(1)^2 + SDPost(1)^2 - 2 * CorrPrePost * SDPre(1) * SDPost(1)}$

$SDChange(2) = \sqrt{SDPre(2)^2 + SDPost(2)^2 - 2 * CorrPrePost * SDPre(2) * SDPost(2)}$

$SDChangePooled = \sqrt{((n(1) - 1) * SDChange(1)^2 + (n(2) - 1) * SDChange(2)^2) / (n(1) + n(2) - 2)}$

StdChangeDiff = RawDiff / SDChangePooled

$StdChangeDiffSE = \sqrt{1 / n(1) + 1 / n(2) + StdChangeDiff^2 / (2 * (n(1) + n(2)))}$

RawDiff = -0.040 - -0.020 = -0.020

$SDChangePooled = \sqrt{((12 - 1) * 0.031^2 + (10 - 1) * 0.036^2) / (12 + 10 - 2)} = 0.033$

StdChangeDiff = -0.020 / 0.033 = -0.599

$StdChangeDiffSE = \sqrt{1 / 12 + 1 / 10 + -0.599^2 / (2 * (12 + 10))} = 0.438$

### Standardized mean difference corrected for bias (Hedges' g)

The program computes the Standardized mean difference (d) and then multiplies d by a correction factor (J) to compute g.

Correction factor J

$J = 1 - (3 / (4 * df - 1))$

Where df = NTot - 2

$J = 1 - (3 / (4 * 20 - 1)) = 0.962$

Computation of g

$g = d * J$

$StdErr(g) = StdErr(d) * J$

$$\text{Variance}(g) = \text{StdErr}(g) ^ 2$$

$$g = -0.599 * 0.962 = \mathbf{-0.577}$$

$$\text{StdErr}(g) = 0.438 * 0.962 = 0.421$$

$$\text{Variance}(g) = 0.421 ^ 2 = 0.177$$

Final result for 5-m linear sprint Hedges' *g* effect size (ES) calculation for between-group comparisons: **-0.577**.

In this example, a negative value means a reduction in the time needed to complete a 5-m linear sprint, favouring the plyometric group in comparison with the control group.

## **Compliance with Ethical Standards**

### *Funding*

No sources of funding were used to assist in the preparation of this article.

### *Conflicts of Interest*

Rodrigo Ramirez-Campillo, Daniel Castillo, Javier Raya-González, Jason Moran, Eduardo Sáez de Villarreal and Rhodri Lloyd declare that they have no conflicts of interest relevant to the content of this review.

### *Authorship Contributions*

RRC, DC and JRG wrote the first draft of the manuscript. RRC, DC and JRG collected data. RRC and JM analyzed and interpreted the data. ESV and RD revised the original manuscript. All authors read and approved the final manuscript.




## **Electronic Supplementary Material Fig. S1 to S9**

**Article title:** Effects of plyometric jump training on jump and sprint performance in young male soccer players: a systematic review and meta-analysis.

**Journal name:** Sports Medicine.


**Author names:** Rodrigo Ramirez-Campillo, Daniel Castillo, Javier Raya-González Jason Moran, Eduardo Sáez de Villarreal, Rhodri S. Lloyd.

**Affiliation and e-mail of the corresponding author:** Rodrigo Ramirez-Campillo, PhD. Human Performance Laboratory. Department of Physical Activity Sciences. Universidad de Los Lagos. Osorno, Chile. Mail: [r.ramirez@ulagos.cl](mailto:r.ramirez@ulagos.cl)




Electronic Supplementary Material Fig. S1. Forest plot of increases in countermovement jump height performance in young male soccer players participating in plyometric jump training compared to controls. Values shown are effect sizes with 95% confidence intervals (CI).

<u>Study name</u>	<u>Statistics for each study</u>						
	Hedges's g	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value
Ramirez-Campillo et al. (2015) bilateral	0.854	0.399	0.159	0.072	1.636	2.140	0.032
Ramirez-Campillo et al. (2015) combine	0.443	0.434	0.188	-0.407	1.294	1.022	0.307
Ramirez-Campillo et al. (2015) combined	0.579	0.389	0.152	-0.184	1.342	1.488	0.137
Ramirez-Campillo et al. (2015) horizontal	0.145	0.429	0.184	-0.696	0.986	0.338	0.735
Ramirez-Campillo et al. (2015) increase	0.050	0.473	0.224	-0.877	0.977	0.106	0.916
Ramirez-Campillo et al. (2015) no increase	0.036	0.473	0.224	-0.890	0.963	0.077	0.939
Ramirez-Campillo et al. (2015) unilateral	0.327	0.359	0.129	-0.376	1.030	0.912	0.362
Ramirez-Campillo et al. (2015) vertical	0.433	0.434	0.188	-0.417	1.283	0.998	0.318
Ramirez-Campillo et al. (2016)	1.002	0.322	0.104	0.371	1.633	3.114	0.002
Rosas et al. (2016) jump	0.149	0.303	0.092	-0.445	0.743	0.491	0.623
Rosas et al. (2016) jump+halteres	0.327	0.305	0.093	-0.270	0.925	1.073	0.283
Sedano et al. (2011)	1.478	0.467	0.218	0.563	2.393	3.166	0.002
	0.481	0.117	0.014	0.252	0.710	4.120	0.000




Electronic Supplementary Material Fig. S2. Forest plot of increases in countermovement jump with arms height performance in young male soccer players participating in plyometric jump training compared to controls. Values shown are effect sizes with 95% confidence intervals (CI).

Study name	Statistics for each study						
	Hedges's g	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value
Chelly et al. (2010)	1.451	0.456	0.208	0.558	2.344	3.184	0.001
Hammami et al. (2019)	1.440	0.430	0.185	0.597	2.283	3.348	0.001
Jlid et al. (2019)	1.359	0.409	0.168	0.557	2.162	3.320	0.001
Jlid et al. (2020)	0.148	0.374	0.140	-0.585	0.882	0.397	0.691
McKinlay et al. (2018)	0.854	0.391	0.153	0.088	1.621	2.184	0.029
Meylan and Makatesta (2009)	0.634	0.400	0.160	-0.149	1.418	1.587	0.113
Ramirez-Campillo et al. (2015) 24h	0.340	0.192	0.037	-0.035	0.716	1.775	0.076
Ramirez-Campillo et al. (2015) 48h	0.295	0.189	0.036	-0.075	0.665	1.562	0.118
Sedano et al. (2011)	-0.836	0.429	0.184	-1.677	0.005	-1.949	0.051
Vaczi et al. (2013)	1.788	0.349	0.122	1.104	2.472	5.126	0.000
	0.725	0.220	0.048	0.293	1.156	3.292	0.001



Electronic Supplementary Material Fig. S3. Forest plot of increases in squat jump height performance in young male soccer players participating in plyometric jump training compared to controls. Values shown are effect sizes with 95% confidence intervals (CI).

<u>Study name</u>	<u>Statistics for each study</u>						
	Hedges's g	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value
Brito et al. (2014)	-1.588	0.404	0.163	-2.379	-0.797	-3.936	0.000
Chelly et al. (2010)	-0.944	0.426	0.181	-1.779	-0.110	-2.218	0.027
Hammami et al. (2016)	-1.597	0.425	0.181	-2.431	-0.763	-3.755	0.000
Hammami et al. (2019)	-2.235	0.491	0.241	-3.198	-1.273	-4.551	0.000
Nakamura et al. (2012)	1.237	0.548	0.300	0.163	2.311	2.258	0.024
Sohnlein et al. (2014)	-0.577	0.421	0.177	-1.402	0.248	-1.370	0.171
	-0.977	0.433	0.187	-1.825	-0.130	-2.259	0.024




Electronic Supplementary Material Fig. S4. Forest plot of increases in 5-m linear sprint performance in young male soccer players participating in plyometric jump training compared to controls. Values shown are effect sizes with 95% confidence intervals (CI).

**Study name**


**Statistics for each study**

**Hedges's g and 95% CI**

	Hedges's g	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value
Beato et al. (2018)	-0.431	0.449	0.202	-1.312	0.450	-0.959	0.337
Chtara et al. (2017)	-0.784	0.446	0.199	-1.658	0.090	-1.757	0.079
Coratella et al.(2018) body mass	-1.258	0.379	0.144	-2.001	-0.516	-3.322	0.001
Coratella et al.(2018) loaded	-2.388	0.456	0.208	-3.281	-1.494	-5.237	0.000
Hammami et al. (2016)	-1.235	0.403	0.163	-2.026	-0.445	-3.064	0.002
Meylan and Malatesta (2009)	0.138	0.390	0.152	-0.627	0.902	0.353	0.724
Michailidis et al. (2013)	-0.888	0.308	0.095	-1.492	-0.284	-2.881	0.004
Nakamura et al. (2012)	0.929	0.536	0.287	-0.121	1.979	1.733	0.083
Ramirez-Campillo et al. (2015) no increase	0.040	0.473	0.224	-0.887	0.967	0.084	0.933
Ramirez-Campillo et al. (2015) increase	-0.103	0.473	0.224	-1.030	0.825	-0.217	0.828
Sedano et al. (2011)	-0.407	0.415	0.172	-1.219	0.406	-0.980	0.327
Sohnlein et al. (2014)	-0.551	0.420	0.177	-1.374	0.273	-1.311	0.190
	-0.604	0.223	0.050	-1.040	-0.167	-2.711	0.007



Electronic Supplementary Material Fig. S5. Forest plot of increases in 10-m linear sprint performance in young male soccer players participating in plyometric jump training compared to controls. Values shown are effect sizes with 95% confidence intervals (CI).




Electronic Supplementary Material Fig. S6. Forest plot of increases in 20-m linear sprint performance in young male soccer players participating in plyometric jump training compared to controls. Values shown are effect sizes with 95% confidence intervals (CI).

Study name

Statistics for each study


Hedges's g and 95% CI

Study name	Hedges's g	Standard error	Statistics for each study		Z-Value	p-Value	
			Variance	Lower limit			Upper limit
Beato et al. (2018)	-0.159	0.445	0.198	-1.031	0.712	-0.358	0.720
Chtara et al. (2017)	-0.491	0.435	0.189	-1.344	0.362	-1.129	0.259
Coratella et al.(2018) body mass	0.000	0.345	0.119	-0.675	0.675	0.000	1.000
Coratella et al.(2018) loaded	-1.258	0.379	0.144	-2.001	-0.516	-3.322	0.001
Hammani et al. (2016)	-0.892	0.387	0.150	-1.651	-0.134	-2.308	0.021
Michailidis et al. (2013)	-0.500	0.298	0.089	-1.085	0.085	-1.676	0.094
Ramirez-Campillo et al. (2015) bilateral	-0.522	0.388	0.150	-1.282	0.238	-1.345	0.179
Ramirez-Campillo et al. (2015) combine	-0.650	0.440	0.194	-1.513	0.214	-1.475	0.140
Ramirez-Campillo et al. (2015) combined	-0.648	0.391	0.153	-1.415	0.119	-1.655	0.098
Ramirez-Campillo et al. (2015) horizontal	-0.376	0.432	0.187	-1.223	0.472	-0.869	0.385
Ramirez-Campillo et al. (2015) unilateral	-0.901	0.375	0.140	-1.635	-0.167	-2.405	0.016
Ramirez-Campillo et al. (2015) vertical	-0.210	0.430	0.185	-1.052	0.632	-0.490	0.624
Ramirez-Campillo et al. (2016)	-1.928	0.369	0.136	-2.650	-1.205	-5.229	0.000
Ramirez-Campillo et al. (in press) com-surface	-0.865	0.512	0.262	-1.869	0.139	-1.689	0.091
Ramirez-Campillo et al. (in press) s-surface	-0.453	0.494	0.244	-1.422	0.515	-0.918	0.359
Sohnlein et al. (2014)	-0.148	0.429	0.184	-0.988	0.693	-0.344	0.731
	-0.637	0.127	0.016	-0.886	-0.388	-5.019	0.000



Electronic Supplementary Material Fig. S7. Forest plot of increases in 30-m linear sprint performance in young male soccer players participating in plyometric jump training compared to controls. Values shown are effect sizes with 95% confidence intervals (CI).





Electronic Supplementary Material Fig. S8. Forest plot of increases in 40-m linear sprint performance in young male soccer players participating in plyometric jump training compared to controls. Values shown are effect sizes with 95% confidence intervals (CI).

### Compliance with Ethical Standards

#### *Funding*

No sources of funding were used to assist in the preparation of this article.

#### *Conflicts of Interest*

Rodrigo Ramirez-Campillo, Daniel Castillo, Javier Raya-González, Jason Moran, Eduardo Sáez de Villarreal and Rhodri Lloyd declare that they have no conflicts of interest relevant to the content of this review.

#### *Authorship Contributions*

RRC, DC and JRG wrote the first draft of the manuscript. RRC, DC and JRG collected data. RRC and JM analyzed and interpreted the data. ESV and RD revised the original manuscript. All authors read and approved the final manuscript.

Electronic Supplementary Material Appendix S1

**Article title:** **Article title:** Effects of plyometric jump training on jump and sprint performance in young male soccer players: a systematic review and meta-analysis.

**Journal name:** Sports Medicine.

**Author names:** Rodrigo Ramirez-Campillo, Daniel Castillo, Javier Raya-González Jason Moran, Eduardo Sáez de Villarreal, Rhodri S. Lloyd.

**Affiliation and e-mail of the corresponding author:** Rodrigo Ramirez-Campillo, PhD. Human Performance Laboratory. Department of Physical Activity Sciences. Universidad de Los Lagos. Osorno, Chile. Mail: [r.ramirez@ulagos.cl](mailto:r.ramirez@ulagos.cl)

## Countermovement jump additional analysis

Regarding interventions with a frequency of  $\leq 2$  sessions per week (27 study groups; ES = 0.81 [95%CI = 0.57 to 1.05],  $p < 0.001$ ; within-group  $I^2 = 68.6\%$ ) and those with  $> 2$  sessions per week (3 study groups; ES = 0.59 [95%CI = -0.37 to 1.55],  $p = 0.228$ ; within-group  $I^2 = 77.5\%$ ) only the former induced significant improvement on countermovement jump height performance, although without between-group differences (between-group  $p = 0.676$ ).

Regarding interventions with a duration of  $\leq 7$  weeks, also comprising  $\leq 14$  total PJT sessions (17 study groups; ES = 0.71 [95%CI = 0.42 to 0.99],  $p < 0.001$ ; within-group  $I^2 = 68.8\%$ ) and those with  $> 7$  weeks, also comprising  $> 14$  total PJT sessions (13 study groups; ES = 0.89 [95%CI = 0.49 to 1.29],  $p < 0.001$ ; within-group  $I^2 = 68.0\%$ ) both induced a similar (between-group  $p = 0.476$ ) significant improvement on countermovement jump height performance.

Regarding interventions in players with  $\leq 13.2$  years (14 study groups; ES = 0.81 [95%CI = 0.40 to 1.22],  $p < 0.001$ ; within-group  $I^2 = 76.3\%$ ) and those with  $> 13.2$  years (16 study groups; ES = 0.77 [95%CI = 0.49 to 1.04],  $p < 0.001$ ; within-group  $I^2 = 60.3\%$ ), both induced a similar (between-group  $p = 0.855$ ) significant improvement on countermovement jump height performance.

Regarding interventions in players U-17 (22 study groups; ES = 0.85 [95%CI = 0.56 to 1.14],  $p < 0.001$ ; within-group  $I^2 = 73.5\%$ ) and U-20 (5 study groups; ES = 0.66 [95%CI = 0.04 to 1.27],  $p = 0.036$ ; within-group  $I^2 = 64.6\%$ ), all induced a significant improvement on countermovement jump height performance, although without between-group differences ( $p = 0.571$ ).

## Squat jump additional analysis

Regarding interventions in players with  $\leq 13.2$  years (3 study groups; ES = 0.94 [95%CI = 0.49 to 1.39],  $p < 0.001$ ; within-group  $I^2 = 0.0\%$ ) and those with  $> 13.2$  years (7 study groups; ES = 0.64 [95%CI = 0.09 to 1.20],  $p = 0.024$ ; within-group  $I^2 = 83.0\%$ ), both induced a similar (between-group  $p = 0.411$ ) significant improvement on squat jump height performance.

Regarding interventions with a duration of  $\leq 7$  weeks, also comprising  $\leq 14$  total PJT sessions (5 study groups; ES = 0.74 [95%CI = 0.18 to 1.29],  $p = 0.009$ ; within-group  $I^2 = 80.7\%$ ) and those with  $> 7$  weeks, also comprising  $> 14$  total PJT sessions (5 study groups; ES = 0.71 [95%CI = -0.09 to 1.50],  $p = 0.082$ ; within-group  $I^2 = 78.6\%$ ), only the former induced a significant improvement on squat jump height performance, although without between-group differences ( $p = 0.949$ ).

## 10-m linear sprint additional analysis

Regarding interventions with  $\leq 14$  total PJT sessions, also comprising a duration of  $\leq 7$  weeks (5 study groups; ES = 0.11 [95%CI = 0.65 to -0.42],  $p = 0.677$ ; within-group  $I^2 = 39.7\%$ ) and those with  $> 14$  sessions, also comprising a duration of  $> 7$  weeks (7 study groups; ES = 0.93 [95%CI = 1.47

to 0.38],  $p = 0.001$ ; within-group  $I^2 = 71.9\%$ ), only the latter induced a significant improvement on 10-m linear sprint performance, with a significant between-group difference ( $p = 0.038$ ).

Regarding interventions in players with  $\leq 13.2$  years (6 study groups; ES = 0.41 [95%CI = 0.78 to 0.03],  $p = 0.032$ ; within-group  $I^2 = 23.1\%$ ) and those with  $>13.2$  years (6 study groups; ES = 0.83 [95%CI = 1.62 to 0.03],  $p = 0.043$ ; within-group  $I^2 = 81.1\%$ ), both induced a significant improvement on 10-m linear sprint performance, without significant between-group differences ( $p = 0.354$ ).

### **20-m linear sprint additional analysis**

Regarding interventions with a duration of  $\leq 7$  weeks, also comprising  $\leq 14$  total PJT sessions (15 study groups; ES = 0.50 [95%CI = 0.73 to 0.27],  $p < 0.001$ ; within-group  $I^2 = 47.1\%$ ) and those with  $>7$  weeks, also comprising  $>14$  total PJT sessions (6 study groups; ES = 1.09 [95%CI = 2.11 to 0.07],  $p = 0.036$ ; within-group  $I^2 = 89.4\%$ ) both induced significant improvement on 20-m linear sprint performance, without significant between-group differences (between-group  $p = 0.270$ ).

Regarding interventions in players with  $\leq 13.2$  years (10 study groups; ES = 0.40 [95%CI = 0.61 to 0.19],  $p < 0.001$ ; within-group  $I^2 = 0.0\%$ ) and those with  $>13.2$  years (11 study groups; ES = 0.89 [95%CI = 1.40 to 0.38],  $p = 0.001$ ; within-group  $I^2 = 83.8\%$ ), both induced significant improvement on 20-m linear sprint performance, without significant between-group differences (between-group  $p = 0.082$ ).

### **30-m linear sprint additional analysis**

Regarding interventions with a duration of  $\leq 7$  weeks, also comprising  $\leq 14$  total PJT sessions (9 study groups; ES = 0.68 [95%CI = 1.05 to 0.31],  $p < 0.001$ ; within-group  $I^2 = 47.3\%$ ) and those with  $>7$  weeks, also comprising  $>14$  total PJT sessions (7 study groups; ES = 0.57 [95%CI = 0.91 to 0.23],  $p = 0.001$ ; within-group  $I^2 = 26.3\%$ ) both induced a similar (between-group  $p = 0.667$ ) significant improvement on 30-m linear sprint performance.

Regarding interventions in players with  $\leq 13.2$  years (11 study groups; ES = 0.53 [95%CI = 0.77 to 0.29],  $p < 0.001$ ; within-group  $I^2 = 0.0\%$ ) and those with  $>13.2$  years (5 study groups; ES = 0.85 [95%CI = 1.57 to -0.14],  $p = 0.019$ ; within-group  $I^2 = 78.0\%$ ), both induced a significant improvement on 30-m linear sprint performance, although without between-group differences ( $p = 0.395$ ).

Additional analyses, including PJT frequency, PJT duration, and total number of PJT sessions, participant's age, and FIFA age categories were not possible as  $<3$  studies were available for at least one of the moderators.

## **Compliance with Ethical Standards**

### *Funding*

No sources of funding were used to assist in the preparation of this article.

### *Conflicts of Interest*

Rodrigo Ramirez-Campillo, Daniel Castillo, Javier Raya-González, Jason Moran, Eduardo Sáez de Villarreal and Rhodri Lloyd declare that they have no conflicts of interest relevant to the content of this review.

### *Authorship Contributions*

RRC, DC and JRG wrote the first draft of the manuscript. RRC, DC and JRG collected data. RRC and JM analyzed and interpreted the data. ESV and RD revised the original manuscript. All authors read and approved the final manuscript.