

## **Endothelial dysfunction in COVID-19: a position paper of the ESC Working Group for Atherosclerosis and Vascular Biology, and the ESC Council of Basic Cardiovascular Science**

Paul C. Evans<sup>1\*</sup>, G. Ed Rainger<sup>2</sup>, Justin C. Mason<sup>3</sup>, Tomasz J. Guzik<sup>4</sup>, Elena Osto<sup>5</sup>, Zania Stamataki<sup>6</sup>, Desley Neil<sup>2</sup>, Imo E. Hoefer<sup>7</sup>, Maria Fragiadaki<sup>1</sup>, Johannes Waltenberger<sup>8</sup>, Christian Weber<sup>9</sup>, Marie-Luce Bochaton-Piallat<sup>10</sup>, Magnus Bäck<sup>11\*</sup>.

<sup>1</sup>Department of Infection, Immunity and Cardiovascular Disease, University of Sheffield, Sheffield, UK and Insigneo Institute for In Silico Medicine, Sheffield, UK.

<sup>2</sup>Institute of Cardiovascular Sciences, College of Medical and Dental Sciences, University of Birmingham, Birmingham B15 2TT, UK

<sup>3</sup>Vascular Science, National Heart and Lung Institute, Imperial College London and Rheumatology Department, Imperial College Healthcare NHS Trust, Hammersmith Hospital, London, UK.

<sup>4</sup>Institute of Cardiovascular and Medical Sciences, University of Glasgow, Glasgow, UK and Department of Medicine, Jagiellonian University Collegium Medicum, Cracow, Poland

<sup>5</sup>University and University Hospital Zurich, Institute of Clinical Chemistry and University Heart Center, Zurich, Switzerland; Swiss Federal Institute of Technology, Laboratory of Translational Nutrition Biology, Zurich, Switzerland.

<sup>6</sup>Institute of Immunology and Immunotherapy College of Medical and Dental Sciences, University of Birmingham, Birmingham B15 2TT, UK

<sup>7</sup>Central Diagnostic Laboratory, University Medical Centre Utrecht, Netherlands

<sup>8</sup>Department of Cardiovascular Medicine, Medical Faculty, University of Münster, Münster, Germany; and SRH Central Hospital Suhl, Suhl, Germany

<sup>9</sup>Institute for Cardiovascular Prevention (IPEK), Ludwig-Maximilian-Universität (LMU) München and German Center for Cardiovascular Research (DZHK), partner site Munich Heart Alliance, and Munich Cluster for Systems Neurology (SyNergy), Munich, Germany; Department of Biochemistry, Cardiovascular Research Institute Maastricht (CARIM), Maastricht University, Maastricht, the Netherlands.

<sup>10</sup>Department of Pathology and Immunology, Faculty of Medicine, University of Geneva, Geneva, Switzerland.

<sup>11</sup>Center for Molecular Medicine and Department of Cardiology, Karolinska University Hospital, Solna, Stockholm, Sweden, and INSERM U1116, Université de Lorraine, Centre Hospitalier Régional Universitaire de Nancy, 54505 Vandoeuvre les Nancy, France.

\* Address correspondence to:

Professor Magnus Back MD, PhD, FESC  
Center for Molecular Medicine and Department of Cardiology,  
Karolinska University Hospital, M85  
14186 Stockholm, Sweden.  
Email: Magnus.Back@ki.se

Professor Paul Evans PhD, FESC, FRSB  
Department of Infection, Immunity and Cardiovascular Disease,  
Bateson Centre & INSIGNEO Institute,  
University of Sheffield,  
Sheffield S10 2RX, UK.  
Email: paul.evans@sheffield.ac.uk

**Abstract.**

The COVID-19 pandemic is an unprecedented healthcare emergency causing mortality and illness across the world. Although primarily affecting the lungs, the SARS-CoV2 virus also affects the cardiovascular system. In addition to cardiac effects e.g. myocarditis, arrhythmias, and myocardial damage, the vasculature is affected in COVID-19, both directly by the SARS-CoV-2 virus, and indirectly as a result of a systemic inflammatory cytokine storm. This includes the role of vascular endothelium in the recruitment of inflammatory leukocytes where they contribute to tissue damage and cytokine release, which are key drivers of acute respiratory distress syndrome (ARDS), in disseminated intravascular coagulation and cardiovascular complications in COVID-19. There is also evidence linking endothelial cells (EC) to SARS-CoV-2 infection including: (1) the expression and function of its receptor angiotensin converting enzyme (ACE) 2 in the vasculature, (2) the prevalence of a Kawasaki Disease-like syndrome (vasculitis) in COVID-19, and (3) evidence of EC infection with SARS-CoV-2 in patients with fatal COVID-19. Here, the working Group on Atherosclerosis and Vascular Biology together with the Council of Basic Cardiovascular Science of the European Society of Cardiology provide a Position Statement on the importance of the endothelium in the underlying pathophysiology behind the clinical presentation in COVID-19 and identify key questions for future research to address. We propose that endothelial biomarkers and tests of function (e.g. flow mediated dilatation) should be evaluated for their value in the risk stratification of COVID-19 patients. A better understanding of the effects of SARS-CoV-2 on endothelial biology in both the micro- and macro-vasculature is required, and endothelial function testing should be considered in the follow-up of convalescent COVID-19 patients for early detection of long-term cardiovascular complications.

## Introduction

Coronavirus Disease 2019 (COVID-19) is caused by a single strand RNA virus called severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) that likely crossed from bats to humans following a gain-of-function mutation in the spike protein that allows infection of human cells<sup>1</sup>. Though primarily known as a disease affecting the respiratory system, cardiovascular complications are common in COVID-19<sup>2 3</sup>. These include myocarditis and myocardial injury that may lead to heart failure. Myocardial infarction (MI) and Takotsubo syndrome has been reported in patients with COVID-19<sup>4 5</sup>. An increase in troponin levels in COVID-19 is associated with more severe disease and mortality, underlining that myocardial injury as a prognostic factor<sup>6</sup>. Increased levels of N-terminal pro B-type natriuretic peptide (NT-proBNP) is also an independent predictor of in-hospital death in COVID-19 patients, but at lower levels than in heart failure<sup>7</sup>, suggesting that relatively mild alteration in cardiac function may critically determine outcome in COVID-19. In addition, arrhythmias are remarkably prevalent among patients with COVID-19 with an incidence of 16.7% reported in a Chinese cohort<sup>8</sup>. Cardiomyocytes are known to express angiotensin converting enzyme 2 (ACE2)<sup>9</sup>, which is the receptor for SARS-CoV-2, however direct evidence of cardiomyocyte infection is lacking.

In addition, underlying cardiovascular disease may aggravate the clinical course of disease in COVID-19. Meta-analysis of data from Chinese cohorts revealed that fatality rates in hospitalised patients with COVID-19 were elevated in those with cardiovascular disease (10.5%), diabetes (7.3%) or hypertension (6%) compared to patients without comorbidities (0.9%)<sup>10</sup>. Cohorts of patients from Italy and U.S.<sup>11</sup> also show that hypertension, diabetes and obesity are common co-morbidities.

Increased D-dimer (fibrin degradation product) levels is a marker of adverse COVID-19 outcomes<sup>12</sup> and disseminated intravascular coagulation with significant risk of venous thromboembolism (VTE) and ischaemic stroke has been reported<sup>13 14 15</sup>. A study of systematic duplex ultrasound in 26 severe COVID-19 patients reported a cumulative VTE incidence of 69% despite anti-coagulation treatment<sup>16</sup>. Ischemic stroke has been linked to COVID-19 progression because it had a higher incidence in patients with severe disease compared to those with milder symptoms<sup>17</sup>. A report of six ischaemic stroke cases from the UK revealed an association with elevated D-dimers, systemic inflammatory changes, and anti-phospholipid antibodies, which are indicative of hypercoagulability<sup>18</sup>. However, caution should be exercised in attributing stroke to viral infection since there were several potential confounding factors including hypertension and atrial fibrillation, which enhance stroke risk.

We refer the reader to Guzik et al<sup>19</sup> for a more complete description of cardiac effects in COVID-19. The aim of this Position Statement from the Working Group on Atherosclerosis and Vascular Biology together with the Council of Basic Cardiovascular Science of the European Society of Cardiology is to draw attention to the importance of the endothelium in COVID-19 and to encourage research on endothelial dysfunction and biomarkers to tackle the COVID-19 pandemic and its potential long-term cardiovascular complications.

## **Endothelium and COVID-19**

The characteristic hyperinflammatory and pro-coagulatory state of COVID-19 implies a critical role of the endothelium, both as an effector contributing to inflammation and thrombosis, and as a target organ, whose dysfunction may contribute to poor outcome<sup>20</sup>. Of particular note, there is also evidence of SARS-CoV-2 infection of vascular ECs<sup>21-24</sup>.

The vascular endothelium forms a critical interface between the circulatory system and underlying tissues and has vital and ubiquitous roles in cardiovascular homeostasis by regulating the transport of cells, nutrients and metabolites between the circulation and underlying tissues<sup>25</sup>. Several risk factors for cardiovascular disease including diabetes, obesity, dyslipidemia, smoking and disturbed blood flow can induce EC dysfunction which is characterised by a spectrum of phenotypes<sup>26,27,28</sup>. These include loss of integrity (e.g. via apoptosis) which is associated with increased permeability; induction of cytokines and adhesion molecules to capture inflammatory cells from the circulation; metabolic changes; a pro-thrombotic phenotype and de-differentiation<sup>29,26</sup>. This is exemplified in multiple studies showing that quantitation of EC function provides a useful marker for early disease detection and stratification of patients with cardiovascular disease<sup>30,31</sup>. The European Society of Cardiology recently reviewed this subject and called for further investigation of EC pathophysiological states, optimisation and standardisation of methodologies for clinical measurement of EC function as well as larger clinical trials to establish reference values and assess clinical utility<sup>25</sup>.

Emerging evidence indicates that EC dysfunction is a central feature of COVID-19. This is evidenced by the critical role of vascular endothelium in inflammation, which is the key driver of cytokine dysregulation in ARDS as well as multiple cardiovascular pathologies. Additionally, the pro-thrombotic phenotype and disseminated intravascular coagulation observed in COVID-19 reflect dysfunction of EC, which enhances thrombosis by reduced integrity leading to exposure of pro-thrombotic sub-endothelial material, capture of platelets and regulation of clotting cascades, thrombin activation and fibrin production<sup>32</sup>.

There are several other lines of evidence that substantiate the role of endothelium in COVID-19 (Figure 1):

### (1) Recruitment of leukocytes, immune response and tissue injury.

Leukocytes play an important role in the pathogenesis of SARS-CoV-2. The importance of the leukocyte-EC axis is exemplified by the observations that patients with severe disease demonstrate a marked increase in blood neutrophils which is associated with lymphopenia, with both CD4<sup>+</sup> T cells and CD8<sup>+</sup> T cells being lower in severe compared to moderate cases<sup>33</sup>. Moreover, histological examination of a severe case, who died of SARS-CoV-2 demonstrated lung interstitial mononuclear inflammatory infiltrates, dominated by lymphocytes<sup>34</sup>. Through the systemic inflammatory response in COVID-19, referred to as cytokine storm, or cytokine release syndrome, the endothelium will be directly exposed to pro-inflammatory cytokines that initiate transcriptional programmes, that induce adhesion molecules and chemokines

driving leukocyte recruitment and inflammation<sup>35</sup>. This process can also lead to EC death that contributes to increased vascular permeability and end-organ damage. Through an amplification loop of the inflammatory response, the endothelium may constitute a significant source of pro-inflammatory cytokines e.g. interleukin (IL)-1 and IL-6, that characterize the cytokine storm in COVID-19<sup>36 37</sup>. Since the process of lymphocyte trafficking in chronic inflammatory and auto-immune diseases is a major therapeutic target, it is possible that anti-inflammatory therapies developed for these conditions could be repurposed to treat SARS-CoV-2 infection.

It is uncertain whether modulating EC function during the inflammatory response to SARS-CoV-2 would be beneficial, with the dilemma that anti-inflammatory agents might be counterproductive and compromise the immune response against the virus. However, in mouse models of pathogenic influenza and respiratory syncytial virus (RSV) infection, where there is evidence that the EC orchestrate a CD8<sup>+</sup> T cell-mediated cytokine storm, agonists of the S1P<sub>1</sub> receptor are effective in improving survival<sup>38-40</sup>. It is also noteworthy that adiponectin is protective in models of sepsis and ARDS through suppression of EC activation<sup>41</sup>. Such observations raise the question of whether the immune response itself may contribute to chronic inflammatory process in severe SARS-CoV-2 infection. The immunology of SARS-CoV-2 infection has recently been comprehensively reviewed by Vabret et al<sup>42</sup> who noted that in SARS-CoV-1 patients survival was associated with immunity skewed towards a CD8 cytotoxic response<sup>43</sup>. Indeed, there is evidence that expansion of virus specific CD4 T cells and a robust Th2 response (including increases in plasma IL-4, IL-5 and IL-10) are associated with death in SARS-CoV-1<sup>44</sup>. Whether this is the case in SARS-Cov-2 remains to be established. It does however appear that the adaptive immune response is an important arbiter of outcomes in SARS-CoV-2, as there is a reported association between plasma IgA titers and severity of disease<sup>45</sup>. It is possible that in SARS-CoV-2, antibody-dependent enhancement (ADE) of inflammation plays a role in EC activation and pathology. In this phenomenon non-neutralising antibodies facilitate Fc-mediated uptake of virus into macrophages, resulting in their activation and liberation of inflammatory cytokines, leading to the secondary recruitment of inflammatory leukocytes by local EC. The phenomenon of ADE is readily demonstrable in a number of experimental models of SARS infection but there is currently no evidence that ADE contributes to pathology in SARS-Cov-2<sup>42</sup>. Since the process of EC dependent leukocyte trafficking in chronic inflammatory and auto-immune diseases is a major therapeutic target, it is possible that anti-inflammatory therapies developed for these conditions could be repurposed to treat SARS-CoV-2 infection. Indeed, the recent success of low dose dexamethasone in the RECOVERY-trial, and its adoption as a front line therapy for SARS-CoV-2 in the UK, is an exemplar of the utility of such an approach (see below).

## (2) Endothelium and thrombosis

When dysfunctional, the thrombotic and coagulant properties of the endothelium change<sup>35</sup>. In particular, a decreased anti-aggregatory prostacyclin

production from EC and an increased pro-aggregatory thromboxane synthesis from activated platelets<sup>46</sup> may skew the homeostatic situation towards a pro-thrombotic and pro-inflammatory phenotype. Interestingly, under some inflammatory conditions EC express adhesion receptors such as von Willebrand Factor on their surface. Both in vitro and in vivo these conditions have been demonstrated to support the recruitment and activation of platelets to intact endothelial monolayers<sup>47</sup>. This in turn can lead to the platelet dependent secondary recruitment of circulating leukocytes, either by leukocyte interactions with platelets adherent to EC<sup>48</sup>, by the recruitment of circulating heterotypic aggregates of platelets and leukocytes<sup>49</sup>, or by the transfer of platelet borne receptors such as GPIb to the leukocyte membrane by platelet derived microvesicles<sup>50</sup>. It is reasonable to assume that such tri-cellular aggregates (EC-platelet-leukocyte) on the walls of smaller vessel would be sufficient to cause loss of microvascular perfusion in the lungs and other organs. Indeed, compromise of myocardial perfusion could play a role in the elevated levels of troponin evident in many SARS-CoV-2 patients. Intravascular thrombosis and coagulation in addition may further damage the endothelium and contribute to endothelial inflammation and dysfunction<sup>35</sup>. In some patients with COVID-19 severe micro-vascular endothelial injury directly mediated by activation of the alternative and lectin complement pathways has been demonstrated and associated with a pro-coagulant state<sup>51</sup>. It remains however to be established if anti-coagulation or platelet inhibition in COVID-19 improves endothelial function, and if dampened endothelial inflammation would attenuate the pro-coagulant state in COVID-19.

### (3) ACE2 expression and function in endothelium.

ACE2 is intimately linked to cardiovascular physiology as part of the renin-angiotensin-aldosterone system (RAAS), which controls blood pressure by altering vascular tone and function. The related molecule ACE converts angiotensin (Ang) I to Ang II, which promotes vasoconstriction, hypertension and vascular inflammation. Because of these properties, anti-hypertensive drugs have been developed to reduce the production (ACE inhibitors, ACE-i) or downstream effects (Ang II receptor blockers, ARBs) of Ang II. The effects of ACE are opposed by ACE2, which converts Ang II into Ang 1-7 molecules thereby promoting vasodilatation and reducing hypertension<sup>52</sup>. Since ACE2 is expressed in cells of the cardiovascular system<sup>53</sup>, there has been considerable interest in the hypothesis that this class of anti-hypertensive drugs may increase the risk of SARS-CoV-2 infection by increasing the expression of ACE2 in vascular cells<sup>54</sup>. However, population-based studies revealed that ACE inhibitors and ARBs do not enhance the risk of COVID-19 or disease severity<sup>55 56</sup> and the European Society of Cardiology and other learned societies recommend that patients should continue with their usual anti-hypertensive medications during the pandemic ([https://www.escardio.org/Councils/Council-on-Hypertension-\(CHT\)/News/position-statement-of-the-esc-council-on-hypertension-on-ace-inhibitors-and-ang](https://www.escardio.org/Councils/Council-on-Hypertension-(CHT)/News/position-statement-of-the-esc-council-on-hypertension-on-ace-inhibitors-and-ang)).

The spike (S) protein of the coronaviruses mediates viral entry into target cells. Entry depends on physical interactions of the surface unit, S1, of the S protein to a host cell receptor, which facilitates viral attachment to the surface of the target cells. SARS-CoV-2 engages ACE2 as the primary receptor<sup>57, 58 59</sup> and

hence it is plausible that COVID-19 may cause reduced bioavailability of ACE2 due to endosomal/lysosomal processing<sup>52</sup>. Treatment of COVID-19 patients with recombinant ACE2 is currently under clinical trial because this may act as a decoy receptor<sup>60</sup>, hence limiting viral entry. It is notable recombinant ACE2 treatment may have additional beneficial effects, by increasing its bioavailability at the endothelial surface. The S2 domain of the S protein facilitates membrane fusion, which requires conformational flexibility, achieved by proteolytic cleavages. The cleavage event employs the cellular transmembrane protease, serine 2 (TMPRSS2), and has profound implications on virulence. A TMPRSS2 inhibitor, approved for clinical use, was shown to block entry and might constitute a treatment option<sup>57</sup>. It has been reported that TMPRSS2 expression is below detection in microvascular endothelial cells<sup>61</sup> and only upregulated in endothelial cells actively undergoing angiogenic or tubulogenic responses. Further studies are needed to better understand the physiological expression and function of TMPRSS2 in adult endothelial cells. However, the expression of TMPRSS2 alone may not be predictive of its function since serine proteases, such as TMPRSS2, are regulated by nitrosylation<sup>62</sup>. Therefore it is plausible that eNOS activity and subsequent production of NO may influence viral infection of EC by altering TMPRSS2 activity. Taken together the S protein mediates entry by connecting the virus to the plasma membrane, and by catalysing subsequent virus-cell membrane fusions.

#### (4) SARS-CoV-2-induced endothelitis.

In addition to the respiratory tract, SARS-CoV-2 viral load is detected in kidneys, liver, heart, and brain<sup>63</sup>, which are all highly vascularized tissues. Indeed, Monteil et al provided an early indication of SARS-CoV-2 tropism for vascularized tissues by demonstrating that this virus can infect human blood vessel and kidney organoids via ACE2<sup>59</sup>. It was suggested by Varga et al by electron microscopy and histology that SARS-CoV-2 can be detected in EC of the kidney (glomerular capillaries), small bowel, lung and myocardium<sup>21</sup>, but these data were recently disputed<sup>64</sup>. Ackerman et al<sup>22</sup> have shown abnormalities within the pulmonary microvasculature with congestion and micro-thrombi similar to Menter et al<sup>23</sup>, and by electron microscopy there is endothelial injury and congestion with cell fragments and degenerate organelles in the lumen. Scanning electron microscopy of corrosion casts shows microvasculature of larger diameter with an irregular surface, which may be due to endothelial injury and/or platelet aggregates/fibrin. Intriguingly there was also evidence of intussusceptive angiogenesis and the authors speculate that this feature may distinguish the pulmonary pathobiology of COVID-19 from other viral infections. SARS-CoV-2 has also been detected in skin EC<sup>24</sup> and circulating endothelial cells are elevated in patients admitted to hospital with COVID-19<sup>65</sup>.

Taken together, these studies point to endothelial SARS-CoV-2-infection as a possible direct trigger of endothelial adverse effects in COVID-19. Indeed, SARS-CoV-2 infected endothelium has been associated with EC apoptosis, suggesting a possible mechanism through which the endothelium may become dysfunctional in COVID-19<sup>21</sup>. The recent reports of a Kawasaki Disease-like syndrome associated with COVID-19 infection in children highlights the

importance of the virus on the vasculature<sup>66 67</sup>. Kawasaki disease is a systemic vasculitis most commonly seen in children which particularly targets the myocardium and coronary arteries. Although the etiology of Kawasaki disease is unknown, infectious agents including RNA viruses have been previously postulated as the cause and the first link to a coronavirus infection was published in 2005<sup>68</sup>. The current outbreaks following infection by SARS-CoV-2 are the subject of intense investigation. A recently reported group of 58 hospitalized children were diagnosed with pediatric inflammatory multisystem syndrome temporally associated with severe acute respiratory syndrome coronavirus 2 (PIMS-TS). Of note, important differences were noted when compared with Kawasaki disease<sup>69</sup>. Moreover, anecdotal cases with evidence of medium and large vessel vasculitis suggest similar complications may be seen in some adults with COVID-19.

#### (5) Pericytes.

Pericytes are multifunctional mural cells of the microvasculature and are essential for the maintenance of the endothelium integrity<sup>70</sup>. Studies suggest that they are involved in COVID-19-related vasculopathy. Recent single-cell or single-nucleus RNA sequencing analyses have shown that ACE2 is highly expressed in pericytes of various organs, such as the heart, both in mouse and man<sup>71 72</sup>. In alveolar capillary of SARS-CoV-2 infected lung, pericytes are markedly decreased, likely through apoptosis<sup>73</sup>. In a genetically modified mouse model with pericyte deficiency (*pdgf-b<sup>ret/ret</sup>* mouse<sup>74</sup>) induced by deletion of the PDGF-B retention motif, loss of pericytes induced thrombogenic reactions in ECs<sup>72</sup>. Therefore, pericytes by acting as direct target for SARS-CoV-2 infection, could play a crucial role in microvascular dysfunction and coagulopathy. It is suggested that a permeable endothelial barrier as observed in hypertension, diabetes and obesity, which are co-morbidity factors in severe cases of COVID-19, allows the virus to reach the pericytes<sup>72</sup>.

#### **Therapeutic targets.**

It remains to be determined whether the endothelial dysfunction and injury seen in COVID-19 predominantly reflects direct infection of EC by SARS-CoV-2 or indirect bystander injury by factors including cytokines, leukocytes, neutrophil nets and complement activation<sup>75</sup>. Notwithstanding, an important consideration is the effect of current cardiovascular drugs in this setting (Figure 2). On the one hand, they may offer enhanced endothelial protection, while on the other they may increase endothelial susceptibility. To date there are only retrospective data to rely upon and prospective clinical studies in COVID-19 with clearly defined cardiovascular endpoints are required. Drugs including HMG-CoA reductase inhibitors (statins), alpha and beta adrenergic blockers and renin–angiotensin–aldosterone system antagonists are widely prescribed in those with diabetes mellitus, hypertension and coronary artery disease, groups known to be at the highest risk from COVID-19. Understanding how these drugs influence outcomes is therefore essential.

In pre-clinical studies, statins increase expression of ACE2. However, these in disease models this may reflect a return to normal levels. Moreover, enhanced



expression of ACE2 has cardiovascular benefit<sup>76</sup>. Previously observed beneficial effects in influenza and actions of statins including their ability to reduce CD147 expression, optimise lipid raft function, regulate autophagy, minimise endothelial activation, down-regulate pro-thrombotic pathways and enhance anti-thrombotic effects, alongside immunomodulatory actions, suggest that statins may exert important endothelial protective effects both against and during SARS-CoV-2 infection<sup>77-79</sup>. COVID-19 clinical trials incorporating statins (NCT04333407; NCT04348695; NCT04380402) have commenced and the results are awaited with interest.

Reported beneficial effects of beta-adrenergic blockers in ARDS and respiratory failure, alongside a potential ability to reduce viral entry by down-regulating ACE2 has resulted in these drugs being suggested as an adjunct therapy for COVID-19, even in those without a primary indication<sup>80</sup>. Retrospective studies underway will help inform this hypothesis. An intriguing large, retrospective analysis of alpha-1 adrenergic receptor antagonists in patients suffering from ARDS or pneumonia has revealed that those incidentally prescribed alpha-blockers were less likely to require ventilation or to die following onset of ventilator support. In contrast, beta-adrenergic blockers had no effect. The clinical study was inspired by pre-clinical data, which demonstrated that alpha-blockade can prevent the ARDS-associated cytokine storm and death in mice by interfering with a catecholamine loop<sup>81</sup>. These findings suggest that alpha-1 adrenergic receptor antagonists merit retrospective and prospective analysis in COVID-19. The identification of the SARS-CoV-2-associated cytokine storm as a potential therapeutic target is also supported by the early data from the dexamethasone arm of the COVID-19 RECOVERY trial<sup>82</sup>.

The ability of SARS-CoV-2 to utilise ACE2 as a co-receptor for cellular entry has led to significant interest in the impact of angiotensin-converting enzyme inhibitors ACE-i and ARBs. Similarly to statins, these commonly used drugs enhance ACE2<sup>76</sup>. Although clinical data are sparse, ACEi and ARBs are reported to improve outcomes in ARDS<sup>83</sup>. There has also been speculation that a dysfunctional RAAS is important for COVID-19 disease pathogenesis<sup>84</sup>. Initial concern regarding potential susceptibility to SARS-CoV-2 infection and/or exacerbation of its effects in those taking ACEi or ARBs have been allayed by retrospective clinical studies<sup>85-87</sup>. Current advice is to continue taking these drugs prescribed for hypertension, cardiac failure and chronic renal disease, and for physicians to prescribe them for new clinical indications as normal<sup>84</sup>. Prospective studies and clinical trials are now urgently needed. Two clinical trials will study the impact of losartan on COVID-19 (NCT04311177 and NCT04312009). Interest will also focus upon Ang 1-7 peptides, ACE2 itself<sup>76, 84</sup> and monoclonal antibodies that prevent SARS-CoV-2 binding ACE2<sup>88</sup>.

### **Position statements**

Further research is urgently needed to combat the COVID-19 pandemic and we emphasize that the role of vascular endothelium requires close scrutiny. There are today several outstanding questions that need to be addressed to

elucidate more precisely the role of EC in COVID-19 and to investigate potential routes to clinical translation.

1. Endothelial biomarkers and tests of function (e.g. flow-mediated dilation, arterial stiffness) should be monitored in studies of COVID-19 outcome and treatment effects. High-quality data collection is needed, with follow-up studies amongst the survivors of acute infection since there are little or no data available on EC function testing in COVID-19. Indeed, collaborative networks have already been established to analyse RNA biomarkers<sup>89</sup> and arterial stiffness (CARTESIAN STUDY) to assess vascular consequences of COVID-19. This may help to enable stratification of COVID-19 patients with highest pro-thrombotic and cardiovascular risk and allow tailored treatments.

2. The significance of SARS-CoV-2-mediated endocytosis and downregulation of ACE2 on cardiovascular health is uncertain, but data from ongoing clinical trials to test recombinant ACE2 may be instrumental in addressing this question.

3. The principle effects of SARS-CoV-2 on endothelial function should be determined, including studies of EC activation, leukocyte recruitment, platelet activation, turnover, signalling etc. EC from both micro and macrovasculature should be investigated. Aging is an important determinant of COVID-19 outcome thus the influence of cellular senescence, oxidative stress and other features of aging on SARS-CoV-2 infection of endothelium should also be assessed. The influence of gender on endothelial responses to SARS-CoV-2 and how this relates to the susceptibility and outcome of COVID-19 patients should be investigated.

4. The effects of common cardiovascular drugs such as statins and beta blockers on endothelial responses to SARS-CoV-19 should be explored, including their influence on ACE2 expression and viral infectivity.

5. The long-term cardiovascular effects following recovery from COVID-19 must be determined during planned patient follow up so that appropriate preventive measures can be taken in time if needed. Measuring endothelial function in addition to myocardial injury and respiratory function markers in convalescent patients may represent a possible means for early detection of vascular sequelae post COVID-19.

## References

1. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, Si HR, Zhu Y, Li B, Huang CL, Chen HD, Chen J, Luo Y, Guo H, Jiang RD, Liu MQ, Chen Y, Shen XR, Wang X, Zheng XS, Zhao K, Chen QJ, Deng F, Liu LL, Yan B, Zhan FX, Wang YY, Xiao GF, Shi ZL. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 2020;**579**:270-273.
2. Guo T, Fan Y, Chen M, Wu X, Zhang L, He T, Wang H, Wan J, Wang X, Lu Z. Cardiovascular Implications of Fatal Outcomes of Patients With Coronavirus Disease 2019 (COVID-19). *JAMA cardiology* 2020.
3. Driggin E, Madhavan MV, Bikdeli B, Chuich T, Laracy J, Bondi-Zoccai G, Brown TS, Nigoghossian C, Zidar DA, Haythe J, Brodie D, Beckman JA, Kirtane AJ, Stone GW, Krumholz HM, Parikh SA. Cardiovascular Considerations for Patients, Health Care Workers, and Health Systems During the Coronavirus Disease 2019 (COVID-19) Pandemic. *Journal of the American College of Cardiology* 2020.
4. Stefanini GG, Montorfano M, Trabattoni D, Andreini D, Ferrante G, Ancona M, Metra M, Curello S, Maffeo D, Pero G, Cacucci M, Assanelli E, Bellini B, Russo F, Ielasi A, Tespili M, Danzi GB, Vandoni P, Bollati M, Barbieri L, Oreglia J, Lettieri C, Cremonesi A, Carugo S, Reimers B, Condorelli G, Chieffo A. ST-Elevation Myocardial Infarction in Patients with COVID-19: Clinical and Angiographic Outcomes. *Circulation* 2020.
5. Meyer P, Degrauwe S, Van Delden C, Ghadri JR, Templin C. Typical takotsubo syndrome triggered by SARS-CoV-2 infection. *Eur Heart J* 2020;**41**:1860.
6. Tersalvi G, Vicenzi M, Calabretta D, Biasco L, Pedrazzini G, Winterton D. Elevated Troponin in Patients With Coronavirus Disease 2019: Possible Mechanisms. *Journal of cardiac failure* 2020.
7. Gao L, Jiang D, Wen XS, Cheng XC, Sun M, He B, You LN, Lei P, Tan XW, Qin S, Cai GQ, Zhang DY. Prognostic value of NT-proBNP in patients with severe COVID-19. *Respiratory research* 2020;**21**:83.
8. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, Wang B, Xiang H, Cheng Z, Xiong Y, Zhao Y, Li Y, Wang X, Peng Z. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *Jama* 2020.
9. Chen L, Li X, Chen M, Feng Y, Xiong C. The ACE2 expression in human heart indicates new potential mechanism of heart injury among patients infected with SARS-CoV-2. *Cardiovascular Research* 2020;**116**:1097-1100.
10. [The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China]. *Zhonghua liu xing bing xue za zhi = Zhonghua liuxingbingxue zazhi* 2020;**41**:145-151.
11. Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, Barnaby DP, Becker LB, Chelico JD, Cohen SL, Cookingham J, Coppa K, Diefenbach MA, Dominello AJ, Duer-Hefele J, Falzon L, Gitlin J, Hajizadeh N, Harvin TG, Hirschwerk DA, Kim EJ, Kozel ZM, Marrast LM, Mogavero JN, Osorio GA, Qiu M, Zanos TP. Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *Jama* 2020.
12. Bikdeli B, Madhavan MV, Jimenez D, Chuich T, Dreyfus I, Driggin E, Nigoghossian C, Ageno W, Madjid M, Guo Y, Tang LV, Hu Y, Giri J,

- Cushman M, Quere I, Dimakakos EP, Gibson CM, Lippi G, Favaloro EJ, Fareed J, Caprini JA, Tafur AJ, Burton JR, Francese DP, Wang EY, Falanga A, McLintock C, Hunt BJ, Spyropoulos AC, Barnes GD, Eikelboom JW, Weinberg I, Schulman S, Carrier M, Piazza G, Beckman JA, Steg PG, Stone GW, Rosenkranz S, Goldhaber SZ, Parikh SA, Monreal M, Krumholz HM, Konstantinides SV, Weitz JI, Lip GYH. COVID-19 and Thrombotic or Thromboembolic Disease: Implications for Prevention, Antithrombotic Therapy, and Follow-up. *J Am Coll Cardiol* 2020.
13. Gando S, Fujishima S, Saitoh D, Shiraishi A, Yamakawa K, Kushimoto S, Ogura H, Abe T, Mayumi T, Sasaki J, Kotani J, Takeyama N, Tsuruta R, Takuma K, Yamashita N, Shiraishi SI, Ikeda H, Shiino Y, Tarui T, Nakada TA, Hifumi T, Otomo Y, Okamoto K, Sakamoto Y, Hagiwara A, Masuno T, Ueyama M, Fujimi S, Umemura Y. The significance of disseminated intravascular coagulation on multiple organ dysfunction during the early stage of acute respiratory distress syndrome. *Thrombosis research* 2020;**191**:15-21.
  14. Tang N, Li D, Wang X, Sun Z. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *Journal of thrombosis and haemostasis : JTH* 2020;**18**:844-847.
  15. Danzi GB, Loffi M, Galeazzi G, Gherbesi E. Acute pulmonary embolism and COVID-19 pneumonia: a random association? *European heart journal* 2020.
  16. Llitjos JF, Leclerc M, Chochois C, Monsallier JM, Ramakers M, Auvray M, Merouani K. High incidence of venous thromboembolic events in anticoagulated severe COVID-19 patients. *J Thromb Haemost* 2020.
  17. Mao L, Jin H, Wang M, Hu Y, Chen S, He Q, Chang J, Hong C, Zhou Y, Wang D, Miao X, Li Y, Hu B. Neurologic Manifestations of Hospitalized Patients With Coronavirus Disease 2019 in Wuhan, China. *JAMA neurology* 2020.
  18. Beyrouti R, Adams ME, Benjamin L, Cohen H, Farmer SF, Goh YY, Humphries F, Jager HR, Losseff NA, Perry RJ, Shah S, Simister RJ, Turner D, Chandratheva A, Werring DJ. Characteristics of ischaemic stroke associated with COVID-19. *Journal of neurology, neurosurgery, and psychiatry* 2020.
  19. Guzik TJ, Mohiddin SA, Dimarco A, Patel V, Savvatis K, Marelli-Berg FM, Madhur MS, Tomaszewski M, Maffia P, D'Acquisto F, Nicklin SA, Marian AJ, Nosalski R, Murray EC, Guzik B, Berry C, Touyz RM, Kreutz R, Wang DW, Bhella D, Sglicco O, Crea F, Thomson EC, McInnes IB. COVID-19 and the cardiovascular system: implications for risk assessment, diagnosis, and treatment options. *Cardiovascular research* 2020.
  20. Klok FA, Kruip M, van der Meer NJM, Arbous MS, Gommers D, Kant KM, Kaptein FHJ, van Paassen J, Stals MAM, Huisman MV, Endeman H. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thrombosis research* 2020.
  21. Varga Z, Flammer AJ, Steiger P, Haberecker M, Andermatt R, Zinkernagel AS, Mehra MR, Schuepbach RA, Ruschitzka F, Moch H. Endothelial cell infection and endotheliitis in COVID-19. *Lancet (London, England)* 2020;**395**:1417-1418.
  22. Ackermann M, Verleden SE, Kuehnel M, Haverich A, Welte T, Laenger F, Vanstapel A, Werlein C, Stark H, Tzankov A, Li WW, Li VW, Mentzer SJ,

- Jonigk D. Pulmonary Vascular Endothelialitis, Thrombosis, and Angiogenesis in Covid-19. *The New England journal of medicine* 2020.
23. Menter T, Haslbauer JD, Nienhold R, Savic S, Hopfer H, Deigendesch N, Frank S, Turek D, Willi N, Pargger H, Bassetti S, Leuppi JD, Cathomas G, Tolnay M, Mertz KD, Tzankov A. Post-mortem examination of COVID19 patients reveals diffuse alveolar damage with severe capillary congestion and variegated findings of lungs and other organs suggesting vascular dysfunction. *Histopathology* 2020.
  24. Colmenero I, Santonja C, Alonso-Riaño M, Noguera-Morel L, Hernández-Martín A, Andina D, Wiesner T, Rodríguez-Peralto JL, Requena L, Torrelo A. SARS-CoV-2 endothelial infection causes COVID-19 chilblains: histopathological, immunohistochemical and ultrastructural study of 7 paediatric cases. *The British journal of dermatology* 2020.
  25. Alexander Y, Osto E, Schmidt-Trucksass A, Shechter M, Trifunovic D, Duncker DJ, Aboyans V, Back M, Badimon L, Cosentino F, De Carlo M, Dorobantu M, Harrison DG, Guzik TJ, Hofer I, Morris PD, Norata GD, Suades R, Taddei S, Vilahur G, Waltenberger J, Weber C, Wilkinson F, Bochaton-Piallat ML, Evans PC. ENDOTHELIAL FUNCTION IN CARDIOVASCULAR PRECISION MEDICINE : A POSITION PAPER ON BEHALF OF THE EUROPEAN SOCIETY OF CARDIOLOGY. *Cardiovasc Res* 2020.
  26. Souilhol C, Harmsen MC, Evans PC, Krenning G. Endothelial-mesenchymal transition in atherosclerosis. *Cardiovasc Res* 2018;**114**:565-577.
  27. Souilhol C, Serbanovic-Canic J, Fragiadaki M, Chico TJ, Ridger V, Roddie H, Evans PC. Endothelial responses to shear stress in atherosclerosis: a novel role for developmental genes. *Nature reviews Cardiology* 2019.
  28. Mundi S, Massaro M, Scoditti E, Carluccio MA, van Hinsbergh VWM, Iruela-Arispe ML, De Caterina R. Endothelial permeability, LDL deposition, and cardiovascular risk factors-a review. *Cardiovasc Res* 2018;**114**:35-52.
  29. Li X, Sun X, Carmeliet P. Hallmarks of Endothelial Cell Metabolism in Health and Disease. *Cell Metab* 2019;**30**:414-433.
  30. Celermajer DS, Sorensen KE, Gooch VM, Spiegelhalter DJ, Miller OI, Sullivan ID, Lloyd JK, Deanfield JE. Non-invasive detection of endothelial dysfunction in children and adults at risk of atherosclerosis. *Lancet* 1992;**340**:1111-1115.
  31. Kitta Y, Obata JE, Nakamura T, Hirano M, Kodama Y, Fujioka D, Saito Y, Kawabata K, Sano K, Kobayashi T, Yano T, Nakamura K, Kugiyama K. Persistent impairment of endothelial vasomotor function has a negative impact on outcome in patients with coronary artery disease. *J Am Coll Cardiol* 2009;**53**:323-330.
  32. Wang M, Hao H, Leeper NJ, Zhu L. Thrombotic Regulation From the Endothelial Cell Perspectives. *Arteriosclerosis, thrombosis, and vascular biology* 2018;**38**:e90-e95.
  33. Zhang B, Zhou X, Zhu C, Feng F, Qiu Y, Feng J, Jia Q, Song Q, Zhu B, Wang J. Immune phenotyping based on neutrophil-to-lymphocyte ratio and IgG predicts disease severity and outcome for patients with COVID-19. *medRxiv* 2020:2020.2003.2012.20035048.
  34. Xu Z, Shi L, Wang Y, Zhang J, Huang L, Zhang C, Liu S, Zhao P, Liu H, Zhu L, Tai Y, Bai C, Gao T, Song J, Xia P, Dong J, Zhao J, Wang FS.

- Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *The Lancet Respiratory medicine* 2020;**8**:420-422.
35. Chousterman BG, Swirski FK, Weber GF. Cytokine storm and sepsis disease pathogenesis. *Seminars in immunopathology* 2017;**39**:517-528.
  36. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z, Yu T, Xia J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo L, Xie J, Wang G, Jiang R, Gao Z, Jin Q, Wang J, Cao B. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;**395**:497-506.
  37. Chen X, Zhao B, Qu Y, Chen Y, Xiong J, Feng Y, Men D, Huang Q, Liu Y, Yang B, Ding J, Li F. Detectable serum SARS-CoV-2 viral load (RNAemia) is closely correlated with drastically elevated interleukin 6 (IL-6) level in critically ill COVID-19 patients. *Clin Infect Dis* 2020.
  38. Teijaro JR, Walsh KB, Cahalan S, Fremgen DM, Roberts E, Scott F, Martinborough E, Peach R, Oldstone MBA, Rosen H. Endothelial Cells Are Central Orchestrators of Cytokine Amplification during Influenza Virus Infection. *Cell* 2011;**146**:980-991.
  39. Walsh KB, Teijaro JR, Brock LG, Fremgen DM, Collins PL, Rosen H, Oldstone MBA. Animal Model of Respiratory Syncytial Virus: CD8(+) T Cells Cause a Cytokine Storm That Is Chemically Tractable by Sphingosine-1-Phosphate 1 Receptor Agonist Therapy. *Journal of Virology* 2014;**88**:6281-6293.
  40. Walsh KB, Teijaro JR, Wilker PR, Jatzek A, Fremgen DM, Das SC, Watanabe T, Hatta M, Shinya K, Suresh M, Kawaoka Y, Rosen H, Oldstone MBA. Suppression of cytokine storm with a sphingosine analog provides protection against pathogenic influenza virus. *Proceedings of the National Academy of Sciences of the United States of America* 2011;**108**:12018-12023.
  41. Teoh H, Quan A, Bang KWA, Wang GL, Lovren F, Vu V, Haitsma JJ, Szmítko PE, Al-Omran M, Wang CH, Gupta M, Peterson MD, Zhang HB, Chan L, Freedman J, Sweeney G, Verma S. Adiponectin deficiency promotes endothelial activation and profoundly exacerbates sepsis-related mortality. *American Journal of Physiology-Endocrinology and Metabolism* 2008;**295**:E658-E664.
  42. Vabret N, Samstein R, Fernandez N, Merad M, Sinai Immunology Review P. Advancing scientific knowledge in times of pandemics. *Nature Reviews Immunology* 2020;**20**:338-338.
  43. Ng OW, Chia A, Tan AT, Jadi RS, Leong HN, Bertoletti A, Tan YJ. Memory T cell responses targeting the SARS coronavirus persist up to 11 years post-infection. *Vaccine* 2016;**34**:2008-2014.
  44. Li CKF, Wu H, Yan HP, Ma SW, Wang LL, Zhang MX, Tang XP, Temperton NJ, Weiss RA, Brenchley JM, Douek DC, Mongkolsapaya J, Tran BH, Lin CLS, Screaton GR, Hou JL, McMichael AJ, Xu XN. T cell responses to whole SARS coronavirus in humans. *Journal of Immunology* 2008;**181**:5490-5500.
  45. Ma H, Zeng W, He H, Zhao D, Yang Y, Jiang D, Zhou P, Qi Y, He W, Zhao C, Yi R, Wang X, Wang B, Xu Y, Yang Y, Kombe Kombe AJ, Ding C, Xie J, Gao Y, Cheng L, Li Y, Ma X, Jin T. COVID-19 diagnosis and study of serum SARS-CoV-2 specific IgA, IgM and IgG by chemiluminescence immunoanalysis. *medRxiv* 2020:2020.2004.2017.20064907.

46. Petri MH, Tellier C, Michiels C, Ellertsen I, Dogne JM, Bäck M. Effects of the dual TP receptor antagonist and thromboxane synthase inhibitor EV-077 on human endothelial and vascular smooth muscle cells. *Biochem Biophys Res Commun* 2013;**441**:393-398.
47. Tull SP, Anderson SI, Hughan SC, Watson SP, Nash GB, Rainger GE. Cellular pathology of atherosclerosis: smooth muscle cells promote adhesion of platelets to cocultured endothelial cells. *Circ Res* 2006;**98**:98-104.
48. Kuckleburg CJ, Yates CM, Kalia N, Zhao Y, Nash GB, Watson SP, Rainger GE. Endothelial cell-borne platelet bridges selectively recruit monocytes in human and mouse models of vascular inflammation. *Cardiovascular Research* 2011;**91**:134-141.
49. Alharbi A, Thompson JP, Brindle NP, Stover CM. Ex vivo modelling of the formation of inflammatory platelet-leucocyte aggregates and their adhesion on endothelial cells, an early event in sepsis. *Clinical and Experimental Medicine* 2019;**19**:321-337.
50. Chimen M, Evryviadou A, Box CL, Harrison MJ, Hazeldine J, Dib LH, Kuravi SJ, Payne H, Price JMJ, Kavanagh D, Iqbal AJ, Lax S, Kalia N, Brill A, Thomas SG, Belli A, Crombie N, Adams RA, Evans SA, Deckmyn H, Lord JM, Harrison P, Watson SP, Nash GB, Rainger GE. Appropriation of GPIb alpha from platelet-derived extracellular vesicles supports monocyte recruitment in systemic inflammation. *Haematologica* 2020;**105**:1248-1261.
51. Magro C, Mulvey JJ, Berlin D, Nuovo G, Salvatore S, Harp J, Baxter-Stoltzfus A, Laurence J. Complement associated microvascular injury and thrombosis in the pathogenesis of severe COVID-19 infection: a report of five cases. *Translational research : the journal of laboratory and clinical medicine* 2020.
52. Gheblawi M, Wang K, Viveiros A, Nguyen Q, Zhong JC, Turner AJ, Raizada MK, Grant MB, Oudit GY. Angiotensin-Converting Enzyme 2: SARS-CoV-2 Receptor and Regulator of the Renin-Angiotensin System: Celebrating the 20th Anniversary of the Discovery of ACE2. *Circ Res* 2020;**126**:1456-1474.
53. Nicin L, Abplanalp WT, Mellentin H, Kattih B, Tombor L, John D, Schmitto JD, Heineke J, Emrich F, Arsalan M, Holubec T, Walther T, Zeiher AM, Dimmeler S. Cell type-specific expression of the putative SARS-CoV-2 receptor ACE2 in human hearts. *European heart journal* 2020.
54. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *The Lancet Respiratory medicine* 2020;**8**:e21.
55. Mehra MR, Desai SS, Kuy S, Henry TD, Patel AN. Cardiovascular Disease, Drug Therapy, and Mortality in Covid-19. *The New England journal of medicine* 2020.
56. Mancia G, Rea F, Ludergnani M, Apolone G, Corrao G. Renin-Angiotensin-Aldosterone System Blockers and the Risk of Covid-19. *The New England journal of medicine* 2020.
57. Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S, Schiergens TS, Herrler G, Wu NH, Nitsche A, Muller MA, Drosten C, Pohlmann S. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* 2020;**181**:271-280.e278.
58. Li W, Moore MJ, Vasilieva N, Sui J, Wong SK, Berne MA, Somasundaran M, Sullivan JL, Luzuriaga K, Greenough TC, Choe H, Farzan M.


- Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. *Nature* 2003;**426**:450-454.
59. Monteil V, Kwon H, Prado P, Hagelkruys A, Wimmer RA, Stahl M, Leopoldi A, Garreta E, Hurtado Del Pozo C, Prosper F, Romero JP, Wirnsberger G, Zhang H, Slutsky AS, Conder R, Montserrat N, Mirazimi A, Penninger JM. Inhibition of SARS-CoV-2 Infections in Engineered Human Tissues Using Clinical-Grade Soluble Human ACE2. *Cell* 2020.
  60. Zhang H, Penninger JM, Li Y, Zhong N, Slutsky AS. Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. *Intensive care medicine* 2020;**46**:586-590.
  61. Aimes RT, Zijlstra A, Hooper JD, Ogbourne SM, Sit ML, Fuchs S, Gotley DC, Quigley JP, Antalis TM. Endothelial cell serine proteases expressed during vascular morphogenesis and angiogenesis. *Thrombosis and haemostasis* 2003;**89**:561-572.
  62. Stamler JS, Lamas S, Fang FC. Nitrosylation. the prototypic redox-based signaling mechanism. *Cell* 2001;**106**:675-683.
  63. Puelles VG, Lutgehetmann M, Lindenmeyer MT, Sperhake JP, Wong MN, Allweiss L, Chilla S, Heinemann A, Wanner N, Liu S, Braun F, Lu S, Pfefferle S, Schroder AS, Edler C, Gross O, Glatzel M, Wichmann D, Wiewe T, Kluge S, Puschel K, Aepfelbacher M, Huber TB. Multiorgan and Renal Tropism of SARS-CoV-2. *The New England journal of medicine* 2020.
  64. Goldsmith CS, Miller SE, Martines RB, Bullock HA, Zaki SR. Electron microscopy of SARS-CoV-2: a challenging task. *Lancet (London, England)* 2020;**395**:e99.
  65. Khider L, Gendron N, Goudot G, Chocron R, Hauw-Berlemont C, Cheng C, Rivet N, Pere H, Roffe A, Clerc S, Lebeaux D, Debuc B, Veyer D, Rance B, Gaussem P, Bertil S, Badoual C, Juvin P, Planquette B, Messas E, Sanchez O, Hulot JS, Diehl JL, Mirault T, Smadja DM. Curative anticoagulation prevents endothelial lesion in COVID-19 patients. *Journal of thrombosis and haemostasis : JTH* 2020.
  66. Verdoni L, Mazza A, Gervasoni A, Martelli L, Ruggeri M, Ciuffreda M, Bonanomi E, D'Antiga L. An outbreak of severe Kawasaki-like disease at the Italian epicentre of the SARS-CoV-2 epidemic: an observational cohort study. *Lancet* 2020.
  67. Riphagen S, Gomez X, Gonzalez-Martinez C, Wilkinson N, Theocharis P. Hyperinflammatory shock in children during COVID-19 pandemic. *Lancet* 2020.
  68. Esper F, Shapiro ED, Weibel C, Ferguson D, Landry ML, Kahn JS. Association between a novel human coronavirus and Kawasaki disease. *The Journal of infectious diseases* 2005;**191**:499-502.
  69. Whittaker E, Bamford A, Kenny J, Kafrou M, Jones CE, Shah P, Ramnarayan P, Fraise A, Miller O, Davies P, Kucera F, Brierley J, McDougall M, Carter M, Tremoulet A, Shimizu C, Herberg J, Burns JC, Lyall H, Levin M, Group fIP-TS, EUCLIDS, Consortia P. Clinical Characteristics of 58 Children With a Pediatric Inflammatory Multisystem Syndrome Temporally Associated With SARS-CoV-2. *JAMA* 2020.
  70. Attwell D, Mishra A, Hall CN, O'Farrell FM, Dalkara T. What is a pericyte? *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism* 2016;**36**:451-455.



71. Chen L, Li X, Chen M, Feng Y, Xiong C. The ACE2 expression in human heart indicates new potential mechanism of heart injury among patients infected with SARS-CoV-2. *Cardiovasc Res* 2020;**116**:1097-1100.
72. He L, Mäe MA, Sun Y, Muhl L, Nahar K, Liébanas EV, Fagerlund MJ, Oldner A, Liu J, Genové G, Pietilä R, Zhang L, Xie Y, Leptidis S, Mocci G, Stritt S, Osman A, Anisimov A, Hemanthakumar KA, Räsänen M, Björkegren J, Vanlandewijck M, Blomgren K, Hansson E, Mäkinen T, Peng X-R, Arnold TD, Alitalo K, Eriksson LI, Lendahl U, Betsholtz C. Pericyte-specific vascular expression of SARS-CoV-2 receptor ACE2 – implications for microvascular inflammation and hypercoagulopathy in COVID-19 patients. *bioRxiv* 2020:2020.2005.2011.088500.
73. Cardot-Leccia N, Hubiche T, Dellamonica J, Burel-Vandenbos F, Passeron T. Pericyte alteration sheds light on micro-vasculopathy in COVID-19 infection. *Intensive care medicine* 2020:1-2.
74. Lindblom P, Gerhardt H, Liebner S, Abramsson A, Enge M, Hellstrom M, Backstrom G, Fredriksson S, Landegren U, Nystrom HC, Bergstrom G, Dejana E, Ostman A, Lindahl P, Betsholtz C. Endothelial PDGF-B retention is required for proper investment of pericytes in the microvessel wall. *Genes Dev* 2003;**17**:1835-1840.
75. Gustafson D, Raju S, Wu R, Ching C, Veitch S, Rathnakumar K, Boudreau E, Howe KL, Fish JE. Overcoming Barriers: The Endothelium As a Linchpin of Coronavirus Disease 2019 Pathogenesis? *Arteriosclerosis, thrombosis, and vascular biology* 2020:Atvbaha120314558.
76. South AM, Diz DI, Chappell MC. COVID-19, ACE2, and the cardiovascular consequences. *American journal of physiology Heart and circulatory physiology* 2020;**318**:H1084-h1090.
77. Rodrigues-Diez RR, Tejera-Muñoz A, Marquez-Exposito L, Rayego-Mateos S, Sanchez LS, Marchant V, Santamaria LT, Ramos AM, Ortiz A, Egido J, Ruiz-Ortega M. Statins: Could an old friend help the fight against COVID-19? *Br J Pharmacol* 2020.
78. Bifulco M, Gazzero P. Statin therapy in COVID-19 infection: much more than a single pathway. *European Heart Journal - Cardiovascular Pharmacotherapy* 2020.
79. Castiglione V, Chiriaco M, Emdin M, Taddei S, Vergaro G. Statin therapy in COVID-19 infection. *European Heart Journal - Cardiovascular Pharmacotherapy* 2020.
80. Vasanthakumar N. Can beta-adrenergic blockers be used in the treatment of COVID-19? *Med Hypotheses* 2020;**142**:109809-109809.
81. Staedtke V, Bai RY, Kim K, Darvas M, Davila ML, Riggins GJ, Rothman PB, Papadopoulos N, Kinzler KW, Vogelstein B, Zhou S. Disruption of a self-amplifying catecholamine loop reduces cytokine release syndrome. *Nature* 2018;**564**:273-277.
82. Horby P, Lim WS, Emberson J, Mafham M, Bell J, Linsell L, Staplin N, Brightling C, Ustianowski A, Elmahi E, Prudon B, Green C, Felton T, Chadwick D, Rege K, Fegan C, Chappell LC, Faust SN, Jaki T, Jeffery K, Montgomery A, Rowan K, Juszczak E, Baillie JK, Haynes R, Landray MJ. Effect of Dexamethasone in Hospitalized Patients with COVID-19: Preliminary Report. *medRxiv* 2020:2020.2006.2022.20137273.
83. Karram T, Abbasi A, Keidar S, Golomb E, Hochberg I, Winaver J, Hoffman A, Abassi Z. Effects of spironolactone and eprosartan on cardiac remodeling

- and angiotensin-converting enzyme isoforms in rats with experimental heart failure. *American journal of physiology Heart and circulatory physiology* 2005;**289**:H1351-1358.
84. Sanchis-Gomar F, Lavie CJ, Perez-Quilis C, Henry BM, Lippi G. Angiotensin-Converting Enzyme 2 and Antihypertensives (Angiotensin Receptor Blockers and Angiotensin-Converting Enzyme Inhibitors) in Coronavirus Disease 2019. *Mayo Clin Proc* 2020;**95**:1222-1230.
  85. Reynolds HR, Adhikari S, Pulgarin C, Troxel AB, Iturrate E, Johnson SB, Hausvater A, Newman JD, Berger JS, Bangalore S, Katz SD, Fishman GI, Kunichoff D, Chen Y, Ogedegbe G, Hochman JS. Renin-Angiotensin-Aldosterone System Inhibitors and Risk of Covid-19. *N Engl J Med* 2020;**382**:2441-2448.
  86. Mancia G, Rea F, Ludergnani M, Apolone G, Corrao G. Renin-Angiotensin-Aldosterone System Blockers and the Risk of Covid-19. *N Engl J Med* 2020;**382**:2431-2440.
  87. Bromage DI, Cannata A, Rind IA, Gregorio C, Piper S, Shah AM, McDonagh TA. The impact of COVID-19 on heart failure hospitalization and management: report from a Heart Failure Unit in London during the peak of the pandemic. *European journal of heart failure* 2020.
  88. Wu Y, Wang F, Shen C, Peng W, Li D, Zhao C, Li Z, Li S, Bi Y, Yang Y, Gong Y, Xiao H, Fan Z, Tan S, Wu G, Tan W, Lu X, Fan C, Wang Q, Liu Y, Zhang C, Qi J, Gao GF, Gao F, Liu L. A noncompeting pair of human neutralizing antibodies block COVID-19 virus binding to its receptor ACE2. *Science* 2020;**368**:1274-1278.
  89. Emanuelli C, Badimon L, Martelli F, Potočnjak I, Carpusca I, Robinson EL, Devaux Y, Action obotE-CC. Call to action for the cardiovascular side of COVID-19: A call for cooperative action from the EU-CardioRNA COST Action. *European Heart Journal* 2020;**41**:1796-1797.

**FIGURE 1**



**FIGURE 2**

