



University of Dundee

Fit to WHO weight standard of European infants over time

Levin, Daniel; Marryat, Louise; Cole, Tim J.; McColl, John; Harjunmaa, Ulla; Ashorn, Per

Published in:
Archives of Disease in Childhood

DOI:
[10.1136/archdischild-2015-309594](https://doi.org/10.1136/archdischild-2015-309594)

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Levin, D., Marryat, L., Cole, T. J., McColl, J., Harjunmaa, U., Ashorn, P., & Wright, C. (2016). Fit to WHO weight standard of European infants over time. *Archives of Disease in Childhood*, 101(5), 455-460.
<https://doi.org/10.1136/archdischild-2015-309594>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



OPEN ACCESS

Fit to WHO weight standard of European infants over time

Daniel Levin,¹ Louise Marryat,¹ Tim J Cole,² John McColl,¹ Ulla Harjunmaa,^{3,4} Per Ashorn,^{3,4} Charlotte Wright⁵

¹School of Mathematics and Statistics, University of Glasgow, Glasgow, UK

²Population, Policy and Practice Programme, UCL Institute of Child Health, London, UK

³Department of International Health, University of Tampere School of Medicine, Tampere, Finland

⁴Department of Paediatrics, Tampere University Hospital, Tampere, Finland

⁵Department of Child Health, University of Glasgow, Queen Elizabeth University Hospital, Glasgow, UK

Correspondence to

Louise Marryat, School of Mathematics and Statistics, University of Glasgow, 15 University Gardens, Glasgow G12 8QW, UK; louise.marryat@glasgow.ac.uk

Received 17 August 2015

Revised 20 January 2016

Accepted 21 January 2016

Published Online First

16 February 2016

ABSTRACT

Objectives The 2006 WHO growth charts were created to provide an international standard for optimal growth, based on healthy, breastfed populations, but it has been suggested that Northern European children fit them poorly. This study uses infant weight data spanning 50 years to determine how well-nourished preschool children from different eras fit the WHO standard, and discuss the implications of deviations.

Design Four longitudinal datasets from the UK and one from Finland were used comprising over 8000 children born between 1959 and 2003. Weights from birth to 2 years were converted to age–sex-adjusted Z scores using the WHO standard and summarised using Generalized Additive Models for Location, Scale and Shape.

Results Weights showed a variable fit to the WHO standard. Mean weights for all cohorts were above the WHO median at birth, but dipped by up to 0.5 SD to a nadir at 8 weeks before rising again. Birth weights increased in successive cohorts and the initial dip became slightly shallower. By age 1 year, cohorts were up to 0.75 SD above the WHO median, but there was no consistent pattern by era.

Conclusions The WHO standard shows an acceptable, but variable fit for Northern European infants. While birth weights increased over time, there was, unexpectedly, no consistent variation by cohort beyond this initial period. Discrepancies in weight from the standard may reflect differences in measurement protocol and trends in infant feeding practice.

INTRODUCTION

Growth charts are widely used in child health to identify undernutrition and overnutrition.¹ Many countries produce their own charts describing how local children grow. However, these do not necessarily characterise healthy growth; in some countries average weight will be low due to undernutrition, while in high-income countries higher average weights reflect obesity. In the past, charts were based predominantly on bottle-fed infants and did not adequately reflect the growth of breastfed infants who should be the physiological norm.² Thus, the WHO developed new growth charts based on healthy breastfed children living in optimal circumstances in six world regions.³ The WHO showed that linear growth differed little between the six datasets⁴ and thus argued that they could be used to define how all children aged 0–5 years *should* grow, whatever their ethnic origin. However, weights by country for the WHO dataset have never been published. Since the publication of the standard, studies

What is already known on this topic

- ▶ Infants in high-income countries tend to fit the WHO 2006 growth standard well for length.
- ▶ Infants do, however, become heavier than the standard after 6 months.

What this study adds

- ▶ Northern European infants demonstrate a largely adequate fit to the WHO weight standard after the first 2 weeks.
- ▶ Infants born recently fit the WHO standard at birth and in the early weeks better than earlier cohorts.
- ▶ All the cohorts were heavier than the WHO standards by age 12 months, with no trend over time, suggesting that this cannot be explained by increasing rates of population obesity.

in unselected, healthy, non-deprived populations such as the UK,⁵ USA,⁶ Canada,⁷ Norway, Belgium,⁸ Italy, Argentina⁹ and Denmark¹⁰ have generally found a close fit for length, but a tendency for children to become heavier than the WHO standard after the first 6 months. Some authors have argued that these discrepancies represent a fundamental difference that renders the standard unsuitable for high-income countries.⁸ Others have suggested that variations in fit are to be expected if the WHO charts represent optimal rather than average growth, as few children will be breastfed to age 1 year¹¹ and that the higher weights reflect rising rates of obesity at all ages.⁵

If the mismatch is due to obesity, it ought to be less evident in historic datasets from eras when obesity was less common. Conversely, if it reflects the nature, prevalence and duration of formula feeding, the fit should be better in more recent cohorts as breastfeeding has increased.

Aims

Our aims are to explore a portfolio of weight datasets from the past 50 years (i) to determine how well real populations of well-nourished preschool children from different eras fit the 2006 WHO growth standard at different ages during infancy;



Open Access
Scan to access more
free content



CrossMark

To cite: Levin D, Marryat L, Cole TJ, et al. *Arch Dis Child* 2016;**101**:455–460.

and (ii) to explore how trends differ by era of the cohort and associated infant feeding patterns.

METHODS

Datasets

Data came from existing longitudinal growth studies, retrieved mainly from routine records. They had already been cleaned, checked and analysed for other purposes, with four studies already published. Details of the five studies are as follows:

Widdowson study (1959)

In a study set up by Dr Elsie Widdowson, routine weights of 1094 babies born in 1959–1965 were obtained contemporaneously from the records of 10 Cambridge Child Welfare Clinics. Weights were measured by clinical staff approximately monthly in the first year of life, with a maximum of 13 weights per child. Of the possible 14 222 measurements all but 864 (6%) were collected. Although the data were cleaned and analysed at the time of collection the results were never published.

Cambridge Infant Growth study (1984)

Cambridge Infant Growth Study (CIGS) was a research cohort of 255 babies recruited in 1984–1987. Infants were sampled in four cohorts from lists of Cambridge city mothers booked to deliver in particular months, with some filtering by midwives. Measurements were taken mainly by one highly trained auxologist every 4 weeks from 4 to 52 weeks, then at 15, 18, 24, 30, 36 and 48 months. Weight, length, head and arm circumferences, triceps and subscapular skinfolds were measured at each visit: 223 (87%) had all 15 measurements from birth to 2 years.^{12–14}

Newcastle Growth and Development study (1987)

This dataset comprises the routine weights of a birth cohort of 3418 children born at term in Newcastle upon Tyne between June 1987 and May 1988. Up to 11 weights measured by clinical staff in infancy were retrieved from baby clinic records, and 3060 of the babies had at least two weights.^{15 16}

Gateshead Millennium study (1999)

Gateshead Millennium Study (GMS) is a birth cohort of 1029 babies (923 term) born in Gateshead in 1999–2000, representing 81% of eligible births during the recruitment period. Routine weights were retrieved from baby clinic records. There was a mean of 13 weights per child in the first year. Research nurses measured 830 infants at 13 months.^{17–19}

Tampere study, Finland (2003)

This dataset comprises the routine heights and weights of 2809 children aged 0–4 years born between October 2003 and September 2004 who attended child health clinics in Tampere. Children were weighed by clinical staff on electronic scales. Up to 16 scheduled events were recorded per child at birth, 1–2 weeks, 6–8 weeks, 2, 3, 4, 5, 6, 8, 10, 12, 18, 24, 36 and 48 months. There was a mean of 12 measurements per child.²⁰

Analysis

The datasets were cleaned and weights converted into Z scores relative to the WHO growth standard. Data beyond 2 years were excluded, when numbers were low and bias was likely. The measurements at birth and 6–8 weeks, 3, 6, 9, 12, 18 and 24 months were summarised by age, sex and cohort (table 1). The mean Z score, SD, skewness and kurtosis were modelled as functions of age for each dataset by sex, using Generalized Additive Models for Location, Scale and Shape (GAMLSS) as implemented in the GAMLSS package in R V.3.1.1. Multiple GAMLSS models were fitted using different hyperparameters and distribution families. In the final models, based on the Box-Cox power exponential family (BCPE), the mean Z score was allowed to vary with age in each of the datasets, whereas the SD, skewness and kurtosis were constrained to be constant as exploratory analyses showed this made little difference to the model fit as determined using the Bayesian Information Criterion. The skewness adjustment ensured that the mean and median were effectively the same. The BCPE requires values to be positive, so all Z scores had 10 added to them prior to analysis, and 10 was then subtracted from the mean curves. The

Table 1 Number of observations at target ages by dataset

Target age (weeks)	Age range	Sex	Number of observations				
			Widdowson (1959)	CIGS (1984)	GDS (1987)	GMS (1999)	Tampere (2003)
0	Birth	M	573	142	1589	484	929
		F	519	122	1582	477	889
7	5–9 weeks	M	452	135	1201	388	1202
		F	410	116	1217	379	1145
13	2–4 months	M	573	137	1357	392	1220
		F	520	118	1365	389	1189
26	5–7 months	M	570	135	1186	240	1192
		F	516	117	1212	247	1174
39	8–10 months	M	569	134	898	222	993
		F	515	116	893	233	962
52	11–14 months	M	501	131	1068	308	1161
		F	476	116	1109	300	1134
78	15–21 months	M	–	125	642	61	1132
		F	–	116	670	66	1098
104	22–30 months	M	–	124	122	–	1076
		F	–	115	115	–	1068

CIGS, Cambridge Infant Growth Study; GDS, Growth and Development Study; GMS, Gateshead Millennium Study.

Table 2 Mean (SD) weight Z scores at target ages by dataset

Target age (weeks)	Sex	Mean (SD) weight Z score of children by age, sex and study cohort, Z-score units				
		Widdowson (1959)	CIGS (1984)	GDS (1987)	GMS (1999)	Tampere (2003)
0	M	0.03 (1.11)	0.06 (0.97)	0.12 (1.01)	0.18 (1.05)	0.37 (1.08)
	F	0.10 (1.01)	0.30 (0.89)	0.08 (1.07)	0.17 (1.1)	0.40 (1.1)
7	M	-0.40 (1.02)	-0.28 (0.99)	-0.25 (0.95)	-0.16 (0.96)	0.14 (1.05)
	F	-0.30 (0.83)	-0.19 (0.83)	-0.30 (0.91)	-0.17 (0.9)	0.12 (0.95)
13	M	-0.33 (1.07)	-0.34 (0.97)	-0.21 (0.98)	-0.16 (0.97)	0.17 (1.03)
	F	-0.17 (0.83)	-0.22 (0.86)	-0.29 (0.9)	-0.18 (0.9)	0.12 (0.91)
26	M	0.28 (0.99)	-0.09 (0.98)	0.15 (1.01)	0.30 (0.99)	0.46 (1.07)
	F	0.42 (0.82)	0.05 (0.87)	0.05 (0.91)	0.21 (0.93)	0.38 (0.93)
39	M	0.62 (1)	0.05 (0.97)	0.38 (1.05)	0.56 (1.01)	0.56 (1.09)
	F	<i>0.67 (0.8)</i>	0.21 (0.86)	0.23 (0.93)	0.43 (0.92)	0.48 (0.96)
52	M	<i>0.69 (0.97)</i>	0.16 (0.97)	0.53 (1.01)	0.55 (1.06)	<i>0.60 (1.01)</i>
	F	<i>0.76 (0.81)</i>	0.26 (0.83)	0.38 (0.89)	0.51 (0.91)	0.52 (0.91)
78	M	-	0.20 (0.94)	0.46 (1)	0.44 (0.97)	<i>0.62 (0.96)</i>
	F	-	0.23 (0.76)	0.27 (0.95)	0.41 (1.01)	0.56 (0.89)
104	M	-	0.04 (0.96)	0.38 (1.12)	-	0.52 (0.96)
	F	-	0.18 (0.71)	0.12 (1.12)	-	0.53 (0.91)

Fit: excellent (≤ 0.17 SD), poor (> 0.67 SD).

CIGS, Cambridge Infant Growth Study; GDS, Growth and Development Study; GMS, Gateshead Millennium Study.

models for each cohort were plotted as mean Z score versus age in boys and girls, along with the constant SD, skewness and kurtosis, to compare how well the cohorts fitted the WHO 2006 standard. A mean Z score of 0, SD 1, skewness 1 and kurtosis 2 indicates a perfect fit, whereas a mean Z score above 0 is heavier than the WHO standard, and below is lighter.

There is no standard definition of what constitutes a good or poor fit to a growth reference. In *this* study fit was defined in terms of mean weight Z score, measured in fractions of a centile channel width relative to the WHO median, where one channel width = 0.67 SDs.²¹ An excellent fit was defined as a difference of no more than ¼ of a channel width (0.17 SD) and a poor fit as greater than a channel width (0.67 SD).

RESULTS

Mean birthweight Z scores in the five datasets were all positive and close to zero (table 2). They were progressively higher in later years, particularly in the boys, rising from 0.03 in 1959 to 0.37 in 2003. For all cohorts the fit was adequate in early infancy, mostly staying within half a channel width of the median, but by 1 year boys and girls in Widdowson and boys in Tampere fitted poorly, more than a channel width above the median.

The smoothed curves of mean Z score versus age from the GAMLSS models were plotted by cohort on separate charts by sex, along with the constant SD (sigma), skewness (nu) and kurtosis (tau) (figure 1). The curves provide a visual assessment of the fit to the standard by cohort. All datasets showed consistent differences relative to the standard. The mean Z scores all tended to fall in the early weeks, with the four UK datasets ending up below zero. This steep fall was followed by a slightly less steep rise, creating a 'trough' in each curve at around 8 weeks. Boys and girls followed broadly the same pattern.

The depth of the trough, like birthweight, became progressively shallower over time. Mean Z score fell by between 0.2 and 0.5 SDs from birth to 7 weeks, the fall tending to be greater in the earlier cohorts (table 3). The earliest dataset (Widdowson) also showed the steepest rise in Z score after the

trough, so that by 52 weeks it was the heaviest, around a channel width above the median. In contrast, the second earliest dataset (CIGS) rose least and remained close to the median. Mean Z scores in the remaining datasets rose by around 0.5 SD and tended to level out after 52 weeks. It should be noted that the Widdowson and GMS datasets had only limited data in later infancy, and thus the trajectory curves are incomplete. Patterns for the girls and boys were similar (figure 1). The fitted constant SDs showed a good fit relative to the WHO standard, being close to 1 throughout. Skewness was less consistent, with negative skewness (nu < 1) in CIGS and positive skewness (nu > 1) in the other datasets. Kurtosis was near 2 as expected.


DISCUSSION

The strengths of this study are that the five datasets represent weight gain before and during the obesity epidemic. The datasets are of high quality and all but CIGS are representative of their populations, with high recruitment rates. The GAMLSS modelling approach makes maximum use of the available data and controls for potential bias. A weakness is that the most contemporary study is from Finland, not the UK, where growth patterns may differ systematically.

Overall, most cohorts were within a channel width of the WHO median and many showed an excellent fit in the early weeks, though less so beyond this point. It seems unlikely that these later differences could be simply genetic, since two of the six WHO datasets were of predominantly Northern European origin and they showed no consistent differences in stature.²² The lack of variation by era argues against obesity being the explanation.

The increase in birthweight over time is consistent with research in other high-income countries.^{3 22 23} This is generally thought to reflect less maternal cigarette smoking and more maternal obesity and gestational diabetes.²⁴ It has been suggested that the higher birthweight seen for all these cohorts compared with the WHO standard may be explained by prior maternal undernutrition in some of the cohorts used to develop the WHO standard.²⁵ However, a recent study suggested that

Figure 1 Mean weight Z scores by age in five Northern European cohorts (top boys; bottom girls). CIGS, Cambridge Infant Growth Study; GDS, Growth and Development Study; GMS, Gateshead Millennium Study.



birth weights across diverse countries, including some of those included in the WHO Multicentre Growth Reference Study (MGRS) cohorts, were not significantly different by country.⁵

Table 3 Change in mean (SD) weight Z scores between target ages by dataset

Target age (weeks)	Sex	Change in SDS				
		Widdowson (1959)	CIGS (1984)	GDS (1987)	GMS (1999)	Tampere (2003)
0–7	M	−0.43	−0.34	−0.37	−0.34	−0.23
	F	−0.40	−0.49	−0.38	−0.34	−0.28
7–52	M	1.09	0.44	0.78	0.71	0.46
	F	1.06	0.45	0.68	0.68	0.40

CIGS, Cambridge Infant Growth Study; GDS, Growth and Development Study; GMS, Gateshead Millennium Study.

Though born heavier, the infants here initially lost weight relative to the WHO standard and then regained it, causing a trough to appear in the Z score growth curve. This has been described in other cohorts,^{5–8} and was influential in the UK rejecting the use of the WHO standard at birth.²⁶ This pattern is the mirror image of the way breastfed infants used to grow on charts based mainly on bottle-fed infants and though seen in all datasets, its depth decreases over time.^{2–5} We do not have information about breastfeeding rates in the routine cohorts but most will have been lower than the WHO sample.²⁷ However, breastfeeding initiation rates have risen in the UK over the past 50 years from 36% in 1970 to over 70% in the early 21st century (UK)^{28–30} while when the Tampere cohort were born, 93% of Finnish infants were initially breastfed.²⁰ The CIGS dataset, a more selective research sample, showed the shallowest trough and had higher rates of initial breastfeeding (75%) and breastfeeding to 6 months (48%).³¹ This pattern by era may also reflect changes in formula milk composition which has increasingly mirrored the nutritional content of breast milk.³²

In contrast, the later rise relative to the standard did not show the same trend over time, suggesting that it may be unrelated to breastfeeding. While observational studies have found an association between use of breast milk substitutes and faster weight gain later in infancy^{2 31} this effect was not seen in the Belarus trial of breastfeeding promotion.³³ The sampling process used to construct the WHO standard could in principle mean that the MGRS cohorts were lighter than the general population. Children who were either outliers or not breastfed to 12 months were excluded from the sample¹ and it has been shown that larger children are more likely to cease breastfeeding early.^{33 34} However, the WHO MGRS group did not find any systematic difference in size between those included and excluded (personal communication De Onis, email communication, 2006).

Measurement error might also play a role, since the MGRS cohorts were measured using the same research protocols, where all clothes were removed or adjusted for.^{4 35} The present cohorts mainly used routine data collected by health professionals; UK and Finnish guidelines recommend weighing infants naked up to the age of 2 years,³⁶ but parents of older infants may be reluctant to comply, leading to higher weights at later ages. The CIGS cohort where all infants were weighed naked to a research protocol tracks closest to the WHO standard beyond 6 months. However, another study that compared routinely collected measurements with research measurements of the same children found little difference in weight.³⁷

CONCLUSIONS

The overall fit to the WHO weight standard of these cohorts ranged from excellent to adequate, so that use of the WHO standard is unlikely to introduce major bias in the assessment of individual children. Some of the more subtle variation in fit is likely to reflect variations in the levels of breast and formula feeding, as well as the composition of formula milk, in different eras. However, the lack of a consistent trend for weight gain after the initial weeks suggests that the higher average weight gain seen in North European cohorts cannot simply be explained by increasing rates of obesity in later childhood and adulthood.

Twitter Follow Louise Marryat at @LMarryat

Acknowledgements This study would not have been possible without the hard work of many researchers who collected and processed the growth data for the cohorts included here, as well as the parents who participated in the Cambridge Infant Growth Study.

Contributors DL cleaned the data, carried out the analyses, drafted the initial manuscript, reviewed further drafts of the manuscript and approved the final manuscript. LM carried out further analyses of the data, revised the manuscript and approved the final manuscript. TJC conceptualised and designed the analysis, provided access to the Cambridge Infant Growth Study and Widdowson datasets, reviewed and revised the manuscript and approved the final manuscript. JM conceptualised and designed the study, reviewed and revised the manuscript and approved the final manuscript. UH designed the Finnish data collection and collected the Finnish data, reviewed and revised the manuscript and approved the final manuscript. PA coordinated a study that yielded the Finnish database used in this study, revised the manuscript and approved the final manuscript. CW was PI for the Gateshead Millennium Study and Newcastle Growth and Development Study, conceptualised and designed this analysis, reviewed and revised the manuscript and approved the final manuscript.

Funding Chief Scientist Office, Scotland. TJC is funded by MRC grant MR/M012069/1.

Competing interests TJC and CW were involved in the decision to adopt the WHO standard in the UK.

Provenance and peer review Not commissioned; internally peer reviewed.

Data sharing statement This paper uses secondary data analysis from five studies. To the authors' knowledge none are currently publicly available.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- de Onis M, Wijnhoven TMA, Onyango AW. Worldwide practices in child growth monitoring. *J Pediatr* 2004;144:461–5.
- Dewey KG, Heinig MJ, Nommsen LA, et al. Growth of breast-fed and formula-fed infants from 0 to 18 months: the DARLING study. *Pediatrics* 1992;89:1035–41.
- de Onis M, Garcia C, Onyango AW, et al. WHO Child Growth Standards based on length/height, weight and age. *Acta Paediatr Suppl* 2006;450:76.
- de Onis M. Assessment of differences in linear growth among populations in the WHO Multicentre Growth Reference Study. *Acta Paediatr Suppl* 2006;95 (S450):56–65.
- Wright C, Lakshman R, Emmett P, et al. Implications of adopting the WHO 2006 Child Growth Standard in the UK: two prospective cohort studies. *Arch Dis Child* 2008;93:566–9.
- Mei Z, Ogden CL, Flegal KM, et al. Comparison of the Prevalence of Shortness, Underweight, and Overweight among US Children Aged 0 to 59 Months by Using the CDC 2000 and the WHO 2006 Growth Charts. *J Pediatr* 2008;153:622–8.
- van Dijk CE, Innis SM. Growth-Curve Standards and the Assessment of Early Excess Weight Gain in Infancy. *Pediatrics* 2009;123:102–8.
- Júlíusson PB, Roelants M, Hoppenbrouwers K, et al. Growth of Belgian and Norwegian children compared to the WHO growth standards: prevalence below –2 and above +2 standard deviations and the effect of breastfeeding. *Arch Dis Child* 2011;96:916–21.
- Onyango AW, de Onis M, Caroli M, et al. Field-Testing the WHO Child Growth Standards in Four Countries. *J Nutr* 2007;137:149–52.
- Tinggaard J, Aksglaede L, Sørensen K, et al. The 2014 Danish references from birth to 20 years for height, weight and body mass index. *Acta Paediatr* 2014;103:214–24.
- Norris SA, Griffiths P, Pettifor JM, et al. Implications of adopting the WHO 2006 Child Growth Standards: Case study from urban South Africa, the Birth to Twenty cohort. *Ann Hum Biol* 2009;36:21–7.
- Wright CM, Waterston A, Aynsley-Green A. Comparison of the use of Tanner and Whitehouse, NCHS, and Cambridge standards in infancy. *Arch Dis Child* 1993;69:420–2.
- Paul AA, Cole TJ, Ahmed EA, et al. The need for revised standards for skinfold thickness in infancy. *Arch Dis Child* 1998;78:354–8.
- Cole TJ, Paul AA, Eccles M, et al. The use of a multiple growth standard to highlight the effects of diet and infection on growth. In: Perspectives in the science of growth and development. *Auxology* 1989;88:91–100.
- Drewett RF, Corbett SS, Wright CM. Cognitive and educational attainments at school age of children who failed to thrive in infancy: a population-based study. *J Child Psychol Psychiatry* 1999;40:551–61.
- Drewett RF, Corbett SS, Wright CM. Physical and emotional development, appetite and body image in adolescents who failed to thrive as infants. *J Child Psychol Psychiatry* 2006;47:524–31.
- Wright CM, Matthews JNS, Waterston A, et al. What is a normal rate of weight gain in infancy? *Acta Paediatr* 1994;83:351–6.
- Wright CM, Cheetham TD. The strengths and limitations of parental heights as a predictor of attained height. *Arch Dis Child* 1999;81:257–60.
- Parkinson K, Wright C, Drewett R. The Gateshead Millennium Baby Study: A Prospective Study of Feeding and Growth. *Int J Soc Res Methodol* 2007;10:335–47.
- Harjunmaa U. *Current growth patterns of Finnish children aged 0 to 4 years*. Master's Thesis. University of Tampere, 2009. <http://tampub.uta.fi/handle/10024/81098>
- Cole TJ. Do growth chart centiles need a face lift? *BMJ* 1994;308:641.
- Chike-Obi U, David RJ, Coutinho R, et al. Birth Weight Has Increased Over a Generation. *Am J Epidemiol* 1996;144:563–9.
- Hadfield RM, Lain SJ, Simpson JM, et al. Are babies getting bigger? An analysis of birth weight trends in new south wales, 1990–2005. *Obstet Gynecol Surv* 2009;64:445–7.
- Ananth CV, Wen SW. Trends in fetal growth among singleton gestations in the United States and Canada, 1985 through 1998. *Semin Perinatol* 2002;26:260–7.
- Kramer MS, Morin I, Yang H, et al. Why are babies getting bigger? Temporal trends in fetal growth and its determinants. *J Pediatr* 2002;141:538–42.
- Wright CM, Williams AF, Elliman D, et al. PRACTICE POINTER: Using the new UK-WHO growth charts. *BMJ* 2010;647–50.
- Whitehead RG, Paul AA, Cole TJ. Diet and the growth of healthy infants. *J Hum Nutr Diet* 1989;2:73–84.
- Denny K. Breastfeeding predicts handedness. *Laterality* 2012;17:361–8.

- 29 McAndrew F, Thompson J, Fellows L, *et al*. *Infant feeding survey 2010*. Leeds: Health and Social Care Information Centre, 2012.
- 30 Sacker A, Kelly Y, Iacovou M, *et al*. Breast feeding and intergenerational social mobility: what are the mechanisms? *Arch Dis Child* 2013;98:666–71.
- 31 Cole TJ, Freeman JV, Preece MA. Body mass index reference curves for the UK, 1990. *Arch Dis Child* 1995;73:25–9.
- 32 Heird WC. Progress in Promoting Breast-Feeding, Combating Malnutrition, and Composition and Use of Infant Formula, 1981–2006. *J Nutr* 2007;137:499S–502S.
- 33 Kramer MS, Guo T, Platt RW, *et al*. Breastfeeding and infant growth: biology or bias? *Pediatrics* 2002;110:343–7.
- 34 Wright CM, Parkinson K, Scott J. Breast-feeding in a UK urban context: who breast-feeds, for how long and does it matter? *Public Health Nutr* 2006;9:686–91.
- 35 de Onis M, Onyango AW, Van den Broeck J, *et al*. Measurement and standardization protocols for anthropometry used in the construction of a new international growth reference. *Food Nutr Bull* 2004;25(1 Suppl):S27–36.
- 36 Wright CM, Parkinson KN, Drewett RF. Why are babies weaned early? Data from a prospective population based cohort study. *Arch Dis Child* 2004;89:813–16.
- 37 Coles LN, Bates K, Phillips K, *et al*. Standards for the weighing of infants, children and young people in the acute healthcare settings. In: Nursing RCo, ed. London: RCN, 2013. https://www2.rcn.org.uk/__data/assets/pdf_file/0009/351972/003828.pdf