

How to create coats for all seasons: elucidating antigenic variation in African trypanosomes

Cher-Pheng Ooi and Gloria Rudenko

Department of Life Sciences, Sir Alexander Fleming Building, Imperial College London, South Kensington, London SW7 2AZ, U.K.

Correspondence: Gloria Rudenko (gloria.rudenko@imperial.ac.uk)



Extracellular parasites of the mammalian bloodstream face considerable challenges including incessant assault by the immune system. African trypanosomes are consummate survivors in this inclement environment and are renowned for their supremely sophisticated strategy of antigenic variation of their protective surface coat during the course of chronic infections. Recent developments are making us realize how complex this antigenic machinery is and are allowing us to tackle previously intractable problems. However, many of the simplest (and arguably the most important) questions still remain unanswered!

How is the extraordinary heterogeneity of variant surface glycoprotein variants during a chronic infection generated?

The bloodstream form of *Trypanosoma brucei* is coated with a dense covering of $\sim 10^7$ variant surface glycoprotein (VSG) molecules, which extend as antigenically variable rods connected to the cell surface via glycosylphosphatidylinositol (GPI) anchors [1]. This GPI attachment allows VSGs to move freely over the cell surface [2]. Tight packing of these VSGs produces a protective barrier shielding invariant surface proteins from recognition by host antibodies and preventing trypanosome lysis by the alternative pathway of the complement system [3]. Extraordinarily high rates of recycling of surface VSG allow selective removal of host cell molecules including antibodies and complement from the trypanosome surface, thereby functioning as a ‘coat-cleaning’ machine [4]. This prolongs trypanosome survival in rising antibody titres and facilitates escape from macrophages [5].

Each trypanosome has a vast repertoire of VSG genes [6], differing in number and content between different strains, presumably as a consequence of diversifying forces. For example, the ‘VSGnome’ of *T. brucei* 427 contains more than 2500 different VSG genes, of which more than 80% are not immediately functional [7]. Switching the active VSG can entail transcriptional switches between the ~ 15 telomeric VSG expression site (ES) transcription units (Figure 1). More importantly, DNA rearrangements and predominantly gene conversions can copy new VSGs (or segments of VSGs) into the active ES, thereby resulting in an antigenic switch [8] (Figure 1). This extraordinary ability to construct novel patchwork, or mosaic, VSGs is key for trypanosome survival [9]. The continuous recombination of various permutations and combinations of sections of VSG genes (or pseudogenes) allows the trypanosome to construct an almost infinite number of antigenically distinct VSG coat types.

A given wave of trypanosome parasitaemia is normally predominantly composed of a major VSG variant, although it has long been known that minor VSG variants are also present within each wave. However, there is possibly phenomenal heterogeneity within these infection waves, as next-generation RNA sequencing of RNA transcripts has documented an extraordinary variety of minor VSG variants [10]. These experiments argue that a considerable percentage of the available VSG repertoire is displayed relatively early in the trypanosome infection. This highlights the necessity of the continuous generation of mosaic VSGs for long-term trypanosome survival. However, this also gives rise to several questions. Where are these new VSGs made? Within a silent ES? How then does the

Received: 11 October 2017

Revised: 8 November 2017

Accepted: 10 November 2017

Version of Record published:
22 December 2017




Figure 1. Mechanisms for switching VSG.

VSG switching can be mediated through DNA rearrangements (gene conversion or telomere exchange) or transcriptional control. Duplicative gene conversion involves copying an intact VSG (A) or segments of a VSG (B) into the active VSG ES resulting in loss of the old VSG. After a telomere exchange (C), there is no loss of genetic information. A transcriptional switch entails activating a new ES and silencing the old one (D). ESs are shown with the promoters indicated with flags, VSG genes with coloured boxes, and transcription of the active ES with an arrow.

trypanosome ‘test’ that a new VSG cobbled together from different pieces is indeed functional? Within an active ES? However, VSG is essential for the bloodstream-form trypanosome, and blocking its synthesis triggers a precytokinesis arrest [11]. This does not give much time for multiple successive gene conversions to occur,

unless this process is transiently stimulated. What is the DNA recombination machinery that allows the trypanosome to construct these novel VSG mosaics? Is this specifically dedicated to the generation of VSG diversity?

Key questions remain unanswered. However, a major complication with analyzing VSG switching in chronic infections is that establishing the order of VSG switching is often not possible. Major variants following each other in successive waves of parasitaemia might not necessarily have been generated from each other, but can be the consequence of the emergence of minor variants that have arisen much earlier. Unravelling this complexity will be seriously non-trivial.

How is VSG expression controlled in a monoallelic fashion?

To allow frugal use of the VSG repertoire, only a single VSG is highly expressed per cell from one of ~15 highly similar telomeric VSG ESs [12]. ES transcription is mediated by RNA polymerase I (Pol I), which also transcribes the ribosomal DNA [13]. The use of Pol I to transcribe protein-coding genes is unprecedented in eukaryotes, where protein-coding genes are normally exclusively transcribed by Pol II. The use of an unconventional RNA polymerase makes it likely that ES control will involve some very novel features, which yet remain to be unravelled. The active ES is located in a unique non-nucleolar Pol I-containing structure called the expression site body (ESB) [14]. Low levels of transcripts are generated from 'silent' ESs which are not located in the ESB [15], but these RNAs are predominantly non-polyadenylated [16]. This could argue that the ESB is a 'factory' which contains components allowing efficient transcription initiation and elongation at the active ES, as well as RNA processing including trans-splicing and polyadenylation. The ESB appears to be Pol I-transcription nucleated, as it disappears exceptionally rapidly after the specific blocking of Pol I transcription using chemical Pol I inhibitors [17].

Whole-genome RNAi libraries are now allowing the development of functional screens to identify essential components involved in basic biochemical processes in African trypanosomes [18]. A whole-genome RNAi screen designed to select for derepression of a silent *T. brucei* telomere has allowed the identification of the first ESB-specific marker, a protein called VEX1 [19]. The identification of additional ESB-specific components, as well as determination of their respective functions, should give us insights into the functioning of this unique subnuclear body.

A single trypanosome transcribes a single ES at a time, with the remaining 14 ESs remaining essentially inactive. What mediates this singularity of ES expression [20]? Using drug selection pressure, it is possible to force the trypanosome to simultaneously activate a second ES; however, this state is not well tolerated by the cell. The resulting 'double-expresser' state is highly unstable, and the trypanosomes appear to rapidly switch back and forth between the two ESs [21]. More detailed analysis of these 'double-expresser' strains should give us insights into the epigenetic modifications which facilitate this perturbation of the monoallelic exclusion machinery. However, central questions regarding the molecular restrictions mediating this striking singular ES expression still remain unanswered.

How do epigenetic factors affect VSG expression and switching?

DNA in eukaryotes is wrapped around nucleosomes composed of the core histones, with chromatin remodeling allowing the cell to modulate the accessibility of DNA regulatory sequences to the transcriptional machinery. Specific post-translational modifications of the histones, particularly at their N-termini, allow the cell to functionally define different types of chromatin. African trypanosomes do not appear to regulate their Pol II transcription, and the bulk of the genome is constitutively transcribed in extensive polycistronic transcription units [22]. This has led to a relatively late appreciation of the role of chromatin in the organization and transcription of the *T. brucei* genome.

Next-generation sequencing is now allowing the investigation of nucleosome positioning in unprecedented detail across the entire *T. brucei* genome. The position of individual nucleosomes across all of the core chromosomes has now been determined in both bloodstream- and procyclic-form *T. brucei* [23,24]. This has allowed us to determine the chromatin state at different types of transcription units. Pol II promoters in *T. brucei* are diffuse and lack distinctive sequence elements [25]. Instead, Pol II promoters appear to be predominantly functionally defined by discrete chromatin structure around the strand switch regions where Pol II transcription initiates [24]. In addition, chromatin structure appears to play a role in the repression of the silent VSG genes,

and positioned nucleosomes are found flanking the silent *VSG* gene arrays [23]. The chromatin structure in the border regions of these *VSG* arrays could represent barrier regions which repress cryptic transcription initiation or readthrough into the *VSG* arrays, keeping these *VSGs* silent.

In contrast, the active ES has an unusually open chromatin structure compared with most of the *T. brucei* genome and is stripped bare of most nucleosomes, as well as the linker histone H1 (Figure 2) [26–28]. This strikingly open chromatin state is characteristic of Pol I-transcription units and presumably facilitates the exceptionally high rate of transcription initiation seen here [29]. In place of nucleosomes, the architectural chromatin protein TDP1 coats the active *VSG* ES [30]. TDP1 is a high mobility group protein which is likely to be analogous to HMO1 in yeast [31]. HMO1 coats active Pol I-transcription units with a mutually exclusive distribution to the linker histone H1 and appears to stabilize chromatin depleted of nucleosomes [29,32]. Consistent with this, in *T. brucei*, TDP1 appears to help maintain an active chromatin state at the ES, even if ES transcription itself is inducibly blocked [33].

In contrast, silent ESs are marked with an unusual modified glucosylated nucleotide called Base J, which is incorporated in a gradient that increases towards the telomere end and could play some role in silencing these telomeric *VSGs* [34,35]. Additionally, different histone variants and modifications have also been shown to play a role in the down-regulation of ESs (Figure 2). The histone variant H3.V is enriched around the termination region of Pol II transcription units [25]. Knockdown of histone variant H3.V not only results in the disruption of Pol II transcription termination but also causes the derepression of the silent telomeric ESs [34,35]. Histone modifications also play an important role in ES control. Knockdown of the H3K76 histone methyltransferase DOT1B results in derepression of silent ESs and slows down the kinetics of transcriptional switching between ESs [36]. However, histone modifications, which are unique for ESs, have yet to be identified. Mass spectrometry analysis of *T. brucei* histones should allow us to identify additional modifications which could prove to be crucial for understanding epigenetic control.

It is now clear that a wide range of epigenetic factors play a role in the control of *VSG* expression, and a long list of chromatin remodellers, histone-modifying enzymes, chromatin ‘readers’ of histone modifications and chromatin architectural proteins have been shown to be important [37]. It is vital that more are identified, as the connections between these different puzzle pieces still remain to be elucidated.

How does repressed chromatin suppress *VSG* switching mediated through DNA recombination?

ESs are extensive telomeric transcription units, normally ranging 40–60 kb [12]. Repressive chromatin domains or gradients appear to control ESs from both ends [20]. An inhibitory chromatin state appears to restrict transcription initiation and elongation at *VSG* ES promoters, as knockdown of chromatin remodellers including ISWI and FACT results in derepression of silent ESs [38,39]. In addition, a repressive chromatin gradient mediated by the chromatin protein RAP1 extends up from the telomere end, and knockdown of RAP1 results in derepression of silent ESs [40].

However, in addition to having an impact on transcription, repressed chromatin states also appear to suppress excessively high rates of DNA recombination at the ES telomere. RAP1 knockdown results not only in ES depression but also in an increase in double-strand DNA breaks at the ES telomere, with a concurrent increase in *VSG* switching mediated by DNA recombination [41]. Intriguingly, this appears to occur through transcriptional regulation of a non-coding RNA. Knockdown of RAP1 results in the depression of transcription of the telomeric non-coding RNA: TERRA. TERRA forms DNA/RNA hybrids at the telomeres which stimulate the generation of double-strand breaks, thereby facilitating DNA recombination-mediated *VSG* switching. Similarly, the telomere-binding proteins TRF and TIF2 also appear to play a role in suppressing DNA recombination at *T. brucei* telomeres, with their knockdown resulting in increased double-strand breaks and *VSG* switching [42,43]. However, it still remains to be determined if these proteins directly establish a repressed chromatin state at the *T. brucei* telomeres.

How is the *VSG* switch mediated? Is there a link between antigenic variation and differentiation?

It has recently been proposed that antigenic variation is linked with differentiation in *T. brucei* [44,45]. Overexpression of a second ectopic *VSG* from an inducible promoter can result in transient silencing of the active ES proceeding upwards from the telomere. This transcriptional attenuation requires DOT1B-mediated




Figure 2. Schematic of VSG expression site control mediated via opposing repressive chromatin gradients.

ESs in different activation states are shown with flags indicating the promoters, transcription with a red arrow and telomere repeats with arrow heads. Chromatin proteins and remodellers (indicated with coloured circles) help maintain a silenced chromatin state at inactive ES promoters. Telomere-binding proteins including RAP1 and TRF help maintain a repressed state at the chromosome end. These operate in a gradient extending upwards from the chromosome end, in combination with the histone methyltransferase DOT1B. The linker histone H1 coats inactivate ESs (blue circles), but upon ES activation is replaced with TDP1 (open balls), which maintains an open chromatin state. VEX1 is a marker for the active ES.

histone methylation [36,46]. This provides evidence for the presence of a reversible chromatin-mediated silencing gradient extending up from the telomere end, and thereby having an impact on transcriptional elongation at the ES telomere. One possibility is that upon overexpression of the ectopic VSG, the trypanosome down-regulates the endogenous VSG in order to maintain homeostasis of VSG levels [47].

However, overexpression of the ectopic VSG also results in the generation of non-replicating ‘stumpy form’ parasites. These non-proliferative trypanosomes are pre-adapted to life in the tsetse fly and typically arise when cells sense that densities have risen above a given threshold. The authors therefore additionally postulate that this ES silencing is a ‘sensor’ that triggers stumpy differentiation and propose that this is initiated through silencing of *ESAG* genes located at the ES telomere [44].

However, an alternative interpretation for the results of Batram et al. [44] is that the overexpression of ectopic VSG (or knockdown of *ESAGs*) produces stress responses which trigger the generation of the stumpy form. One of the first events that occurs upon differentiation to the stumpy form is progressive attenuation of transcription of the active ES. This silencing gradient extends up from the telomere along the active ES, whereby the telomeric VSG is the first gene to be silenced [48]. The concurrent attenuation of VSG transcription at the active ES could therefore be a side effect of this stumpy form differentiation rather than directly causing it. This issue is discussed in ref. [45], but remains controversial. However, despite lack of clarity regarding the exact mechanistic details, these intriguing observations all argue that there is an intricate interplay between antigenic variation and trypanosome differentiation which still needs to be further unravelled.

In summary, new developments have highlighted the unprecedented complexity underlying antigenic variation in African trypanosomes. New experimental approaches are allowing us to tackle these issues at the whole-genome level using high-throughput sequencing of entire genomes or transcriptomes. Similarly, high-end proteomic approaches are allowing us to determine the interactomes of key players in the antigenic variation machinery. Whole-genome RNAi libraries are enabling identification of novel components using functional screens. Effective as these new approaches are, they are raising many more questions than answers regarding how African trypanosomes survive so well in the mammalian bloodstream.

Outstanding questions

- How are chimeric VSG genes assembled? Is there a dedicated DNA recombination pathway for this?
- Where are chimeric VSGs assembled?
- What epigenetic factors mark silent vs. active VSG ESs?
- How do these components regulate the monoallelic expression of a single ES?
- How does antigenic variation have an impact on trypanosome differentiation and *vice versa*?

Abbreviations

ES, expression site; ESB, expression site body; GPI, glycosylphosphatidylinositol; Pol I, polymerase I; VSG, variant surface glycoprotein.

Funding

G.R. is a Wellcome Senior Fellow in the Basic Molecular Sciences [095161] and the research was funded by the Wellcome Trust.

Acknowledgements

We apologize to those who were not cited due to space constraints. We are grateful to Calvin Tiengwe, Carys Davies, James Budzak, Aris Aristodemou, Louise Kerry and Eloise Lockyer for their comments on the manuscript.

Competing Interests

The Authors declare that there are no competing interests associated with the manuscript.

References

- 1 Cross, G.A.M. (1975) Identification, purification and properties of clone-specific glycoprotein antigens constituting the surface coat of *Trypanosoma brucei*. *Parasitology* **71**, 393–417 <https://doi.org/10.1017/S003118200004717X>
- 2 Bulow, R., Overath, P. and Davoust, J. (1988) Rapid lateral diffusion of the variant surface glycoprotein in the coat of *Trypanosoma brucei*. *Biochemistry* **27**, 2384–2388 <https://doi.org/10.1021/bi00407a020>
- 3 Bartossek, T., Jones, N.G., Schäfer, C., Cvitković, M., Glogger, M., Mott, H.R. et al. (2017) Structural basis for the shielding function of the dynamic trypanosome variant surface glycoprotein coat. *Nat. Microbiol.* **2**, 1523–1532 <https://doi.org/10.1038/s41564-017-0013-6>
- 4 Engstler, M., Pfohl, T., Herminghaus, S., Boshart, M., Wiegertjes, G., Heddergott, N. et al. (2007) Hydrodynamic flow-mediated protein sorting on the cell surface of trypanosomes. *Cell* **131**, 505–515 <https://doi.org/10.1016/j.cell.2007.08.046>
- 5 Cheung, J.L.Y., Wand, N.V., Ooi, C.-P., Ridewood, S., Wheeler, R.J. and Rudenko, G. (2016) Blocking synthesis of the variant surface glycoprotein coat in *Trypanosoma brucei* leads to an increase in macrophage phagocytosis due to reduced clearance of surface coat antibodies. *PLoS Pathog.* **12**, e1006023 <https://doi.org/10.1371/journal.ppat.1006023>
- 6 Marcello, L. and Barry, J.D. (2007) Analysis of the VSG gene silent archive in *Trypanosoma brucei* reveals that mosaic gene expression is prominent in antigenic variation and is favored by archive substructure. *Genome Res.* **17**, 1344–1352 <https://doi.org/10.1101/gr.6421207>
- 7 Cross, G.A.M., Kim, H.-S. and Wickstead, B. (2014) Capturing the variant surface glycoprotein repertoire (the VSGnome) of *Trypanosoma brucei* Lister 427. *Mol. Biochem. Parasitol.* **195**, 59–73 <https://doi.org/10.1016/j.molbiopara.2014.06.004>
- 8 Horn, D. (2014) Antigenic variation in African trypanosomes. *Mol. Biochem. Parasitol.* **195**, 123–129 <https://doi.org/10.1016/j.molbiopara.2014.05.001>

- 9 Hall, J.P.J., Wang, H. and Barry, J.D. (2013) Mosaic VSGs and the scale of *Trypanosoma brucei* antigenic variation. *PLoS Pathog.* **9**, e1003502 <https://doi.org/10.1371/journal.ppat.1003502>
- 10 Mugnier, M.R., Cross, G.A.M. and Papavasiliou, F.N. (2015) The in vivo dynamics of antigenic variation in *Trypanosoma brucei*. *Science* **347**, 1470–1473 <https://doi.org/10.1126/science.aaa4502>
- 11 Sheader, K., Vaughan, S., Minchin, J., Hughes, K., Gull, K. and Rudenko, G. (2005) Variant surface glycoprotein RNA interference triggers a precytokinesis cell cycle arrest in African trypanosomes. *Proc. Natl Acad. Sci. U.S.A.* **102**, 8716–8721 <https://doi.org/10.1073/pnas.0501886102>
- 12 Hertz-Fowler, C., Figueiredo, L.M., Quail, M.A., Becker, M., Jackson, A., Bason, N. et al. (2008) Telomeric expression sites are highly conserved in *Trypanosoma brucei*. *PLoS ONE* **3**, e3527 <https://doi.org/10.1371/journal.pone.0003527>
- 13 Gunzl, A., Bruderer, T., Laufer, G., Schimanski, B., Tu, L.-C., Chung, H.-M. et al. (2003) RNA polymerase I transcribes procyclin genes and variant surface glycoprotein gene expression sites in *Trypanosoma brucei*. *Eukaryot. Cell* **2**, 542–551 <https://doi.org/10.1128/EC.2.3.542-551.2003>
- 14 Navarro, M. and Gull, K. (2001) A pol I transcriptional body associated with VSG mono-allelic expression in *Trypanosoma brucei*. *Nature* **414**, 759–763 <https://doi.org/10.1038/414759a>
- 15 Kassem, A., Pays, E. and Vanhamme, L. (2014) Transcription is initiated on silent variant surface glycoprotein expression sites despite monoallelic expression in *Trypanosoma brucei*. *Proc. Natl Acad. Sci. U.S.A.* **111**, 8943–8948 <https://doi.org/10.1073/pnas.1404873111>
- 16 Vanhamme, L., Poelvoorde, P., Pays, A., Tebabi, P., Van Xong, H. and Pays, E. (2000) Differential RNA elongation controls the variant surface glycoprotein gene expression sites of *Trypanosoma brucei*. *Mol. Microbiol.* **36**, 328–340 <https://doi.org/10.1046/j.1365-2958.2000.01844.x>
- 17 Kerry, L.E., Pegg, E.E., Cameron, D.P., Budzak, J., Poortinga, G., Hannan, K.M. et al. (2017) Selective inhibition of RNA polymerase I transcription as a potential approach to treat African trypanosomiasis. *PLoS Neglected Trop. Dis.* **11**, e0005432 <https://doi.org/10.1371/journal.pntd.0005432>
- 18 Glover, L., Alsford, S., Baker, N., Turner, D.J., Sanchez-Flores, A., Hutchinson, S. et al. (2015) Genome-scale RNAi screens for high-throughput phenotyping in bloodstream-form African trypanosomes. *Nat. Protoc.* **10**, 106–133 <https://doi.org/10.1038/nprot.2015.005>
- 19 Glover, L., Hutchinson, S., Alsford, S. and Horn, D. (2016) VEX1 controls the allelic exclusion required for antigenic variation in trypanosomes. *Proc. Natl Acad. Sci. U.S.A.* **113**, 7225–7230 <https://doi.org/10.1073/pnas.1600344113>
- 20 Günzl, A., Kirkham, J.K., Nguyen, T.N., Badjatia, N. and Park, S.H. (2015) Mono-allelic VSG expression by RNA polymerase I in *Trypanosoma brucei*: expression site control from both ends? *Gene* **556**, 68–73 <https://doi.org/10.1016/j.gene.2014.09.047>
- 21 Chaves, I., Rudenko, G., Dirks-Mulder, A., Cross, M. and Borst, P. (1999) Control of variant surface glycoprotein gene-expression sites in *Trypanosoma brucei*. *EMBO J.* **18**, 4846–4855 <https://doi.org/10.1093/emboj/18.17.4846>
- 22 Berriman, M., Ghedin, E., Hertz-Fowler, C., Blandin, G., Renauld, H., Bartholomeu, D.C. et al. (2005) The genome of the African trypanosome *Trypanosoma brucei*. *Science* **309**, 416–422 <https://doi.org/10.1126/science.1112642>
- 23 Maree, J.P., Povelones, M.L., Clark, D.J., Rudenko, G. and Patterson, H.-G. (2017) Well-positioned nucleosomes punctuate polycistronic pol II transcription units and flank silent VSG gene arrays in *Trypanosoma brucei*. *Epigenet. Chromatin* **10**, 14 <https://doi.org/10.1186/s13072-017-0121-9>
- 24 Wedel, C., Forstner, K.U., Derr, R. and Siegel, T.N. (2017) GT-rich promoters can drive RNA pol II transcription and deposition of H2A.Z in African trypanosomes. *EMBO J.* **36**, 2581–2594 <https://doi.org/10.15252/embj.201695323>
- 25 Siegel, T.N., Hekstra, D.R., Kemp, L.E., Figueiredo, L.M., Lowell, J.E., Fenyo, D. et al. (2009) Four histone variants mark the boundaries of polycistronic transcription units in *Trypanosoma brucei*. *Genes Dev.* **23**, 1063–1076 <https://doi.org/10.1101/gad.1790409>
- 26 Stanne, T.M. and Rudenko, G. (2010) Active VSG expression sites in *Trypanosoma brucei* are depleted of nucleosomes. *Eukaryot. Cell* **9**, 136–147 <https://doi.org/10.1128/EC.00281-09>
- 27 Figueiredo, L.M. and Cross, G.A.M. (2010) Nucleosomes are depleted at the VSG expression site transcribed by RNA polymerase I in African trypanosomes. *Eukaryot. Cell* **9**, 148–154 <https://doi.org/10.1128/EC.00282-09>
- 28 Povelones, M.L., Gluenz, E., Dembek, M., Gull, K. and Rudenko, G. (2012) Histone H1 plays a role in heterochromatin formation and VSG expression site silencing in *Trypanosoma brucei*. *PLoS Pathog.* **8**, e1003010 <https://doi.org/10.1371/journal.ppat.1003010>
- 29 Merz, K., Hondele, M., Goetze, H., Gmelch, K., Stoekli, U. and Griesenbeck, J. (2008) Actively transcribed rRNA genes in *S. cerevisiae* are organized in a specialized chromatin associated with the high-mobility group protein Hmo1 and are largely devoid of histone molecules. *Genes Dev.* **22**, 1190–1204 <https://doi.org/10.1101/gad.466908>
- 30 Narayanan, M.S. and Rudenko, G. (2013) TDP1 is an HMG chromatin protein facilitating RNA polymerase I transcription in African trypanosomes. *Nucleic Acids Res.* **41**, 2981–2992 <https://doi.org/10.1093/nar/gks1469>
- 31 Hall, D.B., Wade, J.T. and Struhl, K. (2006) An HMG protein, Hmo1, associates with promoters of many ribosomal protein genes and throughout the rRNA gene locus in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **26**, 3672–3679 <https://doi.org/10.1128/MCB.26.9.3672-3679.2006>
- 32 Panday, A. and Grove, A. (2017) Yeast HMO1: linker histone reinvented. *Microbiol. Mol. Biol. Rev.* **81**, e00037-16 <https://doi.org/10.1128/MMBR.00037-16>
- 33 Aresta-Branco, F., Pimenta, S. and Figueiredo, L.M. (2016) A transcription-independent epigenetic mechanism is associated with antigenic switching in *Trypanosoma brucei*. *Nucleic Acids Res.* **44**, 3131–3146 <https://doi.org/10.1093/nar/gkv1459>
- 34 Schulz, D., Zaringhalam, M., Papavasiliou, F.N. and Kim, H.-S. (2016) Base J and H3.V regulate transcriptional termination in *Trypanosoma brucei*. *PLoS Genet.* **12**, e1005762 <https://doi.org/10.1371/journal.pgen.1005762>
- 35 Reynolds, D., Hofmeister, B.T., Cliffe, L., Alabady, M., Siegel, T.N., Schmitz, R.J. et al. (2016) Histone H3 variant regulates RNA polymerase II transcription termination and dual strand transcription of siRNA loci in *Trypanosoma brucei*. *PLoS Genet.* **12**, e1005758 <https://doi.org/10.1371/journal.pgen.1005758>
- 36 Figueiredo, L.M., Janzen, C.J. and Cross, G.A.M. (2008) A histone methyltransferase modulates antigenic variation in African trypanosomes. *PLoS Biol.* **6**, e161 <https://doi.org/10.1371/journal.pbio.0060161>
- 37 Duraisingh, M.T. and Horn, D. (2016) Epigenetic regulation of virulence gene expression in parasitic protozoa. *Cell Host Microbe* **19**, 629–640 <https://doi.org/10.1016/j.chom.2016.04.020>
- 38 Hughes, K., Wand, M., Foulston, L., Young, R., Harley, K., Terry, S. et al. (2007) A novel ISWI is involved in VSG expression site downregulation in African trypanosomes. *EMBO J.* **26**, 2400–2410 <https://doi.org/10.1038/sj.emboj.7601678>
- 39 Denninger, V., Fullbrook, A., Bessat, M., Ersfeld, K. and Rudenko, G. (2010) The FACT subunit TbSpt16 is involved in cell cycle specific control of VSG expression sites in *Trypanosoma brucei*. *Mol. Microbiol.* **78**, 459–474 <https://doi.org/10.1111/j.1365-2958.2010.07350.x>

- 40 Yang, X., Figueiredo, L.M., Espinal, A., Okubo, E. and Li, B. (2009) RAP1 is essential for silencing telomeric variant surface glycoprotein genes in *Trypanosoma brucei*. *Cell* **137**, 99–109 <https://doi.org/10.1016/j.cell.2009.01.037>
- 41 Nanavaty, V., Sandhu, R., Jehi, S.E., Pandya, U.M. and Li, B. (2017) *Trypanosoma brucei* RAP1 maintains telomere and subtelomere integrity by suppressing TERRA and telomeric RNA:DNA hybrids. *Nucleic Acids Res.* **45**, 5785–5796 <https://doi.org/10.1093/nar/gkx184>
- 42 Jehi, S.E., Li, X., Sandhu, R., Ye, F., Benmerzouga, I., Zhang, M. et al. (2014) Suppression of subtelomeric VSG switching by *Trypanosoma brucei* TRF requires its TTAGGG repeat-binding activity. *Nucleic Acids Res.* **42**, 12899–12911 <https://doi.org/10.1093/nar/gku942>
- 43 Jehi, S.E., Nanavaty, V. and Li, B. (2016) *Trypanosoma brucei* TIF2 and TRF suppress VSG switching using overlapping and independent mechanisms. *PLoS ONE* **11**, e0156746 <https://doi.org/10.1371/journal.pone.0156746>
- 44 Batram, C., Jones, N.G., Janzen, C.J., Markert, S.M. and Engstler, M. (2014) Expression site attenuation mechanistically links antigenic variation and development in *Trypanosoma brucei*. *eLife* **3**, e02324 <https://doi.org/10.7554/eLife.02324>
- 45 Zimmermann, H., Subota, I., Batram, C., Kramer, S., Janzen, C.J., Jones, N.G. et al. (2017) A quorum sensing-independent path to stumpy development in *Trypanosoma brucei*. *PLoS Pathog.* **13**, e1006324 <https://doi.org/10.1371/journal.ppat.1006324>
- 46 Janzen, C.J., Hake, S.B., Lowell, J.E. and Cross, G.A.M. (2006) Selective di- or trimethylation of histone H3 lysine 76 by two DOT1 homologs is important for cell cycle regulation in *Trypanosoma brucei*. *Mol. Cell.* **23**, 497–507 <https://doi.org/10.1016/j.molcel.2006.06.027>
- 47 Ridewood, S., Ooi, C.P., Hall, B., Trenaman, A., Wand, N.V., Sioutas, G. et al. (2017) The role of genomic location and flanking 3'UTR in the generation of functional levels of variant surface glycoprotein in *Trypanosoma brucei*. *Mol. Microbiol.* **106**, 614–634 <https://doi.org/10.1111/mmi.13838>
- 48 Amiguet-Vercher, A., Pérez-Morga, D., Pays, A., Poelvoorde, P., Van Xong, H., Tebabi, P. et al. (2004) Loss of the mono-allelic control of the VSG expression sites during the development of *Trypanosoma brucei* in the bloodstream. *Mol. Microbiol.* **51**, 1577–1588 <https://doi.org/10.1111/j.1365-2958.2003.03937.x>