


Journal Pre-proof

Bovine mastitis is a polymicrobial disease requiring a polydiagnostic approach

Angelopoulou Angeliki, Ross Holohan, Mary C. Rea, Alicja K. Warda, Colin Hill, R. Paul Ross



PII: S0958-6946(19)30168-2

DOI: <https://doi.org/10.1016/j.idairyj.2019.104539>

Reference: INDA 104539

To appear in: *International Dairy Journal*

Received Date: 27 March 2019

Revised Date: 21 July 2019

Accepted Date: 22 July 2019

Please cite this article as: Angeliki, A., Holohan, R., Rea, M.C., Warda, A.K., Hill, C., Ross, R.P., Bovine mastitis is a polymicrobial disease requiring a polydiagnostic approach, *International Dairy Journal*, <https://doi.org/10.1016/j.idairyj.2019.104539>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Published by Elsevier Ltd.

1 **Bovine mastitis is a polymicrobial disease requiring a polydiagnostic approach**

2

3

4

5

6

7

8 Angelopoulou Angeliki^{a,b}, Ross Holohan^b, Mary C. Rea^{b,c}, Alicja K. Warda^b, Colin Hill^{a,b}, R.

9 Paul Ross^{a,b,c}

10

11

12

13

14

15 ^a *School of Microbiology, University College Cork, Cork, Ireland*

16 ^b *APC Microbiome Ireland, University College Cork, Cork, Ireland*

17 ^c *Teagasc Food Research Centre, Moorepark, Fermoy, Co. Cork, Ireland*

18

19

20

21

22

23 *Corresponding author. Tel.: +353 214901322

24 *E-mail address: p.ross@ucc.ie (R. P. Ross)*

25

26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

ABSTRACT

Bovine mastitis, an inflammation of the udder, is associated with increases in milk somatic cell count usually resulting from bacterial infection. We analysed 50 mastitic milk samples via cultivation, 16S rRNA sequencing and a combination of the two (culturomics) to define the complete microbial content of the milk. Most samples contained over 10,000 cfu mL⁻¹ total bacterial counts including isolates that were haemolysin positive (n = 36). Among colonies isolated from blood agar plates, *Streptococcus uberis* was dominant (11/50) followed by *Streptococcus dysgalactiae* (6/50), *Pseudomonas* (6/50), *Enterococcus faecalis* (6/50), *Escherichia coli* (6/50), *Staphylococcus argenteus* (4/50), *Bacillus* (4/50) and *Staphylococcus aureus* (2/50). 16S rRNA profiling revealed that amplicons were dominated by *Rhodococcus*, *Staphylococcus*, *Streptococcus* and *Pseudomonas*. A higher inter-sample diversity was noted in the 16S rRNA readouts, which was not always reflected in the plating results. The combination of the two methods highlights the polymicrobial complexity of bovine mastitis.

51 1. Introduction

52

53 Mastitis is an inflammation of the cow's udder and is a disease of high frequency and
54 economic significance due to depleted milk production, discarded milk, premature culling
55 and treatment costs (Bar et al., 2008; Halasa, Huijps, Osteras, & Hogeveen, 2007; Hertl et al.,
56 2010). A large volume of milk is processed to a variety of dairy products and apart from the
57 risk of bacterial contamination, alterations in the composition of mastitic milk can negatively
58 affect the quality of these products (Merin et al., 2008). For example, it is known that the
59 somatic cell count (SCC) level negatively correlates with cheese yield due to slower
60 coagulation properties of the milk (Le Maréchal, Thiéry, Vautor, & Le Loir, 2011).

61 Mastitis can be classified into clinical or subclinical subgroups, with the latter being
62 indicated by an escalation in SCC in the absence of overt symptoms (Vanderhaeghen et al.,
63 2015). Milk is classified as being clinical or subclinical based on SCC, with a SCC of
64 200,000 cells mL⁻¹ generally being accepted as an indicator of the presence of mastitis
65 infection (IDF, 1997) and the SCC threshold for milk purchasers being 400,000 cells mL⁻¹
66 according to EU regulations (Regulation (EC) No. 853 of 2004). Furthermore, mastitis-
67 causing bacteria have been grouped as contagious or environmental based on their
68 distribution and interplay with the teat and teat duct (Smith & Hogan, 1993). The disease is
69 normally the result of bacterial intramammary infection, and the most commonly associated
70 causative agents are staphylococci, streptococci and coliforms (Bradley, Leach, Breen,
71 Green, & Green, 2007; Vanderhaeghen et al., 2015). However, up to 200 different microbial
72 species have been documented in mastitic cases. These are primarily bacteria, but can
73 include fungi or even monocellular achlorophylic algae (Cvetnić, Samardžija, Habrun,
74 Kompes, & Benić, 2016).

75 Identification of the microbe driving the disease is of critical importance for clinical
76 resolution. The gold standard method used for the characterisation of microorganisms
77 responsible for mastitis is bacterial culture. Nevertheless, restrictions of culture-dependent
78 techniques include a delay of 24 to 48 h to acquire results, and the fact that roughly 25% of
79 milk samples from clinical mastitis cases are culture negative (Taponen, Salmikivi, Simojoki,
80 Koskinen, & Pyörälä, 2009). This highlights the importance of evaluating culture-
81 independent techniques for mastitis diagnosis. It has been suggested that all mastitis
82 treatments should be evidence-based, which primarily requires the identification of the
83 mastitis-causing organism(s) (Milkproduction.com, 2007).

84 Sequencing and analysis of hypervariable regions within the 16S rRNA gene can
85 furnish comparably expeditious and cost-effective methods for appraising bacterial diversity
86 and abundance and has proved an effective tool for pathogen discovery and identification
87 (Oikonomou, Machado, Santisteban, Schukken, & Bicalho, 2012). These technologies have
88 enabled the investigation of microbial communities in milk without some of the limitations of
89 culture methods (Ganda et al., 2016; Jimenez et al., 2015; Oikonomou et al., 2012). It should
90 be noted that the resulting datasets are compositional (Gloor, Macklaim, Pawlowsky-Glahn,
91 & Egozcue, 2017), failing to provide resolution to species/strain level and do not differentiate
92 between living and dead microorganisms.

93 We employed both culture-dependent and culture-independent methods to identify the
94 major pathogenic species found in milks collected from diseased animals.

95

96 **2. Materials and methods**

97

98 *2.1. Sample collection*

99

100 Fifty mastitic milk samples were collected from 46 cows which had elevated SCC
101 ($\geq 200,000$ cells mL⁻¹) during the period of November 2016 to April 2018. Samples were
102 taken after the first streams of milk were discarded and stored below 4 °C, overnight until
103 they were further processed. Aliquots of 1 mL of fresh milk were subject to culture within 24
104 h of donation. Remaining aliquots were immediately frozen at -20 °C for subsequent DNA
105 extraction.

106

107 2.2. *Determination of SCC*

108

109 Milk samples were analysed for SCC using a Somacount 300 (Bentley Instruments,
110 Inc., Chaska, MN, USA) according to the International Dairy Federation (IDF) guidelines
111 (IDF, 1981).

112

113 2.3. *Microbiological analysis*

114

115 Clotted samples with high SCC ($>10,000,000$ cells mL⁻¹) were homogenised for thirty
116 minutes with the use of a stomacher machine (IUL Instruments, SA) whereas the remaining
117 samples were directly processed. Aliquots of milk sample, 1 mL, were mixed with 9 mL of
118 maximum recovery diluent (Oxoid, Basingstoke, UK) to make an initial 10⁻¹ dilution. Serial
119 dilutions were enumerated by the spread plate method in duplicate onto: (i) de Man, Rogosa,
120 Sharpe (MRS) agar (Oxoid) at 37 °C (pH 5.5) for 3 days in anaerobic jars (gas-pack plus
121 anaerobic system, BBL; BD Diagnostics, USA), which selects for lactobacilli; (ii) blood agar
122 base (Oxoid) supplemented with 7% (v/v) defibrinated sheep blood (Cruinn Diagnostics,
123 Ireland) at 37 °C for 48 h aerobically, which is a non-selective medium; (iii) Baird Parker
124 agar (Oxoid) supplemented with 50 mL egg yolk tellurite emulsion (Oxoid) at 37 °C for 48 h

125 aerobically, which selects for staphylococci; (iv) MacConkey agar (Oxoid) at 37 °C for 24 h
126 aerobically, which selects for enterobacteria; (v) plate count agar (Oxoid) at 30 °C for 72 h,
127 aerobically in which total mesophilic bacteria were counted. Plates were assessed for growth
128 and colony morphology characteristics and the blood agar plates were subsequently analysed
129 for haemolytic characteristics.

130

131 2.4. *Species determination by Sanger sequencing*

132

133 Colony PCR was performed on forty isolated colonies from blood agar plates and
134 forty isolated colonies from Baird Parker plates per sample based on different morphology in
135 the analysed samples (Supplementary material, Table S1). Cells were lysed in 10% Igepal
136 630 (Sigma-Aldrich, Germany) at 95 °C for 10 min. PCR was performed in a total volume of
137 25 µL using 10 µL Phusion Green Hot Start II High Fidelity PCR master mix (ThermoFisher
138 Scientific, Waltham, MA, USA), 10 µL PCR-grade water, 1.5 µL of the nonspecific primers
139 27F and 1495R (primer stocks at 0.1 ng µL⁻¹) (Sigma) and 2 µL of DNA template from lysed
140 cells. Amplification was carried out with reaction conditions as follows: initial denaturation
141 at 98 °C for 30 s, followed by 35 cycles of 98 °C for 10 s, annealing at 55 °C for 30 s and
142 elongation at 72 °C for 30 s with a final extension step at 72 °C for 10 min. Five microlitres
143 of the resulting amplicons from each reaction were electrophoresed in a 1.5% (w/v) agarose
144 gel. A GeneGenius Imaging System (Syngene, Cambridge, UK) was used for visualisation.
145 The PCR products were purified using the GeneJet Gel Extraction Kit (Thermo Fisher
146 Scientific). DNA sequencing of the forward strand was performed by Source BioScience
147 (Tramore, Ireland). The resulting sequences were used for searching sequences deposited in
148 the GenBank database using NCBI BLAST database (<http://www.ncbi.nlm.nih.gov/BLAST/>)
149 and the identity of the isolates was determined on the basis of the highest scores (>98%).

150

151 2.5. *DNA extraction and MiSeq sequencing*

152

153 DNA was purified from milk samples using the DNeasy PowerFood Microbial DNA
154 Isolation Kit (MoBio Laboratories, Carlsbad, USA) with slight modifications. Four mL of the
155 milk samples were centrifuged twice at $4000 \times g$ for 30 min. The top fat layer was removed
156 with a sterile cotton swab. The pellet was washed twice with sterile PBS, re-suspended in 90
157 μL of 50 mg mL^{-1} lysozyme and $25 \mu\text{L}$ of 10 KU mL^{-1} mutanolysin and incubated at $55 \text{ }^\circ\text{C}$
158 for 15 min. Subsequently, $28 \mu\text{L}$ of proteinase K was added and the pellet was incubated at
159 $55 \text{ }^\circ\text{C}$ for 15 min. The supernatant was removed after centrifugation at $13,000 \times g$ at $4 \text{ }^\circ\text{C}$.
160 The remaining steps were performed using the DNeasy PowerFood Microbial DNA Isolation
161 Kit according to manufacturer's instructions with the bead-beating time reduced to 3 min to
162 limit DNA shearing. The microbiota composition of the samples was established by amplicon
163 sequencing of a ~ 460 base pair (bp) fragment of the V3–V4 hypervariable region of the
164 bacterial 16S rRNA gene following the Illumina 16S Metagenomic Sequencing Library
165 Preparation guide. PCR amplification of V3–V4 region was performed using the forward
166 primer 5'-
167 TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCTACGGGNGGCWGCAG-3'
168 and reverse primer 5'-
169 GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGACTACHVGGGTATCTAAC-
170 3'. Each $30 \mu\text{L}$ PCR reaction contained up to $5 \text{ ng } \mu\text{L}^{-1}$ microbial genomic DNA, $6 \mu\text{L}$ of
171 each primer ($1 \mu\text{M}$) and $15 \mu\text{L}$ Phusion High-Fidelity PCR Master Mix (ThermoFisher
172 Scientific). The PCR conditions were as follows: initial denaturation for 30 s at $98 \text{ }^\circ\text{C}$; 25
173 cycles of 10 s at $98 \text{ }^\circ\text{C}$, 15 s at $55 \text{ }^\circ\text{C}$ and 20 s at $72 \text{ }^\circ\text{C}$; and $72 \text{ }^\circ\text{C}$ for 5 min for final
174 extension. The Agencourt AMPure XP system (Beckman Coulter, UK) was used to purify the

175 amplicons. A subsequent limited-cycle amplification step was performed to add multiplexing
176 indices and Illumina sequencing adapters. Amplicons were quantified, normalised and pooled
177 using the Qubit[®] dsDNA HS Assay Kit (Life Technologies, Carlsbad, California, USA).
178 Library preparation was carried out by GATC Biotech prior to 2×300 bp sequencing on the
179 Illumina MiSeq platform.

180

181 2.6. *Bioinformatic analysis of high throughput sequencing data*

182

183 Read quality was assessed using FastQC (v0.11.5)
184 (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) both before and after quality
185 filtering with Trimmomatic (v0.32) (Bolger, Lohse, & Usadel, 2014) where a Phred quality
186 threshold of 20 in a sliding window of size 4 was employed. The leading 15 bases of each
187 read was removed followed by a crop at base 270, all reads greater than or equal to 50 bases
188 in length were retained. Read pairs were merged using FLASH (v1.2.11) (Magoč &
189 Salzberg, 2011) on default settings before being processed using VSEARCH in QIIME2
190 (v2018.8) (Bolyen et al., 2018). To do this, reads were dereplicated and clustered de novo at
191 97% forming OTUs. Chimeric reads were removed in two successive steps, both de novo and
192 reference based against the ChimeraSlayer Gold database. Taxonomic classification was
193 determined using mothur (v.1.38.0, bootstrap ≥ 80) (Schloss et al., 2009) and SPINGO
194 (Allard, Ryan, Jeffery, & Claesson 2015) (v1.3, bootstrap ≥ 0.8 , similarity score ≥ 0.5) using
195 the RDP v11.4 database. Analysis was performed using the R programming language (v3.5.1)
196 (Ihaka & Gentleman, 1996) and visualised using ggplot2 (v3.1.0) (Wickham, 2009). Raw data
197 has been made publicly available in the NCBI's Sequence Read Archive under the accession
198 number: PRJNA509157.

199

200 3. Results and discussion

201

202 The aim of the study was to define the microbial composition of milks from mastitic
203 cows using both culture and high throughput sequencing approaches. Fifty bovine milk
204 samples with elevated SCC were analysed in this study, fluctuating between 221,000 and
205 $>10,000,000$ cells mL^{-1} (Table 1). Based on microbiological culturing, the majority of the
206 samples contained isolates with haemolytic patterns with α -haemolysis being dominant (70%
207 of samples). β -Haemolytic bacteria were also detected in 40% of samples, while γ -haemolysis
208 was less common and found in 20% of the samples. Total mesophilic bacteria were
209 enumerated at an average population of $5.92 \log \text{cfu mL}^{-1}$ on PCA, with four culture negative
210 samples (M29, M32, M33, and M45). Comparable mesophilic counts were demonstrated by
211 Dobranić, Kazazić, Filipović, Mikulec and Zdolec (2016), who found up to $5.39 \log \text{cfu mL}^{-1}$
212 total mesophilic counts in bovine milk samples from animals cured of mastitis. The average
213 population of presumptive lactic acid bacteria (LAB) grown on MRS was $4.30 \log \text{cfu mL}^{-1}$,
214 similar to that reported by Qiao et al. (2015) who enumerated lactobacilli using quantitative
215 PCR (qPCR) in 12 mildly subclinical milk samples and 28 severely subclinical milk samples.
216 In the mild subclinical group (SCC $<500,000$ cells mL^{-1}), the mean counts were $4.83 \log \text{cfu}$
217 mL^{-1} whereas in the severely subclinical group (SCC $>500,000$ cells mL^{-1}), the mean counts
218 for lactobacilli were $4.74 \log \text{cfu mL}^{-1}$.

219 High SCC does not always correlate with a high bacterial load. For example, while
220 samples M7, M21, M37 and M49 had a SCC in excess of 5 million and were clotted in
221 appearance their total mesophilic counts were only 2.5, 2.4, 4.2 and $3.9 \log \text{cfu mL}^{-1}$,
222 respectively. This could be due to a high load of uncultivable microorganisms in these
223 samples. The identity of microorganisms isolated from blood agar and Baird Parker plates
224 was determined by Sanger sequencing (Fig. 1, Supplementary material Table S2 and Fig. 2,

225 Supplementary material Table S3, respectively). Colonies from blood agar plates were
226 dominated by the genus *Streptococcus* (31.6%), and more specifically by *Streptococcus*
227 *uberis* (18.2%) followed by *Streptococcus dysgalactiae* (11.1%), *Streptococcus agalactiae*
228 (2.2%) and *Streptococcus urinalis* (0.1%). *St. uberis*, *St. dysgalactiae* and *St. agalactiae* are
229 well-known mastitic pathogens (Klaas & Zadoks, 2018), while *St. urinalis* belongs to a
230 subgroup of streptococci which cause urinary tract infections in humans and has not been
231 associated with bovine mastitis until now (Peltroche-Llacsahuanga, Frye, & Haase, 2012).

232 Of the isolates from blood agar plates, 18.3% were staphylococci with a relatively
233 even distribution of *Staphylococcus aureus* (3.9%), *Staphylococcus argenteus* (4%),
234 *Staphylococcus sciuri* (3.3%) and *Staphylococcus chromogenes* (2.9%). *S. aureus* is a well-
235 established mastitis pathogen both in cows and humans while *S. argenteus* is a relatively
236 novel species (Tong et al., 2015) that has been isolated from human infections (Jiang et al.,
237 2018), but not from bovine mastitis until now. The results identifying strains as *S. argenteus*
238 were inconclusive as to whether they were *S. argenteus* or *S. aureus*. *S. sciuri* and *S.*
239 *chromogenes*, both coagulase-negative staphylococci, have been previously isolated in
240 bovine mastitis studies (dos Santos et al., 2016; Hosseinzadeh & Dastmalchi Saei, 2014).
241 Members of *Escherichia*, *Enterococcus* and *Pseudomonas* were identified at comparable
242 frequencies of 9.7 %, 8.9 % and 8.2 %, respectively. All *Escherichia* isolates were
243 *Escherichia coli* while almost 98% of the *Enterococcus* belonged to *Enterococcus faecalis*. *E.*
244 *coli* has been identified as one of the major mastitis-causing pathogens (Luoreng, Wang, Mei,
245 & Zan, 2018; Vasquez et al., 2019), while enterococci have also been frequently isolated
246 from mastitic cows (Gomes, Saavedra and Henriques, 2016). Thirteen species of
247 *Pseudomonas* were detected (see Fig 1), but none was identified as *Pseudomonas aeruginosa*,
248 a microbe that is often detected in mastitis (Park et al., 2014). *Pseudomonas lactis*,
249 *Pseudomonas paralactis* (von Neubeck et al., 2017) and *Pseudomonas weihenstephanensis*

250 (von Neubeck et al., 2016) have been previously isolated from cows' milk. *Kocuria* (3.2%),
251 which is usually found in skin and mucous membranes of humans and animals and is an
252 emerging cause of infection (Kandi et al., 2016), was also detected on blood agar plates
253 together with *Trueperella pyogenes* (2.2%), which has been associated with summer mastitis
254 (Pyörälä, Jousimies-Somer, & Mero, 1992).

255 Due to the semi-selective nature of the media, isolated colonies from Baird Parker
256 agar plates were predominantly identified as *Staphylococcus*, particularly as *S. argenteus*
257 (19.5%), *S. aureus* (19.5%), *S. chromogenes* (11.8%), *Staphylococcus epidermidis* (8.3%)
258 and *Staphylococcus haemolyticus* (7.4%) (Fig. 2, Supplementary material Table S2).

259 It is broadly acknowledged that many bacteria are not cultivable on standard
260 microbiological media under standard conditions (Kamagata & Tamaki, 2005; Sekiguchi,
261 2006) and so in parallel we applied high throughput sequencing to characterise uncultivated
262 microbiota (DeLong, 2005).

263 MiSeq sequencing of 16S rRNA amplicons from bovine mastitic milk samples
264 yielded a total of 14,319,524 quality filtered reads, with a median read length of 234 ± 53
265 bases. Following quality control, we recorded an average of 286,391 reads per sample. At
266 phylum level, Actinobacteria had the highest relative abundance in 38% of the bovine
267 mastitic milk samples, while Firmicutes which were most abundant in 36% and
268 Proteobacteria in 24% (Fig. 3). At genus level *Rhodococcus* was the most abundant in most
269 samples (38%), followed by *Pseudomonas* (16%), *Streptococcus* (12%) and *Staphylococcus*
270 (8%) (Fig. 4, Supplementary material Table S4).

271 The culture-based and metagenomics approaches displayed considerable divergence
272 in their output. For example, approximately 50% of Miseq samples had high levels of
273 *Rhodococcus* while three samples had high levels of *Acinetobacter*, but we did not detect
274 either genus using culture dependent methods (discussed in detail below). Based on the

275 results of both methods, we categorised our samples in four groups: Group 1 (M7, M9, M10,
276 M13, M15, M22, M25, M30, M34 and M40) consisted of ten samples for which both 16S
277 rRNA sequencing and 16S Sanger sequencing from blood agar plates resulted in the detection
278 of the same dominant genus. More specifically, samples M7, M9 and M13 were dominated
279 by *Staphylococcus* whereas in samples M10, M15 and M25, the main genus detected was
280 *Streptococcus*. Sample M22 and M40 were dominated by *Pseudomonas*, sample M34 was
281 dominated by *Trueperella*, and sample M30 was dominated by *Escherichia*. However, this
282 was not the case for the other samples. Group 2 consisted of 17 samples (M1, M11, M14,
283 M23, M24, M26, M27, M31, M32, M38, M39, M44–M46 and M48–M50) displaying few
284 similarities between the two data sets while the 18 samples in Group 3 displayed no
285 similarities (M2–M6, M8, M16, M18–M21, M28, M29, M33, M36, M41, M42 and M47).
286 Finally, Group 4 (M12, M17, M34, M35 and M37) comprised of 5 samples that did not give
287 rise to colonies on blood agar plates.

288 Of 36 genera detected in the bovine mastitic milk samples by 16S rRNA analysis,
289 only 8 were found in the culture-dependent analysis; namely *Staphylococcus*, *Bacillus*,
290 *Carnobacterium*, *Escherichia/Shigella*, *Enterococcus*, *Streptococcus*, *Trueperella* and
291 *Pseudomonas*. Moreover, from the culture-based approach we detected *Barnesiella* sp.,
292 *Kocuria*, *Microbacterium* and *Raoultella* sp., which were not detected in the 16S rRNA
293 analysis. Aerococci were detected in higher percentages in Blood agar plates than via
294 sequencing. It should be emphasised that 16S rRNA profiling only provides relative
295 abundances and is not quantitative, albeit that *Corynebacterium* was found in similar relative
296 abundances using the two methods. Furthermore, no colonies were obtained from samples
297 M7, M11 and M24 on Baird Parker plates, even though staphylococci were found on blood
298 agar plates for M7 and M11. This may demonstrate differences in nutritional requirements.

299 16S rRNA profiling indicated relative abundances for *Staphylococcus* at 83.81% (M7),
300 0.01% (M11) and 5.27% (M24).

301 *Pseudomonas* was detected in 6 samples (M1, M4, M5, M22, M41 and M47) by both
302 culture and Sanger Sequencing, while it was predominant in the 16S profiling in 9 other
303 samples (M18, M22, M31, M39, M40, M43, M44, M46 and M48). The detected differences
304 could be due to the fact that *Pseudomonas* either could not grow in the culture conditions
305 used or that it was there in different amounts, albeit 16S rRNA profiling only provides
306 relative abundances and does not differentiate between viable and non-viable bacteria.
307 *Pseudomonas* has been found in previous studies in raw milk (von Neubeck et al., 2015),
308 bulk tank milk (Rodrigues, Lima, Canniatti-Brazaca, & Bicalho, 2017) and is a member of
309 the healthy core microbiome in human milk (Murphy et al., 2017). Nevertheless, individual
310 cases or sporadic outbreaks of mastitis may be caused by *Pseudomonas* sp., *T. pyogenes*,
311 *Serratia* sp., or other unusual pathogens (Harmon, 1994). *Pseudomonas* has also been
312 associated with water contamination, including purified water systems (Ryan, Pembroke, &
313 Adley, 2011; Kuehn et al., 2013). Water could be a significant source of microbial
314 contamination considering that modern milking practices depend heavily on water for
315 cleaning milking units. Indeed, mastitis caused by *P. aeruginosa* has been previously linked
316 with contamination of water systems and teat disinfectants in the milking parlour (Kirk &
317 Bartlett, 1984).

318 Eighteen of fifty samples were dominated by *Rhodococcus* according to 16S rRNA
319 profiling. In particular, 8 samples (M5, M6, M16, M21, M23, M29, M36 and M37) had over
320 80% *Rhodococcus* and were accompanied by low total mesophilic counts (up to 5 log cfu
321 mL⁻¹). *Rhodococcus* was not detected from the colonies grown on Blood agar plates and it is
322 likely that the species detected by the 16S rRNA sequencing were either anaerobic or
323 couldn't grow under the conditions used in this study. *Rhodococcus* was previously

324 misidentified in bovine mastitis milk samples as *Corynebacterium bovis* (Watts, Lowery,
325 Teel, & Rossbach, 2000) while *Rhodococcus equi* was identified as the causative agent in an
326 immunocompromised woman with granulomatous mastitis (Nath, Mathew, Mohan, & Anila,
327 2013). One study has identified *Rhodococcus* sp. as a causative agent in 4 out of 65 paired
328 milk samples, collected from mastitic and healthy quarters of diseased dairy cows (Oultram,
329 Ganda, Boulding, Bicalho, & Oikonomou, 2017).

330 Based on 16S rRNA profiling, 5 samples (M4, M10, M14, M20 and M47) were
331 dominated by streptococci (ranging from 79.93 to 99.68%), however, three of those samples
332 (M4, M20 and M47) were negative for streptococci on blood agar plates, possibly due to their
333 anaerobicity. In other studies, streptococci not only have been linked with high SCC milk
334 samples (Park et al., 2007; Zanardi et al., 2014; Rodrigues et al., 2017), but they were also
335 found in the healthy core microbiome of bovine (Quigley et al., 2013) and human milk
336 (Murphy et al., 2017).

337 16S rRNA profiling identified *Trueperella* in 5 samples (M27, M31, M33, M34, M48
338 and M49 at 0.1%, 4.1%, 21.4%, 49.5% and 0.7%, respectively). Sample M34 was dominated
339 by *Trueperella*, and *T. pyogenes* was the only member of this genus cultured. *T. pyogenes*
340 has been shown to act synergistically with anaerobic bacteria, namely *Fusobacterium*
341 *necrophorum*, *Bacteroides* sp, *Porphyromonas levii* in summer mastitis (Pyörälä et al, 1992).
342 Oikonomou et al. (2012) found that milk samples which were diagnosed as *T. pyogenes*
343 mastitis, had a high prevalence of *F. necrophorum* subsp. *funduliforme*. We were unable to
344 confirm this finding.

345 Samples M2, M8 and M19 were dominated by *Acinetobacter*. *Brochothrix* and
346 *Pseudomonas* were detected in all three samples by 16S rRNA profiling. In sample M8,
347 clostridia were detected while in sample M19 *Bacillus* was identified. Patel et al. (2017)
348 demonstrated that 18 healthy mothers were rich in *Acinetobacter* compared with women with

349 mastitis. Moreover, Kable et al. (2016) showed that *Acinetobacter* belongs to the core milk
350 microbiota while Quigley et al. (2013) reported that *Acinetobacter* is often found in raw milk.
351 *Acinetobacter* has the ability to adapt to various environmental conditions and several
352 emerging pathogens have been described (Gurung et al., 2013). *Acinetobacter* is also known
353 to cause spoilage (Hantsis-Zacharov & Halpern, 2007) and is rarely a primary cause of
354 mastitis (Oliver & Murinda, 2012).

355 Five samples contained *Escherichia/Shigella* at genus level (M15, M26, M30, M38
356 and M48 at 40.6%, 20.4%, 50.2%, 10.8% and 29.6% respectively) which agreed with higher
357 abundances of *Enterobacteriaceae* at family level and high abundances of Proteobacteria
358 (Madigan, Bender, Buckley, Sattley, & Stahl, 2018). These findings are in agreement with
359 previously published studies (Ganda et al., 2016; Lima, Bicalho, & Bicalho, 2018; Vasquez et
360 al., 2019) which also found the same pattern. *E. coli* was not detected in aerobic culture in
361 sample M15, while for sample M26 only 25% of the colonies were identified as *E. coli*.
362 Nonetheless, *E. coli* was the only bacterium recovered from samples M30 and M38 on blood
363 agar plates.

364 16S rRNA profiling of samples M30 and M38 revealed a large diversity of taxonomic
365 families, which is in accordance with previous characterisation of the microbiota of mastitic
366 and healthy human and bovine milk, a finding that supports the possibility of an entero-
367 mammary pathway (Perez et al., 2007; Angelopoulou et al., 2018). This is a pathway in
368 which bacteria from the gastrointestinal lumen reach the mammary gland with the help of
369 dendritic cells and CD18⁺ cells (Macpherson & Uhr, 2004; Rescigno et al., 2001). We
370 detected many families in mastitic milk that are normally present in the gastrointestinal tract
371 (GIT), such as *Ruminococcaceae*, *Clostridiaceae*, *Peptostreptococcaceae* and
372 *Lachnospiraceae*. This is consistent with findings from normally sampled quarters and
373 samples acquired via cannula (Ganda et al., 2016; Jost, Lacroix, Braegger, Rochat, &

374 Chassard, 2014; Oikonomou et al., 2014; Pang et al., 2018; Vasquez et al., 2019; Young,
375 Hine, Wallace, Callaghan, & Bibiloni, 2015). Members of these families have been
376 previously detected in samples from different anatomical parts of the bovine GIT (Lima et al.,
377 2015; Mao, Zhang, Liu, & Zhu, 2015). *Ruminococcaceae*, *Clostridiaceae*,
378 *Peptostreptococcaceae* and *Lachnospiraceae* were also identified in faecal matter from cows;
379 making it possible that their presence represents either contamination of samples or
380 translocation into the udder (Young et al., 2015).

381 It is obvious that there are limitations to both culture-based and culture-independent
382 diagnostics. Not all organisms causing infection can be cultured and/or are recovered on
383 culture while 16S rRNA compositional profiling does not provide sufficient resolution to
384 pinpoint particular species and/or strains and furthermore, cannot differentiate between live
385 and dead bacteria. Additionally, practical considerations such as price, time and labour
386 intensity will influence the choice of method. On one hand culturing bacteria is laborious, has
387 a set price per sample (effect of sample number if limited), and can take up to a week to get
388 results. On the other hand, 16S compositional sequencing is less laborious, its price per
389 sample can be greatly affected by number of samples and can be very time consuming
390 (outsourced sequencing usually takes 6 weeks, followed by data analysis). Nevertheless,
391 metagenomic approaches are increasingly applied to acquire a detailed picture of the bacteria
392 involved in the pathogenesis of mastitis.

393 Both strategies are almost certainly compromised by the low microbial biomass of
394 most milk samples (and other types of samples). Salter et al. (2014) and Glassing, Dowd,
395 Galandiuk, Davis, and Chiodini (2016) have both pointed out the potential for incorrect
396 results in low biomass samples due to contamination or other artefacts that could lead to the
397 discrepancies observed between the different studies examining the mastitic bovine milk
398 microbiota (Kuehn et al., 2013; Lima et al., 2018; Oikonomou et al., 2012; Oikonomou et al.,

399 2014; Oultram et al., 2017; Pang et al., 2018; Rodríguez et al., 2017; Vasquez et al., 2019).
400 However, we feel that this is less likely to be an issue in this study given that the majority of
401 samples have a relatively high bacterial load. Storage conditions are another factor that
402 should be taken into consideration as all samples were stored below 4 °C, overnight until they
403 were further processed. However, psychrotrophs such as *Pseudomonas* sp. could proliferate
404 at low temperature, complicating the interpretation of the finding that 16% of samples were
405 dominated by *Pseudomonas*. However, it should be emphasised that *Pseudomonas* has been
406 found in the core microbiome of healthy human milk samples (Murphy et al., 2017), and
407 therefore it remains possible for samples to be dominated by *Pseudomonas* at the time of
408 sampling. Moreover, *S. aureus*, the main causative agent of bovine mastitis in Ireland has a
409 temperature range for growth of 7–48 °C and so numbers should not increase on
410 refrigeration. As storage conditions could influence the results of both culture-dependent and
411 culture-independent methods, it is essential to minimise effects of sample handling in
412 particular collection method, time until sample processing and sample storage.

413 We identified the microbiota composition of fifty bovine mastitic milks using both
414 culture-dependent and independent approaches with 20% (n = 10) of the tested samples
415 giving similar outputs (Group 1). Group 2 displayed few similarities when 16S rRNA
416 profiling was compared with culturing. Group 3 consisted of samples for which the two
417 approaches were inconclusive and Group 4 samples gave no growth on Blood agar plates,
418 indicating that the culture conditions used were not appropriate for the bacteria present in
419 these samples. A high inter-sample diversity was noted in the 16S rRNA profiling, which
420 wasn't always reflected in plating results. Thus, we suggest that the combination of the two
421 methods sheds light into the microbial complexity of the disease and that symptoms might be
422 driven or exacerbated by more than one insulting organism.

423

424 **Acknowledgement**

425

426 This work was funded by APC Microbiome Ireland, a Centre for Science and
427 Technology (CSET) funded by the Science Foundation Ireland (SFI), grant number
428 SFI/12/RC/2273.

429

430 **References**

431

432 Allard, G., Ryan, F. J., Jeffery, I. B., & Claesson, M. J. (2015). SPINGO: a rapid species-
433 classifier for microbial amplicon sequences. *BMC Bioinformatics*, *16*, Article 324.

434 Angelopoulou, A., Field, D., Ryan, C. A., Stanton, C., Hill, C., & Ross, R. P. (2018). The
435 microbiology and treatment of human mastitis. *Medical Microbiology and Immunology*,
436 *207*, 83–94.

437 Bar, D., Tauer, L. W., Bennett, G., González, R. N., Hertl, J. A., Schukken, Y. H. et al.
438 (2008). The cost of generic clinical mastitis in dairy cows as estimated by using
439 dynamic programming. *Journal of Dairy Science*, *91*, 2205–2214.

440 Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina
441 sequence data. *Bioinformatics*, *30*, 2114–2120.

442 Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C., Al-Ghalith, G. A., et al.
443 (2018). QIIME 2: Reproducible, interactive, scalable, and extensible microbiome data
444 science. *PeerJ Preprints*, *6*, Article 27295v2.

445 Bradley, A. J., Leach, K. A., Breen, J. E., Green, L. E., & Green, M. J. (2007). Survey of the
446 incidence and aetiology of mastitis on dairy farms in England and Wales. *Veterinary*
447 *Record*, *160*, Article 253.

- 448 Cvetnić L., Samardžija M., Habrun B., Kompes G., & Benić M. (2016). Microbiological
449 monitoring of mastitis pathogens in the control of udder health in dairy cows. *Slovenian*
450 *Veterinary Research*, 53, 131–140.
- 451 DeLong, E. F. (2005). Microbial community genomics in the ocean. *Nature Reviews*
452 *Microbiology*, 3, 459–469.
- 453 Dobranić, V., Kazazić, S., Filipović, I., Mikulec, N., & Zdolec, N. (2016). Composition of
454 raw cow's milk microbiota and identification of enterococci by MALDI-TOF MS -
455 short communication. *Veterinarski Arhiv*, 86, 581–590.
- 456 dos Santos, D. C., Lange, C. C., Avellar-Costa, P., Dos Santos, K. R. N., Brito, M. A. V. P.,
457 & Giambiagi-deMarval, M. (2016). *Staphylococcus chromogenes*, a coagulase-negative
458 *Staphylococcus* species that can clot plasma. *Journal of Clinical Microbiology*, 54,
459 1372–1375.
- 460 Ganda, E. K., Bisinotto, R. S., Lima, S. F., Kronauer, K., Decter, D. H., Oikonomou, G. et al
461 (2016). Longitudinal metagenomic profiling of bovine milk to assess the impact of
462 intramammary treatment using a third-generation cephalosporin. *Scientific Reports*, 6,
463 Article 37565.
- 464 Glassing, A., Dowd, S. E., Galandiuk, S., Davis, B., & Chiodini, R. J. (2016). Inherent
465 bacterial DNA contamination of extraction and sequencing reagents may affect
466 interpretation of microbiota in low bacterial biomass samples. *Gut pathogens*, 8, Article
467 24.
- 468 Gloor, G. B., Macklaim, J. M., Pawlowsky-Glahn, V., & Egozcue, J. J. (2017). Microbiome
469 datasets are compositional: and this is not optional. *Frontiers in Microbiology*, 8,
470 Article 2224.

- 471 Gomes, F., Saavedra, M. J., & Henriques, M. (2016). Bovine mastitis disease/pathogenicity:
472 evidence of the potential role of microbial biofilms. *Pathogens and Disease*, 74, Article
473 ftw006.
- 474 Gurung, M., Nam, H. M., Tamang, M. D., Chae, M. H., Jang, G. C., Jung, S. C., et al. (2013).
475 Prevalence and antimicrobial susceptibility of *Acinetobacter* from raw bulk tank milk in
476 Korea. *Journal of Dairy Science*, 96, 1997–2002.
- 477 Halasa, T., Huijps, K., Osteras, O., & Hogeveen, H. (2007). Economic effects of bovine
478 mastitis and mastitis management: a review. *Veterinary Quarterly*, 29, 18–31.
- 479 Hantsis-Zacharov, E., & Halpern, M. (2007). Culturable psychrotrophic bacterial
480 communities in raw milk and their proteolytic and lipolytic traits. *Applied and*
481 *Environmental Microbiology*, 73, 7162–7168.
- 482 Harmon, R. J. (1994). Physiology of mastitis and factors affecting somatic cell counts.
483 *Journal of Dairy Science*, 77, 2103–2112.
- 484 Hertl, J. A., Grohn, Y. T., Leach, J. D., Bar, D., Bennett, G. J., Gonzalez, R. N. et al. (2010).
485 Effects of clinical mastitis caused by gram-positive and gram-negative bacteria and
486 other organisms on the probability of conception in New York State Holstein dairy
487 cows. *Journal of Dairy Science*, 93, 1551–1560.
- 488 Hosseinzadeh, S., & Dastmalchi Saei, H. (2014). Staphylococcal species associated with
489 bovine mastitis in the North West of Iran: emerging of coagulase-negative
490 staphylococci. *International Journal of Veterinary Science and Medicine*, 2, 27–34.
- 491 Ihaka, R., & Gentleman, R. (1996). R: A language for data analysis and graphics. *Journal of*
492 *Computational and Graphical Statistics*, 5, 299–314.
- 493 IDF. (1981). *ISO central secretariat, chemin de blandonnet 8CP 401, 1214 Vernier, Geneva,*
494 *Switzerland. Laboratory methods for use in mastitis work, Bulletin No. 132, pp 17–18.*
495 Brussels, Belgium: International Dairy Federation.

- 496 IDF. (1997). *Recommendation for presentation of mastitis-related data. Bulletin No.*
497 *321/1997.* Brussels, Belgium, International Dairy Federation.
- 498 Jiang, B., You, B., Tan, L., Yu, S., Li, H., Bai, G., et al. (2018). Clinical *Staphylococcus*
499 *argenteus* develops to small colony variants to promote persistent infection. *Frontiers*
500 *in Microbiology, 9*, 1347–1347.
- 501 Jimenez, E., de Andres, J., Manrique, M., Pareja-Tobes, P., Tobes, R., Martinez-Blanch, J. F.
502 et al. (2015). Metagenomic analysis of milk of healthy and mastitis-suffering women.
503 *Journal of Human Lactation, 31*, 406–415.
- 504 Jost, T., Lacroix, C., Braegger, C. P., Rochat, F., & Chassard, C. (2014). Vertical mother-
505 neonate transfer of maternal gut bacteria via breastfeeding. *Environmental*
506 *Microbiology, 16*, 2891–2904.
- 507 Kable, M. E., Srisengfa, Y., Laird, M., Zaragoza, J., McLeod, J., et al. (2016). The core and
508 seasonal microbiota of raw bovine milk in tanker trucks and the impact of transfer to a
509 milk processing facility. *mBio, 7*, e00836-00816.
- 510 Kamagata, Y., & Tamaki, H. (2005). Cultivation of uncultured fastidious microbes. *Microbes*
511 *and Environments, 20*, 85–91.
- 512 Kandi, V., Palange, P., Vaish, R., Bhatti, A. B., Kale, V., Kandi, M. R., et al. (2016).
513 Emerging bacterial infection: identification and clinical significance of *Kocuria*
514 species. *Cureus, 8*, Article 731.
- 515 Kirk, J. H., & Bartlett, P. C. (1984). Nonclinical *Pseudomonas aeruginosa* mastitis in a dairy
516 herd. *Journal of American Veterinary Medical Association, 184*, 671–673.
- 517 Klaas, I. C., & Zadoks, R. N. (2018). An update on environmental mastitis: challenging
518 perceptions. *Transboundary and Emerging Diseases, 65*, 166–185.

- 519 Kuehn, J. S., Gorden, P. J., Munro, D., Rong, R., Dong, Q., Plummer, P. J. et al. (2013).
520 Bacterial community profiling of milk samples as a means to understand culture-
521 negative bovine clinical mastitis. *PLoS ONE* 8, Article 61959.
- 522 Le Maréchal, C., Thiéry, R., Vautor, E., and Le Loir, Y. (2011). Mastitis impact on
523 technological properties of milk and quality of milk products—a review. *Dairy Science,*
524 *& Technology, 91, 247–282.*
- 525 Lima, F. S., Oikonomou, G., Lima, S. F., Bicalho, M. L. S., Ganda, E. K., et al. (2015).
526 Prepartum and postpartum rumen fluid microbiomes: characterization and correlation
527 with production traits in dairy cows. *Applied and Environmental Microbiology, 81,*
528 *1327–1337.*
- 529 Lima, S. F., Bicalho, M. L. d. S., & Bicalho, R. C. (2018). Evaluation of milk sample
530 fractions for characterization of milk microbiota from healthy and clinical mastitis
531 cows. *PLoS ONE, 13, Article 0193671.*
- 532 Luoreng, Z. M., Wang, X. P., Mei, C. G., & Zan, L. S. (2018). Comparison of microRNA
533 profiles between bovine mammary glands infected with *Staphylococcus aureus* and
534 *Escherichia coli*. *International Journal of Biological Sciences, 14, 87–99.*
- 535 Macpherson, A. J., & Uhr, T. (2004). Induction of protective IgA by intestinal dendritic cells
536 carrying commensal bacteria. *Science, 303, 1662–1665.*
- 537 Madigan, M. T., Bender, K. S., Buckley, D. H, Sattley, W. M., & Stahl, D. S. (2018). Biology
538 of microorganisms (15th edn, Chapt. 33). Prentice Hall Inc., Holboken, NJ, USA.
- 539 Magoc, T., & Salzberg, S. L. (2011). FLASH: fast length adjustment of short reads to
540 improve genome assemblies. *Bioinformatics, 27, 2957–2963.*
- 541 Mao, S., Zhang, M., Liu, J., & Zhu, W. (2015). Characterising the bacterial microbiota across
542 the gastrointestinal tracts of dairy cattle: membership and potential function. *Scientific*
543 *Reports, 5, Article 16116.*

- 544 Merin, U., Fleminger, G., Komanovsky, J., Silanikove, N., Bernstein, S., & Leitner, G.
545 (2008). Subclinical udder infection with *Streptococcus dysgalactiae* impairs milk
546 coagulation properties: The emerging role of proteose peptones. *Dairy Science and*
547 *Technology*, 88, 407–419.
- 548 Milkproduction.com. (2007). *Bovine mastitis treatment failure – Milkproduction.com* [online]
549 Available at: [http://www.milkproduction.com/Library/Scientific-articles/Animal-](http://www.milkproduction.com/Library/Scientific-articles/Animal-health/Bovine-mastitis-treatment-failure/)
550 [health/Bovine-mastitis-treatment-failure/](http://www.milkproduction.com/Library/Scientific-articles/Animal-health/Bovine-mastitis-treatment-failure/) [Accessed 15 March 2019].
- 551 Murphy, K., Curley, D., O’Callaghan, T. F., O’Shea, C.-A., Dempsey, E. M., O’Toole, P. W.,
552 et al. (2017). The composition of human milk and infant faecal microbiota over the first
553 three months of life: A pilot study. *Scientific Reports*, 7, Article 40597.
- 554 Nath, S. R., Mathew, A. P., Mohan, A., & Anila, K. R. (2013). *Rhodococcus equi*
555 granulomatous mastitis in an immunocompetent patient. *Journal of Medical*
556 *Microbiology*, 62, 1253–1255.
- 557 Oikonomou, G., Bicalho, M. L., Meira, E., Rossi, R. E., Foditsch, C., Machado, V. S. et al
558 (2014). Microbiota of cow’s milk; distinguishing healthy, sub-clinically and clinically
559 diseased quarters. *PLoS ONE*, 9, Article 85904.
- 560 Oikonomou, G., Machado, V. S., Santisteban, C., Schukken, Y. H., & Bicalho, R.C. (2012).
561 Microbial diversity of bovine mastitic milk as described by pyrosequencing of
562 metagenomic 16s rDNA. *PLoS ONE*, 7, Article 47671.
- 563 Oliver, S. P., & Murinda, S. E. (2012). Antimicrobial resistance of mastitis pathogens.
564 *Veterinary Clinics of North America. Food Animal Practice*, 28, 165–185.
- 565 Oultram, J. W. H., Ganda, E. K., Boulding, S. C., Bicalho, R. C., & Oikonomou, G. (2017).
566 A Metataxonomic approach could be considered for cattle clinical mastitis diagnostics.
567 *Frontiers in Veterinary Science*, 4, Article 36.

- 568 Pang, M., Xie, X., Bao, H., Sun, L., He, T., Zhao, H., et al. (2018). Insights into the bovine
569 milk microbiota in dairy farms with different incidence rates of subclinical mastitis.
570 *Frontiers in Microbiology*, 9, Article 2379.
- 571 Park, H., Hong, M., Hwang, S., Park, Y., Kwon, K., Yoon, J., et al. (2014). Characterisation
572 of *Pseudomonas aeruginosa* related to bovine mastitis. *Acta Veterinaria Hungarica*, 62,
573 1–12.
- 574 Park, Y. K., Koo, H. C., Kim, S. H., Hwang, S. Y., Jung, W. K., Kim, J. M., et al. (2007).
575 The analysis of milk components and pathogenic bacteria isolated from bovine raw
576 milk in Korea. *Journal of Dairy Science*, 90, 5405–5414.
- 577 Patel, S. H., Vaidya, Y. H., Patel, R. J., Pandit, R. J., Joshi, C. G., et al. (2017). Culture
578 independent assessment of human milk microbial community in lactational mastitis.
579 *Scientific Reports*, 7, Article 7804.
- 580 Peltroche-Llacsahuanga, H., Frye, B., & Haase, G. (2012). Isolation of *Streptococcus urinalis*
581 from a human blood culture. *Journal of Medical Microbiology*, 61, 740–742.
- 582 Perez, P. F., Dore, J., Leclerc, M., Levenez, F., Benyacoub, J., Serrant, P., et al. (2007).
583 Bacterial imprinting of the neonatal immune system: lessons from maternal cells?
584 *Pediatrics*, 119, 724–732.
- 585 Pyörälä, S., Jousimies-Somer, H., & Mero, M. (1992). Clinical, bacteriological and
586 therapeutic aspects of bovine mastitis caused by aerobic and anaerobic pathogens.
587 *British Veterinary Journal*, 148, 54–62.
- 588 Qiao, J., Kwok, L., Zhang, J., Gao, P., Zheng, Y., Guo, Z., et al. (2015). Reduction of
589 *Lactobacillus* in the milks of cows with subclinical mastitis. *Beneficial Microbes*, 6,
590 485–490.

- 591 Quigley, L., McCarthy, R., O'Sullivan, O., Beresford, T. P., Fitzgerald, G. F., Ross, R. P. et
592 al. (2013). The microbial content of raw and pasteurized cow milk as determined by
593 molecular approaches. *Journal of Dairy Science*, *96*, 4928–4937.
- 594 Rescigno, M., Urbano, M., Valzasina, B., Francolini, M., Rotta, G., Bonasio, R. et al. (2001).
595 Dendritic cells express tight junction proteins and penetrate gut epithelial monolayers to
596 sample bacteria. *Nature Immunology*, *2*, 361–367.
- 597 Rodrigues, M. X., Lima, S. F., Canniatti-Brazaca, S. G., & Bicalho, R. C. (2017). The
598 microbiome of bulk tank milk: characterization and associations with somatic cell count
599 and bacterial count. *Journal of Dairy Science*, *100*, 2536–2552.
- 600 Ryan, M. P., Pembroke, J. T., & Adley, C. C. (2011). Genotypic and phenotypic diversity of
601 *Ralstonia pickettii* and *Ralstonia insidiosa* isolates from clinical and environmental
602 sources including high-purity water. Diversity in *Ralstonia pickettii*. *BMC*
603 *Microbiology*, *11*, Article 194.
- 604 Salter, S. J., Cox, M. J., Turek, E. M., Calus, S. T., Cookson, W. O., Moffatt, M. F., et al.
605 (2014). Reagent and laboratory contamination can critically impact sequence-based
606 microbiome analyses. *BMC Biology*, *12*, Article 87.
- 607 Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., et al.
608 (2009). Introducing mothur: open-source, platform-independent, community-supported
609 software for describing and comparing microbial communities. *Applied and*
610 *Environmental Microbiology*, *75*, Article 7537.
- 611 Sekiguchi, Y. (2006). Yet-to-be cultured microorganisms relevant to methane fermentation
612 processes. *Microbes and Environments*, *21*, 1–15.
- 613 Smith, K. L., & Hogan, J. S. (1993). Environmental mastitis. *Veterinary Clinics of North*
614 *America. Food Animal Practice*, *9*, 489–498.

- 615 Taponen, S., Salmikivi, L., Simojoki, H., Koskinen, M. T., & Pyörälä, S. (2009). Real-time
616 polymerase chain reaction-based identification of bacteria in milk samples from bovine
617 clinical mastitis with no growth in conventional culturing. *Journal of Dairy Science*, *92*,
618 2610–2617.
- 619 Tong, S. Y., Schaumburg, F., Ellington, M. J., Corander, J., Pichon, B., Leendertz, F., et al.
620 (2015). Novel staphylococcal species that form part of a *Staphylococcus aureus*-related
621 complex: the non-pigmented *Staphylococcus argenteus* sp. nov. and the non-human
622 primate-associated *Staphylococcus schweitzeri* sp. nov. *International Journal of*
623 *Systemic and Evolutionary Microbiology*, *65*, 15–22.
- 624 Vanderhaeghen, W., Piepers, S., Leroy, F., Van Coillie, E., Haesebrouck, F., & De Vliegher,
625 S. (2015). Identification, typing, ecology and epidemiology of coagulase negative
626 staphylococci associated with ruminants. *Veterinary Journal*, *203*, 44–51.
- 627 Vasquez, A. K., Ganda, E. K., Capel, M. B., Eicker, S., Virkler, P. D., Bicalho, R. C., et al.
628 (2019). The microbiome of *Escherichia coli* and culture-negative nonsevere clinical
629 mastitis: characterization and associations with linear score and milk production.
630 *Journal of Dairy Science*, *102*, 578–594.
- 631 von Neubeck, M., Baur, C., Krewinkel, M., Stoeckel, M., Kranz, B., Stressler, T., et al.
632 (2015). Biodiversity of refrigerated raw milk microbiota and their enzymatic spoilage
633 potential. *International Journal of Food Microbiology*, *211*, 57–65.
- 634 von Neubeck, M., Huptas, C., Gluck, C., Krewinkel, M., Stoeckel, M., Stressler, T., et al.
635 (2016). *Pseudomonas helleri* sp. nov. and *Pseudomonas weihenstephanensis* sp. nov.,
636 isolated from raw cow's milk. *International Journal of Systemic and Evolutionary*
637 *Microbiology*, *66*, 1163–1173.
- 638 von Neubeck, M., Huptas, C., Gluck, C., Krewinkel, M., Stoeckel, M., Stressler, T., et al.
639 (2017). *Pseudomonas lactis* sp. nov. and *Pseudomonas paralactis* sp. nov., isolated

- 640 from bovine raw milk. *International Journal of Systemic and Evolutionary*
641 *Microbiology*, 67, 1656–1664.
- 642 Watts, J. L., Lowery, D. E., Teel, J. F., & Rossbach, S. (2000). Identification of
643 *Corynebacterium bovis* and other coryneforms isolated from bovine mammary glands.
644 *Journal of Dairy Science*, 83, 2373–2379.
- 645 Wickham, H. (2009). *ggplot2: Elegant graphics for data analysis*. (1st edn, Chapt. 10). New
646 York NY, USA: Springer-Verlag.
- 647 Young, W., Hine, B. C., Wallace, O. A., Callaghan, M., & Bibiloni, R. (2015). Transfer of
648 intestinal bacterial components to mammary secretions in the cow. *PeerJ Preprints*, 3,
649 Article 888.
- 650 Zanardi, G., Caminiti, A., Delle Donne, G., Moroni, P., Santi, A., Galletti, G., et al. (2014).
651 Short communication: comparing real-time PCR and bacteriological cultures for
652 *Streptococcus agalactiae* and *Staphylococcus aureus* in bulk-tank milk samples.
653 *Journal of Dairy Science*, 97, 5592–5598.

Table 1Heat map of the microbial load in fifty bovine mastitic milk samples. ^a

Samples	SCC	Blood agar			McC	BP	PCA	MRS
		α - haemolysis	β - haemolysis	γ - haemolysis				
M1	>10,000,000	4.6			4.6	5.5	6.0	4.4
M2	>10,000,000	5.0				4.9	5.4	5.0
M3	>10,000,000	5.4			4.9	2.0	5.6	5.3
M4	3,343,000	2.8			1.8	1.5	3.7	1.9
M5	324,000	2.5			1.6	1.0	4.2	2.6
M6	3,917,000			2.0	2.0		4.0	3.0
M7	5,707,000		1.7	1.5	2.4		2.5	2.8
M8	346,000	3.5			2.5	3.0	3.8	2.5
M9	700,000		2.0		4.4	1.5	3.5	2.8
M10	9,422,000	5.0			4.2	3.8	4.3	5.7
M11	8,115,000	5.7					5.2	
M12	644,000				5.3	5.0	5.2	7.2
M13	221,000	5.2		1.0	4.6	5.5	6.3	3.9
M14	4,330,000	3.0	1.0	1.2	1.4	1.5	7.4	6.6
M15	2,502,000	3.1		4.5	4.1		4.6	3.5
M16	9,999,000	0.5	2.0	4.4	0.6	5.0	5.0	1.9
M17	809,000				3.1	5.2	6.3	4.0
M18	1,247,000	8.3			8.3	5.5	9.5	1.8
M19	2,607,000	9.0			7.3	4.6	8.3	5.0
M20	>10,000,000	6.5			5.2	4.1	9.5	4.0
M21	>10,000,000	2.8	2.3		1.0	2.4	4.9	
M22	>10,000,000	8.1					7.7	
M23	>10,000,000	4.5			4.5		4.1	
M24	>10,000,000	7.1			4.0		6.8	
M25	>10,000,000	8.9			3.7	6.1	8.9	
M26	>10,000,000	5.6	5.4		5.5	4.5	5.2	
M27	>10,000,000	5.0	4.6		4.4	4.5	5.9	
M28	>10,000,000	6.0		5.1	4.6	3.0	6.1	
M29	592,000			3.0		3.0		
M30	>10,000,000	5.9			4.9		5.7	
M31	>10,000,000			7.2	6.9		7.9	6.4
M32	>10,000,000	8.8			4.5	3.5		5.1
M33	8,181,000		6.3			3.0		
M34	>10,000,000		5.8				6.1	3.3
M35	934,000	4.6			4.7	5.3	4.1	3.6
M36	1,061,000		4.1		4.6	4.1	4.3	4.1
M37	>10,000,000				3.1		4.2	4.4
M38	>10,000,000		4.7		4.4		4.5	4.3
M39	>10,000,000	5.6	4.5		6.3	5.4	6.3	4.1
M40	>10,000,000	7.2	6.2		7.5	5.3	6.8	4.6
M41	>10,000,000	6.1		8.2	7.7	5.9	6.9	5.2
M42	>10,000,000	7.1	7.0		7.3	7.3	7.0	5.1
M43	>10,000,000				5.8	6.7	7.5	6.4
M44	>10,000,000	6.4	5.7		7.5	2.0	7.1	3.7
M45	>10,000,000	6.4	6.5		5.8	2.2		5.0
M46	>10,000,000		8.1		2.6		7.3	5.5
M47	>10,000,000	4.6	3.3		4.8	3.9	5.6	
M48	>10,000,000	6.2			5.6	2.2	6.8	3.8
M49	>10,000,000		3.8		4.0	4.2	3.9	
M50	>10,000,000	5.6	6.8			2.0	7.1	6.2

^a SCC are also shown. Colour intensity corresponds to the microbial load on various media. Red represents bacterial numbers approaching $10 \log \text{cfu mL}^{-1}$ and white represents absence of microbial growth in the tested media.

Figure legends

Fig. 1. Average species identification on blood agar from fifty mastitic milk samples. Inner circle depicts genus and outer circle indicates species. Results depicting *S. argenteus* were inconclusive as to whether it was *S. argenteus* or *S. aureus*.

Fig. 2. Average species identification on Baird Parker agar plates from fifty mastitic milk samples. Inner circle depicts genus and outer circle indicates species. Results depicting *S. argenteus* were inconclusive as to whether it was *S. argenteus* or *S. aureus*.

Fig. 3. Phylum level assignments of average relative abundances of the microbiota in fifty bovine mastitic milk samples. Phyla with abundances below 1% are grouped as “Other”. Results depicting *S. argenteus* were inconclusive whether it is *S. argenteus* or *S. aureus*.

Fig. 4. Genus level assignments of average relative abundances of the microbiota in bovine mastitic milk samples. Bacterial genera under 1% are grouped as “Other”. The samples are represented in 4 groups based on level of similarity between genus level assignments of relative abundances and results from 16S Sanger sequencing on blood agar plates. Group 1 is comprised of samples where in both cases the dominant genus is identified. Group 2 display some similarities, group 3 have no similarities between the 2 data sets and group 4 did not show any growth on blood agar plates.

