

Thalassas: An International Journal of Marine Sciences

Additional records of Tripletail Lobotes surinamensis (Bloch, 1790), from the eastern Mediterranean --Manuscript Draft--

Manuscript Number:	THAL-D-20-00072R2
Full Title:	Additional records of Tripletail Lobotes surinamensis (Bloch, 1790), from the eastern Mediterranean
Article Type:	Original Article
Funding Information:	
Abstract:	In the Mediterranean Sea, Lobotes surinamensis (Bloch, 1790) is considered as a rare species, albeit an increasing number of individual-based sightings published in the literature. In this study, we present 32 additional records of this thermophilic species from the Greek and Cypriot waters; reinforcing the evidence that the species is becoming more common to the region. All of the records were collected as part of the citizen science project "Is it Alien to you? Share it!!!" by a participatory process involving fishers and taxonomic experts. This work highlights the important role of citizen science as a tool for public engagement and for monitoring species distribution in a changing environment, like the Mediterranean basin.
Corresponding Author:	Vasileios Minasidis iSea, Environmental Organisation for the Preservation of the Aquatic Ecosystems GREECE
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	iSea, Environmental Organisation for the Preservation of the Aquatic Ecosystems
Corresponding Author's Secondary Institution:	
First Author:	Vasileios Minasidis
First Author Secondary Information:	
Order of Authors:	Vasileios Minasidis Nikolaos Doumpas Periklis Kleitou Roxani Naasan Aga Spyridopoulou Paschalis Papadamakis Ioannis Giovos
Order of Authors Secondary Information:	
Author Comments:	
Response to Reviewers:	Reviewer #1: Dear authors, the manuscript was improved and I suggest acceptance in its current form. All the best. Reviewer #2: Comment 1: The authors need to cite all the records of this species from the Mediterranean Sea, Red Sea, Arabian Sea and the Arabian Gulf area. In spite of the authors mentioned that they have included such distribution in the introduction section, I cannot see any addition of a new references being added to the introduction.

Authors' Response:

The references added

[Click here to view linked References](#)

1
2
3
4 **Additional records of Tripletail *Lobotes surinamensis* (Bloch, 1790),**
5
6 **from the eastern Mediterranean**
7

8
9 Vasileios Minasidis^{1*}, Nikolaos Doumpas¹, Periklis Kleitou^{1,2}, Roxani Naasan Aga
10
11 Spryridopoulou¹, Paschalis Papadamakis³, Ioannis Giovos^{1,2}
12

13 ¹*Sea, Environmental Organisation for the Preservation of the Aquatic Ecosystems, Ochi Av. 11, 55438*
14 *Agios Paulos, Thessaloniki, Greece*
15

16
17 ²*Marine and Environmental Research (MER) Lab Ltd., 202 Amathountos Av., Marina Gardens, Block B,*
18 *Offices #13-14, Parekklesia, 4533 Limassol, Cyprus*
19

20
21 ³*Fisheries Research Institute, 64007 Nea Peramos, Kavala, Greece*
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58

59 ***Corresponding author:** vasilis.minasidis@isea.com.gr, +30 2313 090696
60
61
62
63
64
65

1
2
3
4 **Abstract**
5
6

7 In the Mediterranean Sea, *Lobotes surinamensis* (Bloch, 1790) is considered as a rare species, albeit an
8 increasing number of individual-based sightings published in the literature. In this study, we present 32
9 additional records of this thermophilic species from the Greek and Cypriot waters; reinforcing the evidence
10 that the species is becoming more common to the region. All of the records were collected as part of the
11 citizen science project “Is it Alien to you? Share it!!!” by a participatory process involving fishers and
12 taxonomic experts. This work highlights the important role of citizen science as a tool for public
13 engagement and for monitoring species distribution in a changing environment, like the Mediterranean
14 basin.
15
16
17
18
19
20
21
22
23

24 **Keywords: Lobotidae, Eastern Mediterranean Sea, Citizen science, Monitoring, Rare species**
25
26

27 **Introduction**
28
29

30 The Atlantic tripletail or tripletail, *Lobotes surinamensis* (Bloch, 1790), is a marine species found in
31 tropical and subtropical waters of all oceans; in the western Atlantic from New England southward to
32 Argentina (Carpenter 2003) and Falkland Islands (Carpenter and Robertson 2015), in the eastern Atlantic
33 from the Straits of Gibraltar to the Gulf of Guinea and Madeira (Carpenter and Johnson 2016), in the Indo-
34 Pacific from East Africa through all countries of Southeast Asia north to Taiwan Province of China and
35 southern Japan, northern Australia to southern Queensland, New Guinea to New Britain, and south to Fiji,
36 except eastern Pacific (Florida Museum of Natural History 2005). It is a demersal and thermophilic species
37 (Riede 2004) that lives at depths from 0 to 70 meters (Fricke et al. 2011), usually shallower than 10 meters,
38 and with preference to brackish waters (Myers 1999; Kuitert and Tono-zuka 2001). The maximum length
39 reported is 110 cm (Robins and Ray 1986), normally ranging between 40 and 80 cm (Bouhleb 1988). The
40 color of the fish is green-grey, darker in the top and lighter in the belly, with yellowish shades all over the
41 body (Heemstra 1986). Juveniles are mottled with yellowish, brownish and black blotches, while adults can
42 show a more uniform color, dark brown, greyish or blackish. Its common name “tripletail” comes from the
43 dorsal and anal fins, that are more flattened, looking like extra tails (Smith 1997). *L. surinamensis* is often
44 found swimming near the surface alone or in very small schools of two to four individuals (Breder 1949).
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 The species exhibits a well-known plant-mimetic behavior; swimming side by side with floating objects as
5
6 a survival strategy to attract unsuspecting prey, avoid predators, and probably benefit from drifting long
7
8 distances (Breder 1949; Watson 1996; Massuti and Renores 1994).
9

10
11 *Lobotes surinamensis* distribution has been geographically expanded from the Atlantic Ocean into the
12
13 Mediterranean Sea via the Strait of Gibraltar (Deidun et al. 2010). In the Mediterranean Sea, the species
14
15 was first recorded in 1875 (Doderlein 1875). Since then, *L. surinamensis* individuals have been reported
16
17 across the entire region (Palom 1991; Gücü and Bingel 1994; De Pirro et al. 1996; Riera et al. 1999;
18
19 Hemida et al. 2003; Camilleri et al., 2005; Zava et al. 2007; Deidun et al., 2010; Dulčić and Dragičević
20
21 2011; Tiralongo et al., 2016; Tunçer and Önal 2016; Tiralongo et al. 2018; Azzurro et al., 2019; Elbaraasi et
22
23 al., 2019; Licchelli and Denitto, 2020). In Greece, the first documented report of the species dates back in
24
25 1968 (Aegean Sea) (Bini 1968). Since then, few records have been reported from the Greek seas (Ondrias
26
27 1971; Roux in Whitehead et al. 1986; Fischer et al. 1987; Papaconstantinou 1988; Koutsogiannopoulos
28
29 2010), including the Aegean Sea (Economidis 1973; Economidis and Bauchot 1976) and sporadically other
30
31 areas, such as Thermaikos Gulf (Minos and Economidis 2007) and Maliakos Gulf (Kavadas and Bekas
32
33 2014). In Cyprus, *L. surinamensis* was first sighted in 2008 (Coral Bay, Paphos) and subsequently reported
34
35 from Limassol in 2015 (34°33'58.1"N, 33°01'05.1"E), and from Paphos, Peyia - Coral bay port
36
37 (34°51'22.0"N, 32°21'41.6"E), in 2016 (Dulčić et al. 2014; Kleitou and Crocetta 2016).
38
39

40
41 In this work, we utilized citizen science and social media to collect observations of tripletail from the Greek
42
43 and Cypriot marine waters (eastern Mediterranean) providing additional information on basic ecological
44
45 information of the species. Citizen science is currently emerging globally, empowering scientists by
46
47 providing massive and spatially broad data (Newman et al. 2012). The method is currently flourishing in
48
49 the Mediterranean, highly supported by the increasing use of social media and other modern technologies
50
51 like smartphones (Cardoso et al. 2017). Several projects exist that facilitate massive reporting of various
52
53 marine taxa (Giovos et al. 2019 and references within; Tiralongo et al., 2019a; Bargnesi et al. 2020 and
54
55 references within; Naasan Aga Spyridopoulou et al. 2020) and habitats (Gerovasileiou et al. 2016). Our
56
57 data highlight the value of citizen science and social media for supporting marine research in the changing
58
59 environment of the Mediterranean Sea.
60
61
62
63
64
65

1
2
3
4 **Materials and Methods**
5
6

7 In 2016, an online data repository was established by iSea, in which citizen scientists could easily upload
8 photographic material along with information on specimen size (length and/or weight), depth, number of
9 specimens, exact location, date, type of observation (freediving, underwater photography, shore-base
10 fishing, boat-based fishing, spearfishing), substrate (seagrass, rocky substrate, soft substrate) and the
11 adeptness of the reporter (recreational fisher, photographer, naturalist, etc). All pictures are checked for
12 their authenticity and originality using the automatic image recognition tool of Google. All verified pictures
13 are sent to a team of taxonomic experts that identifies the species to the lowest taxonomic level possible
14 and validates the observations before a record is uploaded in the project's database (see Giovos et al. 2019).
15
16

17
18
19
20
21
22
23
24 A group on Facebook was established back in 2017 to facilitate citizen science. The project's Facebook
25 group currently numbers more than 11,500 members, with about 5,000 actively engaged on a daily basis.
26
27
28 The vast majority of the participants are recreational fishers, followed by scuba divers, naturalists and
29 professional fishers (Giovos et al. 2019). Furthermore, the group community includes several experts on
30 marine alien species offering their expertise for the identification of the reported specimens while at the
31 same time influence and engaging public engagement with scientific issues.
32
33
34
35
36

37 **Results and Discussion**
38
39

40 In total, 27 observations of the species were collected in the context of this work between 2007-2019.
41 Specifically, 9 observations were recorded from Cyprus and 18 from Greece (Table 1; Fig. 1). From those
42 observations, 32 individuals were reported. The majority of the observers were recreational fishers
43 (45.00%), followed by scientists (22.00%), free divers (11.00%) and others (e.g. beachgoers, journalists
44 and news reporters) while most records were reported in 2018 (Table 1). About 19% of the records were
45 found in brackish waters (number 4, 5, 16, 24, 25 in Table 1). Interestingly, several records were juvenile
46 individuals exhibiting various coloration patterns.
47
48
49
50
51
52
53

54
55 The present work provides additional information for one uncommon species of fish in the Mediterranean
56 Sea. To date, the citizen science project of iSea "Is it Alien to you? Share it !!!" has gathered a vast amount
57 of information, including several records of alien and rare species from the eastern Mediterranean (Giovos
58
59
60
61
62
63
64
65

1
2
3
4 et al. 2019; Naasan Aga Spyridopoulou et al. 2020) and first records of alien and native species from the
5
6 Mediterranean (e.g. Giovos et al. 2018; Langeneck et al. 2019; Giovos et al. 2020), Greece (e.g. Tiralongo
7
8 et al. 2019b; Tsiamis and Giovos 2019) and Cyprus (e.g. Doumpas and Kleitou 2019; Kleitou et al. 2019).
9

10
11 Our data suggest that *L. surinamensis* might be more frequent than previously thought (Coll et al. 2010
12
13 estimated the mean probability of *L. surinamensis* occurrence 0.36, with 0.40 being the threshold for
14
15 frequent species) at least in the waters of Greece and Cyprus. The species is thermophilic and originally
16
17 was more abundant in the South and East Mediterranean (Akyol and Kara 2012; Bilge et al. 2016). Several
18
19 native species in the Mediterranean are currently displaying a northward expansion like *Sparisoma cretense*
20
21 (Linnaeus, 1758) (Kruschel et al. 2012) or *Alectis alexandrina* (Geoffroy Saint-Hilaire, 1817) (Naasan Aga
22
23 Spyridopoulou et al. 2020) as a result of the increasing sea surface temperatures (Volosciuk et al. 2016;
24
25 Pastor et al., 2018). It could be the case that the species is favored by the climate change as predicted for
26
27 thermophilic species in general (Moullec et al. 2019; Dimitriadis et al. 2020). Interestingly, almost half of
28
29 the observations (48.00%) were reported inside ports/harbors, while approximately 19% of records were
30
31 observed close to estuaries and brackish waters, such as the Evros River, the Strymonas River and the
32
33 Lissos River (Fig. 1; Table 1), indicating a preference of the species for protected, unexposed habitats .
34
35 Furthermore, most of the records were found in shallow waters below 10 meters depth (Table 1). It is
36
37 possible that the increasing number of the species records might be a result of the increasing number of
38
39 citizen science projects in the Mediterranean and the use of social media (Giovos et al., 2019 and all
40
41 references within).
42
43

44
45 The contribution of citizen science is becoming crucial in marine science nowadays. The proper
46
47 communication with scientists is constantly producing new data that can be used to understand the presence
48
49 and distribution of species in the Mediterranean Sea. Citizen science has the potential to address a big gap
50
51 in marine research despite a margin of error that should be taken into account since reports are provided by
52
53 non-scientists (Marshall and Pierce 2012; Katsanevakis and Moustakas 2018; Naasan Aga Spyridopoulou
54
55 et al. 2020). In a shifting climate and a changing basin, the participation and contribution of citizen
56
57 scientists in monitoring shifts in species distributions is critical (Giovos et al., 2019; Azzuro et al., 2019).
58
59
60
61
62
63
64
65

1
2
3
4 Scientists in the Mediterranean must continue monitoring the presence of the species to confirm or reject
5
6 the hypothesis of its fast range expansion due to the increasing sea surface temperatures in the basin.
7
8

9 **Acknowledgements**

10
11 We would like to express our gratitude to the recreational fishers George Aristidou, George Boukis, Elias
12
13 Karach, Yiannis Karavas, Elias Koutsoukanidis, Christos Tsoupas and Stelios Zafeirelis for sharing their
14
15 data and pictures from Greece; Dimitris Georgiou, Christoforos Panagi, Andreas Stavrou and Paris Tkdf for
16
17 Cyprus specimens. We are also grateful to researcher Aris Christidis as well as photographers Giorgos
18
19 Poullos (Greece) and Matt Smith (Cyprus) for sharing photographic evidence and pieces of information.
20
21
22
23
24
25
26
27
28
29
30
31
32

33 **References**

- 34
35
36 Akyol O, Kara A (2012) Record of the Atlantic tripletail, *Lobotes surinamensis* (Bloch, 1790) in the Bay of Izmir,
37
38 northern Aegean Sea. *J Appl Ichthyol* 28:645-646
39
40 Azzurro E, Bariche M, Bolognini L, Cerri J, Gianni F, Sbragaglia V, Souissi JB (2019) Climate change, biological invasions,
41
42 and the shifting distribution of Mediterranean fishes: A large-scale survey based on local ecological
43
44 knowledge. *Global Change Biology*. <https://doi.org/10.31230/osf.io/zgupd>
45
46 Bargnesi F, Ferretti F, Lucrezi S (2019) Opportunities from citizen science for shark conservation, with a focus on the
47
48 Mediterranean Sea. *Eur Zool J* 87(1):20-34
49
50 Bettoso N, Comisso G, Kruzic P (2016) First Record of the Tripletail *Lobotes surinamensis* (Pisces: Lobotidae) in the
51
52 Lagoon of Marano and Grado (Gulf of Trieste, Northern Adriatic Sea). *Annales Series Historia naturalis* 26:209-
53
54 212
55
56 Bilge G, Filiz H, Gülşahin A (2016) Occurrence of *Lobotes surinamensis* (Osteichthyes: Lobotidae) in the
57
58 Mediterranean: Historical and recent data. *Zoology in the Middle East*.
59
60 <https://doi.org/10.1080/09397140.2017.1269392>
61
62
63
64
65

- 1
2
3
4 Bini G (1968) Un pesce perciforme raro per i mari Italiani "Lobotes surinamensis, (Bloch 1790)". Atti della Accademia
5
6 Peloritana dei Pericolanti Classe di Scienze Fisiche, Matematiche e Naturali 14(1/2):3-7
7
8 Bouhlel M (1988) Poissons de Djibouti. Placerville (California, USA): RDA International, Inc p 416
9
10 Breder Jr CM (1949) "On the Behavior of Young Lobotes surinamensis". Copeia(4):237-242
11
12 Camilleri M, Ragonese S, Darmanin M, Rosso B (2005) The discovery of a specimen of Lobotes surinamensis off the
13
14 Maltese Islands (central Mediterranean Sea). Biologia Marina Mediterranea 12:480-483
15
16 Cardoso A, Adriaens T, Copas K, Flevaris S, Galiay P, Gervasini E, Jennings E, Josefsson M, López B, Magan J,
17
18 Marchante E, Montani E, Quintas M, Roy H, Schade S, See L, Tsiamis K, Taucer F, von Schomberg R (2017)
19
20 Citizen Science and Open Data: a model for Invasive Alien Species in Europe. Research Ideas and Outcomes.
21
22 <https://doi.org/10.3897/rio.3.e14811>
23
24 Carpenter KE (2003) Lobotidae. Tripletails. In Carpenter K.E. (ed) The living marine resources of the Western Central
25
26 Atlantic, Vol 3: Bony fishes part 2 (Opistognathidae to Molidae), sea turtles and marine mammals. FAO species
27
28 identification guide for fishery purposes, p 1505
29
30 Carpenter KE, Robertson R (2015) Lobotes surinamensis. The IUCN Red List of Threatened Species 2015:
31
32 e.T198670A16644032. <http://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T198670A16644032.en>
33
34 Carpenter KE, Johnson GD (2016) Lobotidae. Tripletails. In K.E. Carpenter and N. De Angelis (eds) The living marine
35
36 resources of the Eastern Central Atlantic. Vol 4: Bony fishes part 2 (Perciformes to Tetradontiformes) and Sea
37
38 turtles. FAO Species Identification Guide for Fishery Purposes, Rome, FAO, p 2544
39
40 Dimitriadis C, Azzurro E, Castriota L, Corsini-Foka M, Falautano M, Fournari-Konstantinidou I, Galanidi M, Giovos I,
41
42 Issaris Y, Kalimeris A, Karachle P, Katsanevakis S, Kytinou E, Zenetos A (2020) Updating the occurrences of
43
44 Pterois miles in the Mediterranean Sea, with considerations on thermal boundaries and future range
45
46 expansion. Mediterranean Marine Science 21(1):62-69. <https://doi.org/10.12681/mms.2184>
47
48 De Pirro M, Tosi G, Vanni S (1996) Terza catturanei mari italiani di Lobotes surinamensis (Bloch, 1790) (Actinopterygii,
49
50 Perciformes, Lobotidae). Atti della Societa Toscana de Scienze Naturali di Pisa, Serie B 103:113-114
51
52 Deidun A, Sammut R, Sciberras A, Vella P (2010) On the increasing occurrence of Lobotes surinamensis in Maltese
53
54 coastal waters. Aquatic Invasions 5(Supp 1):S113-S116
55
56 Doderlein P (1875) Descrizione di una specie delgenere esotico Lobotes, presa nelle acque dei contornidi Palermo.
57
58 Memoria del socio Prof. Pietro Doderlein Palermo Tipografi a Fraudentfelder, pp 13
59
60
61
62
63
64
65

- 1
2
3
4 Dulčić J, Dragičević B (2011) First record of the Atlantic tripletail, *Lobotes surinamensis* (Bloch, 1790), in the Adriatic
5
6 Sea. *J Appl Ichthyol* 27:1385-1386
7
- 8 Dulčić J, Dragičević B, Lipej L, Štifanić M (2014) Range extension of tripletail *Lobotes surinamensis* (Lobotidae) in the
9
10 Adriatic Sea. A northernmost record in the Mediterranean. *Cybiurn* 38:153-154
11
- 12 Economidis PS (1973) Catalogue de Poisons de la Grèce. *Hellenic Oceanology and Limnology* 11:421-598
13
- 14 Economidis PS, Bauchot ML (1976) Sur une collection de poissons des mers hélléniques (mers Égée et Ionienne)
15
16 déposée au Muséum national d'histoire naturelle. *Bulletin du Museum National d'Histoire Naturelle. 3e Serie.*
17
18 *Zoologie, Paris, 3e ser., No 392, Zool* 274:871-903
19
- 20 Elbaraasi H, Azzurro E, Bashir A, Elaabidi S, Elabar B, Elsilini O, Shakman E (2019) Updated checklist of bony fishes
21
22 along the Libyan coasts (Southern Mediterranean Sea). *Mediterranean Marine Science* 20(1):90-105.
23
24 <https://doi.org/10.12681/mms.15570>
25
- 26 Fischer W, Bauchot ML, Schneider MS (1987) *Fishes FAO d'identification des espèces pour les besoins de la Pêche*
27
28 (Revision 1). Zone de pêche 37. Vol 2, Vertébrés. FAO et CEE Publishers, Rome, pp 761- 1530
29
- 30 Florida Museum of Natural History (2005) Biological profiles: tripletail. Retrieved on 26 August 2005, from
31
32 www.flmnh.ufl.edu/fish/Gallery/Descript/Tripletail/Tripletail.html. Ichthyology at the Florida Museum of
33
34 Natural History: Education-Biological Profiles. FLMNH, University of Florida.
35
- 36 Fricke R, Kulbicki M, Wantiez L (2011) Checklist of the fishes of New Caledonia, and their distribution in the Southwest
37
38 Pacific Ocean (Pisces). *Stuttgarter Beiträge zur Naturkunde A, Neue Serie* 4:341-463
39
- 40 Gerovasileiou V, Arvanitidis C, Dailianis T, Dimitriadis C, Doğan A, Filiopoulou I, Gatti G, Issaris Y, Jean-Pierre F,
41
42 Koutsoubas D, Laure Thierry de Ville d'Avray, Melih Ertan Ç, Michalakis N, Panteri E, Romain D, Salomidi M,
43
44 Sini M (2016) CIGESMED for divers: Establishing a citizen science initiative for the mapping and monitoring of
45
46 coralligenous assemblages in the Mediterranean Sea. *Biodiversity Data Journal* 4:1-23.
47
48 <https://doi.org/10.3897/BDJ.4.e8692>
49
- 50 Giosos I, Crocetta F, Doumpas N, Kaminas A, Kleitou P, Langeneck J, Tiralongo F (2020) First record of the Atlantic
51
52 spadefish *Chaetodipterus faber* (Broussonet, 1782) in the Mediterranean Sea: is it a new aquarium release?
53
54 *BiolInvasions Records* 9(1):89-95
55
- 56 Giosos I, Bernardi G, Kleitou P, Marmara D, Romanidis-Kyriakidis G (2018) First records of the fish *Abudefduf*
57
58 *sexfasciatus* (Lacepède, 1801) and *Acanthurus sohal* (Forsskål, 1775) in the Mediterranean Sea. *BiolInvasions*
59
60 *Records* 7(2):205-210
61
62
63
64
65

- 1
2
3
4 Giovos I, Batjakas I, Bernardi G, Crocetta F, Doumpas N, Kalogirou S, Kampouris T, Katsanevakis S, Keramidas I, Kleitou
5
6 P, Langeneck J, Maximiadi M, Mitsou E, Poursanidis D, Romanidis-Kyriakidis G, Stoilas VO, Tiralongo F, Xentidis
7
8 N, Zenetos A (2019) Citizen-science for monitoring marine invasions and stimulating public engagement: a
9
10 case project from the eastern Mediterranean. *BioInvasions Records* 21:3707-3721
11
12 Gücü AC, Bingel F (1994) Trawlable species assemblages on the continental shelf of the northeastern Levant Sea
13
14 (Mediterranean) with an emphasis on Lessepsian migration. *Acta Adriat* 35:83-100
15
16 Heemstra PC (1986) Smiths' sea fishes. In M.M. Smith and P.C. Heemstra (eds) *Lobotidae*. Springer-Verlag, Berlin, pp
17
18 621-622
19
20 Hemida F, Capape C, Diatta Y, Golani D (2003) On the occurrence of tripletail, *Lobotes surinamensis* (Bloch, 1790)
21
22 (Osteichthyes: Lobotidae) off the coast of Algeria (southern Mediterranean). *Annales Series Historia naturalis*
23
24 13:145-148
25
26 Katsanevakis S, Acar Ü, Ammar I, Balci BA, Bekas P et al (2014) New Mediterranean Biodiversity Records (October
27
28 2014). *Mediterranean Marine Science* 15(3):675-695
29
30 Katsanevakis S, Moustakas A (2018) Uncertainty in Marine Invasion Science. *Front Mar Sci* 5:38
31
32 Kavadas S, Bekas P (2014) New record of *Lobotes surinamensis* (Bloch, 1790) from Maliakos Gulf (Central Aegean Sea,
33
34 Greece). *Mediterranean Marine Science* 15:691
35
36 Kleitou P, Crocetta F (2016) First record of the rare native fish *Lobotes surinamensis* (Bloch, 1790) in Cyprus.
37
38 *Mediterranean Marine Science* 17:813
39
40 Kleitou P, Crocetta F, Giovos I, Wolf W (2019) On the importance of citizen-science: the first record of *Goniobranchus*
41
42 *obsoletus* (Rüppell and Leuckart, 1830) from Cyprus (Mollusca: Gastropoda: Nudibranchia). *BioInvasions*
43
44 *Records* 8(2):252-257
45
46 Koutsogiannopoulos DD (2010) *The fish of Greece. A complete guide of all fish which live in Greek seas*. Athens,
47
48 Greece, pp 494
49
50 Kruschel C, Dahlke S, Schultz ST, Zubak I, (2012). New records of the parrot fish, *Sparisoma cretense*, and the cleaver
51
52 wrasse, *Xyrichtys novacula*, by visual census in the Southern Adriatic. *Annales Series Historia naturalis*
53
54 22(1):47
55
56 Kuitert RH, Tonozuka T (2001) *Pictorial guide to Indonesian reef fishes. Part 2. Fusiliers - Dragonets, Caesionidae -*
57
58 *Callionymidae*. Zoonetics, Australia, pp 304-622
59
60
61
62
63
64
65

- 1
2
3
4 Langeneck J, Boero F, Crocetta F, Doumpas N, Giovos I, Piraino S (2019) First record of the non-native jellyfish
5
6 *Chrysaora cf. achlyos* (Cnidaria: Pelagiidae) in the Mediterranean Sea. *BioInvasions Records* 8(3):608-613
7
8 Licchelli C, Denitto F (2020) First record of *Lobotes surinamensis* (Bloch, 1790) in the North Ionian Sea. *Mediterranean*
9
10 *Marine Science* 21(2):348
11
12 Marshall AD, Pierce SJ (2012) The use and abuse of photographic identification in sharks and rays. *J Fish Biol* 80:1361-
13
14 1379
15
16 Massuti E, Renones O (1994) Observations on the pelagic fish community around floating objects in the open sea off
17
18 Mallorca. *Bol Inst Esp Oceanogr* 10:81-93
19
20 Minos G, Economidis PS (2007) On the occurrence of tripletait, *Lobotes surinamensis* (Bloch, 1790) (Pisces:
21
22 *Lobotidae*), in North Aegean Sea (Greece). p. 242. In: 12th European Congress of Ichthyologists. Cavat
23
24 (Dubrovnik), Croatia, 9-13 September 2007. EIS.
25
26 Moullec F, Barrier N, Guilhaumon F, Marsaleix P, Shin YJ, Somot S (2019). An End-to-End model reveals losers and
27
28 winners in a warming Mediterranean Sea. *Frontiers in Marine Science* 6:345
29
30 Myers RF (1999) *Micronesian reef fishes: a comprehensive guide to the coral reef fishes of Micronesia*, 3rd revised
31
32 and expanded edition. Coral Graphics, Barrigada, Guam, pp 330
33
34 Naasan Aga Spyridopoulou R, Bouziotis D, Giovos I, Kalogirou S, Kleitou P, Langeneck J (2020) Filling the Gap of Data-
35
36 Limited Fish Species in the Eastern Mediterranean Sea: A Contribution by Citizen Science. *J Mar Sci Eng*
37
38 107(2):107. <https://doi.org/10.3390/jmse8020107>
39
40 Newman G, Crall A, Crowston K, Graham E, Newman S, Wiggins A (2012) The future of citizen science: emerging
41
42 technologies and shifting paradigms. *The Ecological Society of America journals. Frontiers in Ecology and the*
43
44 *Environment* 10(6):298-304. <https://doi.org/10.1890/110294>
45
46 Ondrias JC (1971) A list of the fresh and sea water fishes of Greece. *Hellenic Oceanology and Limnology* 10:23-96
47
48 Palom O (1991) Primera cita de *Lobotes surinamensis* (Bloch, 1790) (Pisces, *Lobotidae*) para la ictiofauna iberica.
49
50 *Miscellanea Zoologica* 15:240-242
51
52 Papaconstantinou C (1988) Check-list of marine fishes of Greece. *Fauna Graeciae IV*. National Centre Marine Research
53
54 & Hellenic Zoological Society Publishers, Athens, pp 257
55
56 Pastor F, Valiente JA, Palau JL (2018) Sea Surface Temperature in the Mediterranean: Trends and Spatial Patterns
57
58 (1982–2016). *Pure Appl Geophys* 175:4017–4029
59
60
61
62
63
64
65

- 1
2
3
4 Riede K (2004) Global register of migratory species - from global to regional scales. Final Report of the R&D-Projekt
5
6 808 05 081. Federal Agency for Nature Conservation, Bonn, Germany, pp 329
7
8 Riera F, Grau A, Grau AM, Pastor E, Pou S, Quetglas A (1999) Ichthyofauna associated with drifting floating objects in
9
10 the Balearic Islands (western Mediterranean). *Sci Mar* 63:229-235
11
12 Robins CR, Ray GC (1986) A field guide to Atlantic coast fishes of North America. Houghton Mifflin Company, Boston,
13
14 U.S.A., pp 354
15
16 Smith CL (1997) National Audubon Society field guide to tropical marine fishes of the Caribbean, the Gulf of Mexico,
17
18 Florida, the Bahamas, and Bermuda. Alfred A. Knopf, Inc., New York, pp 720
19
20 Tiralongo F (2016) New record of *Lobotes surinamensis* (Bloch, 1790) from Italian waters (Adriatic Sea). In: Dailianis et
21
22 al. New Mediterranean Marine Biodiversity Records. *Mediterranean Marine Science* 17(2):608-626
23
24 Tiralongo F, Coco S, Lombardo BM, Messina G (2018) On the Presence of a Well-Established Population of *Lobotes*
25
26 *surinamensis* (Bloch, 1790) in the central Mediterranean Sea. *Annales Series Historia naturalis* 28(1):31-36
27
28 Tiralongo F, Lillo AO, Tibullo D, Tondo E, Lo Martire C, D'Agnese R, Macali A, Mancini E, Giovos I, Coco S, Azzurro E
29
30 (2019a) Monitoring uncommon and non-indigenous fishes in Italian waters: one year of results for the
31
32 AlienFish project. *Regional Studies in Marine Science*. <https://doi.org/10.1016/j.rsma.2019.100606>
33
34 Tiralongo F, Crocetta F, Doumpas N, Giovos I, Kleitou P, Langeneck J (2019b) Is the mangrove red snapper *Lutjanus*
35
36 *argentimaculatus* (Forsskål, 1775) established in the eastern Mediterranean Sea? First records from Greece
37
38 through a citizen science project. *BioInvasions Records* 8(4):911-916
39
40 Tsiamis K, Giovos I (2018) First record of the green alga *Acetabularia caliculus* from Greece. *Mediterranean Marine*
41
42 *Science* 19(3):684
43
44 Tunçer S, Önal U (2016) The occurrence of the Atlantic Tripletail, *Lobotes surinamensis* (Bloch, 1790), in the Çanakkale
45
46 Strait. *Mediterranean Marine Science* 17(1):247-248
47
48 Volosciuk C, Gulev SK, Latif M, Maraun D, Semenov VA, Tilinina N (2016) Rising mediterranean sea surface
49
50 temperatures amplify extreme summer precipitation in central Europe. *Scientific reports* 6(1):1-7
51
52 Watson W (1996) *Lobotidae: tripletails*. p. 995-997. In H.G. Moser (ed.) *The early stages of fishes in the California*
53
54 *Current Region. California Cooperative Oceanic Fisheries Investigations (CalCOFI) Atlas No. 33*. Allen Press,
55
56 Inc., Lawrence, Kansas. 1505 p.
57
58 Whitehead PJP, Bauchot ML, Hureau JC, Nielsen J, Tortonese E (1986) *Fishes of the North-eastern Atlantic and the*
59
60 *Mediterranean*. UNESCO (Ed). Vol 2, pp 511-1007; Vol 3, pp 1008-1473
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Zava B, Gianguzza P, Riggio S (2007) New capture of the tripletail *Lobotes surinamensis* (Bloch, 1790) in the southern Tyrrhenian Sea (Osteichthyes: Lobotidae) (in Italian with English abstract). *Biologia Marina Mediterranea* 14:370

1
2
3
4 **Tables**
5

6 **Table 1.** Records of *Lobotes surinamensis* in the eastern Mediterranean Sea with information on number of individuals
7 (N), depth (m), area, coordinates (Latitude: Longitude), date (DD/MM/YYYY), type of observation (T.OB.)
8 (spearfishing, S; boat fishing, BF; shore fishing, SF; shore photography, SP; recreational longlining, RL; swimming
9 animal, SA; underwater photography, UP; hand gathering, HG; nets, N; land observation, LO) and links of the
10 photographic evidence of each observation.
11
12
13
14
15
16

n	N Individuals	Depth	Area	Coordinates	Date	T.OB.	Links
1	1	10	Pirgadikia, Greece	40°20'04.0"N, 23°43'10.2"E	03/07/2007	S	Photo 1
2	1	N/A	Lesvos island, Greece	39°10'04.3"N, 26°09'43.4"E	01/07/2009	BF	Photo 2
3	1	N/A	Skopelos island, Greece	39°07'19.3"N, 23°43'44.7"E	08/07/2015	SP	Photo 3
4	1	0	Argolic Gulf, Greece	37°32'08.2"N, 22°55'07.4"E	27/04/2017	SF	Photo 4
5	1	45	Strimonikos Gulf, Greece	40°45'39.3"N, 23°49'42.5"E	03/09/2017	RL	Photo 5
6	1	N/A	Skopelos island, Greece	39°07'23.2"N, 23°43'48.0"E	In 2017	SA	-
7	1	N/A	Allonisos island, Greece	39°08'39.8"N, 23°51'55.5"E	In 2017	SA	-
8	1	N/A	Lakonikos Gulf, Greece	36°30'26.2"N, 23°03'28.4"E	In 2017	SF	-
9	1	0	Chrysochou Gulf, Cyprus	35°04'16.6"N, 32°20'07.1"E	13/10/2017	UP	Photo 9
10	1	0	Ormos Panagias, Greece	40°13'57.6"N, 23°44'17.9"E	In 2017	SP	Photo 10
11	1	0	Protaras, Cyprus	35°00'00.6"N, 34°04'06.5"E	24/09/2017	HG	Photo 11
12	1	7	Chrysochou Gulf, Cyprus	35°03'08.6"N, 32°22'27.9"E	24/10/2017	UP	Photo 12
13	1	0	Protaras, Cyprus	35°03'05.5"N, 34°01'26.3"E	27/09/2017	S	Photo 13
14	1	2	Evoikos Gulf, Greece	38°20'14.4"N, 23°54'50.4"E	23/05/2018	SF	Photo 14
15	1	N/A	Protaras, Cyprus	35°00'11.5"N, 34°03'50.9"E	12/03/2018	HG	Photo 15
16	1	N/A	Delta Evrou, Greece	40°48'25.3"N, 26°01'32.3"E	09/07/2018	N	Photo 16
17	1	0	Cape Greko, Cyprus	34°59'01.3"N, 34°04'11.9"E	29/08/2018	UP	-
18	2	N/A	Skopelos island, Greece	39°07'26.0"N, 23°43'49.9"E	09/09/2018	SA	Photo 18
19	2	0.1	Ierissos Gulf, Greece	40°22'57.1"N, 23°55'53.0"E	In 2018	SP	Photo 19
20	1	0	Protaras, Cyprus	35°00'00.6"N, 34°04'06.5"E	25/08/2018	HG	Photo 20
21	1	0	Ormidia, Cyprus	34°58'43.4"N, 33°46'04.4"E	30/01/2018	HG	Photo 21
22	1	5	Zygi, Cyprus	34°42'52.9"N, 33°18'25.4"E	04/08/2018	S	Photo 22
23	1	2	Thermaikos Gulf, Greece	40°34'43.0"N, 22°56'17.8"E	01/06/2018	LO	Photo 23
24	1	0.1	Kiparrisiakos Gulf, Greece	37°39'15.9"N, 21°19'30.9"E	01/08/2018	LO	Photo 24
25	1	N/A	Thracian Sea, Greece	40°56'27.5"N, 25°19'58.4"E	08/10/2019	SF	Photo 25
26	1	1	Kastelorizo island, Greece	36°09'01.0"N, 29°35'27.0"E	13/09/2019	SF	-
27	4	0.3	Rhodes island, Greece	35°55'40.3"N, 27°51'27.0"E	22/10/2019	SA	-

57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figures

Fig. 1 Spatial distribution of the observations of the tripletail *Lobotes surinamensis* (Bloch, 1790) reported to the citizen-science project. Detailed information about each observation number is shown in Table 1

Fig. 2 Some of the reported specimens based on the photographic evidence provided by the citizen scientists to the project. Detailed information about each observation number is shown in Table 1

Conflict of Interest: The authors declare no conflict of interest.



