



On some Chebyshev type inequalities for the complex integral

SILVESTRU SEVER DRAGOMIR*

Victoria University, College of Engineering and Science, Melbourne, Australia.

University of the Witwatersrand, School of Computer Science and Applied Mathematics, Johannesburg, South Africa.

Abstract. Assume that f and g are continuous on γ , $\gamma \subset \mathbb{C}$ is a piecewise smooth path parametrized by $z(t)$, $t \in [a, b]$ from $z(a) = u$ to $z(b) = w$ with $w \neq u$, and the *complex Chebyshev functional* is defined by

$$\mathcal{D}_\gamma(f, g) := \frac{1}{w-u} \int_\gamma f(z)g(z) dz - \frac{1}{w-u} \int_\gamma f(z) dz \frac{1}{w-u} \int_\gamma g(z) dz.$$

In this paper we establish some bounds for the magnitude of the functional $\mathcal{D}_\gamma(f, g)$ under Lipschitzian assumptions for the functions f and g , and provide a complex version for the well known Chebyshev inequality.

Keywords: Complex integral, Continuous functions, Holomorphic functions, Chebyshev inequality.

MSC2010: 26D15, 26D10, 30A10, 30A86.

Sobre algunas desigualdades tipo Chebyshev para la integral compleja

Resumen. Sean f y g funciones continuas sobre γ , siendo $\gamma \subset \mathbb{C}$ un camino suave por partes parametrizado por $z(t)$, $t \in [a, b]$ con $z(a) = u$ y $z(b) = w$, $w \neq u$, y el *funcional de Chebyshev complejo* definido por

$$\mathcal{D}_\gamma(f, g) := \frac{1}{w-u} \int_\gamma f(z)g(z) dz - \frac{1}{w-u} \int_\gamma f(z) dz \frac{1}{w-u} \int_\gamma g(z) dz.$$

En este artículo establecemos algunas cotas para la magnitud del funcional $\mathcal{D}_\gamma(f, g)$ bajo condiciones de lipschitzianidad para las funciones f y g , y damos una versión compleja para la conocida desigualdad de Chebyshev.

Palabras clave: Integral compleja, funciones continuas, funciones holomórficas, desigualdad de Chebyshev.

*E-mail: sever.dragomir@vu.edu.au

Received: 21 March 2019, Accepted: 3 July 2019.

To cite this article: S.S. Dragomir, On some Chebyshev type inequalities for the complex integral, *Rev. Integr. temas mat.* 37 (2019), No. 2, 307–317. doi: 10.18273/revint.v37n2-2019006.

1. Introduction

For two Lebesgue integrable functions $f, g : [a, b] \rightarrow \mathbb{C}$, in order to compare the integral mean of the product with the product of the integral means, we consider the *Chebyshev functional* defined by

$$C(f, g) := \frac{1}{b-a} \int_a^b f(t)g(t) dt - \frac{1}{b-a} \int_a^b f(t) dt \frac{1}{b-a} \int_a^b g(t) dt.$$

In 1934, G. Grüss [17] showed that

$$|C(f, g)| \leq \frac{1}{4} (M-m)(N-n), \quad (1)$$

provided m, M, n, N are real numbers with the property that

$$-\infty < m \leq f \leq M < \infty, \quad -\infty < n \leq g \leq N < \infty \quad \text{a.e. on } [a, b]. \quad (2)$$

The constant $\frac{1}{4}$ in (1) is sharp.

Another, however less known result, even though it was obtained by Chebyshev in 1882, [8], states that

$$|C(f, g)| \leq \frac{1}{12} \|f'\|_\infty \|g'\|_\infty (b-a)^2, \quad (3)$$

provided that f', g' exist and are continuous on $[a, b]$ and $\|f'\|_\infty = \sup_{t \in [a, b]} |f'(t)|$. The constant $\frac{1}{12}$ cannot be improved in the general case.

The Chebyshev inequality (3) also holds if $f, g : [a, b] \rightarrow \mathbb{R}$ are assumed to be *absolutely continuous* and $f', g' \in L_\infty[a, b]$, while $\|f'\|_\infty = \text{essup}_{t \in [a, b]} |f'(t)|$.

For other inequality of Grüss' type see [1]-[16] and [18]-[28].

In order to extend Grüss' inequality to complex integral we need the following preparations.

Suppose γ is a smooth path parametrized by $z(t)$, $t \in [a, b]$ and f is a complex valued function which is continuous on γ . Put $z(a) = u$ and $z(b) = w$ with $u, w \in \mathbb{C}$. We define the integral of f on $\gamma_{u,w} = \gamma$ as

$$\int_{\gamma} f(z) dz = \int_{\gamma_{u,w}} f(z) dz := \int_a^b f(z(t)) z'(t) dt.$$

We observe that the actual choice of parametrization of γ does not matter.

This definition immediately extends to paths that are piecewise smooth. Suppose γ is parametrized by $z(t)$, $t \in [a, b]$, which is differentiable on the intervals $[a, c]$ and $[c, b]$; then, assuming that f is continuous on γ , we define

$$\int_{\gamma_{u,w}} f(z) dz := \int_{\gamma_{u,v}} f(z) dz + \int_{\gamma_{v,w}} f(z) dz,$$

where $v := z(c)$. This can be extended for a finite number of intervals.

We also define the integral with respect to arc-length:

$$\int_{\gamma_{u,w}} f(z) |dz| := \int_a^b f(z(t)) |z'(t)| dt,$$

and the length of the curve γ is then

$$\ell(\gamma) = \int_{\gamma_{u,w}} |dz| = \int_a^b |z'(t)| dt.$$

Let f and g be holomorphic in G , an open domain, and suppose $\gamma \subset G$ is a piecewise smooth path from $z(a) = u$ to $z(b) = w$. Then we have the *integration by parts formula*

$$\int_{\gamma_{u,w}} f(z) g'(z) dz = f(w) g(w) - f(u) g(u) - \int_{\gamma_{u,w}} f'(z) g(z) dz. \quad (4)$$

We recall also the *triangle inequality* for the complex integral, namely,

$$\left| \int_{\gamma} f(z) dz \right| \leq \int_{\gamma} |f(z)| |dz| \leq \|f\|_{\gamma,\infty} \ell(\gamma), \quad (5)$$

where $\|f\|_{\gamma,\infty} := \sup_{z \in \gamma} |f(z)|$.

We also define the p -norm with $p \geq 1$ by

$$\|f\|_{\gamma,p} := \left(\int_{\gamma} |f(z)|^p |dz| \right)^{1/p}.$$

For $p = 1$ we have

$$\|f\|_{\gamma,1} := \int_{\gamma} |f(z)| |dz|.$$

If $p, q > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$, then, by Hölder's inequality, we have

$$\|f\|_{\gamma,1} \leq [\ell(\gamma)]^{1/q} \|f\|_{\gamma,p}.$$

Suppose $\gamma \subset \mathbb{C}$ is a piecewise smooth path parametrized by $z(t)$, $t \in [a, b]$ from $z(a) = u$ to $z(b) = w$ with $w \neq u$. If f and g are continuous on γ , we consider the *complex Chebyshev functional* defined by

$$\mathcal{D}_{\gamma}(f, g) := \frac{1}{w-u} \int_{\gamma} f(z) g(z) dz - \frac{1}{w-u} \int_{\gamma} f(z) dz \frac{1}{w-u} \int_{\gamma} g(z) dz.$$

In this paper we establish some bounds for the magnitude of the functional $\mathcal{D}_{\gamma}(f, g)$ under various assumptions for the functions f and g , and provide a complex version for the Chebyshev inequality (3).

2. Chebyshev type results

We start with the following identity of interest:

Lemma 2.1. Suppose $\gamma \subset \mathbb{C}$ is a piecewise smooth path parametrized by $z(t)$, $t \in [a, b]$ from $z(a) = u$ to $z(b) = w$ with $w \neq u$. If f and g are continuous on γ , then

$$\begin{aligned} \mathcal{D}_\gamma(f, g) &= \frac{1}{2(w-u)^2} \int_\gamma \left(\int_\gamma (f(z) - f(w))(g(z) - g(w)) dw \right) dz \quad (6) \\ &= \frac{1}{2(w-u)^2} \int_\gamma \left(\int_\gamma (f(z) - f(w))(g(z) - g(w)) dz \right) dw \\ &= \frac{1}{2(w-u)^2} \int_\gamma \int_\gamma (f(z) - f(w))(g(z) - g(w)) dz dw. \end{aligned}$$

Proof. For any $z \in \gamma$ the integral $\int_\gamma (f(z) - f(w))(g(z) - g(w)) dw$ exists and

$$\begin{aligned} I(z) &:= \int_\gamma (f(z) - f(w))(g(z) - g(w)) dw \\ &= \int_\gamma (f(z)g(z) + f(w)g(w) - g(z)f(w) - f(z)g(w)) dw \\ &= f(z)g(z) \int_\gamma dw + \int_\gamma f(w)g(w) dw - g(z) \int_\gamma f(w) dw - f(z) \int_\gamma g(w) dw \\ &= (w-u)f(z)g(z) + \int_\gamma f(w)g(w) dw - g(z) \int_\gamma f(w) dw - f(z) \int_\gamma g(w) dw. \end{aligned}$$

The function $I(z)$ is also continuous on γ , then the integral $\int_\gamma I(z) dz$ exists and

$$\begin{aligned} \int_\gamma I(z) dz &= \int_\gamma \left[(w-u)f(z)g(z) + \int_\gamma f(w)g(w) dw \right. \\ &\quad \left. - g(z) \int_\gamma f(w) dw - f(z) \int_\gamma g(w) dw \right] dz \\ &= (w-u) \int_\gamma f(z)g(z) dz + (w-u) \int_\gamma f(w)g(w) dw \\ &\quad - \int_\gamma f(w) dw \int_\gamma g(z) dz - \int_\gamma g(w) dw \int_\gamma f(z) dz \\ &= 2(w-u) \int_\gamma f(z)g(z) dz - 2 \int_\gamma f(z) dz \int_\gamma g(z) dz \\ &= 2(w-u)^2 \mathcal{D}_\gamma(f, g), \end{aligned}$$

which proves the first equality in (6).

The rest follows in a similar manner and we omit the details. ✓

Suppose $\gamma \subset \mathbb{C}$ is a piecewise smooth path from $z(a) = u$ to $z(b) = w$ and $h : \gamma \rightarrow \mathbb{C}$ a continuous function on γ . Define the quantity:

$$\begin{aligned}\mathcal{P}_\gamma(h, \bar{h}) &= \frac{1}{\ell(\gamma)} \int_\gamma |h(z)|^2 |dz| - \left| \frac{1}{\ell(\gamma)} \int_\gamma h(z) |dz| \right|^2 \\ &= \frac{1}{\ell(\gamma)} \int_\gamma \left| h(v) - \frac{1}{\ell(\gamma)} \int_\gamma h(z) |dz| \right|^2 |dv| \geq 0.\end{aligned}\quad (7)$$

We say that the function $f : G \subset \mathbb{C} \rightarrow \mathbb{C}$ is L - h -Lipschitzian on the subset G if

$$|f(z) - f(w)| \leq L |h(z) - h(w)|$$

for any $z, w \in G$. If $h(z) = z$, we recapture the usual concept of L -Lipschitzian functions on G .

Theorem 2.2. Suppose $\gamma \subset \mathbb{C}$ is a piecewise smooth path parametrized by $z(t)$, $t \in [a, b]$ from $z(a) = u$ to $z(b) = w$ with $w \neq u$, $h : \gamma \rightarrow \mathbb{C}$ is continuous, f and g are L_1 , L_2 - h -Lipschitzian functions on γ ; then

$$|\mathcal{D}_\gamma(f, g)| \leq L_1 L_2 \frac{\ell^2(\gamma)}{|w - u|^2} \mathcal{P}_\gamma(h, \bar{h}). \quad (8)$$

Proof. Taking the modulus in the first equality in (6), we get

$$\begin{aligned}|\mathcal{D}_\gamma(f, g)| &= \frac{1}{2|w - u|^2} \left| \int_\gamma \left(\int_\gamma (f(z) - f(w))(g(z) - g(w)) dw \right) dz \right| \\ &\leq \frac{1}{2|w - u|^2} \int_\gamma \left| \int_\gamma (f(z) - f(w))(g(z) - g(w)) dw \right| |dz| \\ &\leq \frac{1}{2|w - u|^2} \int_\gamma \left(\int_\gamma |(f(z) - f(w))(g(z) - g(w))| |dw| \right) |dz| \\ &\leq \frac{L_1 L_2}{2|w - u|^2} \int_\gamma \left(\int_\gamma |h(z) - h(w)|^2 |dw| \right) |dz| =: A.\end{aligned}\quad (9)$$

Now, observe that

$$\begin{aligned}
& \int_{\gamma} \left(\int_{\gamma} |h(z) - h(w)|^2 |dw| \right) |dz| \\
&= \int_{\gamma} \left(\int_{\gamma} \left(|h(z)|^2 - 2\operatorname{Re} \left(h(z) \overline{h(w)} \right) + |h(w)|^2 \right) |dw| \right) |dz| \\
&= \int_{\gamma} \left(\ell(\gamma) |h(z)|^2 - 2\operatorname{Re} \left(h(z) \int_{\gamma} \overline{h(w)} |dw| \right) + \int_{\gamma} |h(w)|^2 |dw| \right) |dz| \\
&= \ell(\gamma) \int_{\gamma} |h(z)|^2 |dz| - 2\operatorname{Re} \left(\int_{\gamma} h(z) |dz| \int_{\gamma} \overline{h(w)} |dw| \right) \\
&\quad + \ell(\gamma) \int_{\gamma} |h(w)|^2 |dw| \\
&= 2\ell(\gamma) \int_{\gamma} |h(z)|^2 |dz| - 2\operatorname{Re} \left(\int_{\gamma} h(z) |dz| \overline{\left(\int_{\gamma} h(w) |dw| \right)} \right) \\
&= 2 \left[\ell(\gamma) \int_{\gamma} |h(z)|^2 |dz| - \left| \int_{\gamma} h(z) |dz| \right|^2 \right] = 2\ell^2(\gamma) \mathcal{P}_{\gamma}(h, \bar{h}).
\end{aligned} \tag{10}$$

Therefore, by (10) we get

$$A = L_1 L_2 \frac{\ell^2(\gamma)}{|w-u|^2} \mathcal{P}_{\gamma}(h, \bar{h}),$$

and by (9) we get the desired result (8). \square

Further, for $\gamma \subset \mathbb{C}$ a piecewise smooth path parametrized by $z(t)$, and by taking $h(z) = z$ in (7), we can consider the quantity

$$\begin{aligned}
\mathcal{P}_{\gamma} &:= \frac{1}{\ell(\gamma)} \int_{\gamma} |z|^2 |dz| - \left| \frac{1}{\ell(\gamma)} \int_{\gamma} z |dz| \right|^2 \\
&= \frac{1}{\ell(\gamma)} \int_{\gamma} \left| v - \frac{1}{\ell(\gamma)} \int_{\gamma} z |dz| \right|^2 |dv| \\
&= \frac{1}{2\ell^2(\gamma)} \int_{\gamma} \left(\int_{\gamma} |z-w|^2 |dw| \right) |dz| \geq 0.
\end{aligned} \tag{11}$$

Corollary 2.3. Suppose $\gamma \subset \mathbb{C}$ is a piecewise smooth path parametrized by $z(t)$, $t \in [a, b]$ from $z(a) = u$ to $z(b) = w$ with $w \neq u$ and f and g are L_1 , L_2 -Lipschitzian functions on γ ; then

$$|\mathcal{D}_{\gamma}(f, g)| \leq L_1 L_2 \frac{\ell^2(\gamma)}{|w-u|^2} \mathcal{P}_{\gamma}. \tag{12}$$

Remark 2.4. Assume that f is L -h-Lipschitzian on γ . For $g = f$ we have

$$\mathcal{D}_{\gamma}(f, f) = \frac{1}{w-u} \int_{\gamma} f^2(z) dz - \left(\frac{1}{w-u} \int_{\gamma} f(z) dz \right)^2, \tag{13}$$

and by (8) we get

$$|\mathcal{D}_\gamma(f, f)| \leq L^2 \frac{\ell^2(\gamma)}{|w-u|^2} \mathcal{P}_\gamma(h, \bar{h}). \quad (14)$$

For $g = \bar{f}$ we have

$$\mathcal{D}_\gamma(f, \bar{f}) = \frac{1}{w-u} \int_\gamma |f(z)|^2 dz - \frac{1}{w-u} \int_\gamma f(z) dz \frac{1}{w-u} \int_\gamma \overline{f(z)} dz, \quad (15)$$

and by (8) we get

$$|\mathcal{D}_\gamma(f, \bar{f})| \leq L^2 \frac{\ell^2(\gamma)}{|w-u|^2} \mathcal{P}_\gamma(h, \bar{h}). \quad (16)$$

If f is L -Lipschitzian on γ , then

$$|\mathcal{D}_\gamma(f, f)| \leq L^2 \frac{\ell^2(\gamma)}{|w-u|^2} \mathcal{P}_\gamma \quad (17)$$

and

$$|\mathcal{D}_\gamma(f, \bar{f})| \leq L^2 \frac{\ell^2(\gamma)}{|w-u|^2} \mathcal{P}_\gamma. \quad (18)$$

If the path γ is a segment $[u, w]$ connecting two distinct points u and w in \mathbb{C} , then we write $\int_\gamma f(z) dz$ as $\int_u^w f(z) dz$.

Now, if f and g are L_1, L_2 -Lipschitzian functions on $[u, w] := \{(1-t)u + tw, t \in [0, 1]\}$, then by (12) we have

$$|\mathcal{D}_\gamma(f, g)| \leq L_1 L_2 \mathcal{P}_{[u, w]},$$

where

$$\begin{aligned} \mathcal{P}_{[u, w]} &= \frac{|w-u|^2}{2|w-u|^2} \int_0^1 \left(\int_0^1 |(1-t)u + tw - (1-s)u - sw|^2 dt \right) ds \\ &= \frac{1}{2} |w-u|^2 \int_0^1 \left(\int_0^1 (t-s)^2 dt \right) ds = \frac{1}{12} |w-u|^2. \end{aligned}$$

Therefore,

$$\begin{aligned} &\left| \frac{1}{w-u} \int_\gamma f(z) g(z) dz - \frac{1}{w-u} \int_\gamma f(z) dz \frac{1}{w-u} \int_\gamma g(z) dz \right| \\ &\leq \frac{1}{12} |w-u|^2 L_1 L_2, \end{aligned} \quad (19)$$

if f and g are L_1, L_2 -Lipschitzian functions on $[u, w]$.

If f is L -Lipschitzian on $[u, w]$, then

$$\left| \frac{1}{w-u} \int_\gamma f^2(z) dz - \left(\frac{1}{w-u} \int_\gamma f(z) dz \right)^2 \right| \leq \frac{1}{12} |w-u|^2 L^2 \quad (20)$$

and

$$\begin{aligned} &\left| \frac{1}{w-u} \int_\gamma |f(z)|^2 dz - \frac{1}{w-u} \int_\gamma f(z) dz \frac{1}{w-u} \int_\gamma \overline{f(z)} dz \right| \\ &\leq \frac{1}{12} |w-u|^2 L^2. \end{aligned} \quad (21)$$

3. Examples for circular paths

Let $[a, b] \subseteq [0, 2\pi]$ and the circular path $\gamma_{[a,b],R}$ centered in 0 and with radius $R > 0$:

$$z(t) = R \exp(it) = R(\cos t + i \sin t), \quad t \in [a, b].$$

If $[a, b] = [0, \pi]$, then we get a half circle, while for $[a, b] = [0, 2\pi]$ we get the full circle.

Since

$$\begin{aligned} |e^{is} - e^{it}|^2 &= |e^{is}|^2 - 2\operatorname{Re}(e^{i(s-t)}) + |e^{it}|^2 \\ &= 2 - 2\cos(s-t) = 4\sin^2\left(\frac{s-t}{2}\right) \end{aligned}$$

for any $t, s \in \mathbb{R}$, then

$$|e^{is} - e^{it}|^r = 2^r \left| \sin\left(\frac{s-t}{2}\right) \right|^r \quad (22)$$

for any $t, s \in \mathbb{R}$ and $r > 0$. In particular,

$$|e^{is} - e^{it}| = 2 \left| \sin\left(\frac{s-t}{2}\right) \right|$$

for any $t, s \in \mathbb{R}$.

If $u = R \exp(ia)$ and $w = R \exp(ib)$, then

$$\begin{aligned} w - u &= R[\exp(ib) - \exp(ia)] = R[\cos b + i \sin b - \cos a - i \sin a] \\ &= R[\cos b - \cos a + i(\sin b - \sin a)]. \end{aligned}$$

Since

$$\cos b - \cos a = -2 \sin\left(\frac{a+b}{2}\right) \sin\left(\frac{b-a}{2}\right)$$

and

$$\sin b - \sin a = 2 \sin\left(\frac{b-a}{2}\right) \cos\left(\frac{a+b}{2}\right),$$

hence

$$\begin{aligned} w - u &= R \left[-2 \sin\left(\frac{a+b}{2}\right) \sin\left(\frac{b-a}{2}\right) + 2i \sin\left(\frac{b-a}{2}\right) \cos\left(\frac{a+b}{2}\right) \right] \\ &= 2R \sin\left(\frac{b-a}{2}\right) \left[-\sin\left(\frac{a+b}{2}\right) + i \cos\left(\frac{a+b}{2}\right) \right] \\ &= 2Ri \sin\left(\frac{b-a}{2}\right) \left[\cos\left(\frac{a+b}{2}\right) + i \sin\left(\frac{a+b}{2}\right) \right] \\ &= 2Ri \sin\left(\frac{b-a}{2}\right) \exp\left[\left(\frac{a+b}{2}\right)i\right]. \end{aligned}$$

If $\gamma = \gamma_{[a,b],R}$, then the *circular complex Chebyshev functional* is defined by

$$\begin{aligned} & \mathcal{C}_{[a,b],R}(f, g) \\ &:= \mathcal{D}_{\gamma_{[a,b],R}}(f, g) \\ &= \frac{1}{2 \sin(\frac{b-a}{2}) \exp[(\frac{a+b}{2})i]} \int_a^b f(R \exp(it)) g(R \exp(it)) \exp(it) dt \\ &\quad - \frac{1}{4 \sin^2(\frac{b-a}{2}) \exp[2(\frac{a+b}{2})i]} \\ &\quad \times \int_a^b f(R \exp(it)) \exp(it) dt \int_a^b g(R \exp(it)) \exp(it) dt. \end{aligned} \tag{23}$$

If $\gamma = \gamma_{[a,b],R}$, then

$$\mathcal{P}_\gamma := \frac{1}{2\ell^2(\gamma)} \int_\gamma \left(\int_\gamma |z-w|^2 |dw| \right) |dz| \tag{24}$$

$$\begin{aligned} &= \frac{R^4}{2R^2(b-a)^2} \int_a^b \left(\int_a^b |e^{is} - e^{it}|^2 dt \right) ds \\ &= \frac{R^4}{2R^2(b-a)^2} \int_a^b \left(\int_a^b [2 - 2 \cos(s-t)] dt \right) ds \\ &= \frac{R^2}{(b-a)^2} \int_a^b \left(\int_a^b [1 - \cos(s-t)] dt \right) ds \\ &= \frac{R^2}{(b-a)^2} \int_a^b (b-a - \sin(b-s) - \sin(s-a)) ds \\ &= \frac{R^2}{(b-a)^2} [(b-a)^2 - 1 + \cos(b-a) + \cos(b-a) - 1] \\ &= \frac{R^2}{(b-a)^2} [(b-a)^2 - 2(1 - \cos(b-a))] \\ &= \frac{R^2}{(b-a)^2} \left[(b-a)^2 - 4 \sin^2 \left(\frac{b-a}{2} \right) \right] \\ &= \frac{4R^2}{(b-a)^2} \left[\left(\frac{b-a}{2} \right)^2 - \sin^2 \left(\frac{b-a}{2} \right) \right]. \end{aligned} \tag{25}$$

We have the following result:

Proposition 3.1. *Let $\gamma_{[a,b],R}$ be a circular path centered in 0, with radius $R > 0$ and $[a, b] \subset [0, 2\pi]$. If f and g are L_1 , L_2 -Lipschitzian functions on $\gamma_{[a,b],R}$, then*

$$|\mathcal{C}_{[a,b],R}(f, g)| \leq \frac{R^2}{\sin^2(\frac{b-a}{2})} \left[\left(\frac{b-a}{2} \right)^2 - \sin^2 \left(\frac{b-a}{2} \right) \right] L_1 L_2. \tag{26}$$

Acknowledgement. The author would like to thank the anonymous referee for valuable suggestions that have been implemented in the final version of the paper.

References

- [1] Alomari M.W., “A companion of Grüss type inequality for Riemann-Stieltjes integral and applications”, *Mat. Vesnik* 66 (2014), No. 2, 202–212.
- [2] Andrica D. and Badea C., “Grüss’ inequality for positive linear functionals”, *Period. Math. Hungar.* 19 (1988), No. 2, 155–167.
- [3] Baleanu D., Purohit S.D. and Uçar F., “On Grüss type integral inequality involving the Saigo’s fractional integral operators”, *J. Comput. Anal. Appl.* 19 (2015), No. 3, 480–489.
- [4] Cerone P., “On a Čebyšev-type functional and Grüss-like bounds”, *Math. Inequal. Appl.* 9 (2006), No. 1, 87–102.
- [5] Cerone P. and Dragomir S.S., “A refinement of the Grüss inequality and applications”, *Tamkang J. Math.*, 38 (2007), No. 1, 37–49.
- [6] Cerone P. and Dragomir S.S., “Some new Ostrowski-type bounds for the Čebyšev functional and applications”, *J. Math. Inequal.* 8 (2014), No. 1, 159–170.
- [7] Cerone P., Dragomir S.S. and Roumeliotis J., “Grüss inequality in terms of Δ -seminorms and applications”, *Integral Transforms Spec. Funct.* 14 (2003), No. 3, 205–216.
- [8] Chebyshev P.L., “Sur les expressions approximatives des intégrals définis par les autres prises entre les même limites”, *Proc. Math. Soc. Charkov*, 2 (1882), 93–98.
- [9] Cheng X.L. and Sun J., “A note on the perturbed trapezoid inequality”, *JIPAM. J. Inequal. Pure Appl. Math.* 3 (2002), No. 2, Art. 29, 7 pp.
- [10] Dragomir S.S., “A generalization of Grüss’s inequality in inner product spaces and applications”, *J. Math. Anal. Appl.* 237 (1999), No. 1, 74–82.
- [11] Dragomir S.S., “A Grüss’ type integral inequality for mappings of r -Hölder’s type and applications for trapezoid formula”, *Tamkang J. Math.* 31 (2000), No. 1, 43–47.
- [12] Dragomir S.S., “Some integral inequalities of Grüss type”, *Indian J. Pure Appl. Math.* 31 (2000), No. 4, 397–415.
- [13] Dragomir S.S., “Integral Grüss inequality for mappings with values in Hilbert spaces and applications”, *J. Korean Math. Soc.* 38 (2001), No. 6, 1261–1273.
- [14] Dragomir S.S. and Fedotov I.A., “An inequality of Grüss’ type for Riemann-Stieltjes integral and applications for special means”, *Tamkang J. Math.* 29 (1998), No. 4, 287–292.
- [15] Dragomir S.S. and Gomm I., “Some integral and discrete versions of the Grüss inequality for real and complex functions and sequences”, *Tamsui Oxf. J. Math. Sci.* 19 (2003), No. 1, 67–77.
- [16] Fink A.M., “A treatise on Grüss’ inequality”, in *Analytic and Geometric Inequalities and Applications* (eds. RassiasHari T. and Srivastava M.), Math. Appl. 478, Kluwer Acad. Publ. (1999), 93–113.

- [17] Grüss G., “Über das Maximum des absoluten Betrages von $\frac{1}{b-a} \int_a^b f(x)g(x)dx - \frac{1}{(b-a)^2} \int_a^b f(x)dx \int_a^b g(x)dx'$ ”, *Math. Z.* 39 (1935), No. 1, 215–226.
- [18] Jankov Maširević D. and Pogány T.K., “Bounds on Čebyšev functional for $C_\varphi[0, 1]$ function class”, *J. Anal.* 22 (2014), 107–117.
- [19] Liu Z., “Refinement of an inequality of Grüss type for Riemann-Stieltjes integral”, *Soochow J. Math.* 30 (2004), No. 4, 483–489.
- [20] Liu Z., “Notes on a Grüss type inequality and its application”, *Vietnam J. Math.* 35 (2007), No. 2, 121–127.
- [21] Lupaş A., “The best constant in an integral inequality”, *Mathematica* (Cluj, Romania), 15 (38) (1973), 219–222.
- [22] Mercer A.Mc.D. and Mercer P.R., “New proofs of the Grüss inequality”, *Aust. J. Math. Anal. Appl.* 1 (2004), No. 2, Art. 12, 6 pp.
- [23] Minculete N. and Ciurdariu L., “A generalized form of Grüss type inequality and other integral inequalities”, *J. Inequal. Appl.* 2014, 2014:119, 18 pp.
- [24] Ostrowski A.M., “On an integral inequality”, *Aequationes Math.* 4 (1970), No. 3, 358–373.
- [25] Pachpatte B.G., “A note on some inequalities analogous to Grüss inequality”, *Octagon Math. Mag.* 5 (1997), No. 2, 62–66.
- [26] Pečarić J. and Ungar Š., “On a inequality of Grüss type”, *Math. Commun.* 11 (2006), No. 2, 137–141.
- [27] Sarikaya M. Z. and Budak H., “An inequality of Grüss like via variant of Pompeiu’s mean value theorem”, *Konuralp J. Math.* 3 (2015), No. 1, 29–35.
- [28] Ujević N., “A generalization of the pre-Grüss inequality and applications to some quadrature formulae”, *JIPAM. J. Inequal. Pure Appl. Math.* 3 (2002), No. 1, Art. 13, 9 pp.