Letter to the Editor



Received: January 12, 2018 Accepted: February 5, 2018 Published online: March 14, 2018

Skin Appendage Disord 2018;4:347–348 DOI: 10.1159/000487493

Do Not Hurt the Nail Matrix: Safe Technique for **Proximal Nail Fold Incision**

Nilton Di Chiacchio^a Leandro Fonseca Noriega^a Cristina Diniz Borges Figueira de Mellob Nilton Gioia Di Chiacchio^{a, c} Jorge Ocampo-Garza^{c, d}

^a Dermatology Clinic, Hospital do Servidor Público Municipal de São Paulo, São Paulo, Brazil; ^bDermatology Department, Universidade Estadual de Campinas, Campinas, Brazil; ^cDermatology Department, Faculdade de Medicina do ABC, São Paulo, Brazil; ^dDermatology Department, University Hospital Dr. José Eleuterio González, Universidad Autónoma de Nuevo León, Monterrey, Mexico

Dear Editor,


For a better view of the nail matrix during nail surgery, the proximal nail fold (PNF) must be incised and reclined. Two oblique incisions at 45° at the junction of the proximal and lateral nail fold are performed [1, 2]. As the nail matrix is just below the PNF, there is a considerable risk of hurting the matrix, and consequently nail dystrophy [1]. How to avoid this risk when both incisions are performed?

The detachment of PNF from the nail plate is performed creating a space between them (Fig. 1a). The blade is placed horizontally into this space (Fig. 1b, c). By a rotation movement, the sharp blade portion is placed upward on the ventral portion of the PNF, while the dorsal part of the blade (not sharp) touches the nail plate and nail matrix (Fig. 2). Then, the incision is performed from the ventral to the dorsal part of the PNF, without any risk for the nail matrix.









sion from the ventral to the dorsal portion



of the PNF.

Statement of Ethics

The authors have no ethical conflicts to disclose.

Disclosure Statement

The authors have no conflicts of interest to declare.

References

- 1 Di Chiacchio N, Belda W Jr, Criado PR: Unhas: o que há de novo. Rio de Janeiro, Atheneu, 2015.
- 2 Göktay F, Güneş P, Yaşar Ş, Güder H, Aytekin S: New observations of intraoperative dermoscopic features of the nail matrix and bed in longitudinal melanonychia. Int J Dermatol 2015;54:1157–1162.

Skin Appendage Disord 2018;4:347–348 DOI: 10.1159/000487493