

Mathematical analysis of plankton population dynamics

Fatin Nadiah Yussof^a, Normah Maan^a, Nadzri Reba^{b,*}

^a Department of Mathematics, Faculty of Science, Universiti Teknologi Malaysia, 81310 UTM Johor Bahru, Johor, Malaysia

^b Faculty of Geoinformation and Real Estate, Universiti Teknologi Malaysia, 81310 UTM Johor Bahru, Johor, Malaysia

* Corresponding author: normahmaan@utm.my

Article history

Received 2 January 2019

Revised 12 March 2019

Accepted 2 May 2019

Published Online 2 February 2020

Abstract

Harmful algal blooms (HABs) event that causes enormous economic loss and health effect raises concerns among environmentalists. In this paper, a mathematical model of interaction between nutrient, toxin-producing phytoplankton (TPP), non-toxic phytoplankton (NTP), zooplankton, and toxic chemicals is proposed to study on how the process of these HABs occurred. The model of interaction is represented by Ordinary Differential Equations (ODEs) and stability analysis of the model is conducted. Several conditions for the system to be stable around trivial and interior equilibrium point are obtained. From the analysis, it is observed that under nutrient limitation, the amounts of toxic chemicals secreted out by the TPP are increased. As a result, NTP population and zooplankton population are affected by the situation. If this situation is prolonged, this will result in the extinction of both populations. Overall, this study shows that TPP release more toxic chemicals when the nutrient is limited and gives a better understanding on the occurrence of HABs event.

Keywords: Stability analysis, harmful algal blooms, toxin-producing phytoplankton, nutrient limitation

© 2020 Penerbit UTM Press. All rights reserved

INTRODUCTION

In the marine ecosystem, there are two forms of plankton community which are plant and animal. Phytoplankton is plant forms of plankton community while zooplankton is an animal form of plankton which feeds on phytoplankton. Generally, phytoplankton can be divided into two, either it is toxin-producing phytoplankton (TPP) or non-toxic phytoplankton (NTP). Toxic chemicals released by TPP cause tremendous effect to the marine organism such as shellfish and fish where the toxic will accumulate in the filter feeder and cause food poisoning to the human [3,17]. Toxic chemicals that usually produced by TPP are okadaic acid (OA) and dinophysistoxin (DTX-1) [1]. Over time, the secretion of these toxic chemicals by TPP may result in harmful algal blooms which gives adverse effect to the marine ecosystem and human health.

For microalgae, [2] concluded that the growth of competing algae can be controlled through the toxin production. [5] conducted a field study as well as analyzing mathematical model and claimed that the survival of weak competitors is enhanced by the toxin allelopathy. [4] have shown that TPP population can control the bloom of NTP. Some dinoflagellates-released toxin such as *Alexandrium tamarense* and *Alexandrium fundyense* decrease the production of toxin under deficient nitrogen condition while under phosphorus limited condition the dinoflagellates increase in production of toxin [6-8]. Besides, other studies have shown that dinoflagellates such as *Procentrum lima* and *Dinophysis acuminata* increase the production of toxin when there are limited nitrogen and phosphorus condition [9-11].

It is important to model on plankton communities especially on HAB and co-existence of species since it is lacking in the literature. Mathematical models can give better understanding of the interaction between the nutrient and plankton population. The complexity of an ecosystem response to variant conditions also can be recognized through the developed model with ordinary differential equations. Several mathematical models that described these methods were

recently developed [4-5,12,18-21]. In this paper, the developed model is modified from [4] model where the model study the effect of nutrient limitation on toxic production. However, the model does not include and study the interaction of the developed model with zooplankton population. Therefore, our concern is to include the zooplankton population into the model system and investigate the interaction of zooplankton in the system.

The flow of this paper is as follows: Section 2 presents the model; Section 3 shows all the conditions for the existence of the equilibrium points in natural phenomena; Section 4 analyzes the local stability of each equilibrium points and the analysis is implemented using Mathematica software; Section 5 presents the numerical simulations of the developed model; Section 6 discusses on the result obtained; and finally, Section 7 concludes this paper.

MATHEMATICAL MODEL

This model system is comprised of five ordinary differential equations which are concentration of nutrient levels $N(t)$, TPP populations $P_1(t)$, NTP populations $P_2(t)$, zooplankton populations $P_3(t)$, and toxin concentration $Z(t)$ present at time t . Let A be the constant rate of nutrient flows into the system while d is the rate of nutrient loss from the system. Assumptions are made in which all the parameter values are defined as follows:

- m_1 = maximum nutrient uptake rates for TPP
- m_2 = maximum nutrient uptake rates for NTP
- a = mortality rates of zooplankton due to toxin
- a_1 = the half saturation constant
- a_2 = the half saturation constant
- r_1 = mortality rates of NTP due to toxin

- r_2 = mortality rates of zooplankton due to toxin
- k = toxic production rate
- k_1 = conversion factor from TPP to zooplankton
- k_2 = conversion factor from NTP to zooplankton
- δ = natural death rate of TPP
- γ = natural death rate of NTP
- θ = toxic washout rate
- β_1 = maximum zooplankton ingestion rate on TPP
- β_2 = maximum zooplankton ingestion rate on NTP
- ε = natural death rate of zooplankton

$$N^{(1)} = \frac{A}{d}$$

- The NTP and zooplankton free equilibrium $E_2=(N^{(2)},P_1^{(2)},0,0,Z^{(2)})$ always exists if

$$N^{(2)} = -\frac{a_1\delta}{\delta - m_1}, P_1^{(2)} = \frac{A\delta + a_1d\delta - Am_1}{\delta(\delta - m_1)},$$

$$Z^{(2)} = \frac{-A\delta k - a_1d\delta k + Akm_1}{a_1m_1\delta\theta}$$

- Michaelis-Menten functions $\frac{N}{a_1 + N}$ and $\frac{N}{a_2 + N}$ are used to model the nutrient uptake for growth of TPP and NTP, respectively, in order to illustrate that nutrient concentration level influences the phytoplankton nutrient uptake rate [15]. For growth and nutrient-limited uptake, the equations involved follow the standard representation form [16].
- The functional response of zooplankton that feeds on TPP and NTP is described by the law of predation of Holling type-II [13,14].
- The amount of toxic chemicals released by TPP are higher when the concentration of nutrient becomes limited. Therefore, this phenomenon is described by a monotonic decreasing function of nutrient, $\frac{kP_1}{a + N}$.
- Toxic chemicals released by TPP gives adverse effect to the existence of NTP and zooplankton population where the toxic chemicals caused the population die.

Local stability analysis of equilibrium points

A system is locally asymptotically stable if all the solutions approaching the equilibrium points as $t \rightarrow \infty$. The local stability of the equilibrium points is investigated by finding the eigenvalues of the associated Jacobian matrices at system (1).

The Jacobian matrix at $E_1=(N^{(1)},0,0,0,0)$ is

$$J_1 = \begin{bmatrix} -d & -\frac{m_1N^{(1)}}{a_1 + N^{(1)}} & -\frac{m_2N^{(1)}}{a_2 + N^{(1)}} & 0 & 0 \\ 0 & -\delta + \frac{m_1N^{(1)}}{a_1 + N^{(1)}} & 0 & 0 & 0 \\ 0 & 0 & -\gamma + \frac{m_2N^{(1)}}{a_2 + N^{(1)}} & 0 & 0 \\ 0 & 0 & 0 & -\varepsilon & 0 \\ 0 & \frac{k}{a_1 + N^{(1)}} & 0 & 0 & -\theta \end{bmatrix}$$

Eigenvalues of J_1 is given by:

$$\lambda_N = -d, \lambda_{P_1} = -\delta + \frac{m_1N^{(1)}}{a_1 + N^{(1)}}, \lambda_{P_2} = -\gamma + \frac{m_2N^{(1)}}{a_2 + N^{(1)}},$$

$$\lambda_{P_3} = -\varepsilon, \lambda_Z = -\theta$$

The equilibrium point E_1 is locally asymptotically stable if the condition along with the following conditions are satisfied:

$$\frac{m_1N^{(1)}}{a_1 + N^{(1)}} < \delta$$

$$\frac{m_2N^{(1)}}{a_2 + N^{(1)}} < \gamma$$

Biological interpretation

If the respective maximal growth rates $\frac{m_1N^{(1)}}{a_1 + N^{(1)}}$ and $\frac{m_2N^{(1)}}{a_2 + N^{(1)}}$

of TPP and NTP populations are less than their respective corresponding natural removal rates (δ and γ) and the amount of nutrient is stabilized at $N^{(1)}$, both TPP and NTP populations will undergo extinction.

The Jacobian matrix at $E_2=(N^{(2)}, P_1^{(2)}, 0, 0, Z^{(2)})$ is

The model suggested in this work is as follows:

$$\begin{aligned} \frac{dN}{dt} &= A - dN - \frac{m_1NP_1}{a_1 + N} - \frac{m_2NP_2}{a_2 + N} \\ \frac{dP_1}{dt} &= \frac{m_1NP_1}{a_1 + N} - \delta P_1 - \frac{\beta_1 P_1 P_3}{a_1 + P_1} \\ \frac{dP_2}{dt} &= \frac{m_2NP_2}{a_2 + N} - \gamma P_2 - \frac{\beta_2 P_2 P_3}{a_2 + P_2} - r_1 P_2 Z \\ \frac{dP_3}{dt} &= \frac{k_1\beta_1 P_1 P_3}{a_1 + P_1} + \frac{k_2\beta_2 P_2 P_3}{a_2 + P_2} - \varepsilon P_3 - r_2 P_3 Z \\ \frac{dZ}{dt} &= \frac{kP_1}{a_1 + N} - \theta Z \end{aligned} \tag{1}$$

System (1) is subject to the following initial conditions:

$$N(0) > 0, P_1(0) > 0, P_2(0) > 0, P_3(0) > 0, Z(0) > 0.$$

Existence of equilibrium points

The system (1) possesses the following equilibrium points:

- The plankton and toxin free equilibrium $E_1=(N^{(1)},0,0,0,0)$ always exists if

$$J_2 = \begin{bmatrix} -d + \frac{m_1 N^{(2)} P_1^{(2)}}{(a_1 + N^{(2)})^2} - \frac{m_1 P_1^{(2)}}{a_1 + N^{(2)}} & -\frac{m_1 N^{(2)}}{a_1 + N^{(2)}} & -\frac{m_2 N^{(2)}}{a_2 + N^{(2)}} & 0 & 0 \\ -\frac{m_1 N^{(2)} P_1^{(2)}}{(a_1 + N^{(2)})^2} + \frac{m_1 P_1^{(2)}}{a_1 + N^{(2)}} & -\delta + \frac{m_1 N^{(2)}}{a_1 + N^{(2)}} & 0 & -\frac{\beta_1 P_1^{(2)}}{a_1 + P_1^{(2)}} & 0 \\ 0 & 0 & -\gamma + \frac{m_2 N^{(2)}}{a_2 + N^{(2)}} - r_1 Z^{(2)} & 0 & 0 \\ 0 & 0 & 0 & \frac{\beta_1 k_1 P_1^{(2)}}{a_1 + P_1^{(2)}} - \varepsilon - r_2 Z^{(2)} & 0 \\ -\frac{k P_1^{(2)}}{(a_1 + N^{(2)})^2} & \frac{k}{a_1 + N^{(2)}} & 0 & 0 & -\theta \end{bmatrix}$$

The characteristic equation of this Jacobian matrix is given by

$$\left(-\gamma + \frac{m_2 N^{(2)}}{a_2 + N^{(2)}} - r_1 Z^{(2)} - \lambda_{2P_2}\right) \left(-\varepsilon + \frac{\beta_1 k_1 P_1^{(2)}}{a_1 + P_1^{(2)}} - r_2 Z^{(2)} - \lambda_{2P_3}\right) \left(-\theta - \lambda_{2Z}\right) \left(\lambda_{2N}^2 + A_4 \lambda_{2N} + A_5\right) = 0$$

Therefore,

$$\lambda_{2P_1, 2N} = \frac{-dw - \delta w + y + m_1 N^{(2)2}}{2w} \pm \sqrt{(dw + \delta w - y)^2 - 4w(d\delta w - \delta y - dm_1 N^{(2)2})}$$

where

$$w = a_1^2 + 2a_1 N^{(2)} + N^{(2)2}$$

$$y = a_1 m_1 N^{(2)} - a_1 m_1 P_1^{(2)}$$

Since all the eigenvalues have negative real parts, then the equilibrium point E_2 is locally asymptotically stable if the existence condition satisfies the following conditions:

$$\frac{m_2 N^{(2)}}{a_2 + N^{(2)}} < \gamma + r_1 Z^{(2)}$$

$$\frac{\beta_1 k_1 P_1^{(2)}}{a_1 + P_1^{(2)}} < \varepsilon + r_2 Z^{(2)}$$

Biological interpretation

When the amount of nutrient and toxic chemicals are stabilized at $N^{(2)}$ and $Z^{(2)}$, respectively, NTP and zooplankton populations become extinct if their maximal growth rates $\frac{m_2 N^{(2)}}{a_2 + N^{(2)}}$ and $\frac{\beta_1 k_1 P_1^{(2)}}{a_1 + P_1^{(2)}}$ are less than total loss rate $\gamma + r_1 Z^{(2)}$ and $\varepsilon + r_2 Z^{(2)}$.

The Jacobian matrix at $E^*=(N^*, P_1^*, P_2^*, P_3^*, Z^*)$ is

$$J(N^*, P_1^*, P_2^*, P_3^*, Z^*) = \begin{bmatrix} A_1 & A_2 & A_3 & 0 & 0 & 1 \\ A_4 & A_5 & 0 & A_6 & 0 & 2 \\ A_7 & 0 & A_8 & A_9 & -P_2^* r & \\ 0 & A_{10} & A_{11} & A_{12} & -P_3^* r & \\ A_{13} & A_{14} & 0 & 0 & -\theta & \end{bmatrix}$$

where

$$A_1 = -d + \frac{m_1 N^* P_1^*}{(a_1 + N^*)^2} - \frac{m_1 P_1^*}{a_1 + N^*} + \frac{m_2 N^* P_2^*}{(a_2 + N^*)^2} - \frac{m_2 P_2^*}{a_2 + N^*}$$

$$A_2 = -\frac{m_1 N^*}{a_1 + N^*}$$

$$A_3 = -\frac{m_2 N^*}{a_2 + N^*}$$

$$A_4 = -\frac{m_1 N^* P_1^*}{(a_1 + N^*)^2} + \frac{m_1 P_2^*}{a_1 + N^*}$$

$$A_5 = -\delta + \frac{m_1 N^*}{a_1 + N^*} + \frac{\beta_1 P_1^* P_3^*}{(a_1 + P_1^*)^2} - \frac{\beta_1 P_3^*}{a_1 + P_1^*}$$

$$A_6 = -\frac{\beta_1 P_1^*}{a_1 + P_1^*}$$

$$A_7 = -\frac{m_2 N^* P_2^*}{(a_2 + N^*)^2} + \frac{m_2 P_2^*}{a_2 + N^*}$$

$$A_8 = -\gamma + \frac{m_2 N^*}{a_2 + N^*} + \frac{\beta_2 P_2^* P_3^*}{(a_2 + P_2^*)^2} - \frac{\beta_2 P_3^*}{a_2 + P_2^*} - r_1 Z^*$$

$$A_9 = -\frac{\beta_2 P_2^*}{a_2 + P_2^*}$$

$$A_{10} = -\frac{\beta_1 k_1 P_1^* P_3^*}{(a_1 + P_1^*)^2} + \frac{\beta_1 k_1 P_3^*}{a_1 + P_1^*}$$

$$A_{11} = -\frac{\beta_2 k_2 P_2^* P_3^*}{(a_2 + P_2^*)^2} + \frac{\beta_2 k_2 P_3^*}{a_2 + P_2^*}$$

$$A_{12} = \frac{\beta_1 k_1 P_1^*}{a_1 + P_1^*} + \frac{\beta_2 k_2 P_2^*}{a_2 + P_2^*} - \varepsilon - r_2 Z^*$$

$$A_{13} = -\frac{k P_1^*}{(a_1 + N^*)^2}$$

$$A_{14} = \frac{k}{a_1 + N^*}$$

$$\det|J - \lambda I| = \begin{vmatrix} A_1 - \lambda & A_2 & A_3 & 0 & 0 \\ A_4 & A_5 - \lambda & 0 & A_6 & 0 \\ A_7 & 0 & A_8 - \lambda & A_9 & -P_2^* r_1 \\ 0 & A_{10} & A_{11} & A_{12} - \lambda & -P_3^* r_2 \\ A_{13} & A_{14} & 0 & 0 & -\theta - \lambda \end{vmatrix} = 0$$

Characteristic equation

$$\lambda^5 + B_1 \lambda^4 + B_2 \lambda^3 + B_3 \lambda^2 + B_4 \lambda + B_5 = 0$$

where

$$B_1 = -A_1 - A_3 - A_5 - A_8 + \theta$$

$$B_2 = A_1 A_3 - A_{10} A_4 + A_1 A_5 + A_3 A_5 - A_{11} A_6 - A_{12} A_7 + A_1 A_8 + A_3 A_8 + A_5 A_8 - A_2 A_9 - A_1 \theta - A_3 \theta - A_5 \theta - A_8 \theta$$

$$B_3 = A_{10} A_3 A_4 - A_1 A_3 A_5 + A_1 A_{11} A_6 + A_{11} A_5 A_6 + A_1 A_{12} A_7 + A_{12} A_3 A_7 - A_1 A_3 A_8 + A_{10} A_4 A_8 - A_1 A_5 A_8 - A_3 A_5 A_8 + A_2 A_5 A_9 + A_2 A_8 A_9 + A_1 A_3 \theta - A_{10} A_4 \theta + A_1 A_5 \theta + A_3 A_5 \theta - A_{11} A_6 \theta - A_{12} A_7 \theta + A_1 A_8 \theta + A_3 A_8 \theta + A_5 A_8 \theta - A_2 A_9 \theta + A_{10} A_{13} P_2^* r_1 + A_{11} A_{14} P_3^* r_2$$

$$\begin{aligned}
 B_4 = & -A_{10}A_{12}A_2A_6 + A_{10}A_{11}A_4A_6 - A_1A_{11}A_5A_6 - A_1A_{12}A_3A_7 \\
 & - A_{10}A_3A_4A_8 + A_1A_3A_5A_8 + A_{12}A_2A_7A_9 - A_{11}A_4A_7A_9 \\
 & - A_2A_5A_8A_9 + A_{10}A_3A_4\theta - A_1A_3A_5\theta + A_1A_{11}A_6\theta + A_{11}A_5A_6\theta \\
 & + A_1A_{12}A_7\theta + A_{12}A_3A_7\theta - A_1A_3A_8\theta + A_{10}A_4A_8\theta - A_1A_5A_8\theta \\
 & - A_3A_5A_8\theta + A_2A_5A_9\theta + A_2A_8A_9\theta + A_2A_8A_9\theta + A_{10}A_{14}A_2P_2^*r_1 \\
 & - A_{10}A_{13}A_3P_2^*r_1 + A_{11}A_{14}A_7P_2^*r_1 - A_{10}A_{13}A_8P_2^*r_1 + A_{10}A_{12}A_{13}P_3^*r_2 \\
 & - A_1A_{11}A_{14}P_3^*r_2 - A_{11}A_{14}A_5P_3^*r_2 + A_{11}A_{13}A_9P_3^*r_2
 \end{aligned}$$

$$\begin{aligned}
 B_5 = & -A_{10}A_{12}A_2A_6\theta + A_{10}A_{11}A_4A_6\theta - A_1A_{11}A_5A_6\theta \\
 & - A_1A_{12}A_3A_7\theta - A_{10}A_3A_4A_8\theta + A_1A_3A_5A_8\theta + A_{12}A_2A_7A_9\theta \\
 & - A_{11}A_4A_7A_9\theta - A_2A_5A_8A_9\theta - A_{10}A_{11}A_{13}A_6P_2^*r_1 \\
 & - A_1A_{11}A_{14}A_7P_2^*r_1 - A_{10}A_{14}A_2A_8P_2^*r_1 - A_{10}A_{14}A_2A_8P_2^*r_1 \\
 & + A_{10}A_{13}A_3A_8P_2^*r_1 + A_{10}A_{13}A_3A_8P_2^*r_1 + A_{11}A_{13}A_7A_9P_2^*r_1 \\
 & + A_{10}A_{12}A_{14}A_2P_3^*r_2 - A_{10}A_{12}A_{13}A_3P_3^*r_2 - A_{10}A_{11}A_{14}A_4P_3^*r_2 \\
 & + A_1A_{11}A_{14}A_5P_3^*r_2 - A_{11}A_{13}A_5A_9P_3^*r_2
 \end{aligned}$$

A system is stable if all the eigenvalues are negative real parts. Thus, by following Routh-Hurwitz criterion, all the eigenvalues must satisfy the following conditions:

$$B_i > 0$$

$$B_1B_2B_3 > B_3^2 + B_1^2B_4$$

$$(B_1B_4 - B_5)(B_1B_2B_3 - B_3^2 - B_1^2B_4) > B_5(B_1B_2 - B_3)^2 + B_1B_5^2$$

Biological interpretation

If all the roots satisfy the above conditions, then the equilibrium point of E^* is stable, which indicates that there is no occurrence of HABs.

NUMERICAL SIMULATIONS


In this section, numerical simulations for system (1) is performed to study the dynamical behavior of the system. By varying the parameter value of A , and the other values of parameter are as shown in Table 1. The simulation results are obtained and it is observed that, system (1) has an equilibrium point at $E^* = (N^*, P_1^*, P_2^*, P_3^*, Z^*)$ as shown in Figure 1 when the parameter value is $A=1.0$.

Table 1 Parameter values used in numerical analysis.


Name	Description	Values
d	Amount of nutrient loss	0.1
a_1	Half saturation constant	0.07
a_2	Half saturation constant	0.2
m_1	Uptake rate of TPP	1.0
m_2	Uptake rate of NTP	0.7
k	Toxin production rate	0.07
k_1	Conversion rate of TPP	0.01
k_2	Conversion rate of NTP	0.01
r_1	Mortality rate of NTP due to toxin	0.8
r_2	Mortality rate of zooplankton due to toxin	0.8
ϵ	Per capita natural death rate of zooplankton	0.02
β_1	Maximum zooplankton ingestion rate on TPP	0.01
β_2	Maximum zooplankton ingestion rate on NTP	0.02
θ	Washout rate of toxin	0.8
γ	Per capita natural death rate of NTP	0.01
δ	Per capita natural death rate of TPP	0.5

If the parameter value is $A=1.5$ and the other values of parameter are as shown in Table 1, hence system (1) has an equilibrium point at $E_1=(N^{(1)}, 0,0,0,0)$ as shown in Figure 2. Next, if we consider parameter value of $A=0.5$ and the other values of parameter are as shown in Table 1, then system (1) has an equilibrium point at $E_2=(N^{(2)}, P_1^{(2)}, 0, 0, Z^{(2)})$ as shown in Figure 3. Table 1 shows the parameter values used in this model [4].

a) Nutrient.



b) Toxic phytoplankton, non-toxic phytoplankton, and zooplankton.



c) Toxic chemicals.






Figure 1 From (a-e), the solution goes to components $E_2=(N^*, P_1^*, P_2^*, P_3^*, Z^*)$ when the parameter value is $A = 1.0$.


a) Nutrient




a) Nutrient




b) Toxic phytoplankton, non-toxic phytoplankton, and zooplankton.



b) Toxic phytoplankton, non-toxic phytoplankton, and zooplankton.



c) Toxic chemicals.



c) Toxic chemicals.




Figure 2 From (a-e), the solution goes to components $E_1=(N^{(1)}, 0,0,0,0)$ where the parameter value is $A = 1.5$.

Figure 3 From (a-e), the solution goes to components $E_2=(N^{(2)}, P_I^{(2)}, 0, 0, Z^{(2)})$ where the parameter value is $A = 0.5$.

DISCUSSION

In Figure 1, the solution is asymptotically stable at component $E_2=(N^*, P_1^*, P_2^*, P_3^*, Z^*)$ where the parameter value is $A=1.0$. In this situation, HAB does not occur as all the variables exist in the ecosystem and the system is balance. Next, in Figure 2, it can be seen that, when the parameter value of $A=1.5$, the solution goes to components $E_1=(N^{(1)}, 0, 0, 0, 0)$ where in this situation, only nutrient is present in the water. Therefore, HAB will not occur in this context.

In Figure 3, when the parameter value is $A=0.5$, the solution is asymptotically stable at components $E^*=(N^*, P_1^*, 0, 0, Z^*)$. From the graph, it can be seen that when the nutrient is limited, TPP produces more toxic chemicals and there is no significant change when the nutrient is sufficient. This shows that the environmental stress forced the TPP to produce more toxic chemicals. As the amount of toxic chemicals increase in the water, concentrations of NTP and zooplankton drop tremendously. Hence, it shows that HAB caused other population to die due to the effects of toxic chemicals by TPP in the water.

CONCLUSION

The results obtained demonstrate that when the nutrient is limited, TPP releases more toxic chemicals. This is due to the environmental stress faced by the TPP population since the nutrient is limited forcing them to release more toxic chemicals. When toxic chemicals concentration is higher in the water, this situation affects the co-existence of NTP and zooplankton populations. Toxic chemicals produced by TPP population will harm the NTP and zooplankton populations, causing these populations to cease. Elimination of zooplankton population as the grazer in the system results in ecosystem imbalance, thus allowing the TPP population to dominate the ecosystem. Therefore, the amount of toxic chemicals in the water will increase and this will pose adverse effect to human and also marine organism. Hence, the existence of zooplankton population is important to balance the marine ecosystem as well as to control the algal bloom. The data shown from the model presented enable us to have better understanding on the occurrence of HAB and the importance of population dynamic to prevent ecosystem imbalance.

ACKNOWLEDGEMENT

This work was financially supported by the Universiti Teknologi Malaysia under the Research University Grant and Ministry of Higher Education Malaysia through research grant of vot no. 4L854.

REFERENCES

- [1] Windust, A. J., Wright, J. L. C., and McLachlan, J. L. 1996. The effects of the diarrhetic shellfish poisoning toxins, okadaic acid and dinophysistoxin-1, on the growth of microalgae. *Marine biology*. 126(1): 19-25.
- [2] Hulot, F. D., and Huisman, J. 2004. Allelopathic interactions between phytoplankton species: the roles of heterotrophic bacteria and mixing intensity. *Limnology and Oceanography*. 49(4part2): 1424-1434.
- [3] Lim, H. C., Leaw, C. P., Tan, T. H., Kon, N. F., Yek, L. H., Hii, K. S., Teng, S. T., Razali, R. M., Usup, G., Iwataki, M., and Lim, P. T. 2014. A bloom of *Karlodinium australe* (Gymnodiniales, Dinophyceae) associated with mass mortality of cage-cultured fishes in West Johor Strait, Malaysia. *Harmful Algae*. 40: 51-62.
- [4] Chakraborty, S., Roy, S., and Chattopadhyay, J. 2008. Nutrient-limited toxin production and the dynamics of two phytoplankton in culture media: A mathematical model. *Ecological modelling*. 213(2): 191-201.
- [5] Roy, S., and Chattopadhyay, J. 2007. Toxin-allelopathy among phytoplankton species prevents competitive exclusion. *Journal of Biological Systems*. 15(1): 73-93.
- [6] Anderson, D. M., Kulis, D. M., Sullivan, J. J., and Hall, S. 1990. Analysis of a dengue fever transmission model. *Toxicon*. 28(8): 885-893.
- [7] Boyer, G. L., Sullivan, J. J., Andersen, R. J., Harrison, P. J., and Taylor, F. J. R. 1987. Effects of nutrient limitation on toxin production and composition in the marine dinoflagellate *Protogonyaulax tamarensis*. *Marine Biology*. 96(1): 123-128.
- [8] Hall, S. 1982. Toxins and toxicity of *Protogonyaulax* from the northeast Pacific. PhD thesis. University of Alaska Fairbanks.
- [9] McLachlan, J. L., Marr, J. C., ad Conlon-Keily, A., and Adamson, A. 1994. Effects of nitrogen concentration and cold temperature on DSP-toxin concentrations in the dinoflagellate *Prorocentrum lima* (prorocentrales, dinophyceae). *Natural toxins*. 2(5): 263-270.
- [10] Sohet, K., Pereira, A., and Braekman, J. C., and Houvenaghel, G. 1995. Growth and toxicity of *Prorocentrum lima* (Ehrenberg) dodge in different culture media: Effect of humic acids and organic phosphorus. *in*: Lassus, P (Ed.). *Harmful marine algal blooms: Proceedings of the Sixth International Conference on toxic marine phytoplankton = Proliférations d'algues marines nuisibles: Sixième Conférence Internationale sur le phytoplancton toxique, October 18-22, 1993, Nantes, France*. pp. 669-674.
- [11] Johansson, N., and Graneli, E. 1999. Influence of different nutrient conditions on cell density, chemical composition and toxicity of *Prymnesium parvum* (Haptophyta) in semi-continuous cultures. *Journal of Experimental Marine Biology and Ecology*. 239(2): 243-258.
- [12] Rehim, M., Zhang, Z., and Muhammadhaji, A. 2016. Mathematical analysis of a nutrient-plankton system with delay. *SpringerPlus*. 5(1): 1055.
- [13] Chattopadhyay, J., Sarkar, R. R., and Mandal, S. 2002. Toxin-producing plankton may act as a biological control for planktonic blooms field study and mathematical modelling. *Journal of Theoretical Biology*. 215(3): 333-344.
- [14] Das, K., and Ray, S. 2008. Effect of delay on nutrient cycling in phytoplankton-zooplankton interactions in estuarine system. *Ecological Modelling*. 215(1-3): 69-76.
- [15] Dugdale, R. C. J. 1967. Nutrient limitation in the sea: Dynamics, identification, and significance. *Limnology and Oceanography*. 12(4): 685-695.
- [16] Franks, P. J. S. 2002. NPZ models of plankton dynamics: Their construction, coupling to physics and application. *Journal of Oceanography*. 58(2): 379-387.
- [17] Teen, L. P., Pin, L. C., Gires, U. 2012. Harmful algal blooms in Malaysian waters. *Sains Malaysiana*. 41(12):1509-1515.
- [18] Fan, A., Han, P., and Wang, K. 2013. Global dynamics of a nutrient-plankton system in the water ecosystem. *Applied Mathematics and Computation*. 219(15): 8269-8276.
- [19] Chakraborty, K., and Dasb, K. 2015. Modeling and analysis of a two-zooplankton one phytoplankton system in the presence of toxicity. *Applied Mathematical Modelling*. 39(3-4): 1241-1265.
- [20] Khare, S., Dhar, J., and Misra, O. P. 2010. Role of toxin-producing phytoplankton on a plankton ecosystem. *Nonlinear Analysis: Hybrid Systems*. 4(3): 496-502.
- [21] Chatterjee, A., Pal, S., and Chatterjee, S. 2011. Bottom up and top down effect on toxin-producing phytoplankton and its consequence on the formation of plankton bloom. *Applied Mathematics and Computation*. 218(7): 3387-3398.