



Miskolc Mathematical Notes  
Vol. 20 (2019), No. 2, pp. 997–1011

HU e-ISSN 1787-2413  
DOI: 10.18514/MMN.2019.2489

## SOME NEW INTEGRAL INEQUALITIES FOR $N$ -TIMES DIFFERENTIABLE $R$ -CONVEX AND $R$ -CONCAVE FUNCTIONS

HURIYE KADAKAL, MAHIR KADAKAL, AND İMDAT İŞCAN

*Received 02 January, 2018*

*Abstract.* In this work, by using an integral identity together with both the Hölder and the Power-Mean integral inequality we establish several new inequalities for  $n$ -time differentiable  $r$ -convex and concave functions.

2010 *Mathematics Subject Classification:* 26A51; 26D10; 26D15

*Keywords:* convex function,  $r$ -convex function, Hölder integral inequality and power-mean integral inequality

### 1. INTRODUCTION

A function  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  is said to be convex if the inequality

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y)$$

is valid for all  $x, y \in I$  and  $t \in [0, 1]$ . If this inequality reverses, then  $f$  is said to be concave on interval  $I \neq \emptyset$ . This definition is well known in the literature. Convexity theory has appeared as a powerful technique to study a wide class of unrelated problems in pure and applied sciences. Many articles have been written by a number of mathematicians on convex functions and inequalities for their different classes, using, for example, the last articles [3, 8–15] and the references in these papers.

$f : [a, b] \rightarrow \mathbb{R}$  be a convex function, then the inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a) + f(b)}{2}$$

is known as the Hermite-Hadamard inequality (see [6] for more information). Since then, some refinements of the Hermite-Hadamard inequality on convex functions have been extensively investigated by a number of authors (e.g., [3, 4, 15]). In [20], the first author obtained a new refinement of the Hermite-Hadamard inequality for convex functions. The Hermite-Hadamard inequality was generalized in [17] to an  $r$ -convex positive function which is defined on an interval  $[a, b]$ .

**Definition 1.** A positive function  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  is called  $r$ -convex function on  $[a, b]$ , if for each the  $x, y \in [a, b]$  and  $t \in [0, 1]$ ,

$$f(tx + (1-t)y) \leq \begin{cases} [tf^r(x) + (1-t)f^r(y)]^{\frac{1}{r}}, & r \neq 0, \\ [f(x)]^t [f(y)]^{1-t}, & r = 0. \end{cases}$$

If the equality is reversed, then the function  $f$  is said to be  $r$ -concave.

It is obvious 0-convex functions are simply log-convex functions, 1-convex functions are ordinary convex functions and  $-1$ -convex functions are arithmetically harmonically convex. One should note that if  $f$  is  $r$ -convex on  $[a, b]$ , then the function  $f^r$  is a convex function for  $r > 0$  and  $f^r$  is a concave function for  $r < 0$ . We note that if  $f$  and  $g$  are convex and  $g$  is increasing, then  $g \circ f$  is convex; moreover, since  $f = \exp(\log f)$ , it follows that a log-convex function is convex.

The definition of  $r$ -convexity naturally complements the concept of  $r$ -concavity, in which the inequality is reversed [18] and which plays an important role in statistics.

It is easily seen that if  $f$  is  $r$ -convex on  $[a, b]$ ,

$$f^r\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f^r(x) dx \leq \frac{f^r(a) + f^r(b)}{2}, \quad r > 0 \quad (1.1)$$

$$f^r\left(\frac{a+b}{2}\right) \geq \frac{1}{b-a} \int_a^b f^r(x) dx \geq \frac{f^r(a) + f^r(b)}{2}, \quad r < 0 \quad (1.2)$$

Some refinements of the Hadamard inequality for  $r$ -convex functions could be found in [2, 7, 16, 19, 21]. In [1], Bessenyei studied Hermite-Hadamard-type inequalities for generalized 3-convex functions. In [16], the authors showed that if  $f$  is  $r$ -convex in  $[a, b]$  and  $0 < r \leq 1$ , then

$$\frac{1}{b-a} \int_a^b f(x) dx \leq \frac{r}{r+1} [f^r(a) + f^r(b)]^{\frac{1}{r}}. \quad (1.3)$$

**Theorem 1 ([5]).** Suppose that  $f$  is a positive  $r$ -convex function on  $[a, b]$ . Then

$$\frac{1}{b-a} \int_a^b f(t) dt \leq L_r(f(a), f(b)).$$

If  $f$  is a positive  $r$ -concave function, then the inequality is reversed, where

$$L_r(f(a), f(b)) = \begin{cases} \frac{r}{r+1} \frac{f^{r+1}(a) - f^{r+1}(b)}{f^r(a) - f^r(b)}, & r \neq 0, -1, & f(a) \neq f(b) \\ \frac{f(a) - f(b)}{\ln f(a) - \ln f(b)}, & r = 0, & f(a) \neq f(b) \\ f(a) f(b) \frac{\ln f(a) - \ln f(b)}{f(a) - f(b)}, & r = -1, & f(a) \neq f(b) \\ f(a), & & f(a) = f(b). \end{cases}$$

**Theorem 2.** Let  $f : [a, b] \rightarrow (0, \infty)$  be  $r$ -convex function and  $r \geq 1$ . Then the following inequality holds:

$$\frac{1}{b-a} \int_a^b f(x) dx \leq \left[ \frac{f^r(a) + f^r(b)}{2} \right]^{\frac{1}{r}}.$$

**Lemma 1.** Let  $a \geq 0, b \geq 0$ . Then  $(a+b)^\lambda \leq a^\lambda + b^\lambda, 0 < \lambda \leq 1$ .

Let  $0 < a < b$ , throughout this paper we will use

$$A(a, b) = \frac{a+b}{2}, \quad G(a, b) = \sqrt{ab}$$

$$L_p(a, b) = \left( \frac{b^{p+1} - a^{p+1}}{(p+1)(b-a)} \right)^{\frac{1}{p}}, \quad a \neq b, \quad p \in \mathbb{R}, \quad p \neq -1, 0$$

for the arithmetic, geometric, generalized logarithmic mean, respectively. Also for shortness we will use the following notation:

$$I(a, b, n, f) = \sum_{k=0}^{n-1} (-1)^k \left( \frac{f^{(k)}(b) b^{k+1} - f^{(k)}(a) a^{k+1}}{(k+1)!} \right) - \int_a^b f(x) dx$$

where an empty sum is understood to be nil.

## 2. MAIN RESULTS

We will use the following Lemma for obtain our main results.

**Lemma 2** ([14]). Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be  $n$ -times differentiable mapping on  $I^\circ$  for  $n \in \mathbb{N}$  and  $f^{(n)} \in L[a, b]$ , where  $a, b \in I^\circ$  with  $a < b$ , we have the identity

$$I(a, b, n, f) = \frac{(-1)^{n+1}}{n!} \int_a^b x^n f^{(n)}(x) dx.$$

where an empty sum is understood to be nil.

**Theorem 3.** For  $n \in \mathbb{N}$ ; let  $f : I \subset (0, \infty) \rightarrow \mathbb{R}$  be  $n$ -times differentiable function on  $I^\circ$ ,  $r > 0$  and  $a, b \in I^\circ$  with  $a < b$ . If  $f^{(n)} \in L[a, b]$  and  $|f^{(n)}|^q$  for  $q > 1$  is  $r$ -convex function on  $[a, b]$ , then the following inequality holds:

$$|I(a, b, n, f)| \leq \frac{b-a}{n!} L_{np}^n(a, b) L_{\frac{1}{r}}^{\frac{1}{r}} \left( |f^{(n)}(a)|^{qr}, |f^{(n)}(b)|^{qr} \right)$$

*Proof.* If  $|f^{(n)}|^q$  for  $q > 1$  is  $r$ -convex function on  $[a, b]$  and  $r > 0$ , using Lemma 2, the Hölder integral inequality and

$$|f^{(n)}(x)|^q = \left| f^{(n)} \left( \frac{x-a}{b-a} b + \frac{b-x}{b-a} a \right) \right|^q$$

$$\leq \left[ \frac{x-a}{b-a} |f^{(n)}(b)|^{qr} + \frac{b-x}{b-a} |f^{(n)}(a)|^{qr} \right]^{\frac{1}{r}},$$

we have

$$\begin{aligned} & |I(a, b, n, f)| \\ & \leq \frac{1}{n!} \int_a^b x^n |f^{(n)}(x)| dx \\ & \leq \frac{1}{n!} \left( \int_a^b x^{np} dx \right)^{\frac{1}{p}} \left( \int_a^b |f^{(n)}(x)|^q dx \right)^{\frac{1}{q}} \\ & \leq \frac{1}{n!} \left( \int_a^b x^{np} dx \right)^{\frac{1}{p}} \left( \int_a^b \left[ \frac{x-a}{b-a} |f^{(n)}(b)|^{qr} + \frac{b-x}{b-a} |f^{(n)}(a)|^{qr} \right]^{\frac{1}{r}} dx \right)^{\frac{1}{q}} \\ & = \frac{1}{n!} \left( \int_a^b x^{np} dx \right)^{\frac{1}{p}} \left( \int_{|f^{(n)}(a)|^{qr}}^{|f^{(n)}(b)|^{qr}} u^{\frac{1}{r}} \frac{b-a}{|f^{(n)}(b)|^{qr} - |f^{(n)}(a)|^{qr}} du \right)^{\frac{1}{q}} \\ & = \frac{1}{n!} (b-a) \left( \frac{b^{np+1} - a^{np+1}}{(np+1)(b-a)} \right)^{\frac{1}{p}} \left( \frac{|f^{(n)}(b)|^{qr(\frac{1}{r}+1)} - |f^{(n)}(a)|^{qr(\frac{1}{r}+1)}}{(\frac{1}{r}+1)(|f^{(n)}(b)|^{qr} - |f^{(n)}(a)|^{qr})} \right)^{\frac{1}{q}} \\ & = \frac{1}{n!} (b-a) L_{np}^n(a, b) \left( \frac{|f^{(n)}(b)|^{qr(\frac{1}{r}+1)} - |f^{(n)}(a)|^{qr(\frac{1}{r}+1)}}{(\frac{1}{r}+1)(|f^{(n)}(b)|^{qr} - |f^{(n)}(a)|^{qr})} \right)^{\frac{1}{q}} \\ & = \frac{1}{n!} (b-a) L_{np}^n(a, b) L_{\frac{1}{r}}^{\frac{1}{q}}(|f^{(n)}(a)|^{qr}, |f^{(n)}(b)|^{qr}). \end{aligned}$$

This completes the proof of theorem.  $\square$

*Remark 1.* The results obtained in this paper reduces to the results of [14] in case of  $r = 1$ .

**Corollary 1.** Under the conditions Theorem 3 for  $n = 1$  we have the following inequality:

$$\left| \frac{f(b)b - f(a)a}{b-a} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq L_p(a, b) L_{\frac{1}{r}}^{\frac{1}{q}}(|f'(a)|^{qr}, |f'(b)|^{qr}).$$

**Proposition 1.** Let  $a, b \in (0, \infty)$  with  $a < b$ ,  $q > 1$  and  $m \geq 1$ ,  $r \geq 1$  we have

$$L_{\frac{m}{q}+1}^{\frac{m}{q}+1}(a, b) \leq L_p(a, b) L_{\frac{1}{r}}^{\frac{1}{q}}(a^{mr}, b^{mr}).$$

*Proof.* Under the assumption of the Proposition, let  $f(x) = \frac{q}{m+q}x^{\frac{m}{q}+1}$ ,  $x \in (0, \infty)$ . Then  $|f'(x)|^q = x^m$  is  $r$ -convex on  $(0, \infty)$  and the result follows directly from Corollary 1.  $\square$

*Remark 2.* Under the assumption of the Proposition 2.1, If  $r = 1$ ,  $m = 1$ , then the results obtained in this paper reduces to the results of [14].

**Theorem 4.** For  $n \in \mathbb{N}$ ; let  $f : I \subset (0, \infty) \rightarrow \mathbb{R}$  be  $n$ -times differentiable function on  $I^\circ$ ,  $r > 0$  and  $a, b \in I^\circ$  with  $a < b$ . If  $f^{(n)} \in L[a, b]$  and  $|f^{(n)}|^q$  for  $q \geq 1$  is  $r$ -convex function on  $[a, b]$ , then the following inequality holds:

$$|I(a, b, n, f)| \leq \begin{cases} \frac{1}{n!} (b-a)^{1-\frac{1}{q}-\frac{1}{qr}} L_n^{\left(\frac{q-1}{q}\right)}(a, b) \left[ C_1 |f^{(n)}(b)|^q + C_2 |f^{(n)}(a)|^q \right]^{\frac{1}{q}}, & r \geq 1 \\ \frac{1}{n!} (b-a)^{1-\frac{1}{q}-\frac{1}{qr}} L_n^{\left(\frac{q-1}{q}\right)}(a, b) \left[ C_1^r |f^{(n)}(b)|^{qr} + C_2^r |f^{(n)}(a)|^{qr} \right]^{\frac{1}{qr}}, & r \leq 1 \end{cases}$$

where

$$C_1 = C_1(a, b, r, n) = \int_a^b x^n (x-a)^{\frac{1}{r}} dx, \quad C_2 = C_2(a, b, r, n) = \int_a^b x^n (b-x)^{\frac{1}{r}} dx.$$

*Proof.* From Lemma 2 and Power-Mean integral inequality, we get

$$\begin{aligned} & |I(a, b, n, f)| \\ & \leq \frac{1}{n!} \int_a^b x^n |f^{(n)}(x)| dx \\ & \leq \frac{1}{n!} \left( \int_a^b x^n dx \right)^{1-\frac{1}{q}} \left( \int_a^b x^n |f^{(n)}(x)|^q dx \right)^{\frac{1}{q}} \\ & \leq \frac{1}{n!} \left( \int_a^b x^n dx \right)^{1-\frac{1}{q}} \left( \int_a^b x^n \left[ \frac{x-a}{b-a} |f^{(n)}(b)|^{qr} + \frac{b-x}{b-a} |f^{(n)}(a)|^{qr} \right]^{\frac{1}{r}} dx \right)^{\frac{1}{q}}. \end{aligned}$$

Here, using Lemma 1 we obtain respectively,

For  $r \geq 1$

$$\begin{aligned} & |I(a, b, n, f)| \\ & \leq \frac{1}{n!} \left( \int_a^b x^n dx \right)^{1-\frac{1}{q}} \times \left( \int_a^b x^n \left\{ \left[ \frac{x-a}{b-a} |f^{(n)}(b)|^{qr} \right]^{\frac{1}{r}} + \left[ \frac{b-x}{b-a} |f^{(n)}(a)|^{qr} \right]^{\frac{1}{r}} \right\} dx \right)^{\frac{1}{q}} \\ & = \frac{1}{n!} \left( \frac{1}{b-a} \right)^{\frac{1}{qr}} \left( \frac{b^{n+1} - a^{n+1}}{n+1} \right)^{1-\frac{1}{q}} \end{aligned}$$

$$\begin{aligned}
& \times \left( \int_a^b x^n (x-a)^{\frac{1}{r}} |f^{(n)}(b)|^q dx + \int_a^b x^n (b-x)^{\frac{1}{r}} |f^{(n)}(a)|^q dx \right)^{\frac{1}{q}} \\
& = \frac{1}{n!} \left( \frac{1}{b-a} \right)^{\frac{1}{qr}} (b-a)^{1-\frac{1}{q}} \left( \frac{b^{n+1}-a^{n+1}}{(n+1)(b-a)} \right)^{1-\frac{1}{q}} \\
& \quad \times \left[ C_1(a, b, r, n) |f^{(n)}(b)|^q + C_2(a, b, q, n) |f^{(n)}(a)|^q \right]^{\frac{1}{q}} \\
& = \frac{1}{n!} (b-a)^{1-\frac{1}{q}-\frac{1}{qr}} L_n^{\left(\frac{q-1}{q}\right)}(a, b) \left[ C_1 |f^{(n)}(b)|^q + C_2 |f^{(n)}(a)|^q \right]^{\frac{1}{q}},
\end{aligned}$$

For  $r \leq 1$ , using Minkowski inequality, we have

$$\begin{aligned}
& |I(a, b, n, f)| \\
& \leq \frac{1}{n!} \left( \int_a^b x^n dx \right)^{1-\frac{1}{q}} \left( \frac{1}{b-a} \right)^{\frac{1}{qr}} \\
& \quad \times \left( \int_a^b \left[ x^{nr} (x-a) |f^{(n)}(b)|^{qr} + x^{nr} (b-x) |f^{(n)}(a)|^{qr} \right]^{\frac{1}{r}} dx \right)^{\frac{1}{q}} \\
& = \frac{1}{n!} \left( \frac{b^{n+1}-a^{n+1}}{n+1} \right)^{1-\frac{1}{q}} \\
& \quad \times \left( \left\{ \left[ \int_a^b x^n (x-a)^{\frac{1}{r}} |f^{(n)}(b)|^q dx \right]^r + \left[ \int_a^b x^n (b-x)^{\frac{1}{r}} |f^{(n)}(a)|^q dx \right]^r \right\}^{\frac{1}{r}} \right)^{\frac{1}{q}} \\
& = \frac{1}{n!} (b-a)^{1-\frac{1}{q}-\frac{1}{qr}} L_n^{\left(\frac{q-1}{q}\right)}(a, b) \left[ C_1^r |f^{(n)}(b)|^{qr} + C_2^r |f^{(n)}(a)|^{qr} \right]^{\frac{1}{qr}}.
\end{aligned}$$

This completes the proof of theorem.  $\square$

**Corollary 2.** Under the conditions Theorem 4 for  $n = 1$  we have the following inequalities:

$$|J(a, b, f)| \leq \begin{cases} A^{1-\frac{1}{q}}(a, b) \left\{ \frac{r(b-a)}{2r+1} [ |f'(b)|^q - |f'(a)|^q ] + \frac{r[|f'(b)|^q + |f'(a)|^q]}{r+1} \right\}^{\frac{1}{q}}, & r \geq 1 \\ A^{1-\frac{1}{q}}(a, b) \left\{ \left( \frac{r^2(a+b)+br}{(r+1)(2r+1)} \right)^r |f'(b)|^{qr} + \left( \frac{r^2(a+b)+ar}{(r+1)(2r+1)} \right)^r |f'(a)|^{qr} \right\}^{\frac{1}{qr}}, & r \leq 1 \end{cases}$$

where  $J(a, b, f) = \frac{I(a, b, 1, f)}{b-a}$ .

**Proposition 2.** Let  $a, b \in (0, \infty)$  with  $a < b$ ,  $q \geq 1$  and  $m \geq 1$ , we have the following inequalities:

$$L^{\frac{m}{q}+1}(a, b) \leq \begin{cases} A^{1-\frac{1}{q}}(a, b) \left[ \frac{2rA(a^{m+1}, b^{m+1})}{2r+1} + \frac{2r^2G^2(a, b)A(a^{m-1}, b^{m-1})}{(r+1)(2r+1)} \right]^{\frac{1}{q}}, & r \geq 1 \\ A^{1-\frac{1}{q}}(a, b) \left\{ \left( \frac{r^2(a+b)+br}{(r+1)(2r+1)} \right)^r b^{mr} + \left( \frac{r^2(a+b)+ar}{(r+1)(2r+1)} \right)^r a^{mr} \right\}^{\frac{1}{qr}}, & r \leq 1. \end{cases}$$

*Proof.* The result follows directly from Corollary 2 for function  $f(x) = \frac{q}{m+q}x^{\frac{m}{q}+1}$ ,  $x \in (0, \infty)$ . □

**Corollary 3.** Using Proposition 2 for  $m = 1$ , we have following inequalities:

$$L^{\frac{1}{q}+1}(a, b) \leq \begin{cases} A^{1-\frac{1}{q}}(a, b) \left[ \frac{2r}{2r+1}A(a^2, b^2) + \frac{2r^2}{(r+1)(2r+1)}G^2(a, b) \right]^{\frac{1}{q}}, & r \geq 1 \\ A^{1-\frac{1}{q}}(a, b) \left\{ \left( \frac{r^2(a+b)+br}{(r+1)(2r+1)} \right)^r b^r + \left( \frac{r^2(a+b)+ar}{(r+1)(2r+1)} \right)^r a^r \right\}^{\frac{1}{qr}}, & r \leq 1. \end{cases}$$

**Corollary 4.** Using Proposition 2 for  $q = 1$ , we have following inequalities:

$$L^{m+1}(a, b) \leq \begin{cases} \frac{2rA(a^{m+1}, b^{m+1})}{2r+1} + \frac{2r^2G^2(a, b)A(a^{m-1}, b^{m-1})}{(r+1)(2r+1)}, & r \geq 1 \\ \left\{ \left( \frac{r^2(a+b)+br}{(r+1)(2r+1)} \right)^r b^{mr} + \left( \frac{r^2(a+b)+ar}{(r+1)(2r+1)} \right)^r a^{mr} \right\}^{\frac{1}{r}}, & r \leq 1. \end{cases}$$

**Corollary 5.** Using Corollary 4. for  $m = 1$ , we have following inequalities:

$$L_2^2(a, b) \leq \begin{cases} \frac{2r}{2r+1}A(a^2, b^2) + \frac{2r^2}{(r+1)(2r+1)}G^2(a, b), & r \geq 1 \\ \left\{ \left( \frac{r^2(a+b)+br}{(r+1)(2r+1)} \right)^r b^r + \left( \frac{r^2(a+b)+ar}{(r+1)(2r+1)} \right)^r a^r \right\}^{\frac{1}{r}}, & r \leq 1. \end{cases}$$

**Corollary 6.** Under the conditions Theorem 4 for  $q = 1$  we have the following inequalities:

$$|I(a, b, n, f)| \leq \begin{cases} \frac{1}{n!}(b-a)^{-\frac{1}{r}} \left[ C_1 |f^{(n)}(b)| + C_2 |f^{(n)}(a)| \right], & r \geq 1 \\ \frac{(b-a)^{-\frac{1}{r}}}{n!} \left[ C_1^r |f^{(n)}(b)|^r + C_2^r |f^{(n)}(a)|^r \right]^{\frac{1}{r}}, & r \leq 1 \end{cases}$$

**Theorem 5.** For  $n \in \mathbb{N}$ ; let  $f : I \subset (0, \infty) \rightarrow \mathbb{R}$  be  $n$ -times differentiable function on  $I^\circ$ ,  $r > 0$  and  $a, b \in I^\circ$  with  $a < b$ . If  $f^{(n)} \in L[a, b]$  and  $|f^{(n)}|^q$  for  $q > 1$  is  $r$ -convex function on  $[a, b]$ , then the following inequality holds:

$$|I(a, b, n, f)| \leq \begin{cases} \frac{1}{n!}(b-a)^{\frac{1}{p}-\frac{1}{qr}} \left( |f^{(n)}(b)|^q D_1 + |f^{(n)}(a)|^q D_2 \right)^{\frac{1}{q}}, & r \geq 1 \\ \frac{1}{n!}(b-a)^{\frac{1}{p}-\frac{1}{qr}} \left( |f^{(n)}(b)|^{qr} D_1^r + |f^{(n)}(a)|^{qr} D_2^r \right)^{\frac{1}{qr}}, & r \leq 1 \end{cases}$$

where

$$D_1 = D_1(a, b, r, n, q) = \int_a^b x^{nq}(x-a)^{\frac{1}{r}} dx$$

$$D_2 = D_2(a, b, r, n, q) = \int_a^b x^{nq} (b-x)^{\frac{1}{r}} dx.$$

*Proof.* Since  $|f^{(n)}|^q$  for  $q > 1$  is  $r$ -convex function on  $[a, b]$ , using Lemma 2 and the Hölder integral inequality, we have the following inequality:

$$\begin{aligned} & |I(a, b, n, f)| \\ & \leq \frac{1}{n!} \int_a^b 1 \cdot x^n |f^{(n)}(x)| dx \\ & \leq \frac{1}{n!} \left( \int_a^b 1^p dx \right)^{\frac{1}{p}} \left( \int_a^b x^{nq} |f^{(n)}(x)|^q dx \right)^{\frac{1}{q}} \\ & \leq \frac{1}{n!} \left( \int_a^b 1^p dx \right)^{\frac{1}{p}} \left( \int_a^b x^{nq} \left[ \frac{x-a}{b-a} |f^{(n)}(b)|^{qr} + \frac{b-x}{b-a} |f^{(n)}(a)|^{qr} \right]^{\frac{1}{r}} dx \right)^{\frac{1}{q}}. \end{aligned}$$

Here, using Lemma 1 we obtain respectively,

For  $r \geq 1$ ,

$$\begin{aligned} & |I(a, b, n, f)| \\ & \leq \frac{1}{n!} \left( \int_a^b 1^p dx \right)^{\frac{1}{p}} \left( \int_a^b x^{nq} \left[ \left( \frac{x-a}{b-a} \right)^{\frac{1}{r}} |f^{(n)}(b)|^q + \left( \frac{b-x}{b-a} \right)^{\frac{1}{r}} |f^{(n)}(a)|^q \right] dx \right)^{\frac{1}{q}} \\ & = \frac{(b-a)^{-\frac{1}{qr}}}{n!} (b-a)^{\frac{1}{p}} \left( |f^{(n)}(b)|^q D_1 + |f^{(n)}(a)|^q D_2 \right)^{\frac{1}{q}} \\ & = \frac{1}{n!} (b-a)^{\frac{1}{p} - \frac{1}{qr}} \left( |f^{(n)}(b)|^q D_1 + |f^{(n)}(a)|^q D_2 \right)^{\frac{1}{q}}, \end{aligned}$$

For  $r \leq 1$ , using Minkowski inequality, we have

$$\begin{aligned} & |I(a, b, n, f)| \\ & \leq \frac{1}{n!} (b-a)^{\frac{1}{p} - \frac{1}{qr}} \left( \int_a^b \left[ x^{nqr} (x-a) |f^{(n)}(b)|^{qr} + x^{nqr} (b-x) |f^{(n)}(a)|^{qr} \right]^{\frac{1}{r}} dx \right)^{\frac{1}{q}} \\ & = \frac{1}{n!} (b-a)^{\frac{1}{p} - \frac{1}{qr}} \\ & \quad \times \left\{ \left( |f^{(n)}(b)|^q \int_a^b x^{nq} (x-a)^{\frac{1}{r}} dx \right)^r + \left( |f^{(n)}(a)|^q \int_a^b x^{nq} (b-x)^{\frac{1}{r}} dx \right)^r \right\}^{\frac{1}{qr}} \\ & = \frac{1}{n!} (b-a)^{\frac{1}{p} - \frac{1}{qr}} \left( |f^{(n)}(b)|^{qr} D_1^r + |f^{(n)}(a)|^{qr} D_2^r \right)^{\frac{1}{qr}}. \end{aligned}$$



This completes the proof of theorem.  $\square$

**Corollary 7.** Under the conditions Theorem 5 for  $n = 1$  we have the following inequalities:

$$\begin{aligned} & \left| \frac{f(b)b - f(a)a}{b-a} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \begin{cases} (b-a)^{\frac{1}{p} - \frac{1}{qr} - 1} (|f'(b)|^q D_1 + |f'(a)|^q D_2)^{\frac{1}{q}}, & r \geq 1 \\ (b-a)^{\frac{1}{p} - \frac{1}{qr} - 1} (|f'(b)|^{qr} D_1^r + |f'(a)|^{qr} D_2^r)^{\frac{1}{qr}}, & r \leq 1. \end{cases} \end{aligned}$$

**Proposition 3.** Let  $a, b \in (0, \infty)$  with  $a < b$ ,  $q > 1$  and  $m \geq 1$ , we have

$$L^{\frac{m}{q}+1}_{\frac{m}{q}+1}(a, b) \leq \begin{cases} (b-a)^{\frac{1}{p} - \frac{1}{qr} - 1} (b^m D_1 + a^m D_2)^{\frac{1}{q}}, & r \geq 1 \\ (b-a)^{\frac{1}{p} - \frac{1}{qr} - 1} (b^{mr} D_1^r + a^{mr} D_2^r)^{\frac{1}{q}}, & r \leq 1. \end{cases}$$

*Proof.* The result follows directly from Corollary 7 for  $f(x) = \frac{q}{m+q} x^{\frac{m}{q}+1}$ ,  $x \in (0, \infty)$ .  $\square$

**Corollary 8.** For  $m = 1$  from Proposition 3, we obtain the following inequality:

$$L^{\frac{1}{q}+1}_{\frac{1}{q}+1}(a, b) \leq \begin{cases} (b-a)^{\frac{1}{p} - \frac{1}{qr} - 1} (bD_1 + aD_2)^{\frac{1}{q}}, & r \geq 1 \\ (b-a)^{\frac{1}{p} - \frac{1}{qr} - 1} (b^r D_1^r + a^r D_2^r)^{\frac{1}{q}}, & r \leq 1. \end{cases}$$

**Theorem 6.** For  $n \in \mathbb{N}$ ; let  $f : I \subset (0, \infty) \rightarrow \mathbb{R}$  be  $n$ -times differentiable function on  $I^\circ$  (interior of  $I$ ),  $r > 0$  and  $a, b \in I^\circ$  with  $a < b$ . If  $f^{(n)} \in L[a, b]$  and  $|f^{(n)}|^q$  for  $q > 1$  is  $r$ -convex function on  $[a, b]$ , then the following inequalities holds:

$$\begin{aligned} & |I(a, b, n, f)| \\ & \leq \begin{cases} 2^{\frac{1}{qr}} \frac{b-a}{n!} \left(\frac{r}{r+1}\right)^{\frac{1}{q}} L_{np}^n(a, b) A^{\frac{1}{qr}} \left( |f^{(n)}(a)|^{qr}, |f^{(n)}(b)|^{qr} \right), & 0 < r \leq 1, \\ \frac{b-a}{n!} L_{np}^n(a, b) A^{\frac{1}{qr}} \left( |f^{(n)}(a)|^{qr}, |f^{(n)}(b)|^{qr} \right), & r \geq 0, \\ \frac{1}{n!} (b-a) L_{np}^n(a, b) \left( L_r \left( |f^{(n)}(a)|^q, |f^{(n)}(b)|^q \right) \right)^{\frac{1}{q}}, & r > 0, \end{cases} \end{aligned}$$

*Proof.* For  $0 < r \leq 1$ , since  $|f^{(n)}|^q$  for  $q > 1$  is  $r$ -convex function on  $[a, b]$ , with respect to Hermite-Hadamard inequality we have

$$\int_a^b |f^{(n)}(x)|^q dx \leq (b-a) \frac{r}{r+1} \left[ |f^{(n)}(a)|^{qr} + |f^{(n)}(b)|^{qr} \right]^{\frac{1}{r}}.$$

Using Lemma 2 and the Hölder integral inequality we have

$$|I(a, b, n, f)|$$

$$\begin{aligned}
&\leq \frac{1}{n!} \int_a^b x^n |f^{(n)}(x)| dx \\
&\leq \frac{1}{n!} \left( \int_a^b x^{np} dx \right)^{\frac{1}{p}} \left( \int_a^b |f^{(n)}(x)|^q dx \right)^{\frac{1}{q}} \\
&\leq \frac{1}{n!} \left( \int_a^b x^{np} dx \right)^{\frac{1}{p}} \left( (b-a) \frac{r}{r+1} \left[ |f^{(n)}(a)|^{qr} + |f^{(n)}(b)|^{qr} \right]^{\frac{1}{r}} \right)^{\frac{1}{q}} \\
&= 2^{\frac{1}{qr}} \frac{b-a}{n!} \left( \frac{r}{r+1} \right)^{\frac{1}{q}} \left[ \frac{b^{np+1} - a^{np+1}}{(np+1)(b-a)} \right]^{\frac{1}{p}} \left[ \frac{|f^{(n)}(a)|^{qr} + |f^{(n)}(b)|^{qr}}{2} \right]^{\frac{1}{qr}} \\
&= 2^{\frac{1}{qr}} \frac{b-a}{n!} \left( \frac{r}{r+1} \right)^{\frac{1}{q}} L_{np}^n(a, b) A^{\frac{1}{qr}} \left( |f^{(n)}(a)|^{qr}, |f^{(n)}(b)|^{qr} \right).
\end{aligned}$$

For  $r \geq 1$ , since  $|f^{(n)}|^q$  for  $q > 1$  is  $r$ -convex function on  $[a, b]$ , with respect to Theorem 2 we get

$$\int_a^b |f^{(n)}(x)|^q dx \leq (b-a) \left[ \frac{|f^{(n)}(a)|^{qr} + |f^{(n)}(b)|^{qr}}{2} \right]^{\frac{1}{r}}.$$

Using Lemma 2 and the Hölder integral inequality we have

$$\begin{aligned}
|I(a, b, n, f)| &\leq \frac{1}{n!} \int_a^b x^n |f^{(n)}(x)| dx \\
&\leq \frac{1}{n!} \left( \int_a^b x^{np} dx \right)^{\frac{1}{p}} \left( \int_a^b |f^{(n)}(x)|^q dx \right)^{\frac{1}{q}} \\
&= \frac{b-a}{n!} \left[ \frac{b^{np+1} - a^{np+1}}{(np+1)(b-a)} \right]^{\frac{1}{p}} \left[ \frac{|f^{(n)}(a)|^{qr} + |f^{(n)}(b)|^{qr}}{2} \right]^{\frac{1}{qr}} \\
&= \frac{b-a}{n!} L_{np}^n(a, b) A^{\frac{1}{qr}} \left( |f^{(n)}(a)|^{qr}, |f^{(n)}(b)|^{qr} \right).
\end{aligned}$$

For  $r > 0$ , using Lemma 2, Theorem 1 and the Hölder integral inequality we have

$$|I(a, b, n, f)| \leq \frac{1}{n!} \int_a^b x^n |f^{(n)}(x)| dx$$

$$\begin{aligned}
&\leq \frac{1}{n!} \left( \int_a^b x^{np} dx \right)^{\frac{1}{p}} \left( \int_a^b |f^{(n)}(x)|^q dx \right)^{\frac{1}{q}} \\
&\leq \frac{1}{n!} \left( \int_a^b x^{np} dx \right)^{\frac{1}{p}} \left( (b-a) L_r \left( |f^{(n)}(a)|^q, |f^{(n)}(b)|^q \right) \right)^{\frac{1}{q}} \\
&= \frac{1}{n!} (b-a) L_{np}^n(a,b) \left( L_r \left( |f^{(n)}(a)|^q, |f^{(n)}(b)|^q \right) \right)^{\frac{1}{q}}.
\end{aligned}$$

This completes the proof of theorem.  $\square$

**Corollary 9.** Under the conditions Theorem 6 for  $n = 1$  we have the following inequalities:

$$\begin{aligned}
&\left| \frac{f(b)b - f(a)a}{b-a} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\
&\leq \begin{cases} 2^{\frac{1}{qr}} \left( \frac{r}{r+1} \right)^{\frac{1}{q}} L_p(a,b) A^{\frac{1}{qr}} (|f'(a)|^{qr}, |f'(b)|^{qr}), & 0 < r \leq 1 \\ L_p(a,b) A^{\frac{1}{qr}} (|f'(a)|^{qr}, |f'(b)|^{qr}), & r \geq 1 \\ L_p(a,b) (L_r(|f'(a)|^q, |f'(b)|^q))^{\frac{1}{q}}, & r > 0. \end{cases}
\end{aligned}$$

**Proposition 4.** Let  $a, b \in (0, \infty)$  with  $a < b$ ,  $q > 1$  and  $m \in [0, 1]$ , we have

$$L_{\frac{m}{q}+1}^{\frac{m}{q}+1}(a,b) \leq \begin{cases} 2^{\frac{1}{qr}} \left( \frac{r}{r+1} \right)^{\frac{1}{q}} L_p(a,b) A^{\frac{1}{qr}} (a^{mr}, b^{mr}), & 0 < r \leq 1 \\ L_p(a,b) A^{\frac{1}{qr}} (a^{mr}, b^{mr}), & r \geq 1 \\ L_p(a,b) (L_r(a^m, b^m))^{\frac{1}{q}}, & r > 0 \end{cases}$$

Under the assumption of the Proposition, let  $f(x) = \frac{q}{m+q} x^{\frac{m}{q}+1}$ ,  $x \in (0, \infty)$ . Then  $|f'(x)|^q = x^m$  is  $r$ -convex on  $(0, \infty)$  and the result follows directly from Corollary 9.

**Corollary 10.** For  $m = 1$  from Proposition 4, we obtain the following inequalities:

$$L_{\frac{1}{q}+1}^{\frac{1}{q}+1}(a,b) \leq \begin{cases} 2^{\frac{1}{qr}} \left( \frac{r}{r+1} \right)^{\frac{1}{q}} L_p(a,b) A^{\frac{1}{qr}} (a^r, b^r), & 0 < r \leq 1 \\ L_p(a,b) A^{\frac{1}{qr}} (a^r, b^r), & r \geq 1 \\ L_p(a,b) (L_r(a,b))^{\frac{1}{q}}, & r > 0 \end{cases}$$

**Theorem 7.** For  $n \in \mathbb{N}$ ; let  $f : I \subset (0, \infty) \rightarrow \mathbb{R}$  be  $n$ -times differentiable function on  $I^\circ$ ,  $r > 0$  and  $a, b \in I^\circ$  with  $a < b$ . If  $f^{(n)} \in L[a, b]$  and  $|f^{(n)}|^{\frac{q}{r}}$  for  $q > 1$  is  $r$ -convex function on  $[a, b]$ , then the following inequality holds:

$$|I(a, b, n, f)| \leq \frac{1}{n!} (b-a) L_{np}^n(a,b) A^{\frac{1}{q}} \left( |f^{(n)}(a)|^q, |f^{(n)}(b)|^q \right).$$

*Proof.* If  $|f^{(n)}|^{\frac{q}{r}}$  for  $q > 1$  is  $r$ -convex function on  $[a, b]$  and  $r > 0$ , using (1.1) inequality, Lemma 2 and the Hölder integral inequality respectively, we have

$$\int_a^b |f^{(n)}(x)|^q dx = \int_a^b \left( |f^{(n)}(x)|^{\frac{q}{r}} \right)^r dx \leq (b-a) \frac{|f^{(n)}(a)|^q + |f^{(n)}(b)|^q}{2}$$

and

$$\begin{aligned} |I(a, b, n, f)| &\leq \frac{1}{n!} \int_a^b x^n |f^{(n)}(x)| dx \\ &\leq \frac{1}{n!} \left( \int_a^b x^{np} dx \right)^{\frac{1}{p}} \left( \int_a^b |f^{(n)}(x)|^q dx \right)^{\frac{1}{q}} \\ &= \frac{1}{n!} \left( \frac{b^{np+1} - a^{np+1}}{np+1} \right)^{\frac{1}{p}} \left( (b-a) \frac{|f^{(n)}(a)|^q + |f^{(n)}(b)|^q}{2} \right)^{\frac{1}{q}} \\ &= \frac{1}{n!} (b-a) L_{np}^n(a, b) A^{\frac{1}{q}} \left( |f^{(n)}(a)|^q, |f^{(n)}(b)|^q \right). \end{aligned}$$

This completes the proof of theorem.  $\square$

**Corollary 11.** Under the conditions Theorem 7 for  $n = 1$  we have the following inequality:

$$\left| \frac{f(b)b - f(a)a}{b-a} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq L_p(a, b) A^{\frac{1}{q}} \left( |f'(a)|^q, |f'(b)|^q \right).$$

**Proposition 5.** Let  $a, b \in (0, \infty)$  with  $a < b$ ,  $q > 1$  and  $m \in [0, 1]$ , we have

$$\left| \frac{f(b)b - f(a)a}{b-a} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq L_p(a, b) A^{\frac{1}{q}} (a^{mr}, b^{mr}).$$

*Proof.* Under the assumption of the Proposition, let  $f(x) = \frac{q}{mr+q} x^{\frac{mr}{q}+1}$ ,  $x \in (0, \infty)$ . Then  $|f'(x)|^{\frac{q}{r}} = x^m$  is  $r$ -convex on  $(0, \infty)$  and the result follows directly from Corollary 11.  $\square$

**Corollary 12.** For  $m = 1$  from Proposition 5, we obtain the following inequality:

$$\left| \frac{f(b)b - f(a)a}{b-a} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq L_p(a, b) A^{\frac{1}{q}} (a^r, b^r).$$

**Theorem 8.** For  $n \in \mathbb{N}$ ; let  $f : I \subset (0, \infty) \rightarrow \mathbb{R}$  be  $n$ -times differentiable function on  $I^\circ$ ,  $r > 0$  and  $a, b \in I^\circ$  with  $a < b$ . If  $f^{(n)} \in L[a, b]$  and  $|f^{(n)}|^{\frac{q}{r}}$  for  $q > 1$  is

$r$ -concave function on  $[a, b]$ , then the following inequality holds:

$$|I(a, b, n, f)| \leq \frac{b-a}{n!} L_{np}^n(a, b) \left| f^{(n)}\left(\frac{a+b}{2}\right) \right|.$$

*Proof.* If  $|f^{(n)}|^{\frac{q}{r}}$  for  $q > 1$  is  $r$ -concave function on  $[a, b]$  and  $r > 0$ , using Lemma 2, the Hölder integral inequality and

$$\int_a^b |f^{(n)}(x)|^q dx = \int_a^b \left( |f^{(n)}(x)|^{\frac{q}{r}} \right)^r dx \leq (b-a) \left| f^{(n)}\left(\frac{a+b}{2}\right) \right|^q$$

we have

$$\begin{aligned} |I(a, b, n, f)| &\leq \frac{1}{n!} \int_a^b x^n |f^{(n)}(x)| dx \\ &\leq \frac{1}{n!} \left( \int_a^b x^{np} dx \right)^{\frac{1}{p}} \left( \int_a^b |f^{(n)}(x)|^q dx \right)^{\frac{1}{q}} \\ &\leq \frac{1}{n!} \left( \int_a^b x^{np} dx \right)^{\frac{1}{p}} \left( (b-a) \left| f^{(n)}\left(\frac{a+b}{2}\right) \right|^q \right)^{\frac{1}{q}} \\ &= \frac{1}{n!} (b-a) L_{np}^n(a, b) \left| f^{(n)}\left(\frac{a+b}{2}\right) \right|. \end{aligned}$$

This completes the proof of theorem.  $\square$

**Corollary 13.** Under the conditions Theorem 8 for  $n = 1$  we have the following inequality:

$$\left| \frac{f(b)b - f(a)a}{b-a} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq L_p(a, b) \left| f'\left(\frac{a+b}{2}\right) \right|.$$

**Proposition 6.** Let  $a, b \in (0, \infty)$  with  $a < b$ ,  $q > 1$  and  $m \in [0, 1]$ , we have

$$\left| \frac{f(b)b - f(a)a}{b-a} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq L_p(a, b) A^{\frac{mr}{q}}(a, b).$$

*Proof.* Under the assumption of the Proposition, let  $f(x) = \frac{q}{mr+q} x^{\frac{mr}{q}+1}$ ,  $x \in (0, \infty)$ . Then  $|f'(x)|^{\frac{q}{r}} = x^m$  is  $r$ -concave on  $(0, \infty)$  and the result follows directly from Corollary 13.  $\square$

**Corollary 14.** For  $m = 1$  from Proposition 6, we obtain the following inequality:

$$\left| \frac{f(b)b - f(a)a}{b-a} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq L_p(a, b) A^{\frac{r}{q}}(a, b).$$

## REFERENCES

- [1] M. Bessenyei, “Hermite-Hadamard-type inequalities for generalized 3-convex functions.” *Publ. Math. (Debr.)*, vol. 65, no. 15, pp. 223–232, 2004.
- [2] F. Chen and X. Liu, “Refinements on the Hermite-Hadamard Inequalities for  $r$ -Convex Functions.” *J. Appl. Math.*, vol. 2013, no. 1-2, p. 5, 2013, doi: [10.1155/2013/978493](https://doi.org/10.1155/2013/978493).
- [3] S. Dragomir, “Refinements of the Hermite-Hadamard integral inequality for log-convex functions.” *Aust. Math. Soc. Gaz.*, vol. 28, no. 3, pp. 129–134, 2001.
- [4] S. Dragomir and C. Pearce, *Selected Topics on Hermite-Hadamard Inequalities and Its Applications*. Victoria University: RGMIA Monograph, 2002.
- [5] P. Gill, C. Pearce, and J. Pečarić, “Hadamard’s Inequality for  $r$ -Convex Functions.” *J. Math. Anal. Appl.*, vol. 215, no. AY975645, pp. 461–470, 1997.
- [6] J. Hadamard, “Etude sur les proprietes des fonctions entieres en particulier d’une fonction consideree par Riemann.” *J. Math. Pures Appl.*, no. 58, pp. 171–216, 1893.
- [7] L. Han and L. Liu, “Integral Inequalities of Hermite-Hadamard Type for  $r$ -Convex Functions.” *Appl. Math.*, no. 3, pp. 1967–1971, 2012, doi: [10.4236/am.2012.312270](https://doi.org/10.4236/am.2012.312270).
- [8] I. İşcan, “Ostrowski type inequalities for  $p$ -convex functions.” *New Trends in Mathematical Sciences (Ntmsci)*, vol. 4, no. 3, pp. 140–150, 2016, doi: [10.20852/ntmsci.2016318838](https://doi.org/10.20852/ntmsci.2016318838).
- [9] I. İşcan, H. Kadakal, and M. Kadakal, “Some new integral inequalities for  $n$ -times differentiable log-convex functions.” *New Trends in Mathematical Sciences (Ntmsci)*, vol. 5, no. 2, pp. 10–15, 2017, doi: [10.20852/ntmsci.2017.150](https://doi.org/10.20852/ntmsci.2017.150).
- [10] I. İşcan and M. Kunt, “Hermite-Hadamard-Fejer type inequalities for quasi-geometrically convex functions via fractional integrals.” *J. Math.*, vol. 2016, no. 6523041, p. 7, 2016, doi: [10.1155/2016/6523041](https://doi.org/10.1155/2016/6523041).
- [11] I. İşcan and S. Turhan, “Generalized Hermite-Hadamard-Fejer type inequalities for GA-convex functions via Fractional integral.” *Moroccan J. Pure and Appl. Anal.(MJPA)*, vol. 2, no. 1, pp. 34–46, 2016, doi: [10.7603/s40956-016-0004-2](https://doi.org/10.7603/s40956-016-0004-2).
- [12] H. Kadakal, M. Kadakal, and I. İşcan, “Some new integral inequalities for  $n$ -times differentiable  $s$ -convex and  $s$ -concave functions in the second sense.” *Mathematics and Statistic.*, vol. 5, no. 2, pp. 94–98, 2017, doi: [10.13189/ms.2017.050207](https://doi.org/10.13189/ms.2017.050207).
- [13] M. Kadakal, H. Kadakal, and I. İşcan, “Some new integral inequalities for  $n$ -times differentiable  $s$ -convex functions in the first sense.” *Turkish Journal of Analysis and Number Theory (Tjant)*, vol. 5, no. 2, pp. 63–68, 2017, doi: [10.12691/tjant-5-2-4](https://doi.org/10.12691/tjant-5-2-4).
- [14] M. Maden, H. Kadakal, M. Kadakal, and I. İşcan, “Some new integral inequalities for  $n$ -times differentiable convex and concave functions.” *J. Nonlinear Sci. Appl.*, vol. 10, no. 12, pp. 6141–6148, 2017, doi: [10.22436/jnsa.010.12.01](https://doi.org/10.22436/jnsa.010.12.01).
- [15] B. Mihaly, “Hermite-Hadamard-type inequalities for generalized convex functions.” *J. inequal. pure and appl. math*, vol. 9, no. 3, p. 51, 2008.
- [16] N. Ngoc, N. Vinh, and P. Hien, “Integral inequalities of Hadamard type for  $r$ -convex functions.” *Int. Math. Forum*, vol. 4, no. 35, pp. 1723–1728, 2009.
- [17] C. Pearce, J. Pečarić, and V. Šimić, “Stolarsky means and Hadamard’s inequality.” *J. Math. Anal. Appl.*, vol. 220, no. 1, pp. 99–109, 1998, doi: [10.1006/jmaa.1997.5822](https://doi.org/10.1006/jmaa.1997.5822).
- [18] B. Uhrin, “Some remarks about the convolution of unimodal functions.” *Ann. Probab.*, vol. 12, no. 2, pp. 640–645, 1984, doi: [10.1214/aop/1176993312](https://doi.org/10.1214/aop/1176993312).
- [19] G. Yang and D. Hwang, “Refinements of Hadamard inequality for  $r$ -convex functions.” *Indian J. Pure Appl. Math.*, vol. 32, no. 10, pp. 1571–1579, 2001.
- [20] G. Zabandan, “A new refinement of the Hermite-Hadamard inequality for convex functions.” *J. Inequal. Pure Appl. Math.*, vol. 10, no. 2, p. 7, 2009.

- [21] G. Zabandan, A. Bodagh, and A. Kilicman, "The Hermite-Hadamard inequality for  $r$ -convex functions." *Journal of Inequalities and Applications*, vol. 2012, no. 1, p. 215, 2012, doi: [10.1186/1029-242X-2012-215](https://doi.org/10.1186/1029-242X-2012-215).

*Authors' addresses*

**Huriye Kadakal**

Ministry of Education, Giresun Hamdi Bozbağ Anatolian High School Giresun-TÜRKİYE

*E-mail address:* huriyekadakal@hotmail.com

**Mahir Kadakal**

Department of Mathematics, Faculty of Sciences and Arts, Giresun University, Giresun-TÜRKİYE

*E-mail address:* mahirkadakal@gmail.com

**İmdat İşcan**

Department of Mathematics, Faculty of Sciences and Arts, Giresun University, Giresun-TÜRKİYE

*E-mail address:* imdat.iscan@giresun.edu.tr