



## ON HERMITE-HADAMARD TYPE INEQUALITIES FOR MULTIPLICATIVE FRACTIONAL INTEGRALS

H. BUDAK AND K. ÖZÇELİK

*Received 17 November, 2019*

*Abstract.* In this study, we first establish two Hermite-Hadamard type inequality for multiplicative (geometric) Riemann-Liouville fractional integrals. Then, by using some properties of multiplicative convex function, we give some new inequalities involving multiplicative fractional integrals.

2010 *Mathematics Subject Classification:* 26D07; 26D10; 26D15; 26A33

*Keywords:* Hermite-Hadamard inequality, integral inequalities, bounded functions

### 1. INTRODUCTION

The inequalities discovered by C. Hermite and J. Hadamard for convex functions are considerable significant in the literature (see, e.g., [8], [11], [18, p.137]). These inequalities state that if  $f : I \rightarrow \mathbb{R}$  is a convex function on the interval  $I$  of real numbers and  $a, b \in I$  with  $a < b$ , then

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a)+f(b)}{2}. \quad (1.1)$$

Both inequalities hold in the reversed direction if  $f$  is concave. We note that Hadamard's inequality may be regarded as a refinement of the concept of convexity and it follows easily from Jensen's inequality.

Over the last twenty years, the numerous studies have focused on to obtain new bound for left hand side and right and side of the inequality (1.1). For some examples, please refer to ([3], [5], [6], [7], [8], [9], [10], [12], [13], [14], [15], [16], [17], [19], [20], [21], [22], [23], [24], [25]).

#### 1.1. *Multiplicative Calculus*

Recall that the concept of the multiplicative integral called \* integral is denoted by  $\int_a^b (f(x))^{dx}$  which introduced by Bashirov et al. in [4]. While the sum of the terms of product is used in the definition of a classical Riemann integral of  $f$  on  $[a, b]$ , the

product of terms raised to power is used in the definition multiplicative integral of  $f$  on  $[a, b]$ .

There is the following relation between Riemann integral and multiplicative integral [4]:

**Proposition 1.** *If  $f$  is Riemann integrable on  $[a, b]$ , then  $f$  is multiplicative integrable on  $[a, b]$  and*

$$\int_a^b (f(x))^{dx} = e^{\int_a^b \ln(f(x)) dx}.$$

Moreover, Bashirov et al [4] show that multiplicative integrable has the following results and properties:

**Proposition 2.** *If  $f$  is positive and Riemann integrable on  $[a, b]$ , then  $f$  is  $*$ integrable on  $[a, b]$  and*

- (i)  $\int_a^b ((f(x))^p)^{dx} = \int_a^b ((f(x))^{dx})^p,$
- (ii)  $\int_a^b (f(x)g(x))^{dx} = \int_a^b (f(x))^{dx} \cdot \int_a^b (g(x))^{dx},$
- (iii)  $\int_a^b \left(\frac{f(x)}{g(x)}\right)^{dx} = \frac{\int_a^b (f(x))^{dx}}{\int_a^b (g(x))^{dx}},$
- (iv)  $\int_a^b (f(x))^{dx} = \int_a^c (f(x))^{dx} \cdot \int_c^b (f(x))^{dx}, \quad a \leq c \leq b.$
- (v)  $\int_a^a (f(x))^{dx} = 1$  and  $\int_a^b (f(x))^{dx} = \left(\int_b^a (f(x))^{dx}\right)^{-1}.$

On the other hand, Abdeljawed and Grossman [1] introduce the following Multiplicative Riemann-Liouville fractional integrals.

**Definition 1.** The multiplicative left Riemann-Liouville fractional integral  $({}_a I_*^\alpha f)(x)$  of order  $\alpha \in \mathbb{C}$ ,  $Re(\alpha) > 0$  starting from  $\alpha$  is defined by

$$({}_a I_*^\alpha f)(x) = e^{(J_{a^+}^\alpha (\ln \circ f))(x)}$$

and the multiplicative right one is defined by

$$({}_* I_b^\alpha f)(x) = e^{(J_{b^-}^\alpha (\ln \circ f))(x)},$$

Here  $J_{a^+}^\alpha f(x)$  and  $J_{b^-}^\alpha f(x)$  denote the left and right Riemann-Liouville fractional integral, defined by

$$J_{a^+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, x > a$$

and

$$J_{b-}^{\alpha} f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt, x < b$$

respectively.

### 1.2. Hermite-Hadamard inequality and convexity

For the our main results we need to following definition.

**Definition 2.** A non-empty set  $K$  is said to be convex, if for every  $a, b \in K$  we have

$$a + \mu(b - a) \in K, \forall \mu \in [0, 1].$$

**Definition 3.** A function  $f$  is said to be convex function on set  $K$ , if

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y), \forall t \in [0, 1].$$

**Definition 4.** A function  $f$  is said to be *log* or multiplicatively convex function on set  $K$ , if

$$f(tx + (1-t)y) \leq [f(x)]^t [f(y)]^{1-t}, \forall t \in [0, 1].$$

**Proposition 3.** If  $f$  and  $g$  are *log* (multiplicatively) convex functions, then the functions  $fg$  and  $\frac{f}{g}$  are *log* (multiplicatively) convex functions.

The classical Hermite-Hadamard inequality for convex function given by the inequality (1.1).

Hermite-Hadamard inequality for multiplicatively convex function is proved by Ali et al. in [2] as follows:

**Theorem 1.** Let  $f$  be a positive and multiplicatively convex function on interval  $[a, b]$ , then the following inequalities hold

$$f\left(\frac{a+b}{2}\right) \leq \left(\int_a^b (f(x))^{dx}\right)^{\frac{1}{b-a}} \leq G(f(a), f(b)), \quad (1.2)$$

where  $G(.,.)$  is a geometric mean.

The inequality (1.2) is the same result proved by Dragomir in [8, Page 197, inequality (5.3)]

On the other hand, Sarikaya et al. [22] proved following important inequality which is the Hermite-Hadamard inequality for Riemann-Liouville fractional integrals.

**Theorem 2.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be a positive function with  $0 \leq a < b$  and  $f \in L_1[a, b]$ . If  $f$  is a convex function on  $[a, b]$ , then the following inequalities for fractional integrals hold:

$$f\left(\frac{a+b}{2}\right) \leq \frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}} [J_{a+}^{\alpha} f(b) + J_{b-}^{\alpha} f(a)] \leq \frac{f(a) + f(b)}{2} \quad (1.3)$$

with  $\alpha > 0$ .

In this paper, we establish Hermite-Haddamard inequality for multiplicative Riemann-Liouville fractional integrals.

## 2. MAIN RESULTS

In this section we obtain some Hermite-Hadamard type inequalities for multiplicatively convex function via multiplicative Riemann-Liouville fractional integrals.

**Theorem 3.** *Let  $f$  be a positive and multiplicatively convex function on interval  $[a, b]$ , Then we have the following Hermite-Haddamard inequality for multiplicative Riemann-Liouville fractional integrals*

$$f\left(\frac{a+b}{2}\right) \leq [{}_a I_*^\alpha f(b) \cdot {}_b I_*^\alpha f(a)]^{\frac{\Gamma(\alpha+1)}{2(b-a)^\alpha}} \leq G(f(a), f(b)) \quad (2.1)$$

where  $G(.,.)$  is a geometric mean.

*Proof.* Since  $f$  is multiplicatively convex function on interval  $[a, b]$ , then we have

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &= f\left(\frac{at + (1-t)b + (1-t)a + tb}{2}\right) \\ &\leq [f(at + (1-t)b)]^{\frac{1}{2}} [f((1-t)a + tb)]^{\frac{1}{2}}, \end{aligned}$$

i.e.

$$\ln f\left(\frac{a+b}{2}\right) = \frac{1}{2} \ln f(at + (1-t)b) + \frac{1}{2} \ln f((1-t)a + tb) \quad (2.2)$$

Multiplying both sides of (2.2) by  $t^{\alpha-1}$  then integrating the resulting inequality with respect to  $t$  over  $[0, 1]$ , we obtain

$$\int_0^1 t^{\alpha-1} \ln f\left(\frac{a+b}{2}\right) dt \leq \frac{1}{2} \int_0^1 t^{\alpha-1} \ln f(at + (1-t)b) dt + \frac{1}{2} \int_0^1 t^{\alpha-1} \ln f((1-t)a + tb) dt.$$

By using the change variable, we have

$$\frac{1}{\alpha} \ln f\left(\frac{a+b}{2}\right) dt \leq \frac{1}{(b-a)^\alpha} \int_a^b (b-x)^{\alpha-1} \ln f(x) dx + \frac{1}{2} \frac{1}{(b-a)^\alpha} \int_a^b (x-a)^{\alpha-1} \ln f(x) dx.$$

That is,

$$\ln f\left(\frac{a+b}{2}\right) \leq \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{b^-}^\alpha \ln f(a) + J_{a^+}^\alpha \ln f(b)].$$

Thus we get,

$$f\left(\frac{a+b}{2}\right) \leq e^{\frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a^+}^\alpha \ln f(b) + J_{b^-}^\alpha \ln f(a)]}$$

$$\begin{aligned}
&= [e^{J_{a^+}^\alpha \ln f(b)} \cdot e^{J_{b^-}^\alpha \ln f(a)}]_{\frac{\Gamma(\alpha+1)}{2(b-a)^\alpha}} \\
&= [{}_a I_*^\alpha f(b) \cdot {}_b I^\alpha f(a)]_{\frac{\Gamma(\alpha+1)}{2(b-a)^\alpha}}.
\end{aligned}$$

which completes the proof of the first inequality in (2.1).

As  $f$  is multiplicatively convex function on interval  $[a, b]$ , then we have

$$f(at + (1-t)b) \leq [f(a)]^t [f(b)]^{1-t}$$

$$f((1-t)a + tb) \leq [f(a)]^{1-t} [f(b)]^t$$

i.e.

$$\ln f(at + (1-t)b) + \ln f((1-t)a + tb) \tag{2.3}$$

$$\leq t \ln f(a) + (1-t) \ln f(b) + (1-t) \ln f(a) + t \ln f(b)$$

$$= \ln f(a) + \ln f(b)$$

Multiplying both sides of (2.3) by  $t^{\alpha-1}$  then integrating the resulting inequality with respect to  $t$  over  $[0, 1]$ , we establish

$$\int_0^1 t^{\alpha-1} \ln f(at + (1-t)b) dt + \int_0^1 t^{\alpha-1} \ln f((1-t)a + tb) dt \leq \frac{\ln f(a) + \ln f(b)}{\alpha}.$$

Hence,

$$\frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{b^-}^\alpha f(a) + J_{a^+}^\alpha f(b)] \leq \frac{1}{2} \ln[f(a) \cdot f(b)].$$

Thus, we have the inequality

$$e^{[J_{b^-}^\alpha \ln f(a) + J_{a^+}^\alpha \ln f(b)]_{\frac{\Gamma(\alpha+1)}{2(b-a)^\alpha}}} \leq \sqrt{f(a) \cdot f(b)}$$

and

$$[{}_a I_*^\alpha f(b) \cdot {}_b I^\alpha f(a)]_{\frac{\Gamma(\alpha+1)}{2(b-a)^\alpha}} \leq \sqrt{f(a) \cdot f(b)}.$$

The proof is completed □

*Remark 1.* If we choose  $\alpha = 1$  in Theorem 3, then Theorem 3 reduces to the Theorem 1.

**Corollary 1.** *If  $f$  and  $g$  are two positive and multiplicative convex functions, then we have the following inequality*

$$f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \leq [{}_a I_*^\alpha f g(b) \cdot {}_b I_*^\alpha f g(a)]^{\frac{\Gamma(\alpha+1)}{2(b-a)^\alpha}} \leq G(f(a), f(b)) \cdot G(g(a), g(b)). \quad (2.4)$$

*Proof.* Since  $f$  and  $g$  are positive and multiplicative convex functions, then  $fg$  is positive and multiplicative convex function. Thus if we apply Theorem 3 to the function  $fg$ , then we obtain the desired inequality (2.4).  $\square$

*Remark 2.* If we take  $\alpha = 1$  in Corollary 1, then we have the following inequality

$$f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \leq \left(\int_a^b (f(x))^{dx} \cdot \int_a^b (g(x))^{dx}\right)^{\frac{1}{b-a}} \leq G(f(a), f(b)) \cdot G(g(a), g(b)) \quad (2.5)$$

which is given by Ali et al. in [2].

**Theorem 4.** *Let  $f$  be a positive and multiplicatively convex function on interval  $[a, b]$ , Then we have the following Hermite-Hadamard inequality for multiplicative Riemann-Liouville fractional integrals*

$$f\left(\frac{a+b}{2}\right) \leq \left[ {}_{\frac{a+b}{2}} I_*^\alpha f(b) \cdot {}_{\frac{a+b}{2}} I_*^\alpha f(a) \right]^{\frac{2^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^\alpha}} \leq G(f(a), f(b)) \quad (2.6)$$

where  $G(.,.)$  is a geometric mean.

*Proof.* By using the multiplicatively convexity of  $f$ , we have

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &= f\left[\frac{1}{2}\left(\frac{t}{2}a + \frac{2-t}{2}b\right) + \frac{1}{2}\left(\frac{2-t}{2}a + \frac{t}{2}b\right)\right] \\ &\leq \left(f\left(\frac{t}{2}a + \frac{2-t}{2}b\right)\right)^{\frac{1}{2}} \cdot \left(f\left(\frac{2-t}{2}a + \frac{t}{2}b\right)\right)^{\frac{1}{2}}, \end{aligned}$$

i.e.

$$\ln f\left(\frac{a+b}{2}\right) \leq \frac{1}{2} \left[ \ln f\left(\frac{t}{2}a + \frac{2-t}{2}b\right) + \ln f\left(\frac{2-t}{2}a + \frac{t}{2}b\right) \right]. \quad (2.7)$$

Multiplying both sides of (2.7) by  $t^{\alpha-1}$  then integrating the resulting inequality with respect to  $t$  over  $[0, 1]$ , we get

$$\frac{1}{\alpha} \ln f\left(\frac{a+b}{2}\right) \leq \frac{1}{2} \int_0^1 t^{\alpha-1} \ln f\left(\frac{t}{2}a + \frac{2-t}{2}b\right) dt + \int_0^1 t^{\alpha-1} \ln f\left(\frac{2-t}{2}a + \frac{t}{2}b\right) dt$$

$$\begin{aligned}
&= \frac{\Gamma(\alpha)}{2} \left[ \int_{\frac{a+b}{2}}^b \left( \frac{2(b-x)}{b-a} \right)^{\alpha-1} \ln f(x) \frac{2}{b-a} dx \right. \\
&\quad \left. + \int_a^{\frac{a+b}{2}} \left( \frac{2(x-a)}{b-a} \right)^{\alpha-1} \ln f(x) \cdot \frac{2}{b-a} dx \right] \\
&= \frac{2^{\alpha-1} \Gamma(\alpha)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^+}^\alpha \ln f(b) + J_{\left(\frac{a+b}{2}\right)^-}^\alpha \ln f(a) \right].
\end{aligned}$$

Then it follows that,

$$\begin{aligned}
f\left(\frac{a+b}{2}\right) &\leq e^{\frac{2^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^\alpha}} \left[ J_{\left(\frac{a+b}{2}\right)^+}^\alpha \ln f(b) + J_{\left(\frac{a+b}{2}\right)^-}^\alpha \ln f(a) \right] \\
&= \left[ {}_{\frac{a+b}{2}} I_*^\alpha f(b) \cdot {}_{\frac{a+b}{2}} I_*^\alpha f(a) \right]^{\frac{2^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^\alpha}}.
\end{aligned}$$

This completes the proof of first inequality in the inequality (2.6).

On the other hand, since  $f$  is multiplicatively convex function, we get

$$f\left(\frac{t}{2}a + \frac{2-t}{2}b\right) + f\left(\frac{2-t}{2}a + \frac{t}{2}b\right) \leq [f(a)]^{\frac{t}{2}} + [f(b)]^{\frac{2-t}{2}} + [f(a)]^{\frac{2-t}{2}} + [f(b)]^{\frac{t}{2}}.$$

Thus, we have

$$\ln f\left(\frac{t}{2}a + \frac{2-t}{2}b\right) + \ln f\left(\frac{2-t}{2}a + \frac{t}{2}b\right) \leq \ln f(a) + \ln f(b) \quad (2.8)$$

Multiplying both sides of (2.8) by  $t^{\alpha-1}$  then integrating the resulting inequality with respect to  $t$  over  $[0, 1]$ , we have

$$\frac{2^\alpha}{(b-a)^\alpha} \Gamma(\alpha) \left[ J_{\left(\frac{a+b}{2}\right)^+}^\alpha \ln f(b) + J_{\left(\frac{a+b}{2}\right)^-}^\alpha \ln f(a) \right] \leq \frac{\ln f(a) + \ln f(b)}{\alpha}$$

i.e.

$$\frac{2^{\alpha-1}}{(b-a)^\alpha} \Gamma(\alpha+1) \left[ J_{\left(\frac{a+b}{2}\right)^+}^\alpha \ln f(b) + J_{\left(\frac{a+b}{2}\right)^-}^\alpha \ln f(a) \right] \leq \frac{\ln [f(a) \cdot f(b)]}{2}.$$

Hence, we get the inequality

$$\left[ {}_{\frac{a+b}{2}} I_*^\alpha f(b) \cdot {}_{\frac{a+b}{2}} I_*^\alpha f(a) \right]^{\frac{2^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^\alpha}} \leq \sqrt{f(a)f(b)}.$$

The proof is completed  $\square$

**Corollary 2.** *If  $f$  and  $g$  are positive and multiplicative convex function, then we have the following inequality*

$$\begin{aligned} & f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \\ & \leq \left[ \left(\frac{a+b}{2}\right) I_*^\alpha f \cdot g(b) \cdot {}_* I_{\left(\frac{a+b}{2}\right)}^\alpha f \cdot g(a) \right]^{\frac{\Gamma(\alpha+1)}{2(b-a)^\alpha}} \leq G(f(a), f(b))G(g(a), g(b)). \end{aligned} \quad (2.9)$$

*Proof.* The proof is similar to the proof of Corollary 1. □

**Remark 3.** If we take  $\alpha = 1$  in Corollary 2, then the inequality (2.9) reduces to the inequality (2.5).

**Corollary 3.** *If  $f$  and  $g$  are positive and multiplicative convex function, then we get the following inequality*

#### REFERENCES

- [1] T. Abdeljawad and M. Grossman, “On geometric fractional calculus,” *J. Semigroup Theory Appl.*, vol. 2016, pp. Article–ID, 2016.
- [2] M. A. Ali, M. Abbas, Z. Zhang, I. B. Sial, and R. Arif, “On integral inequalities for product and quotient of two multiplicatively convex functions,” *Asian Research Journal of Mathematics*, pp. 1–11, 2019.
- [3] A. G. Azpeitia, “Convex functions and the hadamard inequality,” *Rev. Colombiana Mat.*, vol. 28, no. 1, pp. 7–12, 1994.
- [4] A. E. Bashirov, E. M. Kurpinar, and A. Özyapıcı, “Multiplicative calculus and its applications,” *Journal of Mathematical Analysis and Applications*, vol. 337, no. 1, pp. 36–48, 2008.
- [5] H. Chen and U. N. Katugampola, “Hermite–hadamard and hermite–hadamard–fejér type inequalities for generalized fractional integrals,” *Journal of Mathematical Analysis and Applications*, vol. 446, no. 2, pp. 1274–1291, 2017.
- [6] J. de la Cal, J. Cárcamo, and L. Escauriaza, “A general multidimensional hermite–hadamard type inequality,” *Journal of mathematical analysis and applications*, vol. 356, no. 2, pp. 659–663, 2009.
- [7] S. Dragomir and R. Agarwal, “Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula,” *Applied Mathematics Letters*, vol. 11, no. 5, pp. 91–95, 1998.
- [8] S. Dragomir and C. Pearce, “Selected topics on hermite-hadamard inequalities and applications, rgmia monographs, victoria university, 2000,” *Online: <http://www.staff.vu.edu.au/RGMIA/monographs/hermite-hadamard.html>*, 2004.
- [9] G. FARID, A. U. REHMAN, and M. ZAHRA, “On hadamard-type inequalities for  $k$ -fractional integrals,” *Konuralp Journal of Mathematics*, vol. 4, no. 2, pp. 79–86, 2016.
- [10] G. Farid, A. Rehman, and M. Zahra, “On hadamard inequalities for  $k$ -fractional integrals,” *Non-linear Funct. Anal. Appl.*, vol. 21, no. 3, pp. 463–478, 2016.
- [11] J. Hadamard, “Étude sur les propriétés des fonctions entières et en particulier d’une fonction considérée par riemann,” *Journal de mathématiques pures et appliquées*, pp. 171–216, 1893.
- [12] M. Iqbal, S. Qaisar, and M. Muddassar, “A short note on integral inequality of type hermite–hadamard through convexity,” *J. Comput. Anal. Appl.*, vol. 21, no. 5, pp. 946–953, 2016.
- [13] İ. İşcan, “Generalization of different type integral inequalities for  $s$ -convex functions via fractional integrals,” *Applicable Analysis*, vol. 93, no. 9, pp. 1846–1862, 2014.



- [14] İ. İşcan and S. Wu, "Hermite-hadamard type inequalities for harmonically convex functions via fractional integrals," *Applied Mathematics and Computation*, vol. 238, pp. 237–244, 2014.
- [15] M. Jleli and B. Samet, "On hermite-hadamard type inequalities via fractional integrals of a function with respect to another function," *J. Nonlinear Sci. Appl.*, vol. 9, no. 3, pp. 1252–1260, 2016.
- [16] U. S. Kirmaci, "Inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula," *Applied Mathematics and Computation*, vol. 147, no. 1, pp. 137–146, 2004.
- [17] M. A. Noor and M. U. Awan, "Some integral inequalities for two kinds of convexities via fractional integrals," *Trans. J. Math. Mech.*, vol. 5, no. 2, pp. 129–136, 2013.
- [18] J. E. Pejčarić and Y. L. Tong, *Convex functions, partial orderings, and statistical applications*. Academic Press, 1992.
- [19] A. Saglam, M. Z. Sarikaya, and H. Yildirim, "Some new inequalities of hermite-hadamard's type," *arXiv preprint arXiv:1005.0750*, 2010.
- [20] M. Z. Sarikaya, A. Akkurt, H. Budak, M. E. Yıldırım, and H. Yıldırım, "Hermite-hadamard's inequalities for conformable fractional integrals," *An International Journal of Optimization and Control: Theories & Applications (IJOCTA)*, vol. 9, no. 1, pp. 49–59, 2019.
- [21] M. Z. Sarikaya and H. Budak, "Generalized hermite-hadamard type integral inequalities for fractional integrals," *Filomat*, vol. 30, no. 5, pp. 1315–1326, 2016.
- [22] M. Z. Sarikaya, E. Set, H. Yaldiz, and N. Başak, "Hermite-hadamard's inequalities for fractional integrals and related fractional inequalities," *Mathematical and Computer Modelling*, vol. 57, no. 9-10, pp. 2403–2407, 2013.
- [23] M. Z. Sarikaya and H. Yildirim, "On hermite-hadamard type inequalities for riemann-liouville fractional integrals," *Miskolc Mathematical Notes*, vol. 17, no. 2, pp. 1049–1059, 2016.
- [24] E. Set, M. E. Ozdemir, and M. Z. Sarikaya, "New inequalities of ostrowski's type for s-convex functions in the second sense with applications," *arXiv preprint arXiv:1005.0702*, 2010.
- [25] Y. Zhang and J. Wang, "On some new hermite-hadamard inequalities involving riemann-liouville fractional integrals," *Journal of Inequalities and Applications*, vol. 2013, no. 1, p. 220, 2013.

#### *Authors' addresses*

##### **H. Budak**

Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce-Turkey  
*E-mail address:* hsyn.budak@gmail.com

##### **K. Özçelik**

Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce-Turkey  
*E-mail address:* kubilayozcelik@windowslive.com