

Title: Molecular evolutionary analysis of nematode *Zona Pellucida* (ZP) modules reveals disulfide-bond reshuffling and standalone ZP-C domains

Author: Cameron J. Weadick

Affiliation: Department of Biosciences, University of Exeter, Stocker Road, Exeter EX4 4QD, United Kingdom

Email: c.weadick@exeter.ac.uk

Abstract

Zona pellucida (ZP) modules mediate extracellular protein-protein interactions and contribute to important biological processes including syngamy and cellular morphogenesis. While some biomedically-relevant ZP modules are well-studied, little is known about the protein family's broad-scale diversity and evolution. The increasing availability of sequenced genomes from "non-model" systems provides a valuable opportunity to address this issue, and to use comparative approaches to gain new insights into ZP module biology. Here, through phylogenetic and structural exploration of ZP module diversity across the nematode phylum, I report evidence that speaks to two important aspects of ZP module biology. First, I show that ZP-C domains—which in some modules act as regulators of ZP-N domain-mediated polymerization activity, and which have never before been found in isolation—can indeed be found as standalone domains. These standalone ZP-C domain proteins originated in independent (paralogous) lineages prior to the diversification of extant nematodes, after which they evolved under strong stabilizing selection, suggesting the presence of ZP-N domain-independent functionality. Second, I provide a much-needed phylogenetic perspective on disulfide bond variability, uncovering evidence for both convergent evolution and disulfide-bond reshuffling. This result has implications for our evolutionary understanding and classification of ZP module structural diversity and highlights the usefulness of phylogenetics and diverse sampling for protein structural biology. All told, these findings set the stage for broad-scale (cross-phyla) evolutionary analysis of ZP modules and position *Caenorhabditis elegans* and other nematodes as important experimental systems for exploring the evolution of ZP modules and their constituent domains.

Keywords

gene family evolution, supradomain, domain architecture, cysteine connectivity, nematode cuticle, cuticlin

Introduction

Secreted proteins help cells withstand, react to, and shape external conditions (Agrawal, et al. 2010; Naba, et al. 2016; Cuesta-Astroz, et al. 2017). The extracellular environment can be variable and stressful, and in order to properly function under such challenging conditions, secreted proteins often employ specialized domains that can be repurposed to different ends by being recombined into different protein architectures (Bork, et al. 1996; Martin, et al. 1998). Obtaining an appreciation of the structural diversity of secreted proteins is key to understanding the many biological processes that extend beyond the cellular membrane. In many cases, however, insights into the biology of secreted protein families derive from restricted and potentially non-representative sets of model proteins (e.g., those linked to particular biomedical conditions, those expressed in already established model systems, and those that can be collected at high levels). Taking a broad, comparative view can uncover important but otherwise overlooked aspects of secreted protein structure and function.

The zona pellucida (ZP) module is a key component of many secreted proteins (Bork and Sander 1992; Plaza, et al. 2010; Litscher and Wassarman 2015; Bokhove and Jovine 2018). Named after the mammalian egg coat (from which the first family-members were found), ZP modules mediate extracellular protein-protein interactions. Through these actions, ZP-module bearing proteins (hereafter referred to simply as 'ZPD proteins', following (Litscher and Wassarman 2015)) contribute to a variety of critical cellular and developmental processes, including regulating sperm-egg interactions (Raj, et al. 2017), acting as a ligand co-receptor in the TGF β /BMP signalling pathway (Lin, et al. 2011; Saito, et al. 2017), and promoting dendrite elongation during neurogenesis (Heiman and Shaham 2009). Knowledge of ZP module structural

biology has increased considerably over the last few years, particularly for ZPD proteins linked to human health (Bokhove and Jovine 2018); mutations in these proteins underlie several human diseases, including hearing loss and renal failure (Verhoeven, et al. 1998; Devuyst, et al. 2017). However, ZP modules are found throughout the animal kingdom, from mammals to jellyfish (Matveev, et al. 2007), and there is still much to learn about their structural and functional diversity, particularly from an evolutionary perspective. For example, their role in gametic interactions implies a link to the evolution of species boundaries (Killingbeck and Swanson 2018) and their role in modulating cell shape suggests a connection to the evolution of morphological diversity (Fernandes, et al. 2010).

For most ZPD proteins studied to date, the primary purpose of the ZP module is to polymerize and trigger the formation of fibrous extracellular matrices (Jovine, et al. 2002; Jovine, et al. 2006). Understanding the mechanics of ZP module polymerization is an area of active research, particularly with regard to the roles played by the two domains that comprise a ZP module: ZP-N and ZP-C (named for their respective N- and C-terminal positions) (Bokhove and Jovine 2018).

Notably, it has been shown that isolated ZP-N domains can spontaneously polymerize into filaments *in vitro* (Jovine, et al. 2006). However, for a complete ZP module to polymerize it must first be activated. Studies of a few biomedically-relevant ZPD proteins such as uromodulin and ZP₃ indicate that cleavage of the ZP-C domain's C-terminal tail is critical to the activation process (Jovine, et al. 2004; Schaeffer, et al. 2009). First, cleavage severs the connection to the membrane, leading to extracellular release. Second, cleavage disrupts inhibitory interactions within the ZP-C domain that prevent polymerization: post-cleavage dissociation exposes an activating 'internal hydrophobic patch' (IHP) that is otherwise buried and suppressed by an 'external hydrophobic patch' (EHP) situated within the now-cleaved C-terminal tail (Jovine,

et al. 2004). These findings led to the notion that the ZP-N domain is the primary agent of protein-protein binding activity, and that the ZP-C domain is a regulator of ZP-N that acts to prevent ill-timed polymerization (Jovine, et al. 2006). Under the strictest form of this hypothesis, ZP-C domains serve no independent function and, consequently, would not be expected to be found on their own. Thus far, comparative data support this prediction: ZP-N domains have been found in isolation whereas ZP-C domains have not (Jovine, et al. 2006; Callebaut, et al. 2007). However, this model of domain functionality cannot directly apply to ZPD proteins that remain membrane-bound and do not polymerize (e.g., the BMP co-receptor endoglin (Saito, et al. 2017)). Moreover, ZP-C domains are capable of folding independently *in vitro* (Lin, et al. 2011; Diestel, et al. 2013; Bokhove, et al. 2016) and they contribute to protein-protein binding interfaces in some ZPD proteins (Han, et al. 2010; Lin, et al. 2011; Diestel, et al. 2013; Okumura, et al. 2015). These points combine to suggest that standalone ZP-C domains could in theory prove functional on their own and exist in nature.

ZP modules are characterized by the presence of multiple intradomain disulfide bonds (Bork and Sander 1992). However, the number of cysteine residues found per module varies and this has led to contrary views about how the cysteines connect and whether this variation has any functional effect (Jovine, et al. 2005; Yonezawa 2014). ZP modules have often been classified as either Type I or Type II based on the number of cysteines found within the ZP-C domain; these two groups were alleged to have non-nested connectivity patterns, and to differ functionally, with Type II but not Type I modules able to homopolymerize (Boja, et al. 2003; Darie, et al. 2004; Kanai, et al. 2008). However, in light of the solved structures of a few ZP modules and isolated ZP-C domains, it was subsequently argued that there is no reliable distinction between these groups, and that polymerization tendencies are unrelated to cysteine connectivity patterns (Bokhove and Jovine 2018).

Rather, Bokhove et al proposed that ZP-C domains typically have a standard set of three disulfide bonds (Cys5-Cys7, Cys6-Cys8, and CysA-CysB), with cysteine variation among ZPD proteins resulting primarily from lineage-specific gains and losses of disulfide pairs.

For example, the ZP module component of the BMP co-receptor endoglin lacks the Cys6-Cys8 and CysA-CysB disulfides found in uromodulin (Saito, et al. 2017), whereas additional disulfides associated with lineage-specific insertions have been found in some vertebrate egg-coat proteins (e.g., trout VE α / β and chicken ZP₃; (Darie, et al. 2004; Han, et al. 2010)). The case of ZP₃ is an interesting example, as this family of egg-coat proteins possesses a ZP-C subdomain that introduces four additional cysteine residues that are closely situated both along the sequence and in 3D space. Through protein crystallography of chicken ZP₃, Han et al. (Han, et al. 2010) showed that disulfide bonds covalently link the ZP-C core to its subdomain. By contrast, the results of earlier mass spectrometric analysis of other vertebrate ZP₃ proteins (but not including chicken ZP₃) indicated several cases where the subdomain's cysteines paired only amongst each other (Boja, et al. 2003; Darie, et al. 2004; Kanai, et al. 2008). If true, this pattern would be consistent with disulfide bond evolution via cysteine swapping, which is believed to be generally rare in nature (Thornton 1981; Rubinstein and Fiser 2008). However, mass spectrometry and crystallography provided contradictory results with regard to cysteine connectivity in mouse ZP₂ (Boja, et al. 2003; Bokhove, et al. 2016), suggesting that an artefactual explanation for the apparent cysteine swapping pattern seen amongst ZP₃ proteins cannot be ruled out. Regardless, the larger-scale comparison of ZP₃ with other ZPD proteins provides clear evidence for expanded cysteine connectivity beyond the core set of bonds defined by Bokhove et al. (Bokhove, et al. 2016). Finally, ZPD proteins may vary in the presence/absence of individual cysteines that contribute to

intermolecular bonds, as well, such as those involved in endoglin dimerization (Saito, et al. 2017). These studies have largely attempted to make sense of variation in ZPD cysteine connectivity through visual inspection of aligned protein sequences or structures without explicit regard to phylogeny. However, employing a phylogenetic approach may prove useful, for example by providing insights into the identification of ancestral versus derived states.

The diversity of ZPD modules found across the animal kingdom derive from a lengthy and complex history of speciation and duplication events that repeatedly provided new opportunities for unexpected structural features to arise. Efforts to test for the presence of isolated ZP-C domains and clear instances of disulfide-bond reshuffling in ZP modules would therefore benefit by taking a broad, phylogeny-informed approach. Recent genome sequencing projects for traditionally “non-model” systems provide the data needed for such studies, but thus far this path has not been taken. I set out to address this shortcoming through a molecular evolutionary study of ZP modules in nematodes.

Nematodes are an intriguing group for exploring the evolution and diversity of ZP modules for several reasons. First, the *Caenorhabditis elegans* genome encodes roughly twice as many ZP modules as are found in mammalian and fruit fly genomes, hinting at unexplored structural and functional diversity (Muriel, et al. 2003; Cohen, et al. 2019). Second, the recent sequencing of dozens of nematode genomes (Coghlan, et al. 2019) has provided the raw material needed for a focussed exploration of ZP module diversity within one of the animal kingdom’s most species-rich groups. Finally, given the proven suitability of *C. elegans* for genetics research, there is the potential for any insights gained from comparative analysis to be explored experimentally. Indeed, several ZPD proteins have already been genetically characterized in *C.*

elegans: these proteins are generally referred to as ‘cuticlin’ or CUT proteins on account of their structural roles in the nematode cuticle (Fujimoto and Kanaya 1973; Sebastiano, et al. 1991; Muriel, et al. 2003; Sapio, et al. 2005; Witte, et al. 2015).

However, the majority of ZPD proteins in *C. elegans* are simply annotated as CUT-like or CUTL proteins and little is known about their biology. Not surprisingly, even less is known about ZPD protein biology in nematodes beyond *C. elegans*, though it has been suggested that study of cuticlin proteins may aid efforts to pharmacologically attack the cuticles of nematodes that parasitize humans, livestock, and crops (Lewis, et al. 1994; Ondrovics, et al. 2016).

Through phylogenetic analysis of 1783 ZP modules from 59 nematode species, I found that the diversity of ZP modules present in *C. elegans* largely reflects the retention of subfamilies that originated and diverged prior to the diversification of modern nematodes. Using this phylogenetic framework, I then uncovered evidence for the evolutionary elaboration of ZP-C cysteine connectivity patterns (involving the modification of an otherwise conserved bond via disulfide-bond reshuffling, and the convergent evolution of novel IHP-stabilizing disulfides) and for the replicated loss of ZP-N domains in independent lineages (providing evidence that standalone ZP-C domains exist in nature, contrary to past predictions and observations). By taking a comparative, evolutionary approach, this work provides new insights into ZP module biology that should benefit efforts to determine ZP module structure-function relationships, in particular the functional role of standalone ZP-C domains. More broadly, this work provides a foundation for future phylogenetic studies aimed at providing an evolutionary classification of ZP modules and domains across the animal kingdom.

Materials and Methods

I compiled a data set of *C. elegans* ZPD protein sequences and used these to search for homologs in other nematodes. WormBase.org version 259 (Lee, et al. 2018) lists 45 genes that encode a "Zona pellucida domain" (i.e., linked to INTERPRO-ID IPR001507, PFSCAN-ID PS51034, PFAM-ID PF00100, and/or SMART-ID SM00241), including five *cut* and 29 *cutl* genes. Two of these were dropped from further consideration: *cutl-21* encodes an isolated and highly divergent ZP-N domain (Jovine, et al. 2006) while *r52.6* seems to have been incorrectly annotated (the PFSCAN motif assignment for R52.6 applies only to its first 40 aa, and BLASTP searches did not indicate sequence similarity with other nematode ZPD proteins; results not shown). When multiple isoforms were available, I selected a single variant, choosing whichever introduced the fewest/shortest indels in preliminary alignments of *C. elegans* ZP modules. This approach resulted in a data set of 43 *C. elegans* ZPD proteins (Supplementary Table 1). As unannotated ZPD proteins would have been missed by the above approach, I also conducted BLASTP searches of the *C. elegans* proteome, using, in turn, the ZP modules from each of the 43 annotated ZPD proteins as the query. (Details on the BLASTP search approach are provided below.) Aside from the already-discounted ZP-N-only protein CUTL-21, doing so did not uncover any additional ZPD proteins (results not shown). ZPD proteins often include other domains upstream of the ZP module; I isolated *C. elegans* ZP modules using GISMO (ver 2.0), an alignment program that uses a Bayesian approach to extract and align the homologous core regions of sequences that potentially contain non-homologous flanks and insertions (Neuwald and Altschul 2016). Because GISMO is stochastic, I applied it multiple times ($n=5$); the positions and lengths of inferred insertions and flanking regions varied among replicates, but all targeted the ZP module, retaining the C-terminal consensus cleavage site and excluding upstream domains and the N-terminal trafficking motif. The 43 flank-trimmed (but not insertion-trimmed) sequences from the

GISMO run with the longest conserved core (obtained using seed 28270; Supplementary File 1) were then used as search queries to detect homologs in 58 additional nematode species through similarity searches of whole-genome predicted protein sets (see Supplementary Table 2 for data set sources). The similarity searches were conducted using BLASTP 2.6.0 (Altschul, et al. 1990), with low complexity regions within the query sequences masked using "seg yes -soft_masking true". After removing subjects best matched by CUTL-21 (the divergent ZP-N-only protein), I filtered the results to retain only those subjects with E-values lower than 10^{-10} and total query coverages of at least 75%. Alternative isoforms were filtered to keep only the longest, though this was only possible for species where predicted isoforms were explicitly identified via sequence name suffixes (e.g., 't1', 't2').

The final data set of 1783 full length ZPD protein sequences (Supplementary File 2) was aligned and trimmed using GISMO. One hundred replicate alignments were generated, with key phylogenetic analyses repeated across all replicates; random seeds are provided in Supplementary File 3. To avoid subjective judgement from biasing the results, alignments were not manually adjusted in any way. Conservation patterns in the focal alignment (the top-scoring alignment according to log-likelihood ratio (LLR) score; Supplementary File 4) were visualized using WebLogo (weblogo.berkeley.edu; (Crooks, et al. 2004)). Throughout the paper, site numbering refers to position in the trimmed focal alignment.

Maximum likelihood (ML) phylogenies were estimated using PhyML via the www.atgc-montpellier.fr/phyml/ web server, with automated SMS-AIC model selection, a BioNJ starting tree, and SPR topology rearrangements (Guindon, et al. 2010; Lefort, et al. 2017). Model parameters were fixed at their SMS-AIC estimates during tree search. This process was applied

to all 100 replicate alignments; results were combined by generating a majority-rule consensus tree and using branch recovery proportions (BRPs) to quantify branch support. Because BRPs can be downwardly biased by rogue taxa/lineages, BRPs were supplemented by 'Transfer Bootstrap Expectations' (TBEs), calculated using BOOSTER (booster.pasteur.fr/new/) (Lemoine, et al. 2018)). The typical methods for estimating branch support on ML trees, namely bootstrapping and aLRT SH-like tests, were not employed as these methods ignore uncertainty in the alignment. $N = 100$ sets of ML branch lengths were estimated for the consensus topology via iqTree 1.6.0 (Nguyen, et al. 2015), using, in turn, each replicate alignment and its corresponding SMS-AIC substitution model. Trees were rooted using the Minimal Ancestor Deviation method via mad 2.2 (Tria, et al. 2017); this approach aims to identify the root position that minimizes deviance in root-to-tip lengths, thereby accounting for heterogeneity in evolutionary rate across the tree (which can mislead the simple midpoint rooting approach). Trees were plotted and analyzed using functions from the ape, phytools, and phangorn R packages (Schliep 2011; Popescu, et al. 2012; Revell 2012).

Patterns of sequence loss were explored by calculating the amount of missing data within each replicate alignment and mapping these values onto the phylogeny. Gap proportions were estimated separately for the ZP-N and ZP-C domains, with the approximate domain boundaries determined according to cysteine conservation patterns: using the nomenclature of (Bokhove, et al. 2016), ZP-N was demarcated using Cys₁ and Cys₄ (positions 1 and 80; Figure 1), while the boundaries of ZP-C were defined using a moderately conserved cysteine in ZP-C's β A strand along with Cys₈ (positions 105 and 218; Figure 1).

Based on the results of the missing-data analysis, three subfamilies were selected for codon model analysis, namely the T01D1.8, F46G11.6, and CUTL-19 subfamilies (named according to their respective *C. elegans* members). In each case, untrimmed protein sequences were re-aligned using GISMO and the alignment with the top LLR score (out of $n=10$ replicates) was used to estimate a subfamily-specific phylogeny (via PhyML, as described above) and build a corresponding codon sequence alignment. (The top scoring alignments were obtained with the following random seeds: T01D1.8 = 25393, F46G11.6 = 21134, and CUTL-19 = 4128.) The codon alignments and trees were used to fit codon substitution models via CodeML from the PAML 4.9a package (Yang 2007). The key parameter for codon models is ω , the nonsynonymous (dN) to synonymous (dS) divergence ratio ($= dN/dS$), with values near zero indicating strong purifying selection and values greater than one suggestive of positive selection. I fit three codon models: M8, M8a, and Mo. M8 and M8a are nested models that were used to test for site-specific positive selection ($\omega > 1$) and to estimate among-site variation in the strength of selective constraint (Swanson, et al. 2003); these models were compared via a likelihood ratio test. The simple Mo model assumes that selection is constant across the alignment and was used to obtain overall estimates of the strength of selection (Goldman and Yang 1994) as well as branch-specific estimates of dS , which were used to check for saturation. All three models assume that selection is constant across the phylogeny.

Homology models were estimated for *C. elegans* ZP modules using the RaptorX web server (raptorx.uchicago.edu; (Kallberg, et al. 2012)). In most cases, full length sequences were submitted for analysis: the exceptions were CUTL-19b, T01D1.8b, and F46G11.6 (which are all short, less than 260 aa long; these sequences were trimmed to remove any predicted N- and C-terminal propeptide flanks), and FBN-1a (which is quite long; only the last 2500 aa of this 2799 aa protein were

analysed owing to RaptorX size limits). The 3.2Å resolution structure for human uromodulin (RCSB PDB code 4wrn; (Bokhove, et al. 2016)) was used as the template for each model; justification for using this template structure is provided in the *Results* section. When examining the resulting models, I only considered the ZP-N and ZP-C domains, not the up and downstream regions or the interdomain linker; domain boundaries were determined from each model's RaptorX structural alignment. Homology models were aligned with one another using DeepAlign:3DCOMB v1.18 (Wang, et al. 2011) and then superimposed on the template for visualization and measurement using the 'super' function in PyMOL v1.8.6.0 (github.com/schrodinger/pymol-open-source).

C-terminal R/K cleavage sites and N-terminal signalling motifs were predicted for untrimmed sequences via the ProP 1.0 Server (www.cbs.dtu.dk/services/ProP/; (Duckert, et al. 2004)), using a score cut-off of 0.5 and discounting cleavage sites predicted within the signalling peptide. C-terminal GPI-anchors were predicted using PredGPI (gpcr.biocomp.unibo.it/predgpi/pred.htm; (Pierleoni, et al. 2008)), using the "general model" option and a specificity cut-off of 99.0%. Protein domains were predicted using PfamScan (www.ebi.ac.uk/Tools/pfa/pfamscan/; (Li, et al. 2015)) with default search settings.

Results

Data set and Alignment

A data set of 1783 nematode ZP modules was assembled by BLASTP searching the whole-genome predicted protein sets of 58 nematode species for homologs of 43 *C. elegans* ZP modules. The search set included both free-living and parasitic

species and covered four of the five major nematode clades defined by (Blaxter, et al. 1998) (Supplementary Table 1); by covering such a wide range of species, this approach should hopefully uncover all major nematode ZP module subfamilies regardless of the idiosyncrasies associated with any particular nematode lineage, or the shortcomings associated with any particular genome project. The number of ZP modules per species in the final data set ranged from 15 for *Romanomermis culicivorax* to 58 for *Toxocara canis*, with Clade-I nematodes contributing fewer ZP modules to the final data set (median = 21; IQR = 19–22) than Clade-III/IV/V species (median = 36; IQR = 28–41).

ZP modules were extracted and aligned using GISMO. Alignment uncertainty is a concern given the short target region and the phylogenetic breadth of the data set. I addressed this by leveraging the stochastic nature of the GISMO alignment procedure, repeating key phylogenetic analyses across 100 replicate alignments. Consistent with expectations for ZP modules, the final GISMO-trimmed alignments were 233–269 aa long, with majority rule consensus sequences possessing 11–13 cysteines. The percentage of gaps and ambiguous data ranged from 6.7–8.6% across alignments. Conservation patterns for the focal alignment (the alignment with the highest LLR score) are shown in Figure 1, with the alignment itself available in Supplementary File 4. Most alignment sites were highly variable, with several cysteine residues and the ZP-C domain's R/K-rich consensus cleavage site (CCS) being notable exceptions. The relationship between position numbering in the focal alignment and untrimmed sequence position for *C. elegans* CUT-1 is shown in Supplementary Figure 1.

Phylogenetics

Evolutionary trees were estimated for the 100 replicate alignments using Maximum Likelihood. Alignment variation affected both model selection and the resulting topology. With regard to the substitution model, VT+I+G was favoured for 77 alignments, LG+I+G for one, and WAG+I+G+F for the remainder, with the top model receiving an AIC weight of 1.000 in 98/100 cases (Supplementary File 3). With regard to the resulting phylogenies, normalized Robinson-Foulds (RF) distances ranged from 0.31 to 0.43 between pairs of trees (where 0 corresponds to topologically identical trees and 1 corresponds to completely contradictory trees). However, this method ignores branch lengths; weighting by branch lengths reduced the pairwise RF distances considerably (range = 0.12–0.23), indicating that many disagreements involved only small-scale differences. The individual trees, including branch lengths and aLRT SH-like partition support values, are provided in Supplementary File 5.

Rather than focussing on the individual ML trees, I constructed a majority-rule consensus tree (Figure 2a), sacrificing resolution for robustness in the face of alignment uncertainty. Doing so reduced the number of internal branches from 1780 to 1266 via the formation of 185 polytomies. Most of the retained branches were relatively well supported, with just over half having Branch Recovery Proportions (BRPs) of at least 0.95, though 21% had BRPs below 0.70; BRPs are analogous to bootstrap support values but quantify the degree of support for a given branch across replicated estimates of the actual alignment as opposed to bootstrap pseudo-alignments. Phylogenetically unstable branches seem likely for a data set of this size and these will tend to reduce recovery frequencies for otherwise robust clades. I therefore also estimated 'Transfer Bootstrap Expectations' (TBEs); this approach calculates how frequently each branch is recovered among replicate trees, but it does so in a manner that accounts for rogue branches (Lemoine, et al. 2018). Reassuringly, branch

support values increased substantially when considering TBE supports, particularly for deeper branches (Supplementary Figures 2,3). Branch length estimates were largely robust to alignment variation: the majority-rule consensus tree drawn in Figure 2a shows branch lengths estimated using the focal alignment, but highly similar results were obtained using any of the other 99 replicate alignments (pairwise Pearson's $r = 0.94-0.98$).

Visual inspection of the nematode ZP module phylogeny revealed three major groups, which I refer to as Type 1, 2, and 3 ZP modules (Figure 2a). These groups are characterized by distinct cysteine conservation patterns that imply alternative ZP-C domain disulfide connectivity patterns (as detailed below via homology modelling) (Figure 2b, Supplementary Figure 4). The branches that define these three groups are well-supported: BRP = 0.90 and TBE = 0.99 for Type 1 vs Type 2/3, and BRP = 0.86 and TBE = 0.99 for Type 2 vs Type 3. The root of the tree was predicted by the Minimal Ancestral Deviation (MAD) method to fall within the Type 1 section of the phylogeny (Figure 2a, Supplementary Figure 5). Notably, the MAD approach is robust to variation in evolutionary rate among lineages, which appears to be important here (note the long branches within the CUTL-19 and CUTL-14 subfamilies, and the shift between the MAD root and the phylogenetic midpoint that is often used to estimate the root position; Figure 2a). This root placement rendered Type 1 modules paraphyletic and therefore suggests that the Type 1 cysteine connectivity pattern is the ancestral state. The Type 2 and 3 modules share a novel pair of ZP-C domain cysteines, and Type 3 modules are further distinguished by the modification of a ZP-C disulfide that remains conserved in Type 1 and 2 modules. Deep relationships within the Type 2 portion of the tree were ambiguous, suggesting a rapid gene family expansion through multiple rounds of duplication and divergence. The consensus tree is inconclusive regarding whether Type 2 and Type 3 modules are sister groups or if Type 2 modules are

paraphyletic, with Type 3 modules representing a derived subclade. The latter scenario is supported by the fully resolved tree obtained using the focal alignment, though the short branch lengths and moderate-to-low recovery frequencies made this conclusion uncertain (Supplementary Figure 3). Notably, Cohen et al. (Cohen, et al. 2019) recently classified *C. elegans* ZP modules into groups based on the number of ZP-C domain cysteine residues present per sequence, and their classification system is broadly congruent with the one provided here (Supplementary Table 3). However, their approach, which was both non-phylogenetic and *C. elegans*-specific, misclassified a few members that independently lost or gained additional disulfides (detailed below), and is equivocal with regard to which cysteine connectivity pattern is ancestral.

The *C. elegans* ZP modules were, with few exceptions, distributed broadly across the phylogeny, and similar patterns were seen for the other species (Figure 2a; Supplementary Figure 6). This pattern indicates that the nematode ZP module phylogeny is characterized by over 40 paralogous subfamilies that originated prior to the diversification of modern nematodes, with the members of each subfamily representing clusters of putative orthologs. Indeed, the lengths of the internal branches that connect the various subfamilies are suggestive of ancient origins, perhaps even predating the origin of the nematode phylum. Follow-up studies would therefore do well to sample broadly (i.e., including closely related phyla), as doing so may uncover deep ZP module conservation between invertebrate groups.

While the tree is largely indicative of stable orthology, occasional lineage-specific gains and losses were also observed. *C. elegans* lacks members of a few subfamilies (i.e., the sister groups to the CUTL-10 and CUTL-23 clades) and the *C. elegans* CUT-1 and CUT-3 modules clearly derive from a recent duplication event. Beyond *C. elegans*, losses appeared particularly

common for Clade 1 nematode ZP modules (Supplementary Figure 6). Adaptive gene loss associated with parasitism likely underlies this pattern (Korhonen, et al. 2016) but data quality issues also play a role: the short intergenic regions typical of Clade 1 nematode genomes can cause false fusion events between neighbouring genes (Pettitt, et al. 2014) and I found that the tandemly arranged *cutl-28* and *dyl-7* genes of Clade 1 (*Trichinella*) species were fused, resulting in only the DYF-7 sequences ending up in the final data set (results not shown). Putting these few departures aside, the overall pattern is consistent with deep conservation of the ZP module complement across the nematode phylum. Assuming that gene duplication is the primary driver of functional divergence within the ZP module family, these results therefore support efforts to leverage knowledge about cuticular biology in the lab model *C. elegans* for use in treating or preventing parasitic nematode infections.

PfamScan analysis identified a total of 2310 domains within 1186 (67%) of the input sequences (Supplementary File 2). Most of the predicted matches (91%) were for domains typical of *C. elegans* ZPD proteins, namely the zona pellucida 'domain' (Pfam:Zona_pellucida; 36%), two types of PAN domain (Pfam: PAN_1 and PAN_4; 31%), two types of Epidermal Growth Factor (EGF)-like domain (Pfam:EGF_CA and EGF_3; 17%), and the von Willebrand factor Type A (vWFA) domain (Pfam:VWA; 7%). The remaining 9% matched 96 different Pfam entries, with none individually accounting for more than 0.7% of the total; these additional domain predictions were not considered further as nearly half derived from Clade 1 nematodes (which, as mentioned, have high incidences of artefactually fused genes (Pettitt, et al. 2014)). The Pfam:Zona_pellucida entry was only returned for 45% of the sequences, indicating that the domain-prediction approach is prone to false negatives, at least for nematode ZP modules. Mapping the PAN, EGF, and vWFA predictions onto the

phylogeny showed that upstream domain predictions within the various subfamilies generally matched expectations, given each clade's respective *C. elegans* member (Supplementary Figure 7). Assuming that domain architecture is conserved within the relevant subfamilies (i.e., that false negatives are more plausible than recurrent domain losses and gains within each subfamily), the majority-rule consensus topology is compatible with single origins for each observed domain architecture. However, the presence of polytomies make this conclusion tentative for PAN+ZP and vWFA+ZP, and the fully resolved topology estimated using the focal alignment (Supplementary Figure 4) actually implies either multiple origins or a single origin followed by multiple losses of the vWFA+ZP arrangement.

Structural Evolution: Sequence Loss

To test for deletions indicative of major structural alterations of the ZP module, I calculated the proportion of missing data for each aligned sequence and mapped these 'gap proportions' onto the phylogeny. This was done separately for the ZP-N and ZP-C domains. Three subfamilies—CUTL-19, T01D1.8, and F46G11.6, named for their respective *C. elegans* members—showed pronounced signatures of ZP-N domain sequence loss (Figure 3a). Averaged across sequences within the respective subfamilies, ZP-N gap proportion ranged from 83–97% for the CUTL-19 subfamily (depending on alignment replicate), 74–96% for the T01D1.8 subfamily, and 83–99% for the F46G11.6 subfamily. Gap proportions tended to be much lower across the rest of the data set, averaging 6–8% depending on the alignment replicate. Some sequences outside the CUTL-19, T01D1.8, and F46G11.6 subfamilies also showed high gap proportions, but these tended to be local outliers and therefore may simply represent artefactual truncations. For the ZP-C domain, the gap proportion was generally quite low (Figure 3b): averaged across sequences, the gap proportion was 3–4% depending on alignment. Cohen

et al (Cohen, et al. 2019) independently noted the apparent lack of the ZP-N domain in *C. elegans* T01D1.8 and F46G11.6 (no results were reported for CUTL-19) but did not explore the issue further.

The phylogenetic distribution of the standalone ZP-C subfamilies indicates that ZP-N loss occurred prior to the emergence of the major nematode lineages, and that it happened at least twice (Figure 3a). The T01D1.8 and F46G11.6 subfamilies are closely related Type 3 modules united by a well-supported branch (BRP = 0.90; TBE = 0.96) and the loss of the ZP-N domain in these subfamilies plausibly represents a single event. The CUTL-19 subfamily, however, is phylogenetically distant, indicating an independent loss of ZP-N within the Type 2 section of the tree. With regard to taxonomic composition, the T01D1.8 subfamily possesses ZP modules from nematodes from all four of the sampled clades (Clades I, III, IV, and V) whereas the F46G11.6 and CUTL-19 subfamilies lack sequences from Clade I nematodes (Supplementary Figure 6). As Clade I nematodes tend to have considerably fewer ZP modules than other nematodes, this difference presumably reflects two instances of Clade I-specific loss.

Codon model analyses were used to estimate the degree of evolutionary constraint experienced within these three subfamilies. Alignment-wide dN/dS under the Mo codon model was $\omega = 0.094$ for T01D1.8, 0.095 for F46G11.6, and 0.135 for CUTL-19, indicating the action of moderately strong purifying selection acting throughout the history of these subfamilies. Selective constraint was generally strongest within the core regions of the ZP-C domain, especially at sites within predicted β strands (Supplementary Figure 8). M8-M8a likelihood ratio tests provided no evidence for site-specific positive selection ($\omega > 1$) in any of the subfamilies (Supplementary Table 4). Under the Mo model, approximately 75% of

branches had $dS < 1$ and 98% had $dS < 3$ in each data set, indicating that saturation is unlikely to have strongly affected these analyses.

N-terminal signal peptides were predicted for most members of all three standalone ZP-C domain subfamilies (73% for To1D1.8, 84% for F46G11.6, and 79% for CUTL-19 versus 66% for the rest of the data set), suggesting that these unusual proteins are still secreted despite the loss of their respective ZP-N domains. However, the three standalone ZP-C subfamilies differed from the norm by generally lacking predicted R/K cleavage sites (30% for To1D1.8, 3% for F46G11.6, and 8% for CUTL-19 versus 66% for the rest). Examination of the subfamily-specific alignments and *C. elegans* homology models showed that the members of the To1D1.8 and F46G11.6 subfamilies tend to possess short C-terminal tails that terminate before the ZP-C domain's final β strand, β G, which contains the regulatory EHP motif (Figure 3c–e). Finally, and unexpectedly, GPI-anchors were predicted for most members of the CUTL-19 subfamily (57%) despite this C-terminal feature being very rare across the rest of the data set (5%, and not found at all in the other two standalone ZP-C subfamilies). Predicted propeptide features for all 1783 sequences are reported in Supplementary File 2.

Structural Evolution: Cysteine Connectivity

Examination of amino acid variability patterns indicated that some cysteine residues were less strongly conserved than others, suggestive of variation in disulfide binding patterns (Figure 1, Supplementary Figure 4). To explore this further, homology models were generated for the 43 *C. elegans* ZPD proteins using RaptorX (Kallberg, et al. 2012). The *C. elegans* ZPD proteins yielded matches to several solved ZP module templates: human uromodulin (RCSB PDB code: 4wrn), chicken

ZP₃ (3nk3), human endoglin (5hzv), mouse ZP₂ ZP-C domain (5bup), and rat betaglycan ZP-C domain (3qw9). I focussed only on homology models generated using the human uromodulin template (Bokhove, et al. 2016). This was done for three reasons: (1) using a common template facilitated aligning and comparing models generated for different sequences; (2) models built using this template were usually the best option according to RaptorX's internal ranking system (first place in 34/43 cases and second place in the rest, and always with highly significant model quality *P* values; Supplementary Table 5); and (3) human uromodulin possesses all three of the putatively typical ZP-C disulfide bonds defined by Bokhove et al (Bokhove, et al. 2016), allowing for evaluation of cysteine connectivity patterns. Homology models and structural alignments are provided in Supplementary File 6.

The ZP-N domain was successfully modelled in 39 of 43 cases, the exceptions being the three standalone ZP-C domain proteins plus CUTL-9, which possesses a long insertion within the ZP-N domain's DE loop that disrupted modelling (Supplementary Table 5). Structural alignment of the models revealed complete conservation of the two disulfides typical of ZP-N domains, namely the Cys1-Cys4 linkage between the β A and β G strands, and the Cys2-Cys3 linkage between the CD and EE' loops (Supplementary Figure 9). These residues correspond to positions 1, 29, 48, and 80 in the focal alignment, all of which are highly conserved (Figure 1). Examining the positions of other cysteine residues in the *C. elegans* models identified a putative β F- β G disulfide specific to CUTL-5 (Supplementary Figure 9). Sequence conservation patterns suggest that this disulfide evolved within the nematode phylum, with the cysteines conserved across Clade III, IV, and V orthologs but not Clade I orthologs (results not shown).

The ZP-C domain was successfully modelled in all 43 cases (Supplementary Table 5; Supplementary Figure 10). The models were generally in good agreement with one another, as expected given the use of a common template structure. However, the C-terminal tails often proved difficult to align and model due to the presence of extended FG loops in the *C. elegans* sequences (as can be seen for CUT-1 in Supplementary Figure 1; note the long unaligned region immediately prior to the consensus cleavage site). In some cases, this led to termination of the model prior to the β G strand (the ZP-C domain's final β strand), even when evidence for it was clearly present in the multiple sequence alignment. In other cases, the β G strand was recovered but connected via a long FG loop that was predicted by RaptorX to have a high propensity for disorder (results not shown). Fortunately, it was still possible to evaluate disulfide binding patterns in most models, as the key cysteine residues are upstream of the poorly modelled region. Doing so revealed clear evidence for large-scale variation in cysteine connectivity among nematode ZP module subfamilies (Figure 2b).

According to Bokhove et al. (Bokhove, et al. 2016), the typical ZP-C domain has three disulfide bonds: Cys5-Cys7, which connects β C to β F; Cys6-Cys8, which connects β C'' to the FG loop, and CysA-CysB, which connects β F to the FG loop. The Cys5-Cys7 disulfide was recovered in nearly all models (Supplementary Figure 10), the only exception being the model for *C. elegans* CUTL-28b. The Cys5 and Cys7 residues are conserved across almost the entire alignment (positions 146 and 201; Figure 1) but both cysteines are absent in the CUTL-28 subfamily (replaced with lysine and alanine, respectively), indicating a subfamily-specific disulfide loss. Cys5-Cys7 loss has also been reported for the *Drosophila* ZPD protein NompA (Bokhove, et al. 2016); whether this represents convergent loss or deep conservation awaits cross-phyla

phylogenetic analysis, though I note that both NompA and CUTL-28 are predicted to have upstream PAN domains (Fernandes, et al. 2010).

The Cys6-Cys8 disulfide was also found to be broadly conserved, though modelling uncertainty makes this conclusion tentative for Type 1 modules. Cys6 mapped to alignment position 164 while Cys8 typically mapped to either 215 or 218 (Supplementary Figure 4a,b), though a single highly conserved Cys8 alignment column was observed for many alignment replicates (results not shown). A disulfide between Cys6 in the $\beta C''$ strand and Cys8 in the FG loop was recovered in all 16 Type 2 modules and in 10 of 12 Type 3 modules (the two exceptions being cases where the unconnected cysteines were placed nearby one another) (Supplementary Figure 10a,b). For Type 1 modules, the Cys6-Cys8 disulfide was recovered (or deemed plausible by proximity) in 7 of 15 models; in the remainder, Cys8 bound or was placed near CysA (Supplementary Figure 10c). While this arrangement could indicate a novel connectivity pattern, the fact that it leaves both Cys6 and CysB (the typical partner of CysA) unbound and distant from one another suggests that it is a consequence of inaccurate modelling of the FG loop; notably, these cysteines were all found to be highly conserved across Type 1 modules (Supplementary Figure 4c). The simplest interpretation is therefore that the Cys6-Cys8 disulfide is conserved in nematode ZPD proteins but is, in some cases, difficult to recover via homology modelling. That said, loss of Cys6-Cys8 has been reported outside of nematodes (e.g., in human endoglin; (Saito, et al. 2017)), indicating that the evolutionary breakdown of the Cys6-Cys8 bond is possible and cannot be conclusively ruled out for all nematode ZPD proteins.

The CysA-CysB disulfide was found to be unexpectedly variable. CysA-CysB, which connects the end of β F strand (position 206) to the beginning of the FG loop (position 210), was recovered in 15 of 16 Type 2 modules (and deemed plausible by proximity in the remaining case) (Supplementary Figure 10b). Whether this linkage is conserved among Type 1 ZP-C domains is unclear given the FG loop modelling uncertainty described above, though the relevant cysteines are highly conserved (Supplementary Figure 4c), and the CysA-CysB linkage was recovered for the DYF-7 and LET-653b models (Supplementary Figure 10c). However, there was a clear loss of the CysA-CysB disulfide in *C. elegans* CUTL-24b; this disulfide has also been lost in some non-nematode ZP proteins (e.g., ZP3; (Han, et al. 2010)) but the example reported here appears to be nematode-specific (shared with Clade III, IV, and V orthologs, but not with orthologs from Clade I nematodes). The CysA-CysB linkage was also lost in Type 3 modules, albeit in an incomplete manner: Type 3 ZP-C domains lack CysA entirely yet surprisingly retain CysB, which is well-positioned to bind a novel cysteine partner in the adjacent β C strand (position 140; median centroid distance of 5.9Å over the 12 Type 3 models; Supplementary Figures 4a, 10a). These findings strongly suggest that the CysA-CysB disulfide was modified via a partner replacement—partially lost, partially conserved.

Beyond the characteristic Cys5-Cys7, Cys6-Cys8, and CysA-CysB disulfides, ZP-C domains sometimes possess additional disulfides, for example the novel Cx-Cy pair found in trout VE α / β egg coat proteins (Darie, et al. 2004) that appears to stabilize a fish-specific expansion of the AB loop, just downstream of the β A-IHP. A few candidates for novel disulfides are apparent in the *C. elegans* ZP-C domain homology models. First, the model for *C. elegans* CUTL-19b included a pair of cysteines that are well placed to link the AB loop and β B (positions 117 and 129; centroid distance = 6.2Å; Figure 3e). These

cysteine residues are both conserved across the CUTL-19 subfamily but are not found beyond it, suggesting that stabilizing the AB loop is particularly important in this standalone ZP-C domain subfamily. The second example, which is more broadly distributed (shared across Type 2 and Type 3 modules), affects the IHP and therefore may be of major functional relevance. Here, cysteines are found at positions 105 (within the IHP motif) and 134 (Supplementary Figure 4a,b). Homology modelling of *C. elegans* ZP-C domains put these cysteines in close proximity: position 105 near the start of β A, and position 134 near the end of β B (median centroid distance of 5.6Å over the 28 Type 2 and Type 3 models; Supplementary Figure 10a,b). Intriguingly, this putative disulfide forms part of a bipartite motif—one divided between the β A and β B strands—that is highly conserved in Type 2 and Type 3 ZP modules. Here, three aromatic residues are projected into the same β A- β B interface bridged by the proposed disulfide bond (Figure 4). Finally, a partially overlapping disulfide appears to have evolved within the early history of the FBN-1 subfamily (a Type 1 module). This putative disulfide is defined by cysteines at alignment positions 105 and, uniquely, 203 (centroid distance = 7.3Å; Supplementary Figure 10c); both of these cysteines are conserved across the FBN-1 family. A disulfide between these residues would anchor β A not to β B (as seems to be the case for the Type 2 and 3 modules), but to β F. This suggests that similar but not identical disulfide bonds have evolved to stabilize the IHP-containing β A strand in different lineages of the nematode ZP module family.

Discussion

The ZP module is a supra-domain (Vogel, et al. 2004)—a combination of structurally independent domains, ZP-N and ZP-C, that function cooperatively and frequently co-occur across a variety of proteins with distinct domain architectures. The co-occurrence between ZP-N and ZP-C is so strong that they were previously considered mere subdomains within a

single “ZP domain” (Monne, et al. 2008; Han, et al. 2010; Bokhove, et al. 2016; Wilburn and Swanson 2017), and while isolated ZP-N domains have been found in a variety of proteins, ZP-C domains have only ever been found within complete modules (Jovine, et al. 2006). This tight but lopsided distribution is consistent with past studies of ZP structure-function relationships that revealed a role for the ZP-C domain as a regulator of ZP-N activity (Litscher and Wassarman 2015; Bokhove and Jovine 2018). Under the assumption that this regulatory role is the ZP-C domain’s primary function (historically, if not currently in each extant ZPD protein), it makes sense that it would only ever be found immediately downstream of a ZP-N domain. However, studies have uncovered non-regulatory (protein-binding) functions for some ZP-C domains (Han, et al. 2010; Lin, et al. 2011; Diestel, et al. 2013; Okumura, et al. 2015; Bokhove, et al. 2016), and this raises questions about the apparent lack of standalone ZP-C domains in nature.

I have shown here that, contrary to expectations, standalone ZP-C domains indeed exist—that they can evolve from pre-existing ZP modules through ZP-N domain loss. My analysis of nematode ZP modules revealed that standalone ZP-C domain proteins originated at least twice, and that they have been maintained over long timeframes—originating prior to the diversification of the major nematode clades and subsequently evolving under strong stabilizing selection. Despite the loss of the upstream ZP-N domain, these standalone ZP-C proteins generally still possess N-terminal signal peptides, suggesting that they remain secreted proteins. Their C-terminal features, by contrast, are atypical: the members of the T01D1.8 and F46G11.6 subfamilies tend to be truncated, indicating that they may be secreted directly without need for proteolytic separation from the membrane, while most members of the CUTL-19 subfamily have predicted GPI-anchor sites (despite this C-terminal feature being rare across the rest of the data set). These findings suggest new dimensions of

functionality for ZP-C domains. One possibility is that these standalone ZP-C domains indeed perform a regulatory role, but as free-agent regulators of unlinked ZP-N domains rather than of physically linked upstream domains; such proteins might prove useful for remodelling ZPD protein-based extracellular matrices. Another is that ZP-C domains are multifunctional, having some uncharacterized non-regulatory function. Some ZP-C domains have been shown to contribute to protein-protein binding (Han, et al. 2010; Lin, et al. 2011; Diestel, et al. 2013; Okumura, et al. 2015) and it may be that these standalone ZP-C domains do likewise. Either way, the finding that standalone ZP-C domains exist in nature will benefit future experimental efforts to explore the ways in which individual domains contribute to higher-level functioning in ZP module-bearing proteins. The ancient origins for the standalone ZP-C proteins suggests that they might be shared with other phyla, but even if standalone ZP-C domains turn out to be restricted to nematodes alone, the mechanistic insights gleaned from their study will likely prove informative in a general sense.

Of the 43 ZPD proteins encoded by the *C. elegans* genome, less than half have been functionally characterized. Aside from DYF-7, which plays a role in neural dendrite elongation (Heiman and Shaham 2009), all of these are cuticular proteins. Several appear to be cuticlins, i.e., non-collageneous structural proteins (Fujimoto and Kanaya 1973; Sebastiano, et al. 1991; Muriel, et al. 2003; Sapio, et al. 2005; Witte, et al. 2015), while others have been linked to cuticular moulting (Frand, et al. 2005) or to the development of various cuticular elaborations and invaginations (Yu, et al. 2000; Kelley, et al. 2015; Gill, et al. 2016; Vuong-Brender, et al. 2017; Cohen, et al. 2019). These cuticular proteins are distributed across the nematode ZP module phylogeny, and cover all four of the major domain architectures (ZP, vWFA+ZP, PAN+ZP, and EGF+ZP), suggesting that many of the uncharacterized ZPD proteins, including the standalone ZP-C domain proteins, probably also

play a role in the cuticle. Consistent with this hypothesis, transcriptome data from Spencer et al. (Spencer, et al. 2011; Lee, et al. 2018) indicate that T01D1.8 and F46G11.6 are both enriched in the epidermis during early development, but that CUTL-19 is enriched in embryonic and larval motor neurons (suggesting a divergent role, perhaps akin to that of DYF-7). A subsequent study found that T01D1.8 is up-regulated in some thermosensitive neurons (Lockhead, et al. 2016), hinting at multiple roles for this standalone ZP-C protein. It will be interesting to see, as more nematode ZPD proteins are characterized, whether phylogeny or domain architecture reliably predict functional role, and whether any of these proteins contribute to the egg coat (as ZPD proteins are known to do in vertebrates and at least some invertebrates; (Killingbeck and Swanson 2018)).

It has been previously shown that artificially isolated ZP-C domains express and fold correctly *in vitro* (Lin, et al. 2011; Diestel, et al. 2013; Bokhove, et al. 2016). The present study provides the first evidence that this experiment has also been performed in nature, with standalone ZP-C domains having evolved from full modules through ZP-N loss. This finding has implications for our understanding of the origin of the ZP module. Two models have been put forth to explain how the original ZP module may have first evolved. The first proposes that the ZP module may have originated via the tandem duplication of a polymerization-capable proto-ZP-N domain, with the C-terminal copy then evolving to form the ZP-C domain (Han, et al. 2010). The second hypothesis suggests that ZP modules may have evolved from antibody light chains polypeptides, as both are composed of 1G-like domains (Bokhove and Jovine 2018). Finding that standalone ZP-C domains are viable in nature suggests it is possible (though unproven) that such proteins could have independently existed in the deep past, and from this admittedly speculative assumption two new possibilities arise: (1) the ZP module could have

evolved through tandem duplication and divergence of an ancient ZP-C-type domain; (2) the ZP module could have formed through the union of pre-existing but independent ZP-N-type and ZP-C-type domains. Given the lack of recognizable sequence-level homology between ZP-N and ZP-C domains, and between either of these domains and their structurally similar counterparts in antibody light chains, distinguishing among these four models will be difficult. Thorough investigation of the diversity of ZP domains in lineages that connect to the deepest nodes in the animal phylogeny (e.g., non-Bilateria, and possibly even closely-related non-animal groups (Swanson, et al. 2011)) will be key to testing these hypotheses.

Identifying highly divergent ZP-C domains will require a good understanding of the domain's sequence conservation patterns. In practical terms, this amounts to an understanding of cysteine conservation patterns, as most sites beyond these disulfide-forming cysteines are highly variable. Bokhove et al. (Bokhove, et al. 2016) argued that cysteine variation in ZP modules largely reflects departures from an otherwise conserved connectivity pattern involving three ZP-C domain disulfides—Cys5-Cys7, Cys6-Cys8, and CysA-CysB—with variation on this theme resulting from occasional losses and gains. This notion is consistent with the general evolutionary patterns observed for disulfide-forming cysteines—that these residues are generally highly conserved, and that they are almost always gained or lost in pairs (Thornton 1981; Rubinstein and Fiser 2008). By contrast, Han et al. (Han, et al. 2010) suggested that the novel ZP-C subdomain found in ZP₃ egg coat proteins accommodate alternative cysteine connectivity patterns in different species (though, as mentioned above, methodological issues might explain this apparent pattern; see Introduction). Extracellular proteins with

numerous, closely situated cysteines, such as ZPD proteins, seem like promising candidates for identifying unusual instances of disulfide reshuffling.

By combining phylogenetic and structural analyses, I found that disulfide variation among nematode ZP-C domains indeed reflects more than just gains and losses: the CysA-CysB disulfide was modified in Type 3 ZP-C domains, with CysA lost and replaced by a novel binding partner in the adjacent β C strand. The CysB- β C disulfide therefore represents a rare case of disulfide-bond reshuffling (Zhang 2007). Importantly, this modified disulfide is not some recently evolved outlier—it is a feature of multiple ZP module subfamilies (covering 12 *C. elegans* paralogs) that are shared across millions of distantly related nematode species. Given the phylogenetic depth of the branch where this reshuffling event is presumed to have occurred, close inspection of ZPD proteins in other invertebrate phyla might plausibly uncover orthologs that share this connectivity pattern. In light of its ancient origin and subsequent conservation across multiple subfamilies, it seems safe to conclude that stabilizing selection has acted to maintain the modified disulfide bond over time. However, it is not obvious whether the modified disulfide's initial origin was adaptive, and whether its evolution resulted in some novel function. For example, the evolution of an extra cysteine residue in the vicinity of CysB could have rendered CysA redundant, allowing for its exchange by drift. Another possibility is that the novel CysB-partner evolved to compensate for the loss of CysA; here, the novel disulfide would be adaptive only in the sense that it corrected some transient maladaptation, with no net change in overall function. Regardless, this finding speaks to the challenges of categorizing proteins using sequence conservation patterns without a robust phylogenetic framework, and to the importance of utilizing new data to update expectations about protein biology.

In contrast, there are several reasons to suspect that the entirely novel disulfide inferred between the β A and β B strands of Type 2 and 3 ZP-C domains is adaptive. First, it occurs in a region of known functional importance: the β A-IHP. Stabilizing the IHP through a disulfide bond could help maintain the tertiary structure of the ZP-C domain upon protein maturation and activation, during which the cleaved C-terminal tail's β G-EHP dissociates from the IHP (Jovine, et al. 2004; Schaeffer, et al. 2009). Second, it is notable that an IHP-stabilizing disulfide evolved independently within the FBN-1 subfamily (a Type 1 module). Convergent evolution is considered one of the strongest forms of observational evidence for adaptation and it seems unlikely that IHP-stabilizing disulfides would evolve repeatedly without providing some benefit to ZP-C domain structure or function. And finally, there is a clear pattern of co-evolution at several nearby sites alongside the same face of the β A and β B strands. These sites are largely fixed for aromatic residues in Type 2 and 3 modules. Fixing aromatic residues along the β A and β B strands may help to stabilize the β A- β B disulfide, act to slow EHP dissociation, or specify a critical interprotein binding surface that is only exposed after EHP release (Bhattacharyya, et al. 2004; Moreira, et al. 2013). Interestingly, it was recently demonstrated that disulfide bonds act as an evolutionary buffer, increasing tolerance for amino acid substitutions that would have ordinarily been structurally disruptive (Feyertag and Alvarez-Ponce 2017); the fixation of several aromatic residues around the novel β A- β B disulfide provides a clear counterexample to this claim. Determining the functional and evolutionary consequences of these convergently evolved disulfide bonds has the potential to provide important insights into the how IHP-EHP interactions affect ZP module activation.

The present study serves as the largest comparative investigation of ZP module evolutionary diversity conducted to date. By combining the newly-estimated nematode ZP module phylogeny with homology modelling of *C. elegans* ZPD proteins, I uncovered evidence for: (1) the parallel loss of the ZP-N domain in at least two lineages, resulting in the unexpected discovery of standalone ZP-C domains; (2) the modification of a highly conserved ZP-C domain disulfide via a rare example of cysteine replacement; and (3) the convergent gain of stabilizing disulfide bonds in the ZP-C domain's regulatory IHP motif. As a purely *in silico* study, it is of course critical that the unusual structural features documented here be confirmed experimentally. Even still, these findings have important implications for our understanding of ZP module structure and function. Moreover, the present study presents a valuable phylogenetic framework for the developmental genetic study of ZPD proteins in nematodes, including the powerful lab model *C. elegans*. Finally, this work sets the stage for future investigation of ZPD protein diversity in the broad sense. Here, the obvious next step will be to bridge the phylogenetic gap between nematodes, other invertebrates and, ultimately, vertebrates.

Acknowledgements

My research is funded by the Royal Society through the Dorothy Hodgkin Fellowship and Enhancement Award programs. I would like to acknowledge the School of Life Sciences at the University of Sussex for providing desk space and IT access during the early stages of this project, and Prof. Ralf Sommer for granting access to the MPI-DevBiol computer cluster. Dr. Christian Rödelsperger, Dr. Gabriel Markov, and three anonymous reviewers provided helpful comments on the manuscript. Finally, I would like to thank Dr. Bonnie Fraser for her support throughout the duration of the project.

Figure Captions

Figure 1: Nematode ZP module amino acid conservation patterns. Residue height indicates its prevalence in the top scoring alignment. Connections between cysteine residues indicate inferred disulfide linkages; also shown are the approximate boundaries of the ZP-N and ZP-C domains, the internal and external hydrophobic patches (IHP/EHP), and the consensus cleavage site (CCS). Non-homologous flanks and insertions were trimmed from the sequences as part of the alignment process; the relationship between alignment numbering and untrimmed sequence position for untrimmed *C. elegans* CUT-1 is provided in Supplementary Figure 1.

Figure 2: Nematode ZP module phylogeny. (a) The majority rule consensus of ML phylogenies estimated for 100 replicate ZP module alignments. Branch recovery proportions (BRPs) are shown using coloured circles; darker/redder circles indicate greater robustness to alignment variation. Branch lengths, drawn in amino acid substitutions per site (see scale bar), were estimated via ML using the top scoring alignment. The labelled arrows indicate the Minimal Ancestor Deviation (MAD) root position and the phylogenetic midpoint. Tip names are shown for *C. elegans* ZP modules; for clarity, CUT-1 was moved slightly to avoid overlap with CUT-3. Three major subtrees are noted (Types 1/2/3), the members of which are defined by different ZP-C domain cysteine connectivity patterns. (b) Cysteine connectivity patterns for Type 1, 2, and 3 ZP-C domains, inferred based on amino acid conservation patterns and homology modelling of *C. elegans* ZPD proteins. The β -strand secondary structure diagram follows that of the human uromodulin ZP-C domain (Bokhove, et al. 2016). The position of the MAD root in (a) suggests that the Type 1 connectivity pattern represents the ancestral state.

Figure 3: ZP-N domain loss and the structure of standalone ZP-C domain proteins. (a,b) Domain-specific gap proportions were calculated for each sequence (averaged over the 100 replicate alignments) and mapped onto the phylogeny: (a) ZP-N domain; (b) ZP-C domain. Gap proportions of 1 (= 100%) indicate cases where the entire domain is missing from the core alignment (coloured circles on the tips of the phylogeny; see legend). Nearly complete signatures of ZP-N-specific domain loss were observed for the To1D1.8, F46G11.6, and CUTL-19 subfamilies. (c-e) Homology models for *C. elegans* proteins with standalone ZP-C domains (pink lines) superimposed on the template structure, human uromodulin (grey cartoon): (c) To1D1.8b, (d) F46G11.6, (e) CUTL-19b. Cysteine residues in the *C. elegans* ZP-C domains are shown in stick format; the three disulfide linkages present in the template (Cys5-Cys7, Cys6-Cys8, and CysA-CysB) are shown as grey dot clouds.

Figure 4: The conserved internal hydrophobic patch (IHP) of Type 2/3 ZP-C domains. Homology models for the ZP-C domains of *C. elegans* CUT-1 (Type 3) and DPY-1a (Type 2) (both shown in pink lines, with key residues shown in stick format) were superimposed on the cartoon structure of the template, human uromodulin (grey cartoon, with the β A and β B strands coloured blue). The residues along the inward facing side of β A comprise the IHP; those residues, and the three adjacent residues in β B, are highly conserved in nematode Type 2 and 3 ZP-C domains and suggest a novel disulfide bond. These same sites are variable in Type 1 ZP-C domains. Conservation patterns for the three ZP-C domain types are shown via sequence logos (extracted from Supplementary Figure 4).

References

- Agrawal GK, Jwa NS, Lebrun MH, Job D, Rakwal R. 2010. Plant secretome: Unlocking secrets of the secreted proteins. *Proteomics* 10:799-827.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403-410.
- Bhattacharyya R, Pal D, Chakrabarti P. 2004. Disulfide bonds, their stereospecific environment and conservation in protein structures. *Protein Eng Des Sel* 17:795-808.
- Blaxter ML, De Ley P, Garey JR, Liu LX, Scheldeman P, Vierstraete A, Vanfleteren JR, Mackey LY, Dorris M, Frisse LM, et al. 1998. A molecular evolutionary framework for the phylum Nematoda. *Nature* 392:71-75.
- Boja ES, Hoodbhoy T, Fales HM, Dean J. 2003. Structural characterization of native mouse zona pellucida proteins using mass spectrometry. *Journal of Biological Chemistry* 278:34189-34202.
- Bokhove M, Jovine L. 2018. Structure of Zona Pellucida Module Proteins. *Curr Top Dev Biol* 130:413-442.
- Bokhove M, Nishimura K, Brunati M, Han L, de Sanctis D, Rampoldi L, Jovine L. 2016. A structured interdomain linker directs self-polymerization of human uromodulin. *Proc Natl Acad Sci U S A* 113:1552-1557.
- Bork P, Downing AK, Kieffer B, Campbell ID. 1996. Structure and distribution of modules in extracellular proteins. *Quarterly Reviews of Biophysics* 29:119-167.
- Bork P, Sander C. 1992. A large domain common to sperm receptors (Zp2 and Zp3) and TGF-beta type III receptor. *FEBS Lett* 300:237-240.
- Callebaut I, Morion JP, Monget P. 2007. Isolated ZP-N domains constitute the N-terminal extensions of Zona Pellucida proteins. *Bioinformatics* 23:1871-1874.
- Coghlan A, Tyagi R, Cotton JA, Holroyd N, Rosa BA, Tsai IJ, Laetsch DR, Beech RN, Day TA, Hallsworth-Pepin K, et al. 2019. Comparative genomics of the major parasitic worms. *Nature Genetics* 51:163-+.
- Cohen JD, Flatt KM, Schroeder NE, Sundaram MV. 2019. Epithelial Shaping by Diverse Apical Extracellular Matrices Requires the Nidogen Domain Protein DEX-1 in *Caenorhabditis elegans*. *Genetics* 211:185-200.
- Crooks GE, Hon G, Chandonia JM, Brenner SE. 2004. WebLogo: a sequence logo generator. *Genome Res* 14:1188-1190.
- Cuesta-Astroz Y, de Oliveira FS, Nahum LA, Oliveira G. 2017. Helminth secretomes reflect different lifestyles and parasitized hosts. *International Journal for Parasitology* 47:529-544.
- Darie CC, Biniossek ML, Jovine L, Litscher ES, Wassarman PM. 2004. Structural characterization of fish egg vitelline envelope proteins by mass spectrometry. *Biochemistry* 43:7459-7478.


- Devuyst O, Olinger E, Rampoldi L. 2017. Uromodulin: from physiology to rare and complex kidney disorders. *Nature Reviews Nephrology* 13:525-544.
- Diestel U, Resch M, Meinhardt K, Weiler S, Hellmann TV, Mueller TD, Nickel J, Eichler J, Muller YA. 2013. Identification of a Novel TGF-beta-Binding Site in the Zona Pellucida C-terminal (ZP-C) Domain of TGF-beta-Receptor-3 (TGFR-3). *PLoS One* 8:e67214.
- Duckert P, Brunak S, Blom N. 2004. Prediction of proprotein convertase cleavage sites. *Protein Eng Des Sel* 17:107-112.
- Fernandes I, Chanut-Delalande H, Ferrer P, Latapie Y, Waltzer L, Affolter M, Payre F, Plaza S. 2010. Zona pellucida domain proteins remodel the apical compartment for localized cell shape changes. *Dev Cell* 18:64-76.
- Feyertag F, Alvarez-Ponce D. 2017. Disulfide Bonds Enable Accelerated Protein Evolution. *Molecular Biology and Evolution* 34:1833-1837.
- Frand AR, Russel S, Ruvkun G. 2005. Functional genomic analysis of *C. elegans* molting. *PLoS Biol* 3:e312.
- Fujimoto D, Kanaya S. 1973. Cuticlin: a noncollagen structural protein from *Ascaris* cuticle. *Arch Biochem Biophys* 157:1-6.
- Gill HK, Cohen JD, Ayala-Figueroa J, Forman-Rubinsky R, Poggioli C, Bickard K, Parry JM, Pu P, Hall DH, Sundaram MV. 2016. Integrity of Narrow Epithelial Tubes in the *C. elegans* Excretory System Requires a Transient Luminal Matrix. *PLoS Genet* 12:e1006205.
- Goldman N, Yang Z. 1994. A codon-based model of nucleotide substitution for protein-coding DNA sequences. *Mol Biol Evol* 11:725-736.
- Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* 59:307-321.
- Han L, Monne M, Okumura H, Schwend T, Cherry AL, Flot D, Matsuda T, Jovine L. 2010. Insights into egg coat assembly and egg-sperm interaction from the X-ray structure of full-length ZP3. *Cell* 143:404-415.
- Heiman MG, Shaham S. 2009. DEX-1 and DYF-7 establish sensory dendrite length by anchoring dendritic tips during cell migration. *Cell* 137:344-355.
- Jovine L, Darie CC, Litscher ES, Wassarman PM. 2005. Zona pellucida domain proteins. *Annu Rev Biochem* 74:83-114.
- Jovine L, Janssen WG, Litscher ES, Wassarman PM. 2006. The PLAC1-homology region of the ZP domain is sufficient for protein polymerisation. *BMC Biochem* 7:11.
- Jovine L, Qi H, Williams Z, Litscher E, Wassarman PM. 2002. The ZP domain is a conserved module for polymerization of extracellular proteins. *Nat Cell Biol* 4:457-461.

- Jovine L, Qi H, Williams Z, Litscher ES, Wassarman PM. 2004. A duplicated motif controls assembly of zona pellucida domain proteins. *Proc Natl Acad Sci U S A* 101:5922-5927.
- Kallberg M, Wang H, Wang S, Peng J, Wang Z, Lu H, Xu J. 2012. Template-based protein structure modeling using the RaptorX web server. *Nat Protoc* 7:1511-1522.
- Kanai S, Kitayama T, Yonezawa N, Sawano Y, Tanokura M, Nakano M. 2008. Disulfide linkage patterns of pig zona pellucida glycoproteins ZP3 and ZP4. *Mol Reprod Dev* 75:847-856.
- Kelley M, Yochem J, Krieg M, Calixto A, Heiman MG, Kuzmanov A, Meli V, Chalfie M, Goodman MB, Shaham S, et al. 2015. FBN-1, a fibrillin-related protein, is required for resistance of the epidermis to mechanical deformation during *C. elegans* embryogenesis. *Elife* 4.
- Killingbeck EE, Swanson WJ. 2018. Egg Coat Proteins Across Metazoan Evolution. *Curr Top Dev Biol* 130:443-488.
- Korhonen PK, Pozio E, La Rosa G, Chang BCH, Koehler AV, Hoberg EP, Boag PR, Tan P, Jex AR, Hofmann A, et al. 2016. Phylogenomic and biogeographic reconstruction of the *Trichinella* complex. *Nature Communications* 7.
- Lee RYN, Howe KL, Harris TW, Arnaboldi V, Cain S, Chan J, Chen WJ, Davis P, Gao S, Grove C, et al. 2018. WormBase 2017: molting into a new stage. *Nucleic Acids Res* 46:D869-D874.
- Lefort V, Longueville JE, Gascuel O. 2017. SMS: Smart Model Selection in PhyML. *Mol Biol Evol* 34:2422-2424.
- Lemoine F, Domelevo Entfellner JB, Wilkinson E, Correia D, Davila Felipe M, De Oliveira T, Gascuel O. 2018. Renewing Felsenstein's phylogenetic bootstrap in the era of big data. *Nature* 556:452-456.
- Lewis E, Sebastiano M, Nola M, Zei F, Lassandro F, Ristoratore F, Cermola M, Favre R, Bazzicalupo P. 1994. Cuticlin Genes of Nematodes. *Parasite* 1:57-58.
- Li W, Cowley A, Uludag M, Gur T, McWilliam H, Squizzato S, Park YM, Buso N, Lopez R. 2015. The EMBL-EBI bioinformatics web and programmatic tools framework. *Nucleic Acids Res* 43:W580-584.
- Lin SJ, Hu Y, Zhu J, Woodruff TK, Jardetzky TS. 2011. Structure of betaglycan zona pellucida (ZP)-C domain provides insights into ZP-mediated protein polymerization and TGF-beta binding. *Proc Natl Acad Sci U S A* 108:5232-5236.
- Litscher E, Wassarman PM. 2015. *Zona pellucida domain proteins*. Hoboken, New Jersey: Wiley.
- Lockhead D, Schwarz EM, O'Hagan R, Bellotti S, Krieg M, Barr MM, Dunn AR, Sternberg PW, Goodman MB. 2016. The tubulin repertoire of *C. elegans* sensory neurons and its context-dependent role in process outgrowth. *Mol Biol Cell*.
- Martin AC, Orengo CA, Hutchinson EG, Jones S, Karmirantzou M, Laskowski RA, Mitchell JB, Taroni C, Thornton JM. 1998. Protein folds and functions. *Structure* 6:875-884.


- Matveev IV, Shaposhnikova TG, Podgornaya OI. 2007. A novel *Aurelia aurita* protein mesoglein contains DSL and ZP domains. *Gene* 399:20-25.
- Monne M, Han L, Schwend T, Burendahl S, Jovine L. 2008. Crystal structure of the ZP-N domain of ZP₃ reveals the core fold of animal egg coats. *Nature* 456:653-657.
- Moreira IS, Martins JM, Ramos RM, Fernandes PA, Ramos MJ. 2013. Understanding the importance of the aromatic amino-acid residues as hot-spots. *Biochim Biophys Acta* 1834:404-414.
- Muriel JM, Brannan M, Taylor K, Johnstone IL, Lithgow GJ, Tuckwell D. 2003. *M14.2.2 (cut-6)*, a novel *Caenorhabditis elegans* matrix gene important for dauer body shape. *Dev Biol* 260:339-351.
- Naba A, Clauser KR, Ding HM, Whittaker CA, Carr SA, Hynes RO. 2016. The extracellular matrix: Tools and insights for the "omics" era. *Matrix Biology* 49:10-24.
- Neuwald AF, Altschul SF. 2016. Bayesian Top-Down Protein Sequence Alignment with Inferred Position-Specific Gap Penalties. *PLoS Comput Biol* 12:e1004936.
- Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 32:268-274.
- Okumura H, Sato T, Sakuma R, Fukushima H, Matsuda T, Ujita M. 2015. Identification of distinctive interdomain interactions among ZP-N, ZP-C and other domains of zona pellucida glycoproteins underlying association of chicken egg-coat matrix. *FEBS Open Bio* 5:454-465.
- Ondrovics M, Gasser RB, Joachim A. 2016. Recent Advances in Elucidating Nematode Moulting - Prospects of Using *Oesophagostomum dentatum* as a Model. *Advances in Parasitology*, Vol 91 91:233-264.
- Pettitt J, Philippe L, Sarkar D, Johnston C, Gothe HJ, Massie D, Connolly B, Muller B. 2014. Operons are a conserved feature of nematode genomes. *Genetics* 197:1201-1211.
- Pierleoni A, Martelli PL, Casadio R. 2008. PredGPI: a GPI-anchor predictor. *BMC Bioinformatics* 9:392.
- Plaza S, Chanut-Delalande H, Fernandes I, Wassarman PM, Payre F. 2010. From A to Z: apical structures and zona pellucida-domain proteins. *Trends Cell Biol* 20:524-532.
- Popescu AA, Huber KT, Paradis E. 2012. ape 3.0: New tools for distance-based phylogenetics and evolutionary analysis in R. *Bioinformatics* 28:1536-1537.
- Raj I, Sadat Al Hosseini H, Dioguardi E, Nishimura K, Han L, Villa A, de Sanctis D, Jovine L. 2017. Structural Basis of Egg Coat-Sperm Recognition at Fertilization. *Cell* 169:1315-1326 e1317.
- Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* 3:217-223.

- Rubinstein R, Fiser A. 2008. Predicting disulfide bond connectivity in proteins by correlated mutations analysis. *Bioinformatics* 24:498-504.
- Saito T, Bokhove M, Croci R, Zamora-Caballero S, Han L, Letarte M, de Sanctis D, Jovine L. 2017. Structural Basis of the Human Endoglin-BMP9 Interaction: Insights into BMP Signaling and HHT1. *Cell Rep* 19:1917-1928.
- Sapio MR, Hilliard MA, Cermola M, Favre R, Bazzicalupo P. 2005. The Zona Pellucida domain containing proteins, CUT-1, CUT-3 and CUT-5, play essential roles in the development of the larval alae in *Caenorhabditis elegans*. *Dev Biol* 282:231-245.
- Schaeffer C, Santambrogio S, Perucca S, Casari G, Rampoldi L. 2009. Analysis of uromodulin polymerization provides new insights into the mechanisms regulating ZP domain-mediated protein assembly. *Mol Biol Cell* 20:589-599.
- Schliep KP. 2011. phangorn: phylogenetic analysis in R. *Bioinformatics* 27:592-593.
- Sebastiano M, Lassandro F, Bazzicalupo P. 1991. cut-1 a *Caenorhabditis elegans* gene coding for a dauer-specific noncollagenous component of the cuticle. *Dev Biol* 146:519-530.
- Spencer WC, Zeller G, Watson JD, Henz SR, Watkins KL, McWhirter RD, Petersen S, Sreedharan VT, Widmer C, Jo J, et al. 2011. A spatial and temporal map of *C. elegans* gene expression. *Genome Res* 21:325-341.
- Swanson WJ, Aagaard JE, Vacquier VD, Monne M, Sadat Al Hosseini H, Jovine L. 2011. The molecular basis of sex: linking yeast to human. *Mol Biol Evol* 28:1963-1966.
- Swanson WJ, Nielsen R, Yang Q. 2003. Pervasive adaptive evolution in mammalian fertilization proteins. *Mol Biol Evol* 20:18-20.
- Thornton JM. 1981. Disulphide bridges in globular proteins. *J Mol Biol* 151:261-287.
- Tria FDK, Landan G, Dagan T. 2017. Phylogenetic rooting using minimal ancestor deviation. *Nature Ecology & Evolution* 1.
- Verhoeven K, Van Laer L, Kirschhofer K, Legan PK, Hughes DC, Schatteman I, Verstreken M, Van Hauwe P, Coucke P, Chen A, et al. 1998. Mutations in the human alpha-tectorin gene cause autosomal dominant non-syndromic hearing impairment. *Nature Genetics* 19:60-62.
- Vogel C, Berzuini C, Bashton M, Gough J, Teichmann SA. 2004. Supra-domains: Evolutionary units larger than single protein domains. *Journal of Molecular Biology* 336:809-823.
- Vuong-Brender TTK, Suman SK, Labouesse M. 2017. The apical ECM preserves embryonic integrity and distributes mechanical stress during morphogenesis. *Development* 144:4336-4349.
- Wang S, Peng J, Xu J. 2011. Alignment of distantly related protein structures: algorithm, bound and implications to homology modeling. *Bioinformatics* 27:2537-2545.


- Wilburn DB, Swanson WJ. 2017. The "ZP domain" is not one, but likely two independent domains. *Mol Reprod Dev* 84:284-285.
- Witte H, Moreno E, Rodelsperger C, Kim J, Kim JS, Streit A, Sommer RJ. 2015. Gene inactivation using the CRISPR/Cas9 system in the nematode *Pristionchus pacificus*. *Dev Genes Evol* 225:55-62.
- Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* 24:1586-1591.
- Yonezawa N. 2014. Posttranslational modifications of zona pellucida proteins. *Adv Exp Med Biol* 759:111-140.
- Yu RY, Nguyen CQ, Hall DH, Chow KL. 2000. Expression of *ram-5* in the structural cell is required for sensory ray morphogenesis in *Caenorhabditis elegans* male tail. *EMBO J* 19:3542-3555.
- Zhang J. 2007. Disulfide-bond reshuffling in the evolution of an ape placental ribonuclease. *Mol Biol Evol* 24:505-512.




(a)




(b)







(c) T01D1.8b




(d) F46G11.6



(e) CUTL-19b






ZP-C type
Type 3

Type 2

Type 1



<http://mc.manuscriptcentral.com/gbe>