

RESEARCH ARTICLE


Psychometric properties of The Baruth Protective Factors Inventory among nursing students

Tayyebeh Ali-Abadi¹ , Siavash Talepasand² , Christopher Boyle³ , Hamid Sharif Nia^{4*} 

1 Department of Nursing, Neyshabur Branch, Islamic Azad University, Neyshabur, Iran, **2** Department of Psychology and Educational Sciences, Semnan University, Semnan, Iran, **3** Education and Psychology, Graduate School of Education, University of Exeter, Exeter, United Kingdom, **4** School of Nursing and Midwifery Amol, Mazandaran University of Medical Sciences, Sari, Iran

 These authors contributed equally to this work.

* pegadis@yahoo.com



OPEN ACCESS

Citation: Ali-Abadi T, Talepasand S, Boyle C, Sharif Nia H (2020) Psychometric properties of The Baruth Protective Factors Inventory among nursing students. PLoS ONE 15(6): e0233760. <https://doi.org/10.1371/journal.pone.0233760>

Editor: César Leal-Costa, Murcia University, SPAIN

Received: August 28, 2019

Accepted: May 12, 2020

Published: June 5, 2020

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0233760>

Copyright: © 2020 Ali-Abadi et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: The author(s) received no specific funding for this work

Abstract

The Baruth Protective Factors Inventory is an instrument which assesses the risk and protective factors behind resilience. However, there is no valid or reliable Persian instrument for measuring resilience amongst nursing students in Iran. This study aimed to evaluate the psychometric properties of the Persian version of The Baruth Protective Factors Inventory among nursing students. This methodological study was done in 2017. The participants were 200 nursing students who were randomly recruited from Neyshabur city. Construct validity was assessed using exploratory and confirmatory factor analysis. Reliability was also assessed through the internal consistency assessment method. Exploratory factor analysis indicated a three-factor structure for the Inventory which accounted for 48.21% of its total variance. Confirmatory factor analysis confirmed the first-order model. The internal consistency values of the Inventory and its domains were good, confirming its great internal consistency and reliability. The Persian version of the Baruth Protective Factors Inventory was found to have acceptable validity and reliability to assess resilience amongst nursing students. Therefore, the Persian version of The Baruth Protective Factors Inventory can help nursing authorities identify non-resilient students, promote their resilience, and thereby improve their academic and clinical performance. Nursing students' improved performance can positively affect both the quality of care and patient outcomes.

Introduction

Nursing students are presented and need to cope with a wide range of challenges that are unique to their profession. For example, care delivery to patients with different physical and mental health problems, patients' death, and the risk of being infected by infectious diseases. These stressors cause nursing students different levels of anxiety, stress, and discomfort [1], also high levels of stress can affect them psychologically and cause sleep disorders, reduction in

Competing interests: The authors have declared that no competing interests exist.

their motivation for clinical activities, and undermine their self-confidence [2]. Simonelli-Muñoz et al (2018) found stress had worse academic performance in nursing students [3].

Resilience is an appropriate strategy for stress management [4]. Students' resilience can be promoted through providing themselves with protective factors such as a caring learning environment, learner-centered education, strong instructor-student relationship, and positive expectations Whereas teaching and learning processes focus merely on information delivery, students do not get psychologically prepared for clinical practice and thus, may be vulnerable to occupational stressors later in their professional lives [5].

However, resilience evaluation necessitates valid and reliable instruments, but due to diversity in resilience definitions, different instruments have been developed for its measurement [6]. These instruments include, but are not limited to, Brief Resilience Scale [7], Connor-Davidson Resilience Scale [8], Baruth Protective Factor Inventory [9], Resilience Scale for Adults [10], and the Brief Resilience Coping Scale [11]. However, as resilience is affected by many factors including context, time, age, and living conditions, its measurement tools also need to be context, time, and age-specific [12].

The conceptual framework of this study was based on the following definition of resilience: "an individualized process of development that occurs through the use of protective factors to successfully navigate perceived stress and adversities" [13]. In this framework, risk factors minimization and protective factors enhancement help promote the ability to cope with challenges in life. As Baruth quoted Wolff (1995) identifying protective stress-causing factors would play a significant role in clinical practices. Individuals with intervene risks, have further resilience while avoiding from environmental problems. When these people got further information about the reasons and protective strategies with the help of expert ones, more resilience and stability in their life would be predictable [9].

One of the research instruments that have been used to measure protective factors specifically in adults is the Baruth Protective Factors Inventory (BPMI). A significant strong point of BPMI theory is the rich theoretical basis for it. This valid and reliable inventory measures protective factors in an adult population through a review of resilience literature (resilience theories [9]). Assessing resilience through BPMI increases the chance of identifying concepts and variables which are important to nursing and health care practice. Therefore these variables will be clarified and important dimensions of the variables that should be measured will be determined [14]. Therefore, BPMI is particularly fascinating in nursing students to quickly identify stress which results in providing interventions to decrease the level of stress.

The BPMI discusses Protective Factors that are repeatedly used by nursing students [1],[15],[16],[5],[17],[18]. The BPMI has been used to examine resilience in various contexts, including among students at a public university in the southwest United States [19], the families of disabled children in Portugal [20], clinical and non-clinical groups [21]. There were no consistent findings regarding the factor structure of BPMI when tested using Confirmatory Factor Analysis (CFA) across different cultures [19, 20]. By considering these findings, previous validation studies in another setting, time and so on, are not warranted to remain valid in another time, culture or context. Therefore, regarding the deficiency of valid tools on nursing student setting the overall aim of this study was evaluation of the psychometric properties of the Persian version of BPMI among nursing students. Identifying the optimal inventory structure may encourage other researchers in this field to utilize this inventory, thus helping to develop the study of resilience in nursing student settings.

Materials and methods

This methodological study was done in 2017 to evaluate the psychometric properties of the Persian BPMI. The study population consisted of all 376 bachelor's students who were studying

nursing in Azad and medical science Universities in Neyshabur, Iran, in the 2017–2018 academic years. A sample of 200 students was recruited via simple random sampling. For instruments with less than forty items, samples of 200 were reported to be adequate [22]. Data collection was performed using BPF1. Baruth et al. developed BPF1 in 2002 based on the existing literature on resilience [9]. As BPF1 has 16 items, they are scored on a five-point Likert-type scale from 1 (“Strongly disagree”) to 5 (“Strongly agree”). Items 1 to 4 are reverse scored. The total BPF1 score varied between 16 and 80, with higher scores indicating more protective factors [21].

Procedure

After obtaining necessary permissions from its developers, the scale was translated from English into a Persian version based on the World Health Organization protocol of the forward-backward translation technique [23]. Two English-Persian translators were requested to independently translate the BPF1. An expert panel, consisting of some of this article’s authors, two professional translators, and a Persian-English bilingual speaker assessed and unified the two translations and constructed a single Persian translation of inventory. Thereafter, a Persian-English translator was asked to back-translate the Persian BPF1 into English. This English version of the Inventory was sent to Persian-English bilingual speaker, for confirmation of the correctness of translations and confirming the similarity of the achieved English BPF1 with its original. Then, the primary Persian BPF1 was piloted on a sample of 28 nursing students in small two to three-person groups. So, face validity assessed this way that Students were asked to comment on the clarity of the items. Their comments required us to revise two items (9 and 10 respectively). Finally, 200 nursing students completed the Persian BPF1 and the collected data were used for construct validity and reliability assessment.

Construct validity

Exploratory factor analysis. The construct validity of the scale was evaluated using Maximum Likelihood Exploratory Factor Analysis (MLEFA) with promax rotation. The Kaiser–Meyer–Olkin (KMO) test and Bartlett’s test of sphericity checked the appropriateness of the study sample and the factor analysis model. The test estimates sampling adequacy for each item in the model and the model as a whole. The indicator is a measure of the proportion of variance among variables that might be common variance [24]. KMO values between 0.8 and 1 indicate the sampling is adequate. In the first step, the latent factors were extracted based on Horn’s Parallel Analysis [25]. The presence of a single item in the factor based on the $CV = 5.152 \div \sqrt{(n-2)}$ formula [26] was estimated to be approximately 0.4 (in the present formula, the CV is the number of extractable factors and n is the sample size of the study). According to the three indicator rule, at least three items must exist for each factor [27]. According to the three-indicator rule, there must be at least three items for each latent variable in the EFA [28]. Items with communalities less than 0.2 were excluded from the EFA [29].

Confirmatory factor analysis. The extracted factors were evaluated by Confirmatory Factor Analysis (CFA) based on the Maximum Likelihood Method. CFA goodness of fit indexes includes the acceptable level of Parsimonious Normed Fit Index (PNFI), Parsimonious Comparative Fit Index (PCFI), Adjusted Goodness of Fit Index (AGFI) indices ($> .5$), Comparative of Fit Index (CFI), Incremental Fit Index (IFI) ($> .9$), Root Mean Square Error of Approximation (RMSEA): $> .08$ and Minimum Discrepancy Function by Degrees of Freedom divided (CMIN/DF) < 3 good [30].

Convergent and divergent validity. In this study for the estimation of convergent and divergent, we used the criterion of Fornell-Larcker [31]. The convergent and divergent validity of the BPF1 was assessed using the average shared squared variance (ASV), the maximum

shared squared variance (MSV) and the average variance extracted (AVE). The convergent validity is established when $AVE > 0.5$ and divergent validity is established when both $MSV < AVE$ and $ASV < AVE$ [31].

Reliability. To assess the internal consistency of the Inventory, coefficients of Cronbach's alpha, Average Inter-Item Correlation (AIC) and Omega McDonald were calculated [32]. Values higher than 0.6 were considered acceptable [33]. In AIC, a good value range .2 to .4 [34]. Construct Reliability (CR), which replaces Cronbach's alpha coefficient in structural equation modeling, was then assessed and CR greater than 0.7 were considered acceptable [35].

Normal distribution, outliers, and missing data. The Univariate and Multivariate normal distribution of data was assessed by skewness (± 3) and kurtosis (± 7) [30]. The presence of any multivariate outlier was investigated by Mahalanobis d-squared ($p < .001$) and multivariate normality by the Mardia coefficient of multivariate kurtosis (< 20) [36]. Missing values calculated by Multiple Imputation technique (which were less than 3%) and were substituted with the mean scores of the corresponding items. All of the statistical procedures were calculated by SPSS- AMOS₂₅ and SPSS R-Menu v2.0.

Ethical considerations. Ethical approval for this study was provided by the Ethics Committee of Semnan University of Medical Sciences, Semnan, Iran (Approval Number: 96.524132). Necessary permissions were obtained from the developers of BPFİ before its translation and psychometric evaluation. The study participants were informed about the study aim. Their agreement in responding to the BPFİ items was considered to be their consent for participation in the study.

Results

A total of 200 nursing student participants were mainly female (47.5%) and single (73%) with a mean age of 21.48 ± 2.86 (Table 1). The mean score of BPFİ score in nursing student was 48.41 ± 4.5 . There was significant difference between female and male nursing student in resilience score ($p = 0.04$).

The minimum score was 23 and the maximum score was 60. After the score reversal of items 9 and 10 in the first stage of face validity, the results showed all items of the inventory to be appropriate, clear, and straightforward.

Construct validity

In this study the KMO was .816 indicating the sample size was adequate for factor analysis and Bartlett's test of sphericity was significant ($\chi^2 = 1283.25$, $df = 105$; $p < .0001$), indicating that

Table 1. Participants' socio-demographic characteristics.

Characteristics		N (%)
Gender	Male	93 (46.5)
	Female	95 (47.5)
Semester	First	55 (27.5)
	Third	30 (15)
	Fifth	49 (24.5)
	Seventh	39 (19.5)
	Other	8 (4)
	Marital status	Single
	Married	43 (21.5)

<https://doi.org/10.1371/journal.pone.0233760.t001>

the correlation matrix was appropriate for factor analysis. The parallel analysis showed a three-factor structure for BPGI which accounted for 48.21% of the total variance (Table 2).

Next the extracted structure was assessed by CFA. The results of the Chi-square test for goodness-of-fit were first obtained as $\chi^2 = 134.420$ ($p < .0001$), and other indices were then assessed for the fit of the model. All the indices (PCFI = .760, CMIN/DF = 1.84, RMSEA = .065, IFI = .948 and CFI = .949) confirmed a good fit of the final model. For a better model, a modification was done to the final factor structure of the BPGI that there was a covary between the measurement errors of items 6 and 15 (Fig 1).

Assessing the ASV, MSV, and AVE, also indicated that the BPGI had good convergent and divergent validity. Also, internal consistency and CR of the BPGI were estimated to be desirable (Table 3).

Discussion

This study represents the first attempt to examine the reliability and validity of a Persian inventory of resilience; the BPGI among nursing students. Previous studies conducted during the development of the BPGI support that the instrument has content, construct, and face validity, as well as reliability in measuring the health-promotion concept of resilience in adults[21].

The study was conducted in three main steps, namely that of 1). Translation and cultural adaptation, 2) data collection, and finally 3). Exploratory and confirmatory factor analysis. The result showed that the BPGI included three distinct and stable factors which explain 48.21% of the total variance in the student nursing population.

However, the original BPGI has a four-factor structure including fewer stressors, adaptable personality, supportive environment, compensating experiences. In primary EFA, three factors were extracted. In this study the factors “Adaptable personality and compensatory experiences” considered as a unit. It is probable that clients in the nonclinical group recalled their families’ adaptive appraisals and compensating experiences as a unit, as well as the familial availability of these sources of competence contributing to their good adaptation[19]. Compensating experiences are defined as a family’s experiences of mastery within the context of

Table 2. Exploratory factor analysis of the Persian version of the BPGI.

Factor	Items	Factor loading	h^2^a	Eigenvalue	% of variance
supportive environment	Q10: I have at least one caring person in my life, (whether in your family or not).	.93	.81	3.91	26.06
	Q9: I have a good relationship with at least one supportive person, (whether in your family or not).	.93	.88		
	Q12: I have at least one person who is interested in my life (whether in your family or not).	.80	.68		
	Q11: I can trust at least one person in my life, (whether in your family or not).	.80	.69		
Compensatory Experience	Q15: I have coped well with one or more major stressors in my life.	.74	.59	3.36	22.40
	Q14: I have control over many (but not all) events in my life.	.74	.56		
	Q13: I have been able to resolve many (but not all) of my problems by my self	.61	.30		
	Q16: I have been able to make “the best out of a bad situation” several times.	.56	.33		
	Q6: I am a creative, resourceful, and independent person.	.49	.33		
	Q8: I am competent and have high self-esteem.	.48	.36		
	Q5: I am optimistic and concentrate on the positives in most situations.	.48	.25		
Fewer Stressor	Q4: I have had more problems than positive experiences with work/school in the past 3 months.	.62	.38	1.67	11.13
	Q2: I have had more problems than positive experiences with my finances in the past 3 months	.55	.33		
	Q3: I have had more problems than positive experiences with my family/friends in the past 3 months	.54	.32		
	Q1: I have had more problems than positive experiences with my health status in the past 3 months.	.53	.30		

^a Communalities




Fig 1. Final model of the first order CFA of the BPF1.

<https://doi.org/10.1371/journal.pone.0233760.g001>

adversity. This mastery includes feelings of positive control over uplifting experiences while having experienced the same situations as hassles [19]. Masten (2001) posited that the influence of family mastery resources as a compensatory factor [37]. As registered nurses, all of the participants had developed compensatory and coping strategies which had enabled them to achieve their qualification and to progress in their careers [38]. However, the Adaptive resources were identified as the ability to plan and the motivation to succeed in the future, behavioral and emotional autonomy, the capacity to handle stressful situations, and access to supportive adults [39].

According to low factor loading, Items 5 and 7 were removed from the Persian version of the scale It is valuable that the differences among different cultures can result in the instability of the factor structure. Studies in the last twenty years indicated that resilience is a multidimensional concept which is affected by context, time, age, gender, cultural origin, and living conditions. In line with our findings, factor analysis in an earlier study of 410 students at a public university in the southwest of the United States revealed a four-factor structure for BPF1 after

Table 3. Convergent and divergent validity, internal consistency, and constructs reliability of BPF1.

Factor	α (CI 95)	AIC	Ω	CR	AVE	MSV	ASV
Supportive Environment	.926 (.908 to .941)	.764	.929	.929	.766	.260	.135
Compensatory Experience	.797 (.750 to .835)	.361	.801	.792	.439	.260	.175
Fewer Stressor	.657 (.572 to .728)	.323	.658	.658	.325	.091	.051

CR: Construct Reliability; AVE: Average Variance Extracted; MSV: Maximum Shared Squared Variance; ASV: Average Shared Squared Variance.

<https://doi.org/10.1371/journal.pone.0233760.t003>

the removal of item 3 [19]. Another study on the families of disabled children in Portugal also reported a four-factor structure for the scale after the removal of item 3 from the fewer stressors domain [20].

According to the final model of the BPFPI, a relationship exists between the measurement error of item 6 (I am a creative, resourceful, and independent person), and 15 (I have coped well with one or more major stressors in my life). This relationship may fall, when the items are not properly understood or due to the conceptual similarity of the sentences. Coping is a protective mechanism that eliminates or modifies the situation, controls the meaning of the experience and manage the emotional consequences of the situation. Coping strategies generally seem to be more effective when an individual has more power or control over a situation [40]. Thus, this is the reason that both items have been put in the compensatory dimension of this study. Compensating experiences have been referred to as a sense of meaning and control over person-life [41] and as a person's experiences of mastery (feelings of positive control over uplifting experiences) within the context of adversity [19]. On the other hand, Compensatory mechanisms include psychological adaptations such as self-efficacy (the confidence a person to their ability to do relevant self-care activities) while the ability to cope with different situations is promoted when a person's self-efficacy is high and when person feels in control during threatening situations [42]. Children and adults who have a less negative assessment of difficult situations or who consider themselves to be in control of certain situations in their lives may be react less negatively to difficult situations and be better equipped to solve problems [43].

The factor "Social support" has been identified as a multidimensional factor that includes family, networks, community, school, employment and Consultant [19]. Social support depict medical students' resilience that experience adversity in their clinical setting [44]. In this inventory, support means a supportive environment aspect from other persons or families. A supportive learning environment promotes the resilience of nursing students, reducing their vulnerability [45, 46]. Staff support for students leads to stay student in the nursing program [44].

The factor "fewer stressors" assesses whether the person perceives more positive or negative experiences in their health status, finances, family and friends, and work or school [21]. ACHA identified stress, especially chronic stress as the number one barrier to college success for most college students [47]. Nursing students inevitably experience stress in their academic education. Resilience is a dynamic process that decreased the effects of stress and prevents the potentially disabling effects of chronic stress [44].

The results of this study also showed that the items of the BPFPI have an appropriate convergent and divergent validity in the final model. The convergent validity exists when the objects of the structure are close together and detect a large variance [35]. On the other hand, it has been stated that divergent validity exists when the factors of the considered structure or the latent extracted factors are completely separate from each other. In other words, we will not have a good convergent validity when the latent factors are not well explained by the extracted items and the items are not related to each other [48]. For example, items 13, 14, and 16 refer to self-efficacy. Students who have higher academic self-efficacy beliefs use positive coping strategies to deal with academic difficulties [49].

The many studies documented that nursing students exposed to different sources of stress during their educational program [13],[18] that include long hours of study and clinical pressures [50], problem with patients or their family, problems in relationship with nurses [51], health problems [52], economic problems [53]. The BPFPI has mentioned these sources of stress (Item1-4). Also the Baruth inventory emphasized protective factors in other items, which has permitted a person to overcome adversity. Researchers have stated that nursing students use these protective factors that include support [15], optimism [17], [1], self- confidence [16], [5], Relationship [5], for coping with stress.

The coefficients of Cronbach's alpha, Omega McDonald, AIC value confirmed the internal consistency of the overall inventory. The fewer stressors dimension presented a low coefficient of internal consistency. This result might be related to the limitation of time (the last three months) and the items in this dimension (personality characteristics or environmental resources that lead to experience greater maladjustment) [20] that interact with other dimension related to personality characteristics and supportive environmental resources that help to prevent maladjustment [54].

Gender is one of the factors influencing the understanding of psychological distress [53]. The number of male nursing students is continuing to increase [55]. The resilience score in this study was different between male and female students, which was consistent with other studies [57, 56]. In these studies, the ratio between females and males were not equal but in this study, the ratio was almost the same. Also data has been collected from multiple samples (Azad university and Neyshabur university of medical science) to fairly generalize the factorial structure and construct validity of this inventory but the present study has several limitations. First, the study relied on self-reported data collected at one time point; thus, follow-up data would be valuable in assessing whether resilience results are consistent in demonstrating dispositional traits of individuals and can help health care authorities understand changes in student nursing resilience over time. Second, the current study did not measure test-retest reliability as a vital indicator of scale reliability. Therefore, future research might consider measuring other indicators of reliability, such as test-retest among undergraduate college students. Despite the study limitations, the findings from the study indicated evidence of the BPFİ reliability and construct validity among undergraduate nursing students and suggest the use of the inventory to effectively assess resilience.

Further psychometric testing with other populations is warranted to enhance the BPFİ's usefulness.

Implications

In this study the first limitation was the lack of access to the sample, thus we had to run on the same sample as EFA and CFA. However we used other tests to increase the validity of the study. With fourteen items in three domains and acceptable validity and reliability, the Persian version of BPFİ can be used to diagnose non-resilience among nursing students. It is shorter than other resilience tools and can be completed in less time. Thus, it can be used for resilience assessment screening purposes and also for evaluating the effects of resilience-promotion interventions among nursing students in Iran.

Supporting information

S1 Data.

(SAV)

Acknowledgments

We deeply appreciate the entire student to participate in this study.

Author Contributions

Conceptualization: Tayyebeh Ali-Abadi.

Formal analysis: Hamid Sharif Nia.

Writing – original draft: Tayyebeh Ali-Abadi.

Writing – review & editing: Siavash Talepasand, Christopher Boyle, Hamid Sharif Nia.

References

1. Thomas LJ, Revell SH. Resilience in nursing students: An integrative review. *Nurs educ today*. 2016; 36:457–62.
2. Pourafzal F, Seyedfatemi N, Inanloo M, Haghani H. Relationship between Perceived Stress with Resilience among Undergraduate Nursing Students. *Hayat*. 2013; 19(1):41–52. (persian).
3. Simonelli-Muñoz AJ, Balanza S, Rivera-Caravaca JM, Vera-Catalán T, Lorente AM, Gallego-Gómez JI. Reliability and validity of the student stress inventory-stress manifestations questionnaire and its association with personal and academic factors in university students. *Nurs educ today*. 2018; 64:156–60.
4. Matua GA, Wal DMVd. Living Under the Constant Threat of Ebola: A Phenomenological Study of Survivors and Family Caregivers During an Ebola Outbreak. *J Nurs Res* 2015; 23(3):217–24. <https://doi.org/10.1097/jnr.000000000000116> PMID: 26270268
5. McAllister M, McKinnon J. The importance of teaching and learning resilience in the health disciplines: a critical review of the literature. *Nurs educ today*. 2009; 29(4):371–9.
6. Xie Y, Peng L, Zuo X, Li M. The Psychometric Evaluation of the Connor-Davidson Resilience Scale Using a Chinese Military Sample. *PLoS One*. 2016; 11(2).
7. Smith BW, Dalen J, Wiggins K, Tooley E, Christopher P, Bernard J. The brief resilience scale: assessing the ability to bounce back. *Int J Behav Med*. 2008; 15(3):194–200. <https://doi.org/10.1080/10705500802222972> PMID: 18696313
8. Connor KM, Davidson JR. Development of a new resilience scale: The Connor-Davidson resilience scale (CD-RISC). *Depress Anxiety*. 2003; 18(2):76–82. <https://doi.org/10.1002/da.10113> PMID: 12964174
9. Baruth KE, Caroll JJ. A formal assessment of resilience: The Baruth Protective Factors Inventory. *J Individ Psychol*. 2002.
10. Friborg O, Hjemdal O, Rosenvinge JH, Martinussen M. A new rating scale for adult resilience: what are the central protective resources behind healthy adjustment? *Int J Methods Psychiatr Res*. 2003; 12(2):65–76. <https://doi.org/10.1002/mpr.143> PMID: 12830300
11. Sinclair VG, Wallston KA. The development and psychometric evaluation of the Brief Resilient Coping Scale. *Assessment*. 2004; 11(1):94–101. <https://doi.org/10.1177/1073191103258144> PMID: 14994958
12. Singh K, Yu X-n. Psychometric evaluation of the Connor-Davidson Resilience Scale (CD-RISC) in a sample of Indian students. *J Psychol*. 2010; 1(1):23–30.
13. Stephens TM. Nursing student resilience: a concept clarification. *Nurs Forum*. 2013; 48(2):125–33. <https://doi.org/10.1111/nuf.12015> PMID: 23600641
14. Waltz CF, Strickland OL, Lenz ER. *Measurement in nursing and health research*: Springer publishing company; 2010.
15. Dugué M DF. Understanding Nursing Student Stress and its Consequences: A Qualitative Study in France. *BAOJ Med Nursing & Health Sciences*. 2018; 4(4):1–8.
16. Sharif F, Masoumi S. A qualitative study of nursing student experiences of clinical practice. *BMC nurs*. 2005; 4(1):6.
17. Snyder C, Rand KL, Sigmon DR. Hope theory. *Handbook of positive psychology*. 2002:257–76.
18. Mason H. The experience of resilience among nursing students: a qualitative study. *New Generat Sci*. 2018; 16(1):62–78.
19. Gardner DL, Huber CH, Steiner R, Vazquez LA, Savage TA. The development and validation of the inventory of family protective factors: A brief assessment for family counseling. *Fam J*. 2008; 16(2):107–17.
20. Augusto CCVCd, Ferreira O, Araújo BR, Rodrigues VMCP, Figueiredo MdCAB. Adaptation and validation of the Inventory of Family Protective Factors for the Portuguese culture. *Rev Lat Am Enfermagem*. 2014; 22(6):1001–8. <https://doi.org/10.1590/0104-1169.3315.2509> PMID: 25591096
21. Baruth KE. The Baruth Protective Factors Inventory as a clinical assessment of resilience 2004.
22. Lin SY, Tseng WT, Hsu MJ, Chiang HY, Tseng HC. A psychometric evaluation of the Chinese version of the nursing home survey on patient safety culture. *J Clin Nurs*. 2017; 26(23–24):4664–74. <https://doi.org/10.1111/jocn.13815> PMID: 28334463
23. World Health Organization. Process of translation and adaptation of instruments 2016. Available from: http://www.who.int/substance_abuse/research_tools/translation/en/.
24. Cerny BA, Kaiser HF. A study of a measure of sampling adequacy for factor-analytic correlation matrices. *Multivariate Behav Res*. 1977; 12(1):43–7. https://doi.org/10.1207/s15327906mbr1201_3 PMID: 26804143
25. Çokluk Ö, Koçak D. Using Horn's Parallel Analysis Method in Exploratory Factor Analysis for Determining the Number of Factors. *Educ Sci Theory Pract*. 2016; 16(2):537–51.

26. Fok D. Development and Testing of a Low Vision Product Selection Instrument (LV-PSI): A Mixed-Methods Approach. 2011.
27. Munro BH. Statistical methods for health care research: lippincott williams & wilkins; 2005.
28. Pett MA, Lackey NR, Sullivan JJ. Making sense of factor analysis: The use of factor analysis for instrument development in health care research: Sage; 2003.
29. Hahs-Vaughn DL. Applied multivariate statistical concepts: Routledge; 2016.
30. Pahlevan Sharif S, Sharif Nia H. Structural Equation Modeling with AMOS. Tehran: Artin Teb; 2018.
31. Fornell C, Larcker DF. Evaluating structural equation models with unobservable variables and measurement error. *J Mark Res.* 1981; 18(1):39–50.
32. Javali SB, Gudaganavar NV, Raj SM. Effect of varying sample size in estimation of coefficients of internal consistency. 2011.
33. Altman DG. Practical statistics for medical research: CRC press; 1990.
34. Michalos AC. Encyclopedia of quality of life and well-being research: Springer Netherlands Dordrecht; 2014.
35. Hair J, Anderson R, Black B, Babin B. Multivariate Data Analysis: Pearson Education; 2016.
36. Esposito Vinzi V, Chin WW, Henseler J, Wang H. Handbook of partial least squares: Concepts, methods and applications: Heidelberg, Dordrecht, London, New York: Springer; 2010.
37. Masten AS. Ordinary magic: Resilience processes in development. *Am Psychol.* 2001; 56(3):227. <https://doi.org/10.1037//0003-066x.56.3.227> PMID: 11315249
38. Major R, Tetley J. Effects of dyslexia on registered nurses in practice. *Nurse Educ Pract.* 2019; 35:7–13. <https://doi.org/10.1016/j.nepr.2018.12.012> PMID: 30633989
39. Maginness A. The development of resilience-a model. 2007.
40. Walton RL. A comparison of perceived stress levels and coping styles of junior and senior students in nursing and social work programs: Citeseer; 2002.
41. Papalia DE, Olds SW, Feldman RD. Human development (81ed.). New York: McGraw Hill; 2001.
42. Robinson-Smith G, Johnston MV, Allen J. Self-care self-efficacy, quality of life, and depression after stroke. *Arch Phys Med Rehabil.* 2000; 81(4):460–4. <https://doi.org/10.1053/mr.2000.3863> PMID: 10768536
43. Vanderbilt-Adriance E, Shaw DS. Conceptualizing and re-evaluating resilience across levels of risk, time, and domains of competence. *Clin Child Fam Psychol Rev.* 2008; 11(1–2):30. <https://doi.org/10.1007/s10567-008-0031-2> PMID: 18379875
44. Reyes AT, Andrusyszyn MA, Iwasiw C, Forchuk C, Babenko-Mould Y. Nursing students' understanding and enactment of resilience: a grounded theory study. *J Adv Nurs.* 2015; 71(11):2622–33. <https://doi.org/10.1111/jan.12730> PMID: 26193950
45. Shiu AT, Chan CW, Lam P, Lee J, Kwong AN. Baccalaureate nursing students' perceptions of peer assessment of individual contributions to a group project: a case study. *Nurs educ today.* 2012; 32(3):214–8.
46. Chow KM, Tang WKF, Chan WHC, Sit WHJ, Choi KC, Sally C. Resilience and well-being of university nursing students in Hong Kong: a cross-sectional study. *BMC med edu.* 2018; 18(1):13.
47. Pope RL, Reynolds AL, Mueller JA. Multicultural competence in student affairs: Jossey-Bass; 2019.
48. Hair J, Anderson RE, Tatham R, Black W. Multivariate data analysis. Saddle River, NJ: Prentice Hall; 1995.
49. Moshtaghi S, Moayed-far H. The role of psychological capital components (Hope, Optimism, Resiliency and Self-Efficacy) and Achievement goals Orientation in predicting Academic Procrastination in Students. *J Cogn Strateg Learn.* 2017; 5(8):61–78.
50. Jimenez C, Navia-Osorio PM, Diaz CV. Stress and health in novice and experienced nursing students. *J Adv Nurs.* 2010; 66(2):442–55. <https://doi.org/10.1111/j.1365-2648.2009.05183.x> PMID: 20423427
51. Humpel N, Caputi P. Exploring the relationship between work stress, years of experience and emotional competency using a sample of Australian mental health nurses. *J Psychiatr Ment Health Nurs.* 2001; 8(5):399–403. <https://doi.org/10.1046/j.1365-2850.2001.00409.x> PMID: 11882159
52. Blake H, Chambers D. Supporting nurse health champions: Developing a 'new generation' of health improvement facilitators. *Health Educ.* 2012; 71(2):205–10.
53. Alvi T, Assad F, Ramzan M, Khan FA. Depression, anxiety and their associated factors among medical students. *J Coll Physicians Surg Pak.* 2010; 20(2):122–6. PMID: 20378041
54. Bolton KW. Development And Validation Of The Resilience Protective Factors Inventory: A Confirmatory Factor Analysis. Ph.D. Dissertations, The University of Texas at Arlington.2013.

55. Hwang E, Shin S. Characteristics of nursing students with high levels of academic resilience: A cross-sectional study. *Nurs educ today*. 2018; 71:54–9.
56. Kim J. Nursing students' relationships among resilience, life satisfaction, psychological well-being, and attitude to death. *Korean J Med Educ*. 2019; 31(3):251. <https://doi.org/10.3946/kjme.2019.135> PMID: 31455054
57. Martin AJ, Marsh HW. Academic resilience and its psychological and educational correlates: A construct validity approach. *Psychol Sch*. 2006; 43(3):267–81.