

Fig. 1. (A) Fundus photography of the macular placoid lesion in the right and left eye. (B) Truncal maculopapular cutaneous erythema at presentation. (C) SD-OCT image of the macula in the right eye on presentation. Disruption of the outer segment structures with grainy, linear hyperdensities and lack of external limiting membrane at presentation. (D) The 'pitchfork sign' (green rectangle, insert) in the right eye. (E) Serous retinal detachment and membrane-like, hyper-reflective subretinal lesion (green rectangle, insert) in the left eye. (F) Seven months after presentation, the outer retinal segment appeared normal in the fovea in the right eye.

Eventually, prompt targeted antibiotic therapy resulted in reconstitution of the outer retinal anatomic landmarks on SD-OCT.

References

De Souza EC, Jalkh AE, Trempe CL, Cunha S & Schepens CL (1988): Unusual central

chorioretinitis as the first manifestation of early secondary syphilis. *Am J Ophthalmol* **105**: 271–276.

Eandi CM, Neri P, Adelman RA, Yannuzzi LA & Cunningham ET Jr (2012): International Syphilis Study Group. Acute syphilitic posterior placoid chorioretinitis: report of a case series and comprehensive review of the literature. *Retina*. **32**: 1915–1941.

Gass JD, Braunstein RA & Chenoweth RG (1990): Acute syphilitic placoid


chorioretinitis. *Ophthalmology* **97**: 1288–1297.

Hoang QV, Cunningham ET Jr, Sorenson JA & Freund KB (2013): The 'pitchfork sign' a distinctive optical coherence tomography finding in inflammatory choroidal neovascularization. *Retina* **33**: 1049–1055.

Correspondence:

Christos Christakopoulos
Department of Ophthalmology
Zealand University Hospital
Ringstedgade 37, 4700, Naestved
Denmark
Tel: +45 56513310
Email: christos@hotmail.dk

Topical levofloxacin, nepafenac and prednisolone acetate medication after cataract surgery in the biggest tertiary eye hospital in Finland during 2015–2018

Sirpa Loukovaara,^{1,2}  Ville Lehtinen,³
Risto Nieminen² and Jukka Moilanen⁴

¹Unit of Vitreoretinal Surgery, Department of Ophthalmology, University of Helsinki and Helsinki University Hospital, Helsinki, Finland; ²Unit of Cataract Surgery, Department of Ophthalmology, Helsinki University Hospital, Helsinki, Finland; ³BCB Medical (bcbmedical.com), Turku, Finland; ⁴Unit of Administration, Department of Ophthalmology, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

doi: 10.1111/aos.14091

Dear Editor,

No general consensus exists regarding the best topical treatment practice after cataract surgery (CS) (Aptel et al. 2017; Juthani et al. 2017). Since no established guidelines are available, practice patterns throughout the world differ. Treatment protocol with topical nonsteroidal anti-inflammatory drugs (NSAIDs) after CS began decades ago (Miyake 1977). Today, NSAIDs such as ketorolac, nepafenac, diclofenac and bromfenac can be used to treat the postoperative

anterior chamber inflammation after CS combined to corticosteroids or as monotherapy.

Our institutional topical treatment regimen was revised in 2014 after topical NSAIDs had shown to offer efficacy comparable to steroids in reducing post-operative inflammation, but with a lower risk for adverse events (Kessel et al. 2014). Using a standardized, computerized platform for patients undergoing CS, we found that during 2015–2018, 60% of the eyes undergoing CS were treated with combination of topical levofloxacin and nepafenac, while 20.3% of eyes were treated with topical levofloxacin and prednisolone acetate (PA; Table 1). Triple combination treatment with topical levofloxacin, nepafenac and PA was used in <10% of our study eyes. Noteworthy, topical PA played an important role, especially in eyes with the most advanced cataract, in the eyes with lowest preoperative visual acuity, and in the eyes with intraoperative complication. Based on our study, we could see seasonal/yearly fluctuation in the use of the most commonly used topical drugs and their combinations in our institution. There was a trend of increased use of topical PA during the study period reflecting partially the instinctive experience of the surgeons to diagnose more inflammatory reaction and cells in the anterior chamber of our patients using nepafenac instead of PA (Grzybowski & Kim 2016). Part of the yearly quartile variation was explained by the fact that nepafenac was temporarily out of stock during the last quartile of the year 2018.

We do acknowledge the following limitations. Our study is retrospective with some inaccuracies in data coding such as missing data. Of note, we could neither assess the postoperative functional visual or refraction outcomes in our study eyes nor patient-reported outcome measures. Neither patient-reported tolerability issues of prescribed medication could be studied (Hovanessian & Holland 2019). In the future, National Cataract Registry and Co-ordination will be highly required in Finland including preoperative risk calculation at least to combination ocular surgeries (Goltz et al. 2018).

Herein, incidence of endophthalmitis was 0.0045%. In 95% of the eyes, we used intracameral cefuroxime shown to be 100% effective in preventing endophthalmitis (Jabbarvand et al. 2016). In

Table 1. Summary of surgical characteristics and actual topical treatment in a large adult cataract surgery cohort operated during 2015–2018.

Basic data of operated eyes		
Operated eyes/patients (n)		21 962/15 994
Female/male		63/37%
ICD-code		
H25.1 normal cataract		20 007 (91.1%)
H26.02 presenile cataract		0.3%
H26.1 traumatic cataract		0.07%
H26.2 related to other eye disease		7.2%
Related to other ICD category		1.3%
Preop risk stratification i.e. estimated severity of cataract		
Green category/yellow/red		37.7/56.3/6.0 (%)
Mode of anaesthesia		
Topical		20 864 (95%)
Peribulbar/general		1098 (5%)
Single/bilateral eye surgery		
Anterior chamber Xylocain + PHNL		20 886 (95.1%)/1076 (4.9%)
Viscoelastic agent		1855 (8.4%)
Discovisc/Viscoat/other		85.8/11.9/2.3 (%)
Trypan blue staining		1059 (4.8%)
Characteristics of anterior segment		
Small pupil		3601 (16.4%)
Exfoliation syndrome+		2174 (9.9%)
Need of pupil expansion device		
Malyugin ring/hook		1405 (6.4%)
Phaco technique		5.9%/0.5%
Stop-chop		9595 (43.7%)
Divide-conquer		8731 (35.8%)
Chop		3278 (14.9%)
Other		355 (1.6%)
Location of inserted IOL		
Posterior chamber		21 304 (97%)
Sulcus fixated		69 (0.3%)
Anterior chamber		24 (0.1%)
Missing info		565 (2.6%)
Intracameral antibiotic 0.1 ml		
Cefuroxime 10 mg/ml		20 832 (94.8%)
Vancomycin 10 mg/ml		352 (1.6%)
Moxifloxacin		54 (0.2%)
Not known		724 (3.3%)
Nylon suture (10-0) to the main wound		219 (1%)
Target refraction		
Emmetropic		18 008 (82%)
-2 to -2.5 D		2525 (11.5%)
Other/missing		4.0%/2.5%
Duration of surgery		
<10 min		1207 (5.5%)
11–15 min		6281 (28.6%)
16–20 min		6083 (27.7%)
>20 min		8367 (38.1%)
Topical medication		
Levofloxacin 5 mg/ml + nepafenac 3 mg/ml		13 349 (60.8%)
Levofloxacin 5 mg/ml + PA 10 mg/ml		4468 (20.3%)
Levofloxacin 5 mg/ml + nepafenac 3 mg/ml + PA 10 mg/ml		1715 (7.8%)
Other, including glaucoma medication		2430 (11.1%)
Use of topical nepafenac and PA in various subgroups		
	VA given in 14	Use of PA and/or N
Preop visual acuity (VA)	792 eyes (67.4%)	
<0.1 on AR	1814 (12.3%)	33.1% used PA + 74.2% used N
0.1–0.5 on AR	11 822 (79.9%)	26.0% used PA + 73.5% used N
>0.5 on AR	1156 (7.8%)	22.7% used PA + 77.3% used N
Risk stratification i.e. Severity of cataract		
Green category (normal)		21.4% of eyes used PA
Yellow (moderate)		28.7% of eyes used PA
Red (advanced)		37% of eyes used PA

Table 1. (Continued)

Intraoperative complications in the whole cohort	400 (1.8%)	
Anterior capsule tear	87 (21.8%)	42.5% used PA + 55.2% used N
Iris damage	118 (29.5%)	47.5% used PA + 59.3% used N
Posterior capsule rupture	110 (27.5%)	48.25 used PA + 59.1% used N
Dropped nucleus	16 (4%)	56.3% used PA + 75% used N
Other	69 (17.2%)	
Changes in PA use during 2015–2018	Operated eyes (<i>n</i>)	Used medication
3rd quartile of the year 2015	903	9% used PA + 75.2% used N
3rd quartile of the year 2016	852	14.3% used PA + 92.7% used N
3rd quartile of the year 2017	994	22.4% used PA + 89.0% used N
3rd quartile of the year 2018	1395	36.3% used PA + 83.1% used N
Postoperative endophthalmitis during 2015–2018		
<i>Propionibacterium acnes</i>	1 eye (0.0045%)	

The eyes with cataract were coded with International Classification of Disease (ICD)-codes. Surgical risk stratification of operated eyes was as follows: normal cataract (green; nuclear sclerosis ++), moderate cataract (yellow; nuclear sclerosis +++), and advanced cataract (red; no red reflex, loose zonules, exfoliation ++, VA < 0.1). The surgical technique was the standard minimally invasive phacoemulsification technique (Infinity/Centurion Vision System, Alcon Laboratories, Inc., Fort Worth, TX, USA) with a 2.4–2.75 mm clear corneal incision. In the uncomplicated cataract surgeries, single-piece acrylic monofocal IOL (AMO Tecnis®, Johnson & Johnson Surgical Vision Inc., Santa Ana, CA, USA) or three-piece aspheric acrylic IOL (Tecnis® ZA9003, Abbott Medical Optics Inc, Santa Ana, CA, USA) was folded and inserted using a lens injector to the capsular bag (posterior chamber IOL). Data were collected from the operating room management system consisting of all cataract operations in the BCB Medical cataract database (BCB cataract database, Turku, Finland) between 1 January 2015 and 31 December 2018. The study represents secondary use of registered surgical data and is without patient contact. AR = autorefractometer, IOL = intraocular lens, N = nepafenac, PA = prednisolone acetate, PHNL = phenylephrine 10%, VA = visual acuity.

previous studies, our mainline strategy to use intracameral cefuroxime together with topical levofloxacin has been shown to lower the chance of endophthalmitis with high-to-moderate quality evidence (Kessel et al. 2015; Gower et al. 2017). We use intracameral antibiotics despite their potential risks and their role being controversial in prophylaxis of endophthalmitis post-CS (Çakır et al. 2015; Witkin et al. 2015; Schwartz et al. 2016). Regimen of topical antibiotic use deserves to be revised next, since resistance issues are an ever-increasing problem globally.

Even though our hospital registry data allow a tool for understanding current clinical practice patterns in our high-volume CS unit, more thorough prospective clinical studies are still needed to answer various specific questions related to ophthalmic outcome parameters (such as final visual function, vision-related quality of life, occurrence of postoperative cystic macular oedema, intraocular lens (IOL) dislocation/tilt, results of combinatorial/post-vitrectomy cataracts surgeries), and issues related to each specific topical therapy. In the future, the structural

data should be linked with systemic medication and type of IOL used. We hope that the actual use of very structured data could, however, stimulate and set standards for other ophthalmic departments.

References

- Aptel F, Colin C, Kaderli S, Deloche C, Bron AM, Stewart MW, Chiquet C; OSIRIS group (2017): Management of postoperative inflammation after cataract and complex ocular surgeries: a systematic review and Delphi survey. *Br J Ophthalmol* **101**: 1–10.
- Çakır B, Celik E, Aksoy NÖ, Bursalı Ö, Uçak T, Bozkurt E & Alagoz G (2015): Toxic anterior segment syndrome after uncomplicated cataract surgery possibly associated with intracameral use of cefuroxime. *Clin Ophthalmol* **17**: 493–497.
- Goltz DE, Baumgartner BT, Politzer CS, DiLallo M, Bolognesi MP & Seyler TM (2018): The American college of surgeons national surgical quality improvement program surgical risk calculator has a role in predicting discharge to post-acute care in total joint arthroplasty. *J Arthroplasty* **33**: 25–29.
- Gower EW, Lindsley K, Tulenko SE, Nanji AA, Leyngold I & McDonnell PJ (2017): Perioperative antibiotics for prevention of acute endophthalmitis after cataract surgery. *Cochrane Database Syst Rev* **2**: CD006364.
- Grzybowski A & Kim SJ (2016): Corticosteroids substituted by nonsteroidal anti-inflammatory drugs: Is it justified by evidence-based medicine? *J Cataract Refract Surg* **42**: 510–511.
- Hovanesian J & Holland E (2019): Tolerability and toxicity of topically applied nepafenac 0.3% compared with generic ketorolac 0.5%. *J Cataract Refract Surg* **45**: 174–180.
- Jabbarvand M, Hashemian H, Khodaparast M, Jouhari M, Tabatabaei A & Rezaei S (2016): Endophthalmitis occurring after cataract surgery: outcomes of more than 480 000 cataract surgeries, epidemiologic features, and risk factors. *Ophthalmology* **123**: 295–301.
- Juthani VV, Clearfield E & Chuck RS (2017): Non-steroidal anti-inflammatory drugs versus corticosteroids for controlling inflammation after uncomplicated cataract surgery. *Cochrane Database Syst Rev* **7**: CD010516.
- Kessel L, Tendal B, Jørgensen KJ, Erngaard D, Flesner P, Andresen JL & Hjortdal J (2014): ost-cataract prevention of inflammation and macular edema by steroid and nonsteroidal anti-inflammatory eye drops: a systematic review. *Ophthalmology* **121**: 1915–1924.
- Kessel L, Flesner P, Andresen J, Erngaard D, Tendal B & Hjortdal J (2015): Antibiotic prevention of postcataract endophthalmitis: a systematic review and meta-analysis. *Acta Ophthalmol* **93**: 303–317.
- Miyake K (1977): Prevention of cystoid macular edema after lens extraction by topical indomethacin (I). A preliminary report. *Albrecht Von Graefes Arch Klin Exp Ophthalmol* **203**: 81–88.
- Schwartz SG, Flynn HW Jr, Grzybowski A, Relhan N & Ferris FL (2016): Intracameral antibiotics and cataract surgery: endophthalmitis rates, costs, and stewardship. *Ophthalmology* **123**: 1411–1413.
- Witkin AJ, Shah AR, Engstrom RE et al. (2015): Postoperative hemorrhagic occlusive retinal vasculitis: expanding the clinical spectrum and possible association with vancomycin. *Ophthalmology* **122**: 1438–1451.

Correspondence:

Sirpa Loukovaara, MD, PhD
Unit of Vitreoretinal Surgery
Department of Ophthalmology
University of Helsinki and Helsinki University Hospital
Haartmaninkatu 4 C, 00290 Helsinki
Finland
Tel: +358 50 4271566
Fax: +358 9 47175100
Email: sirpa.loukovaara@hus.fi

This work was supported by TYH2018127 and TYH2019232 grants (SL).