

# Diversity of freshwater ciliates (Protista) from Argentina

Gabriela C. Küppers<sup>1,\*</sup> & María C. Claps<sup>2</sup>

<sup>1</sup>Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”, CONICET-UNLP, Argentina.

<sup>2</sup>Instituto de Limnología “Dr. R. A. Ringuelet”, CONICET-UNLP, Argentina

\*Corresponding author: [gkoppers@fcnym.unlp.edu.ar](mailto:gkoppers@fcnym.unlp.edu.ar)

**Abstract.** Diversity of ciliates from freshwater and soil were scarcely investigated in Argentina, in spite of their ecological role in these ecosystems and the huge environmental heterogeneity that can be found in this country. In the present study, we describe the morphology of nine species from a temporary pond in Buenos Aires province, by means of live observations and protargol impregnations. *Stentor igneus* Ehrenberg, *Pseudochilodonopsis piscatoris* (Blochmann), *Vorticella halophila* Stiller, *Intranstylum invaginatum* Stokes, and *Epistylis rotans* Švec were recorded for the first time in Argentina, and in most cases, are new for the Neotropical realm as well. In addition, an updated checklist on freshwater ciliates from Argentina is provided, based on ciliates listed for the Salado River basin in the Buenos Aires province, new findings from Rancho Hambre peat bog pools in Tierra del Fuego province and a pond from Misiones province.

**Keywords.** Argentina, Ciliophora, diversity, new records.

## INTRODUCTION

According to the literature and based on own investigations, a total of 208 freshwater ciliate species were recorded in Argentina (KÜPPERS & CLAPS, 2012). Unfortunately; diversity of this group of protists is highly underestimated, in spite of the ecological role of ciliates in microbial food webs and the huge environmental heterogeneity that can be found in Argentina. In marine ecosystems; several investigations on diversity, biogeography, phylogeny, and ecology of plankton ciliates, mostly on tintinnids, were conducted by BALECH (Akselman Cardella, 2008, list of publications), BOLTOVSKOY & ALDER (1989, 1992), BOLTOVSKOY *et al.* (1990),

ALDER & BOLTOVSKOY (1991); and more recently, by THOMPSON *et al.* (1999, 2001, 2005), SANTOFERRARA & ALDER (2009a, b; 2012), and SANTOFERRARA *et al.* (2011, 2012, 2013, 2014, 2015). Brackish water ciliates were mostly investigated by SOUTO (1974), PETTIGROSSO *et al.* (1997), PETTIGROSSO (2003), PETTIGROSSO & POPOVICH (2009), BARRÍA DE CAO (1992), and BARRÍA DE CAO *et al.* (1997, 2013). On the contrary, only few researchers focused on ciliates from freshwater and soil (KÜPPERS & CLAPS, 2012 and citations therein).

The aim of the present study is to characterize biometrically and describe the morphology of nine ciliates from a temporary pond located in Buenos Aires province, by means of live observations and protargol impregnations. These

species were recorded over a 3-year survey along with other species that were described elsewhere and listed in KÜPPERS & CLAPS (2012). In addition, this checklist on freshwater ciliates from Argentina is updated with species listed for the Salado River basin in the Buenos Aires province, new findings from Rancho Hambre peat bog pools in Tierra del Fuego province, and a new record from Misiones province.

## MATERIAL AND METHODS

A temporary pond located near the city of Poblet, Buenos Aires province (35°05'S, 57°48'W), was surveyed during 2003-2005, where plankton and periphyton samples were obtained monthly. This pond went through drought phases mainly in summer and was colonized by hydrophytes like *Ludwigia peploides* (Kunth) Raven and *Alternanthera phylloxeroides* (Martius) Grisebach during desiccation. These macrophytes persisted over the filling phase in autumn, when periphytic samples were obtained. The rest of the hydroperiod was characterized by open waters. For detail description and location of this temporary pond see KÜPPERS *et al.* (2006). During the publication process of the checklist by KÜPPERS & CLAPS (2012), other freshwater ciliates were described or published that are listed in the present work as well (see KÜPPERS & CLAPS, 2010; KÜPPERS *et al.* 2011).

The Salado River is a lowland river located in the Pampean plain from Buenos Aires province, where land floodings over weeks or months occur seasonally; thus influencing the river regime and consequently, its electrical conductivity and transport of dissolved and particulate material (CLAPS *et al.*, 2009). Besides, sedimentary aquifers with high sodium-chloride concentrations in

the headwater of the river influence the water chemistry of streams and lakes that also contribute to the water balance of the Salado River (CLAPS *et al.*, 2009). From the headwater sector through the mouth of the river, the plankton of several tributaries, artificial channels, and backwater ponds were sampled since 1997 by SOLARI *et al.* (2002), GABELLONE *et al.* (2005), and CLAPS *et al.* (2009), where several species of freshwater as well as brackish and marine ciliates were found. For detail information on the geographic location of sampling sites and water characteristics see GABELLONE *et al.* (2008) and previously cited papers.

Rancho Hambre is a dome-shaped, ombrotrophic peat bog located along the southernmost ridges of the Andes in Tierra del Fuego province. Its landscape is dominated by a matrix of *Sphagnum magellanicum* Bridel mosses that holds pools of different size and with nutrient poor, colored, acid waters (ROIG & ROIG, 2004). Plankton samples were taken in five of these pools during a limnological survey in 2008-2010. For detail description of sampling sites, morphometric and physico-chemical characteristics of these pools see GONZÁLEZ GARRAZA *et al.* (2012).

A new ciliate was cited in Misiones province, northeast from Argentina, from a pond formed in the floodplain of the Garupá stream (PESO *et al.*, 2015), which is also included in the list.

Most ciliates described below were found in fresh samples from the pond located near Poblet (Buenos Aires province) and persisted in Petri dishes over 1-2 weeks. Ciliates from Rancho Hambre pools were kept in a culture chamber at 4 °C and 12 hs. light period. Observations were made at magnifications of 10× and 40× under the stereo microscope and at 100×, 400×, and 1000× under the bright field microscope. After observing

ciliates *in vivo*, cells were fixed in Bouin's solution and impregnated with protargol according to WILBERT (1975). Drawings of living cells are free-hand sketches, while impregnated ciliates were illustrated with the aid of a camera lucida. Peritrichs were only studied *in vivo*, because impregnations were too strong. Classification and terminology follow LYNN (2008). Identification of species in the genus *Stentor* Oken were performed according to FOISSNER & WÖLFL (1994), *Vorticella* Linnaeus following WARREN (1986), and specific taxonomic papers cited below.

## RESULTS AND DISCUSSION

Additions to the checklist of freshwater ciliates from Argentina is provided in Table 1. With these findings, the number of freshwater ciliates increases up to 229 species. In the Salado River basin a total of 34 identified species were recorded, from which 5 of the species reported here are new records for Argentina. Some of these ciliates are known to inhabit brackish and / or marine environments. The fluctuating water regime of the river and the chemical nature of the basin cause wide variations in electrical conductivity over wet and dry seasons; thus, probably conditioning ciliate assemblages and allowing the occurrence of euryhaline and halophile species.

Until the investigations of KÜPPERS *et al.* (2011), QUIROGA *et al.* (2013) and unpublished data (KÜPPERS *et al.*, unpubl.), only the peritrich *Epistylis* cf. *umbilicata* was known from lakes in Tierra del Fuego (KÜPPERS & CLAPS, 2012). In the present contribution, a total of 28 identified species are listed in this province, with 13 ciliates being also new records for Argentina.

In Misiones province, PESO *et al.* (2015)

found the peritrich *Ophrydium versatile* (O. F. Müller, 1786) Ehrenberg, 1830 for the first time in a pond formed from the Garupá stream floodplain. This is the only ciliate identified up to species level that was cited for this province.

The following nine species from a temporary pond near the city of Poblet in Buenos Aires province were mentioned in KÜPPERS & CLAPS (2012) but were not biometrically characterized at that time. Five species were new for the freshwater ciliate fauna from Argentina and three of these, were also new for the Neotropical realm.

## Phylum Ciliophora Doflein

### Class Heterotrichea Stein

#### Order Heterotrichida Stein

#### Family Stentoridae Carus

#### ***Stentor igneus* Ehrenberg, 1838 (Figure 1; Table 2)**

Body size *in vivo* 392-490  $\mu\text{m}$  long, trumpet-shaped, with a short mucous lorica. Cortical granules pink-colored, arranged in longitudinal stripes between somatic kineties. Endosymbiotic algae absent. Contractile vacuole anteriorly on the left, with posterior collecting canal (Figure 1a). Nuclear apparatus formed by a single, spherical bead and 3 globular micronuclei. Oral ciliature with 142 membranelles and paroral membrane parallel to the adoral zone. Somatic ciliature composed of 32-36 longitudinal kineties and 7-8 kineties in peristomial region (Figure 1b).

**Occurrence and autoecology.** This species was recorded in autumn 2004, on *Alternanthera philoxeroides* and *Ludwigia peploides*, under the following physico-chemical conditions: conductivity 226.7  $\mu\text{S cm}^{-1}$ , dissolved oxygen concentration 6.3  $\text{mg L}^{-1}$ , temperature 8.6°C, pH 5.4.

**Remarks.** *Stentor igneus* is probably a cosmopolitan

**Table 1.** Additions to the checklist on freshwater ciliates from Argentina, recorded in Buenos Aires province (BA1, BA2, Salado River basin, SRB: S1-S12), Misiones province (M), and Tierra del Fuego province (TF). BA1, temporary pond near Poblet; BA2, temporary pond near Dolores; S1, Salado stream; S2, Piñeiro stream; S3, Junín; S4, Saladillo stream; S4a, Mercante Canal; S4b, Vinculación Canal; S4c, del Este Canal; S5, Achupallas; S6, Ruta 30; S7, Saladillo Vallimanca stream; S8, Roque Pérez; S9, Gorchs; S10, Belgrano; S11, Destino; S12, Guerrero; M, pond of the Garupá stream floodplain; TF, pools from Rancho Hambre peat bog.

Species	BA1	BA2	S1	S2	S3	S4	S4a	S4b	S4c	S5	S6	S7	S8	S9	S10	S11	S12	TF	M	References
<i>Apoamphisella hymenophora</i> (Stokes, 1886) Berger, 1999	+																			Küppers & Claps (2013)
<i>Ascobius lentus</i> Henneguy, 1884																+				New record in SRB
<i>Aspidisca cicada</i> (O. F. Müller, 1786) Claparède & Lachmann, 1858					+		+		+				+							New record in SRB
<i>Aspidisca lynceus</i> (O. F. Müller, 1773) Ehrenberg, 1830					+				+										+	New record in SRB, TF
<i>Balanion planctonicum</i> (Foissner, Oleksiv & Müller, 1990) Foissner, Berger & Kohmann, 1994																				Quiroga <i>et al.</i> (2013)
<i>Bryometopus sphagni</i> (Penard, 1922) Kahl, 1932																			+	New record in Argentina

**Table 1.** Additions to the checklist on freshwater ciliates from Argentina.

<i>Bursariidium pseudobursaria</i> (Fauré-Fremlet, 1924) Kahl, 1927					+	New record in Argentina
<i>Carchesium polypinum</i> (Linnaeus, 1758) Ehrenberg, 1930		+	+			New record in SRB
<i>Cinetochilium margaritaceum</i> (Ehrenberg, 1831) Perty, 1849					+	New record in TF
<i>Clapsiella magnifica</i> Küppers, 2014	+					Küppers (2014)
<i>Colpoda steini</i> Maupas, 1883					+	New record in TF
<i>Condylostoma</i> cf. <i>spatiosum</i>				+		New record in Argentina
<i>Deviata polycirrata</i> Küppers & Claps, 2010	+	+				Küppers & Claps (2010)
<i>Diophrys</i> cf. <i>appendiculata</i>					+	New record in Argentina

Table 1. Additions to the checklist on freshwater ciliates from Argentina.

<i>Epispathidium amphoriforme</i> (Greeff, 1888) Foissner, 1984	+	New record in TF
<i>Fabrea salina</i> Henneguy, 1890	+	New record in Argentina
<i>Frontonia</i> cf. <i>acuminata</i>	+	New record in Argentina
<i>Frontonia atra</i> (Ehrenberg, 1833) Bütschli, 1889	+	New record in TF
<i>Halteria grandinella</i> (O. F. Müller, 1773) Dujardin, 1841	+	Quiroga <i>et al.</i> (2013)
<i>Lembadion lucens</i> (Maskell, 1887) Kahl, 1931	+	New record in Argentina
<i>Limnostrombidium viride</i> (Stein, 1867) Krainer, 1995	+	New record in SRB

**Table 1.** Additions to the checklist on freshwater ciliates from Argentina.

<i>Linostomella vorticella</i> (Ehrenberg, 1833) Aescht in Foissner, Berger & Schaumburg, 1999	+	New record in SRB	
<i>Mesodinium acarus</i> Stein, 1867	+	New record in Argentina	
<i>Ophrydium versatile</i> (O. F. Müller, 1786) Ehrenberg, 1830	+	+	New record in TF, M
<i>Paramecium aurelia</i> complex	+	New record in TF	
<i>Paramecium bursaria</i> (Ehrenberg, 1831) Focke, 1836	+	New record in TF	
<i>Parasterkiella thompsoni</i> (Foissner, 1996) Küppers, da Silva Paiva, do Nascimento Borges, Harada, González Garranza & Mataloni, 2011	+	Küppers <i>et al.</i> (2011)	
<i>Pelagostrombidium fallax</i> (Zacharias, 1896) Krainer, 1991	+	Quiroga <i>et al.</i> (2013)	


**Table 1.** Additions to the checklist on freshwater ciliates from Argentina.

<i>Plagiophyla nasuta</i> Stein, 1860	+	New record in TF
<i>Podophrya fallax</i> Dingfelder, 1961	+	New record in TF
<i>Rimostrombidium hyalinum</i> (Mirabdullaev, 1985) Petz & Foissner, 1992	+	Quiroga <i>et al.</i> (2013)
<i>Stentor coeruleus</i> (Pallas, 1766) Ehrenberg, 1831,	+	New record in TF
<i>Stentor cf. araucanus</i>	+	TF
<i>Stichotricha aculeata</i> Wrzesniowski, 1866	+	New record in Argentina
<i>Strobilidium caudatum</i> (Fromentel, 1876) Foissner, 1987	+	New record in SRB, TF
<i>Strombidium sulcatum</i> Claparède & Lachmann, 1859	+	New record in Argentina
<i>Strombidium cf. stylifer</i>	+	New record in Argentina





species; although, it was not yet found in Australia and Antarctica (FOISSNER *et al.*, 1992). In the Neotropical realm, it was previously recorded in México (ALADRO LUBEL *et al.*, 2006, 2009), Perú (FOISSNER *et al.*, 1992), and Brazil (REGALI-SELEGHIM *et al.*, 2011), while represents a new finding for Argentina. Morphometric features coincide with those provided by FOISSNER *et al.* (1992). The populations studied by SONG & WILBERT (1989) from Bonn, Germany, had a lower number of adoral membranelles (76-96 vs. 142, respectively) and somatic kineties (23-29 vs. 32-36, respectively). According to FOISSNER & WÖLFL (1994), most important features to identify species within the genus *Stentor* are the presence/absence of endosymbionts, morphology of the nuclear apparatus, and color of cortical granules. In this context, the species recorded in this study coincides with *S. igneus*.



**Figure 1.** Morphology of *Stentor igneus* in vivo (a) and after protargol impregnation (b). Arrowhead indicates micronuclei. AM, adoral membranelles; CG, cortical granules; CV, contractile vacuule. Scale bar = 50  $\mu\text{m}$ .

***Stentor roeselii* Ehrenberg, 1835 (Figure 2; Table 2)**

Body size *in vivo* 462-1344  $\mu\text{m}$  in length; trumpet-shaped. With posterior mucous lorica. Cortical granules colorless, arranged in longitudinal stripes between somatic kineties. Endosymbiotic algae absent. Contractile vacuole on left of buccal funnel, with posterior collecting canal (Figure 2a). Nuclear apparatus formed by vermiform macronucleus and 4-8 globular micronuclei. Oral ciliature composed of 142-177 membranelles and paroral membrane parallel to adoral zone. Somatic ciliature composed of 42-63 longitudinal and 9-12 peristomial kineties (Figure 2b).

**Occurrence and autoecology.** This species was found in plankton samples obtained in spring 2003 and 2004, under the following ranges of physico-chemical parameters: conductivity 156.3-226.7  $\mu\text{S cm}^{-1}$ , dissolved oxygen concentration 4.5-7.1  $\text{mg L}^{-1}$ , temperature 8.6-19.2 $^{\circ}\text{C}$ , pH 5.4-9.5. Food vacuoles contained green algae. Occasionally, this species was observed to live in great numbers, in mucous masses formed by individual loricae.

**Remarks.** *Stentor roeselii* is probably cosmopolitan; although, it was not yet found in Australia and Antarctica (FOISSNER *et al.*, 1992). This species was previously recorded in México (ALADRO LUBEL *et al.*, 2006, 2009) and Brazil (PAULETO *et al.*, 2009; REGALI-SELEGHIM *et al.*, 2011; VELHO *et al.*, 2013). In Argentina, MODENUTTI (1991) found *S. roeselii* in lotic environments of del Plata sub-catchment and ZALESKI & CLAPS (1999) in San Miguel del Monte lake (Buenos Aires province), epiphytic on *Myriophyllum quitense* Kunth. Morphometric features provided by different authors are somewhat variable and in some cases are different from those observed in the present study; although, important features to identify species within the genus according to FOISSNER & WÖLFL (1994) are coincident. In

the revision of the species, FOISSNER *et al.* (1992) mentioned a higher number of micronuclei than those observed in the present study (7-20 vs. 4-8, respectively), somatic kineties (about 80 vs. about 50 on average, respectively), and peristomial kineties (14-42 vs. 9-12, respectively). The Argentinean population, however, is similar to that described by SONG & WILBERT (1989) from a stream in Bonn, Germany.

### Class Phyllopharingea de Puytorac *et al.*

### Subclass Phyllopharyngia de Puytorac *et al.*

### Order Chlamyodontida Deroux


### Family Chilodonellidae Deroux

### *Pseudochilodonopsis piscatoris* (Blochmann, 1895)

### Foissner, 1979 (Figure 3; Table 3)

Body size 81-105 × 26-42 μm after protargol

impregnation; dorsoventrally flattened, with a



**Figure 2.** Morphology of *Stentor roeselii* in vivo (a) and after protargol impregnation (b). CV, contractile vacuole; Ma, macronucleus; Mi, micronucleus. Scale bar = 50 μm.

**Table 2.** Morphometric data on *Stentor igneus* (first line) and *S. roeselii* (second line). M, median; Max., maximum observation; Min, minimum observation; n, number of observations; SD, standard deviation. Measurements are in μm and correspond to protargol impregnated specimens, unless indicated.


Character	Mean	M	Min.	Max.	SD	n
Body length <i>in vivo</i>	441.0 919.3	441.0 1015.0	392.0 462.0	490.0 1344.0	69.3 323.4	2 18
Peristome diameter <i>in vivo</i>	---	---	---	---	---	---
Peristome diameter	120.5	112.0	98.0	154.0	15.3	18
Lorica length <i>in vivo</i>	---	---	---	---	---	---
Lorica length	774.2	770.0	630.0	1008.0	110.0	18
Macronuclear nodules number	1 1	1 1	1 1	1 1	0 0	3 18
Macronucleus length	32.7 ---	30.8 ---	28.0 ---	39.2 ---	5.8 ---	3 ---
Macronucleus width	27.8 ---	25.9 ---	22.4 ---	35.0 ---	6.5 ---	3 ---
Micronuclei number	3 5.6	3 5	3 4	3 8	0 1.5	1 5
Micronuclei width	1.8 3.5	1.7 3.8	1.7 2.4	2.1 5.0	0.2 0.9	3 7
Membranelles number	142.0 164.5	142.0 165.5	142.0 142	142.0 177	0 13.1	1 6

Table 2. Continuation.

Somatic kineties number	34 48.7	34 47.0	32 42	36 63	2.8 7.4	2 7
Peristomial kineties number	7.5 10.9	7.5 11.0	7 9	8 12	0.7 1.2	2 7

preoral beak in anterior left margin and posterior end pointed, sometimes rounded. Cytoplasm colorless. Two contractile vacuoles; one located in midbody on right cell margin, and other one in posterior third on left body margin (Figure 3a). With conspicuous dorsal cytoplasmic protrusion or spine posteriorly. Macronucleus ellipsoidal; micronucleus not observed. Cyrtos formed by about 14 rods. Ventral ciliature composed of 5-6 left and 5-6 right kineties, separated by a bare postoral field. One fragmented preoral kinety and 2 circumoral kineties. Preoral kinety consists of 4 fragments; fourth fragment extends around cytopharynx opening (Figure 3b). Dorsal brush arched anteriorly (Figure 3c).

**Occurrence and autoecology.** This species was found in winter-spring 2004 and autumn-spring



**Figure 3.** Morphology of *Pseudochilodonopsis piscatoris* in vivo (a) and after protargol impregnation (b, c). a, b. Ventral view; c. Dorsal view. C, cyrtos; CV, contractile vacuole; DB, dorsal brush; LK, left ventral kineties; Ma, macronucleus; PO, fragmented preoral kinety; RK, Right ventral kineties. Scale bar = 30  $\mu\text{m}$ .

2005, in plankton as well as periphyton samples, under the following ranges of physico-chemical parameters: conductivity 133-243  $\mu\text{S cm}^{-1}$ , dissolved oxygen concentration 4.5-8.0  $\text{mg L}^{-1}$ , temperature 4.0-19.2  $^{\circ}\text{C}$ , pH 5-9.5.

**Remarks.** *Pseudochilodonopsis piscatoris* was previously found in Europe (FOISSNER *et al.*, 1991). For Argentina and the Neotropical realm, it represents a new record. Morphometric features coincide with those observed by FOISSNER (1979) and those provided by FOISSNER *et al.* (1991); although the dorsal spine from Argentinean population is distinctly more conspicuous.

### Class Oligohymenophorea de Puytorac *et al.*

#### Subclass Peritrichia Stein

#### Order Sessilida Kahl

#### Family Vorticellidae Ehrenberg

#### *Vorticella convallaria* Linnaeus, 1758

#### (Figure 4a; Table 4)

Zooid *in vivo* 63-98  $\times$  49-70  $\mu\text{m}$ , inverted bell-shaped. Peristomial disc not elevated above peristome; maximum body width less than peristomial lip diameter. Pellicle finely striated and with normal ribbing between striations. Contractile vacuole below peristome. Contractile stalk with evident thecoplasmic granules on myoneme. Macronucleus J-shaped.

**Occurrence and autoecology.** This species was found in spring 2003 and autumn-winter 2004 on *Alternanthera philoxeroides* and *Ludwigia peploides*, under the following ranges of physico-

**Table 3.** Morphometric data on *Pseudochilodonopsis piscatoris*. Abbreviations are indicated in Table 2. Measurements correspond to protargol impregnated specimens.

Character	Mean	M	Min.	Max.	SD	n
Body length	94.5	94.0	81.0	105.0	8.2	8
Body width	37.6	39.5	26.0	42.0	5.6	8
Macronucleus length	25.4	27.0	20.0	28.0	2.8	7
Macronucleus width	13.0	13.0	11.0	15.0	1.5	7
Dorsal spine length	14.1	14.0	10.5	17.0	2.6	5
Dorsal spine width	2.8	2.8	2.8	2.8	0	1
Circumoral kineties number	2	2	2	2	0	7
Kineties in left ventral field number	5.2	5.0	5	6	0.4	6
Kineties in right ventral Field number	5.2	5.0	5	6	0.4	5

chemical parameters: conductivity 145-227  $\mu\text{S cm}^{-1}$ , dissolved oxygen concentration 5.5-8.8  $\text{mg L}^{-1}$ , temperature 2.4-24.1  $^{\circ}\text{C}$ , pH 5.0-8.4.

**Remarks.** *Vorticella convallaria* belongs to a cosmopolitan species complex that occur in Europe, Asia, Antarctica, and America (FOISSNER *et al.*, 1992). In South America, it was mentioned by GUILLÉN *et al.* (2003) in swamps from Villa in Perú, in freshwater environments from México (ALADRO LUBEL *et al.*, 2009) and Brazil (REGALI-SELEGHIM *et al.*, 2011; VELHO *et al.*, 2013). In Argentina, this species was previously recorded in Buenos Aires province by CELA (1972) in ponds from Berisso, by CLAPS (1984) on *Scirpus californicus* (Meyer) Steud. from Atalaya (Río de la Plata estuary), by CLAPS & MODENUTTI (1984) on *S. californicus* from San Miguel del Monte Lake, on *Stuckenia striata* (Ruiz & Pav.) Holub from Chascomús Lake, and on *Pistia stratiotes* L. from Punta Lara (Río de la Plata estuary); finally, MODENUTTI (1987) found *V. convallaria* in plankton samples of Rodríguez

stream from La Plata city. Morphometric features coincide with those mentioned by WARREN (1986) and FOISSNER *et al.* (1992).


#### ***Vorticella pulchella* Sommer, 1951 (Figure 4b; Table 4)**

Zooid size *in vivo* 30.0-43.3  $\mu\text{m} \times$  26.6-33.3  $\mu\text{m}$ . Peristomial disc prominently elevated above peristome and maximum body width approximately equal to peristomial lip diameter; body constricted below peristomial lip. Pellicle conspicuously striated and with convex ribbing between striation. Contractile vacuole in the center of body. Contractile stalk with inconspicuous thecoplastic granules on myoneme, but evident upon contraction. Macronucleus C-shaped, located horizontally below peristome.

**Occurrence and autoecology.** *Vorticella pulchella* was found in winter 2004, on *L. peplodes*, under the following ranges of physico-chemical parameters:

conductivity 163.3-227.0  $\mu\text{S cm}^{-1}$ , dissolved oxygen concentration 5.5-8.8  $\text{mg L}^{-1}$ , temperature 2.4-7.6  $^{\circ}\text{C}$ , pH 5-6.

**Remarks.** *Vorticella pulchella* is distributed in Switzerland (SOMMER, 1951) and Hungary (STILLER, 1971). In Argentina, it was previously recorded by ZALESKI & CLAPS (2001) on *Myriophyllum quitense* and suspended detritus from San Miguel del Monte Lake, Buenos Aires province. STILLER (1971) found *V. pulchella* as epibiont on copepod crustaceans. Morphometric features are coincident with those mentioned by WARREN (1986) and those observed by ZALESKI & CLAPS (2001).



**Figure 4.** Morphology of *Vorticella convallaria* (a) and *V. pulchella* (b) in vivo. Arrowhead in (b) indicate conspicuously striated pellicle and convex ribbing between striations. CV, contractile vacuole; Ma, macronucleus. Scale bars = 20  $\mu\text{m}$ .

### *Vorticella striata* Dujardin, 1841

#### (Figure 5a,b; Table 4)

Zooid size *in vivo* 56-84  $\mu\text{m} \times 35$ -49  $\mu\text{m}$ . Peristomial disc not elevated above peristome and maximum body width less than peristomial lip diameter.


Pellicle conspicuously striated and with convex ribbing between striations. Contractile vacuole just below peristome. Contractile stalk with evident thecoplastic granules on myoneme. Macronucleus C-shaped, located horizontally below peristome.

**Occurrence.** This species was recorded in summer 2005, in rewetted soil samples obtained from the pond bed during a drought period.

**Remarks.** *Vorticella striata* belongs to the *Vorticella aquadulcis* complex, which species were recorded in Eurasia and America (FOISSNER *et al.*, 1992). In 2008, KOVALCHUK redescribed *V. striata* based on a population from Ukraine. In Argentina, this species was previously found by CLAPS & MODENUTTI (1984) on *Azolla filiculoides* Lam. in Las Víboras stream from Magdalena and on *A. filiculoides*, *Ricciocarpus natans* L., *Lemna* sp., and *Wolfiella* sp. in Chascomús Lake, Buenos Aires province. The occurrence of this peritrich on different type of aquatic macrophytes indicates its low specificity for the substrate. Moreover, this species was also observed as epibiont on subitaneous eggs of the social rotifer *Sinantherina semibullata* (Thorpe) in temporary ponds from Punta Lara and Magdalena, Buenos Aires province (KÜPPERS, unpub.). In agreement with our observations, WARREN (1986) mentioned that this species could occasionally be found as epibiont. FERNÁNDEZ-LEBORANS & TATO-PORTO (2000) mentioned *V. striata* on cladocerans and copepods and it was also found on a crayfish crustacean from México (MAYÉN ESTRADA & ALADRO LUBEL, 2002). In the present study, the finding of this species in rewetted soil samples indicate that it developed from resting cysts, which were also observed (Figure 5b). Morphometric features are in agreement with those mentioned by WARREN (1986) and those observed by CLAPS & MODENUTTI (1984).

***Vorticella halophila* Stiller, 1941**

(Figure 5c; Table 4)



**Figure 5.** Morphology of *Vorticella striata* (a, b) and *V. halophila* (c) in vivo. a, c. Zooids. b. Cyst with pulsating contractile vacuole. Arrowheads indicate pellicle conspicuously striated (a) and theoplasmic granules (c). CV, contractile vacuole; Ma, macronucleus. Scale bars = 20 µm.

Zoid *in vivo* 63-102.5 µm × 35-56 µm. Peristomial disc not elevated above peristome and maximum body width less than peristomial lip diameter. Body constricted below peristome. Pellicle conspicuously striated and with convex ribbing between striations. Two contractile vacuoles, located below peristome. Contractile stalk with scarce colorless theoplasmic granules on myoneme. Macronucleus J-shaped.

**Occurrence and autoecology.** This species was found in summer 2005, in rewetted soil samples obtained from the dried pond bed. It was also observed in spring 2005 in plankton samples, under the following physico-chemical conditions: conductivity 206.7 µS cm<sup>-1</sup>, dissolved oxygen

concentration 4.5 mg L<sup>-1</sup>, temperature 11.6 °C, pH 8.7.

**Remarks.** *Vorticella halophila* was recorded in Hungary (STILLER, 1971). In Argentina and the rest of South America, it represents a new finding. The development of this species from rewetted soil samples from the dried bed of the pond, indicate that it is able to form resting cysts. Moreover, this is the first time *V. halophila* is recorded in soil samples. Unfortunately, resting cysts were not observed. Morphometric characteristics coincide; in general, with those mentioned by WARREN (1986); although, specimens from the present study had greater size (40-50 µm vs. 63-102 µm, respectively).

***Intranstylum invaginatum* Stokes, 1886**

(Figure 6; Table 4)

Zooid size *in vivo* 70-98 µm × 21-28 µm; inverted elongated bell-shaped. With peristomial lip. Peristomial disc slightly convex, with a central button-like protuberance, and slightly elevated above peristome (Figure 6a). Maximum body width less than peristomial lip diameter. Upon contraction, with an anterior snout-like protuberance and slightly folded posteriorly (Figure 6b). Pellicle smooth, without evident striation. Contractile vacuole on the peristome. Macronucleus C-shaped, horizontally located in anterior third of body. Stalk very short, with inconspicuous myoneme and posterior adhesive disc.


**Occurrence and autoecology.** This species was found in winter 2004, epibiont on ostracods in the genus *Cypris* O. F. Müller (Fig. 6c), under the following water conditions: conductivity 227 µS cm<sup>-1</sup>, dissolved oxygen concentration 5.5 mg L<sup>-1</sup>, temperature 2.4°C, pH 5.

**Remarks.** *Intranstylum invaginatum* was recorded in North America and Germany on the ostracods

*Cypris* and *Candona* Baird and also on cladocerans in Germany, England, Finland, and Italy (KAHL, 1935; PÄTSCH, 1974; FERNÁNDEZ-LEBORANS & TATO-PORTO, 2000; CHATTERJEE *et al.*, 2013). In Argentina and the rest of South America, it represents a new finding. Morphology is coincident with specimens recorded on *Cypris* by other authors; although, stalk is much shorter in the Argentinean population. On the contrary, epibionts on *Candona* have greater stalk length and those specimens found on cladocerans lack the button-like protuberance on center of peristomial disc (KAHL, 1935). To our knowledge, this species has rarely been described morphologically after its original description by STOKES (1886) and the observations of KAHL (1935). Unfortunately, protargol impregnations were very strong in order to observe its infraciliature.


### Family Epistylididae Kahl

#### *Epistylis rotans* Švec, 1897 (Figure 7; Table 4)



**Figure 6.** Morphology of *Intranstylum invaginatum* in vivo. a. Zooid with button-like protuberance in center of peristomial disc (arrowhead). b. Zooid upon contraction. c. Basibiont ostracod (*Cypris* sp.), showing zooids in posterior region of the valves. CV, contractile vacuole; Ma, macronucleus. Scale bar = 20 µm.

Colony size *in vivo* about 4 mm high; arbustive, with principal stalk noncontractile and dichotomously branched. Zooid *in vivo* 196-252 µm × 77-98 µm; inverted bell-shaped, with anterior end of body curved or inclined. With peristomial lip; peristomial disc slightly convex, not elevated above peristome. Maximum body width less than peristomial lip diameter. Pellicle conspicuously striated, with convex ribbing between striations, and with small warts that become evident in peristomial zone. Upon contraction, zooid globular-shaped and peristomial lip covers the peristome completely. Contractile vacuole on the peristome. Macronucleus



**Figure 7.** Morphology of *Epistylis rotans* in vivo. a. Zooid. b. Zooid upon contraction. c. General view of the colony. CV, contractile vacuole. Scale bar = 20 µm



C-shaped, horizontally located with respect to cell major axis. Stalk smooth and iridescent at low magnification (less than 40×), colonized by epibiont choanoflagellates.

**Occurrence and autoecology.** This species was found in autumn 2004; on *A. philoxeroides*, *L. peploides*, and suspended organic matter, under the following water characteristics: conductivity

**Table 4.** Morphometric data on *Vorticella convallaria* (Vc), *V. pulchella* (Vp), *V. striata* (Vs), *V. halophila* (Vh), *Intranstylum invaginatum* (Ii), and *Epistylis rotans* (Er) *in vivo*. Spp., species; other abbreviations as in Table 2.

Character	Spp.	Mean	M	Min.	Max.	n
Zooid length	Vc	80.5	80.5	63.0	98.0	20
	Vp	38.3	40.0	30.0	43.3	7
	Vs	63.0	59.5	56.0	84.0	8
	Vh	82.3	84.0	63.0	102.5	15
	Ii	89.3	91.0	70.0	98.0	11
	Er	231.5	238.0	196.0	252.0	15
Zooid width	Vc	56.7	56.0	49.0	70.0	20
	Vp	28.2	28.0	26.6	33.3	7
	Vs	39.4	38.5	35.0	49.0	8
	Vh	47.2	49.0	35.0	56.0	12
	Ii	26.1	28.0	21.0	28.0	11
	Er	85.9	84.0	77.0	98.0	15
Peristome diameter	Vc	73,8	70	63	84	20
	Vp	23.6	23.3	20.0	28.0	5
	Vs	32.4	31.5	28.0	42.0	8
	Vh	57.2	56.0	42.0	70.0	12
	Ii	30.5	28.0	28.0	35.0	11
	Er	117.1	112.0	105.0	126.0	15
Stalk length	Vc	276,8	266	210	378	20
	Vp	162.5	156.5	86.6	223.0	5
	Vs	244.0	266.0	140.0	294.0	7
	Vh	223.6	210.0	168.0	350.0	5
	Ii	---	---	---	---	---
	Er*	2000.0	2000.0	2000.0	2000.0	1
Stalk width	Vc	---	---	---	---	---
	Vp	2.7	3.3	1.7	3.3	5
	Vs	---	---	---	---	---
	Vh	---	---	---	---	---
	Ii	---	---	---	---	---
	Er*	21.0	---	---	---	---
Adhesive disc diameter	Ii	24.5	24.5	21.0	28.0	2

226.7  $\mu\text{S cm}^{-1}$ , dissolved oxygen concentration 6.3  $\text{mg L}^{-1}$ , temperature 8.6°C, pH 5.4.

**Remarks.** *Epistylis rotans* was recorded by other authors in Hungary, Switzerland, Germany, Czech Republic, and Russia (KAHL, 1935; NENNINGER, 1944-48; STILLER, 1971). This species was also found in México (ALADRO LUBEL *et al.*, 2006), while it

represents a new record in Argentina. KAHL (1935) and STILLER (1971) found this peritrich in plankton samples, while NENNINGER (1944-48) recorded it on bryozoans and *Lemna* sp. Morphometric characteristics coincide with those observed by other authors from different geographic locations (KAHL, 1935; NENNINGER, 1944-48), mainly in the

typical shape of zooids. The iridescent coloration of the main stalk of colony was not mentioned by other authors.

### ACKNOWLEDGEMENTS

We are grateful to Dr. Analía Díaz, from Instituto de Limnología Dr. R. A. Ringuelet, for identification of ostracods on which *I. invaginatum* was found and to a Brazilian friend MSc. Paulo *Callithrix* Pinheiro, from Instituto de Biologia UNESP, for the writing of the abstract in Portuguese. Consejo Nacional de Investigaciones Científicas y Técnicas is acknowledged for financial support (PIP 035).

### REFERENCES

- ALADRO-LUBEL, M.A.; MAYÉN-ESTRADA, R. & REYES-SANTOS, M. 2006. **Listados faunísticos de México. XI. Registro actualizado de ciliados (Agosto, 2004)**. México, Instituto de Biología, Universidad Nacional Autónoma de México, 97p.
- ALADRO LUBEL, M.A.; REYES SANTOS, M. & OLVERA BAUTISTA, F. 2009. Diversidad de los protozoos ciliados, pp. 61-68. *In*: Lot, A. & Cano-Santana, Z. (eds.). **Biodiversidad del Pedregal de San Ángel**, Reserva Ecológica del Pedregal de San Ángel y Coordinación de la Investigación Científica, Universidad Nacional Autónoma de México, México, 253 p.
- ALDER, V. & BOLTOVSKOY, D. 1991. The ecology and biogeography of tintinnid ciliates in the Atlantic Sector of the Southern Ocean. **Marine Chemistry** **35**: 337-346
- AKSELMAN CARDELLA, R. 2008. In Memoriam Enrique Balech. **Revista del Instituto de Desarrollo Pesquero** **19**: 81-88.
- BARRÍA DE CAO, M.S. 1992. Abundance and species composition of tintinnina (Ciliophora) in Bahía Blanca Estuary, Argentina. **Estuarine, Coastal and Shelf Science** **34**: 295-303.
- BARRÍA DE CAO, M.S.; LOPEZ ABBATE, C.; PETTIGROSSO, R. & HOFFMEYER, M.S. 2013. The planktonic ciliate community and its relationship with the environmental conditions and water quality in two bays of the Beagle Channel, Argentina. **Journal of the Marine Biological Association of the United Kingdom** **93**: 1753-1760.
- BARRÍA DE CAO, M.S.; PETTIGROSSO, R.E. & POPOVICH, C. 1997. Planktonic ciliates during a diatom bloom in Bahía Blanca Estuary, Argentina. II. Tintinnids. **Oebalia** **23**: 21-31.
- BOLTOVSKOY, D. & ALDER, V. 1989. Summer Weddell Sea microplankton: assemblage structure, distribution and abundance, with special emphasis on the Tintinnina. **Polar Biology** **9**: 447-456.
- BOLTOVSKOY, D. & ALDER, V. 1992. Microzooplankton and tintinnid species-specific assemblage structures: patterns of distribution and year-to-year variations in the Weddell Sea (Antarctica). **Journal of Plankton Research** **14**: 1405-1423.
- BOLTOVSKOY, D.; DINOFRIO, E. & ALDER, V. 1990. Environmental influence on intraspecific variability in Antarctic tintinnids: *Cymatocylis affinis convallaria* in Weddell Sea waters. **Journal of Plankton Research** **12**: 403-413.
- CELA, A.M. 1972. Algunos ciliados vinculados a la vegetación flotante. **Physis** **31**: 559-577.
- CHATTERJEE, T.; KOTOV, A.A. & FERNÁNDEZ-LEBORANS, G. A checklist of epibiotic ciliates (Peritrichia and Suctorina) on the cladoceran crustaceans. **Biologia** **68**: 439-447.

- CLAPS, M.C. 1984. Zooperifiton en *Scirpus* (*Schoenoplectus*) *californicus* (Meyer) Steud. (Río de la Plata – Punta Atalaya). **Neotrópica** **30**: 79-88.
- CLAPS, M.C. & MODENUTTI, B.E. 1984. Contribución al conocimiento de los ciliados (Ciliophora Peritricha) dulceacuícolas de Argentina. II. **Limnobiós** **2**: 581-585.
- CLAPS, M.C.; GABELLONE, N.A. & NESCHUK, N.C. 2009. Influence of regional factors on zooplankton structure in a saline lowland river: the Salado River (Buenos Aires, Argentina). **River Research and Applications** **25**: 453-471.
- FERNÁNDEZ-LEBORANS, G. & TATO-PORTO, M.L. 2002. A review of the species of protozoan epibionts on crustaceans. I. Peritrich ciliates. **Crustaceana** **73**: 643-683.
- FOISSNER, W. 1979. Ökologische und systematische Studien über das Neuston alpiner Kleingewässer, mit besonderer Berücksichtigung der Ciliaten. **Internationale Revue der Gesamten Hydrobiologie** **64**: 99-140.
- FOISSNER, W.; BERGER, H. & KOHMANN, F. 1992. **Taxonomische und ökologische Revision der Ciliaten des Saprobien-systems - Band II: Peritrichia, Heterotrichida, Odontostomatida**. Informationsberichte des Bayer, Landesamtes für Wasserwirtschaft 5/92, 502p.
- FOISSNER, W.; BERGER, H. & SCHAUMBURG, J. 1999. **Identification and ecology of limnetic plankton ciliates**. Informationsberichte des Bayer, Landesamtes Wasserwirtschaft, 3/99, 793p.
- FOISSNER, W. & WÖLFL, S. 1994. Revision of the genus *Stentor* Oken (Protozoa, Ciliophora) and description of *S. araucanus* nov. spec. from South American lakes. **Journal of Plankton Research** **16**: 255-289.
- GABELLONE, N.A.; CLAPS, M.C.; SOLARI, L.C. & NESCHUK, N.C. 2005. Nutrients, conductivity and plankton in a landscape approach to a pampean lowland river (Salado River, Argentina). **Biogeochemistry** **75**: 455-477.
- GONZÁLEZ GARRAZA, G.; MATALONI, G.; ITURRASPE, R.; LOMBARDO, R.; CAMARGO, S. & QUIROGA, M.V. 2012. The limnological character of bog pools in relation to meteorological and hydrological features. **Mires and Peat** **10**: 1-14.
- GUILLÉN G.; MORALES, E. & SEVERINO, R. 2003. Adiciones a la fauna de protozoarios de los pantanos de Villa, Lima, Perú. **Revista Peruana de Biología** **10**: 175-182.
- KAHL, A. 1935. **Urtiere oder Protozoa I: Wimpertiere oder Ciliata (Infusoria)** 4. Peritricha und Chonotricha; Nachtrag I, pp. 651-886. In: Dahl, F. (ed.). Die Tierwelt Deutschlands 30, Gustav Fischer, Jena, 886p.
- KÜPPERS, G.C.; LOPRETTO, E.C. & CLAPS, M.C. 2006. Morphological aspects and seasonal changes of some planktonic ciliates (Protozoa, Ciliophora) from a temporary pond, Buenos Aires province, Argentina. **Pan-American Journal of Aquatic Sciences** **1**: 74-90.
- KÜPPERS, G.C. & CLAPS, M.C. 2010. Morphology and notes on morphogenesis during cell division of *Deviata polycirrata* n. sp. and of *Deviata bacilliformis* (Gelei, 1954) Eigner, 1995 (Ciliophora: Kahliellidae) from Argentina. **Journal of Eukaryotic Microbiology** **57**: 273-284.
- KÜPPERS, G.C.; PAIVA, T.S.; BORGES, B.; HARADA, M.L.; GONZÁLEZ GARRAZA, G. & MATALONI, G. 2011. An Antarctic ciliate, *Parasterkiella thompsoni* (Foissner) nov. gen., nov. comb., recorded

- in Argentinean peat-bogs: morphology, morphogenesis, and molecular phylogeny. **European Journal of Protistology** **47**: 103-123.
- KÜPPERS, G.C. & CLAPS, M.C. 2012. Freshwater ciliates (Protozoa, Ciliophora) from Argentina: an annotated and updated compilation, pp. 61–100. *In*: Thangadurai, D.; Busso, C.A.; Abarca Arenas, L.G. & Jayabalan, S.(Eds.). **Frontiers in Biodiversity Studies**. New Delhi, India, IK International Publishing, 440p.
- KÜPPERS, G.C. & CLAPS, M.C. 2013. Hypotrichous ciliates(Protozoa:Ciliophora)froma temporary pond in Argentina, with redescription of *Apoamphisiella hymenophora* (Stokes, 1886) Berger, 1999. **Zootaxa** **3626**: 55-76.
- KÜPPERS, G.C. 2014. Morphology of a new hypotrichous ciliate, *Clapsiella magnifica* gen. n, sp. n., with a curious dorsal ciliary pattern. **European Journal of Protistology** **50**: 373-381.
- LYNN, D.H. 2008. The Ciliated Protozoa. Characterization, Classification, and Guide to the Literature, 3rd ed. Dordrecht, The Netherlands, Springer, 605p.
- MAYÉN ESTRADA, R. & ALADRO LUBEL, M.A. 2002. Distribution and prevalence of 15 species of epibiont peritrich ciliates on the crayfish *Cambarellus patzcuarensis* Villalobos, 1943 in lake Pátzcuaro, Michoacán, Mexico. **Crustaceana** **74**: 1213-1224.
- MODENUTTI, B.E. 1987. Variaciones en la diversidad del zooplancton del arroyo Rodríguez (prov. de Buenos Aires). **Revista de la Asociación de Ciencias Naturales del Litoral** **18**: 61-70.
- MODENUTTI, B.E. 1991. Zooplancton de ambientes lóticos de la subcuenca delta del río Paraná, Buenos Aires, Argentina. **Iheringia, Série Zoologia** **71**: 67-80.
- NENNINGER, U. 1944-1948. Die Peritrichen der Umgebung von Erlangen mit besonderer Berücksichtigung ihrer Wirtsspezifität. **Zoologische Jahrbücher** **77**: 169-266.
- PÄTSCH, B. 1974. Die Aufwuchsciliaten des Naturlehrparks Haus Wildenrath. Monographische Bearbeitung der Morphologie und Ökologie. **Arbeiten aus dem Institut für landwirtschaftliche Zoologie und Bienenkunde** **1**: 1-82.
- PAULETO, G.M.; VELHO, L.F.M.; BUOSI, P.R.B.; BRÃO, A.F.S.; LANSAC-TÔHA, F.A. & BONECKER, C.C. 2009. Spatial and temporal patterns of ciliate species composition (Protozoa: Ciliophora) in the plankton of the Upper Paraná River floodplain. **Brazilian Journal of Biology** **69**: 517-527.
- PESO, J.G.; MEICHTRY DE ZABURLÍN, N.R.; VOGLER, R.E. & MOLINA, M.J. 2015. First record of *Ophrydium versatile* (Müller, 1786) Ehrenberg, 1838 (Protista: Peritrichida: Ophrydiidae) from the High Paraná River influence area (Misiones province, Argentina). **Check List** **11**: 1967.
- PETTIGROSSO, R.E. 2003. Planktonic ciliates Choreotrichida and Strombidiida from the inner zone of Bahía Blanca estuary, Argentina. **Iheringia**, **93**: 117-126.
- PETTIGROSSO, R.E.; BARRÍA DE CAO, M.S. & POPOVICH, C.A. 1997. Planktonic ciliates during a diatom bloom in Bahía Blanca Estuary, Argentina. I Aloricate. **Oebalia** **23**: 21-31.
- PETTIGROSSO, R.E. & POPOVICH, C.A. 2009. Phytoplankton-aloricate ciliate community in the Bahía Blanca Estuary (Argentina): seasonal patterns and trophic groups. **Brazilian Journal of Oceanography** **57**: 215-227.
- QUIROGA, M.V.; UNREIN, F.; GONZÁLEZ GARRAZA, G.;

- KÜPPERS, G.; LOMBARDO, R.; MARINONE, M.C.; MENÚ MARQUE, S.; VINOCUR, A. & MATALONI, G. 2013. The plankton communities from peat bog pools: structure, temporal variation and environmental factors. **Journal of Plankton Research** **35**: 1234-1253.
- REGALI-SELEGHIM, M.H.; GODINHO, M.J.L. & MATSUMURA-TUNDISI, T. Checklist dos "protozoários" de água doce do Estado de São Paulo, Brasil. **Biota Neotropical** **11(Supl.1)**: 389-426.
- ROIG, C. & ROIG, F.A. 2004. Consideraciones generales, pp. 5-21. *In*: Blanco, D.E. & de la Balze, V.M. (eds.). **Los Turbales de la Patagonia, Bases para su Inventario y la Conservación de su Biodiversidad**, Fundación Humedales, Wetlands International, Buenos Aires, 149 p.
- SANTOFERRARA, L. & ALDER, V.A. 2009a. Abundance trends and ecology of planktonic ciliates of the south-western Atlantic (35-63° S): A comparison between neritic and oceanic environments. **Journal of Plankton Research** **31**: 837-851.
- SANTOFERRARA, L. & ALDER, V.A. 2009b. Morphological variability, spatial distribution and abundance of *Helicostomella* species (Ciliophora, Tintinnina) in relation to environmental factors (Argentine shelf; 40-55°). **Scientia Marina** **73**: 701-716.
- SANTOFERRARA, L. & ALDER, V.A. 2012. Abundance and diversity of tintinnids (planktonic ciliates) under contrasting levels of productivity in the Argentine Shelf and Drake Passage. **Journal of Sea Research** **71**: 25-30.
- SANTOFERRARA, L.F.; GOMEZ, M.I. & ALDER, V.A. 2011. Bathymetric, latitudinal and vertical distribution of protozooplankton in a cold-temperate shelf (southern Patagonian waters) during winter. **Journal of Plankton Research** **33**: 457-468.
- SANTOFERRARA, L.F.; MCMANUS, G.B. & ALDER, V.A. 2012. Phylogeny of the Order Tintinnida (Ciliophora, Spirotrichea) Inferred from Small- and Large-Subunit rRNA Genes. **Journal of Eukaryotic Microbiology** **59**: 423-426.
- SANTOFERRARA, L.F.; MCMANUS, G. & ALDER, V.A. 2013. Utility of genetic markers and morphology for species discrimination within the order Tintinnida (Ciliophora, Spirotrichea). **Protist** **164**: 23-36.
- SANTOFERRARA, L.; GRATTEPANACHE, J.-D.; KATZ, L.A. & MCMANUS, G.B. 2014. Pyrosequencing for assessing diversity of eukaryotic microbes: analysis of data on marine planktonic ciliates and comparison with traditional methods. **Environmental Microbiology** **16**: 2752-2763.
- SANTOFERRARA, L.F.; TIAN, M.; ALDER, V.A. & MCMANUS, G.B. 2015. Discrimination of closely related species in tintinnid ciliates: new insights on crypticity and polymorphism in the genus *Helicostomella*. **Protist** **166**: 78-92.
- SOLARI, L.C.; CLAPS, M.C. & GABELLONE, N.A. River backwater pond interactions in the lower basin of Salado River (Buenos Aires, Argentina). **Archiv für Hydrobiologie** **141**: 99-119.
- SOMMER, G. 1951. Die Peritrichen Ciliaten des grossen Plöner Sees. **Archiv für Hydrobiologie** **44**: 349-440.
- SONG, W. & WILBERT, N. 1989. Taxonomische Untersuchungen an Aufwuchsciliaten (Protozoa, Ciliophora) im Poppelsdorfer Weiher, Bonn. **Lauterbornia** **3**: 1-222.
- SOUTO, S. 1974. Tintínnidos del río de la Plata y su

- zona de influencia (Protozoa, Ciliata). **Physis B 33**: 201-205.
- STILLER, J. 1971. Szájkoszorús Csillósdk-Peritricha. **Fauna Hungariae 105**: 1-245.
- STOKES, A.C. 1886. Notices of new American fresh-water infusoria. **Proceedings of the American Philosophical Society 23**: 562-568.
- THOMPSON, G.A. & ALDER, V.A. 2005. Patterns in tintinnid species composition and abundance in relation to hydrological conditions of the southwestern Atlantic during austral spring. **Aquatic Microbial Ecology 40**: 85-101.
- THOMPSON, G.; ALDER, V. & BOLTOVSKOY, D. 2001. Tintinnids (Ciliophora) and other net microzooplankton (>30 µm) in southwestern Atlantic Shelf Break waters. **Marine Ecology Pubblicazioni della Stazione Zoologica di Napoli I 22**: 343-355.
- THOMPSON, G.; ALDER, V.; BOLTOVSKOY, D. & BRANDINI, F. 1999. Abundance and Biogeography of Tintinnids (Ciliophora) and associated microzooplankton in the Southwestern Atlantic Ocean. **Journal of Plankton Research 21**: 1265-1298.
- VELHO, L.F.M.; LANSAC-TÔHA, F.M.; BUOSI, P.R.B.; RAMOS DE MEIRA, B.; CABRAL, A.F. & LANSAC-TÔHA, F.A. 2013. Structure of planktonic ciliates community (Protist, Ciliophora) from an urban lake of southern Brazil. **Acta Scientiarum 35**: 531-539.
- WARREN, A. 1986. A revision of the genus *Vorticella* (Ciliophora: Peritrichida). **Bulletin of the British Museum of Natural History (Zool.) 50**: 1-57.
- WILBERT, N. 1975. Eine verbesserte Technik der Protargolimprägation für Ciliaten. **Mikrokosmos 64**: 171-179.
- ZALESKI, M. & CLAPS, M.C. 1999. First records of epiphytic limnetic ciliates from Argentina. **Natura Neotropicalis 30**: 77-84.
- ZALESKI, M. & CLAPS, M.C. 2001. First record of some peritrichs ciliates for San Miguel del Monte pond (Buenos Aires, Argentina). **Gayana 65**: 39-49.