

## The giant African snail, *Achatina fulica* (Gastropoda: Achatinidae): Using bioclimatic models to identify South American areas susceptible to invasion

Roberto E. Vogler<sup>1,2</sup>, Ariel A. Beltramino<sup>1,3</sup>, Mariano M. Sede<sup>2,4</sup>, Diego E. Gutiérrez Gregoric<sup>1,2</sup>, Verónica Núñez<sup>1,2</sup>, and Alejandra Rumi<sup>1,2</sup>

<sup>1</sup>División Zoología Invertebrados, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Paseo del Bosque s/n (B1900FWA), La Plata, Argentina

<sup>2</sup>Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina

<sup>3</sup>Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT), Argentina

<sup>4</sup>Instituto de Investigaciones Biomédicas en Retrovirus y SIDA, Facultad de Medicina, Universidad de Buenos Aires, Paraguay 2155 (C1121ABG), Buenos Aires, Argentina

Correspondence, R.E. Vogler: robertovogler@fcnym.unlp.edu.ar; robertovogler@yahoo.com.ar

**Abstract.** The best way to reduce problems related to invasive species is by preventing introductions into potentially susceptible areas. The purpose of this study was to create distribution models for the invasive gastropod *Achatina fulica* Bowdich, 1822 in South America in order to evaluate its potential geographic distribution and identify areas at potential risk. This mollusc, considered one of the 100 world's worst invasive alien species, is the focus of intense concern due to its impact on agriculture, human health, and native fauna. We tested two commonly used ecological niche modeling methods: Genetic Algorithm for Rule-Set Prediction (GARP) and Maximum Entropy (MaxEnt). Models were run with occurrence points obtained from several sources, including the scientific literature, international databases, governmental reports and newspapers, WorldClim bioclimatic variables, and altitude. Models were evaluated with the threshold-independent Receiver Operating Characteristic (ROC) and Area Under the Curve (AUC). Both models had consistent performances with similar areas predicted as susceptible, including areas already affected and new potentially susceptible areas in both tropical and temperate regions of South America.

**Key words:** land mollusc, bioinvasion, distribution, GARP, MaxEnt

In recent decades, South America has been seriously affected by invasive mollusc species (Letelier *et al.* 2007, Gutiérrez Gregoric and Vogler 2010, Rumi *et al.* 2010). Some of these introductions have been the result of unintentional events, such as the cases of *Theba pisana* Müller, 1774 and *Melanooides tuberculata* Müller, 1774 (Rumi *et al.* 2010, Peso *et al.* 2011). In contrast, other introductions have been intentional as a consequence of diverse interests, including biological control and ornamental, medicinal, and economic purposes (Cowie and Robinson 2003, Fernandez *et al.* 2003). The giant African snail *Achatina (Lissachatina) fulica* Bowdich, 1822, native to eastern Africa, is an example of the latter.

Considered one of the 100 world's worst invasive alien species (Lowe *et al.* 2000, Raut and Barker 2002), *Achatina fulica* was first introduced in South America at an agricultural fair in Brazil in the late 1980's as a breeding animal in order to compete with *Cornu aspersum* (Müller, 1774), the "real" escargot (Teles and Fontes 2002, Thiengo *et al.* 2007). Due to the lack of snail-eating habits of the Brazilian population, facilities where the species used to be reared were abandoned and the animals were released into the wild (Teles and Fontes 2002, Thiengo *et al.* 2007, Zenni and Ziller 2010).

These events were followed by a rapid and expansive dispersal in Brazil with its occurrence confirmed in 25 of 26 states in the country (Maldonado Júnior *et al.* 2010, Salgado 2011).

Recent reports suggest that *Achatina fulica* has expanded its range and become widespread throughout several South American countries, including Argentina, Colombia, Ecuador, Paraguay, Peru and Venezuela (Correoso Rodríguez 2006, Martinez-Escarbassiere *et al.* 2008, Borrero *et al.* 2009, Correoso and Coello 2009, Paraguay Biodiversidad 2010, Gutiérrez Gregoric *et al.* 2011). This broad distribution may not only be attributed to active dispersal from established populations (Correoso and Coello 2009) but also to the repeated transportation opportunities provided by people, including fishing (*i.e.* using the individuals as bait), heliculture, sold as a "pet", and the inadvertent transport of specimens (*e.g.*, with waste, building materials, plants). Together these factors have contributed to a rapid geographical range expansion of this invasive species (Borrero *et al.* 2009, Colley 2010, Gutiérrez Gregoric *et al.* 2011). In this context, bioclimatic models (also known as species distribution models and ecological niche models; Jeschke and Stryer 2008) constitute important tools to predict future changes in the species geographic ranges,

since they may be used to identify potential areas of suitable habitat where invasive species may actually be present but not yet detected and favorable regions where they may disperse in the future (Nyári *et al.* 2006, Baldwin 2009, Fukasawa *et al.* 2009, Robinson *et al.* 2010).

Predictive modeling characterizing the distribution of *Achatina fulica* in South America, especially in those countries where the species introduction has been recent, represents a valuable tool for developing effective control policy in order to manage the ongoing invasion. An extensive list of methods to generate spatial distribution models of plants, animals, and viruses at different geographical scales, have been described in the literature (see review by Elith and Leathwick 2009) and include GARP (*Genetic Algorithm for Rule-Setting Prediction*; Stockwell and Peters 1999) and MaxEnt (*Maximum Entropy*; Phillips *et al.* 2004, 2006). Using MaxEnt and 13 environmental variables, Borrero *et al.* (2009) developed a potential distribution model for the giant African snail in South America; however, this model focused on the Andean region. This model indicated low to medium probability that the species would occur in southern South America; however, due to recent findings of the species in subtropical areas of Argentina and Paraguay, we hypothesize that the potential southern distribution area would be larger than previously predicted. In this paper, we generate two high-resolution bioclimatic models using GARP and MaxEnt in order to re-evaluate threat of *A. fulica* in the southernmost region of South America. The models generated will be useful in planning control strategies, prioritizing areas for surveillance, and for taking preventive actions in order to limit the spread of *A. fulica*.

## MATERIAL AND METHODS

### Study area and species records

The study area included all South American countries. Presence data ( $N = 490$ ) for *Achatina fulica* were obtained from several sources, including the scientific literature, international databases, government reports and newspapers (Table 1). For those localities where coordinates were not provided by the source, a georeferenced position was obtained from the gazetteer GEOLocate Web Application (<http://www.museum.tulane.edu/geolocate/web/webgeoref.aspx>).

### Environmental variables

Nineteen bioclimatic variables and a topographic variable (altitude) (Table 2), at a spatial resolution of 30 arc seconds (approximately 1 km<sup>2</sup>), were used as predictors. These data, obtained from WorldClim v. 1.4 (<http://www.worldclim.org>), were derived from weather station

data spanning 1950–2000 (Hijmans *et al.* 2005). Given the heterogeneous habitats that *Achatina fulica* can occupy and its polyphagous diet, we chose to not include vegetation cover in our models, as was done in De Meyer *et al.* (2008).

### Modeling of susceptible areas

Models of the potential distribution of *Achatina fulica* were generated by using two algorithms: GARP (DesktopGarp v. 1.1.6; Stockwell and Noble 1992, Stockwell and Peters 1999, Scachetti-Pereira 2002, Scachetti-Pereira and Siqueira 2007) and MaxEnt (MaxEnt v. 3.3.3a; Phillips *et al.* 2004, 2006). These algorithms model the species' ecological niche (a set of ecological conditions habitable for a species) by examining the relationship between locations of known species presence and the environmental characteristics of that area and then extrapolating from this the areas where similar conditions occur in the study area (Paredes-García *et al.* 2011). In both models, 75% of presence records were randomly selected and used in the model training while the remaining 25% were used in the model testing.

For GARP, we generated 100 models with a convergence limit of 0.05, a maximum of 100 iterations, and all four rules (atomic, range, negated range, and logistic regression). The 20 best models were selected by hand following the recommendations of Anderson *et al.* (2003). The predictions of the best subset were arithmetically combined in ArcGIS v.9.1 (ESRI, Redlands, California) to create a composite prediction of the species potential distribution. This potential distribution was interpreted as a surface of densities related to the probability of suitable conditions for *Achatina fulica*. The consensus GARP map was reclassified into four categories: areas highly susceptible (pixels where the 20 models predicted presence), areas of susceptibility medium-high (pixels where 19/20 models predicted presence), areas of susceptibility medium-low (pixels where 18/20 models predicted presence), and areas of low susceptibility (pixels which converge up to 16 of the 20 models).

For MaxEnt, the model was computed as "logistic". This output returns a continuous map with an estimated probability of presence between 0 (no probability of the species presence) and 1 (high probability of presence), which permits fine distinctions between the suitability of different areas modeled (Giovanelli *et al.* 2008). All other parameters were used by default settings. Increased presence probability areas for *Achatina fulica* were considered to be more susceptible to invasion by the species.

To determine which variables contributed most to the development of the model, the MaxEnt program was configured to calculate the significance of variables using the jackknife procedure. This procedure produces three types of models: models created with an omitted variable and all other

**Table 1.** Sources of *Achatina fulica* occurrence in South America used in the two bioclimatic models.

Country	Consulted sources
Argentina	Gutiérrez Gregoric <i>et al.</i> (2011)
Brazil	Teles <i>et al.</i> (1997) Vasconcellos and Pile (2001) Teles and Fontes (2002) Carvalho <i>et al.</i> (2003) Barçante <i>et al.</i> (2005) Colley <i>et al.</i> (2005) Eston <i>et al.</i> (2006) Sacramento <i>et al.</i> (2006) Berto and Bogéa (2007) Oliveira <i>et al.</i> (2007) Paula and Lopes (2007) Thiengo <i>et al.</i> (2007) Agudo-Padrón (2008) Schiffler <i>et al.</i> (2008) Thiengo <i>et al.</i> (2008) Albuquerque <i>et al.</i> (2009) Colley and Fischer (2009) Ohlweiler <i>et al.</i> (2010) Thiengo <i>et al.</i> (2010) <b>Web sites</b> (November 2011): Base de dados sobre especies exóticas invasoras I3N-Brasil: <a href="http://www.institutohorus.org.br">http://www.institutohorus.org.br</a> ClickSergipe (2010): <a href="http://www.clicksergipe.com.br">http://www.clicksergipe.com.br</a> Conquiologistas do Brasil (2010): <a href="http://www.conchasbrasil.org.br">http://www.conchasbrasil.org.br</a> Gazeta do Triângulo (2010): <a href="http://www.gazetadotriangulo.com.br/novo/">http://www.gazetadotriangulo.com.br/novo/</a> Injipa (2010): <a href="http://injipa.com.br">http://injipa.com.br</a> Portal Agora Maranhão (2010): <a href="http://www.agoramaranhao.com">http://www.agoramaranhao.com</a> Portal de São Francisco do Sul (2010): <a href="http://www.sfs.com.br">http://www.sfs.com.br</a> Portal O Dia (2010): <a href="http://www.portalodia.com">http://www.portalodia.com</a> Prefeitura de Aracruz (2010): <a href="http://www.aracruz.es.gov.br">http://www.aracruz.es.gov.br</a> Prefeitura de Bela Vista (2010): <a href="http://www.belavista.go.gov.br">http://www.belavista.go.gov.br</a> Prefeitura de Jataí (2010): <a href="http://www.jatai.go.gov.br">http://www.jatai.go.gov.br</a> Rede Globo (2010): <a href="http://www.globo.com">http://www.globo.com</a> Rondonia Digital (2010): <a href="http://rondoniadigital.com">http://rondoniadigital.com</a>
Bolivia	Correoso and Coello (2009)
Colombia	Borrero <i>et al.</i> (2009) Correoso and Coello (2009) Base de datos sobre especies exóticas invasoras I3N-Colombia (November 2011): <a href="http://ef.humboldt.org.co">http://ef.humboldt.org.co</a> .
Ecuador	Borrero <i>et al.</i> (2009) Correoso and Coello (2009)
Paraguay	Paraguay Biodiversidad (2010)
Peru	Borrero <i>et al.</i> (2009) Correoso and Coello (2009)
Venezuela	Carvalho <i>et al.</i> (2003) Martínez-Escarbassiere <i>et al.</i> (2008)

variables included, models including only one variable, and a model created with all variables. The variables considered most important for model development are those reducing training gain when are not included in the model and show gain when the model is developed involving just the single

variable (Colacicco-Mayhugh *et al.* 2010, Torres and Jayat 2010).

Finally, comparison of the resulting models from both algorithms was done through Receiver Operating Characteristic (ROC curves analyses; Fielding and Bell 1997) representing

**Table 2.** Bioclimatic variables used in models development. Temperatures are expressed in °C \*10, precipitations in mm, and elevation in m above sea level.

Variable	Description
alt	Altitude
bio1	Annual mean temperature
bio2	Mean diurnal range (monthly mean, T° max-T° min)
bio3	Isothermality (bio2/bio7) x 100
bio4	Temperature seasonality (standard deviation x 100)
bio5	Maximum temperature of warmest month
bio6	Minimum temperature of coldest month
bio7	Temperature annual range (bio5-bio6)
bio8	Mean temperature of wettest quarter
bio9	Mean temperature of driest quarter
bio10	Mean temperature of the warmest quarter
bio11	Mean temperature of coldest quarter
bio12	Annual precipitation
bio13	Precipitation of wettest month
bio14	Precipitation of driest month
bio15	Precipitation seasonality (coefficient of variation)
bio16	Precipitation of wettest quarter
bio17	Precipitation of driest quarter
bio18	Precipitation of the warmest quarter
bio19	Precipitation of the coldest quarter

the distribution predictions vs. independent presence and absence data and, by calculating the Area Under the Curve (AUC). The AUC is a threshold independent index commonly used to assess prediction maps (Pearce and Ferrier 2000, Giovanelli *et al.* 2008), which can take values between 0.5 (no predictability) and 1 (perfect prediction). According to Loo *et al.* (2007), a value above 0.8 indicates a strong prediction. For GARP it is only possible to calculate a single value of AUC due to the nature of the algorithm (Larson *et al.* 2010), whereas for MaxEnt two values, one for model training and one for model testing are calculated. However, since the strength of the predictions of both methods cannot be compared directly (Phillips *et al.* 2006), identification of areas susceptible to invasion by *Achatina fulica* in South America was interpreted separately but in a comparative manner. In addition, to increase the level of detail within each country, the primary political division (*e.g.*, state, province) was incorporated into the models for a better interpretation of results.

## RESULTS

In general terms, both models had very good performance and agreed that a large region of South America is highly susceptible to invasion by *Achatina fulica* (Fig. 1). The

AUC calculated from the ROC curve generated for GARP was 0.805. Meanwhile, MaxEnt showed the best performance for the analyzed dataset, with AUC values of 0.944 for training data and 0.921 for test data, with a standard deviation of 0.010. The MaxEnt model showed a larger area of susceptibility to invasion by the giant African snail, showing in all countries a greater number of affected states (also called departments, regions, provinces or districts according to the country) than GARP (Figs. 2–4, Table 3). According to GARP, countries such as Chile and Uruguay, would not be susceptible to invasion, but would be according to MaxEnt, although the area affected was small (Fig. 4).


The jackknife test of variable importance in MaxEnt showed that temperature seasonality (bio4) and mean temperature of coldest quarter (bio11) were the variables that most influenced the development of the model when these variables were used alone (Fig. 5). Temperature seasonality (bio4), isothermality (bio3), and altitude (alt) exhibited modest reductions in training gain when they were removed from the model, thus indicating that contain information necessary for the model (Fig. 5). The remaining variables contributed less to model development.

## DISCUSSION

In this study, models generated with GARP and MaxEnt produced similar results and consistent prediction maps. Although the predictive power of both algorithms cannot be directly compared (Phillips *et al.* 2006), parallel interpretation of the graphical output was useful in identifying area susceptible to *Achatina fulica* invasion.

Our results are in agreement with other previous studies, which suggest that even though the species exhibits a wide environmental tolerance, it prefers warm habitats (Raut and Barker 2002, Albuquerque *et al.* 2009). Specifically, temperature seasonality and mean temperature of coldest quarter were identified as the two variables having the most important effect on the potential distribution of *Achatina fulica*. These findings would explain the absence of species in southern South America, which is characterized by marked seasonality and low temperatures.

Our models matched many of the predictions made by Borrero *et al.* (2009) and Correoso and Coello (2009) with susceptible areas identified in Guyana, French Guiana, Suriname, Peru, Venezuela, Ecuador and Colombia; however, by including the southernmost records of the species for model training and employing a higher spatial resolution, the potential distribution areas in the South America temperate regions were expanded. According to our results, Chile and Uruguay appear to be the least susceptible countries, with the exception of southern Chile, which might be susceptible to invasion




**Figure 1.** Bioclimatic models of susceptible areas: **A**, GARP and **B**, MaxEnt. Lighter areas indicate low susceptibility regions and darker areas indicate high susceptible regions. Points indicate occurrences records used in study.

due to the existence of favorable environmental conditions. Specifically, this region of Chile contains “Valdivian Temperate Rainforest”, which has been reported to contain another exotic snail, *Cornu aspersum*, and an exotic slug, *Arión intermedius* Normand, 1852 (Cádiz and Gallardo 2007, Artacho and Nespolo 2009), suggesting that environmental conditions exist to support invasions of other mollusc species, including *Achatina fulica*.

In Argentina, the northeast region shows varying degrees of susceptibility but the overall trend is towards a medium susceptibility level. Even the north central area could be identified as susceptible, although with a low probability. Nonetheless, it is important to maintain vigilance of this area because it is comprised of Subtropical Cloudforest (Yungas), which shares some environmental conditions with the Paranaense Rainforest (located in northeastern Argentina, southern Brazil, and East of Paraguay), where the species is already present (Miranda and Cuzzo 2010, Gutiérrez Gregoric *et al.* 2011).

In Paraguay, despite the fact that the majority of the eastern portion of the country can be considered as a highly susceptible area, *Achatina fulica* has only been reported in the one southern location, the Ayolas (Table 1). Due to the large number of occurrences of *A. fulica* in Brazil along the Paraguay border (Thiengo *et al.* 2007, Colley and Fischer 2009), it can be assumed that the species may occur in other locations of Paraguay but its presence has not yet been reported.

The majority of Bolivia is highly susceptible to *Achatina fulica* invasion, with the exception of the southwest region where susceptibility appears to be low. We were unable to locate confirmed locations of the species presence; however, Correoso and Coello (2009) reported that the species was being commercialized in the city of La Paz. This fact, combined with the country’s shared border with Brazil and Paraguay (Paula and Lopes 2007, Thiengo *et al.* 2007) suggests that the species is inevitably present in Bolivia but surveys are needed to confirm its distribution.



**Figure 2.** Areas susceptible to invasion in northern South America (Venezuela, Guyana, French Guiana and Suriname), according to **A**, GARP and, **B**, MaxEnt.





Figure 3. Areas susceptible to invasion in northwestern and central South America (Colombia, Ecuador, Peru, and Brazil), according to A, GARP and, B, MaxEnt.



**Figure 4.** Areas susceptible to invasion in southern South America (Argentina, Chile, Uruguay, Paraguay and Bolivia), according to **A**, GARP and, **B**, MaxEnt.




**Table 3.** Primary political division of South American countries susceptible to invasion according to each model. The summary for each political division includes the total number of units per country, as well as the number of susceptible units identified with GARP and MaxEnt. † Total political units per country; ‡ Political units susceptible according to GARP; § Political units susceptible according to MaxEnt.

Country	Primary political division		
	Total†	GARP‡	MaxEnt§
Argentina	23	6	15
Bolivia	9	7	7
Brazil	26	26	26
Chile	15	0	13
Colombia	32	29	31
Ecuador	24	18	23
French Guiana	2	2	2
Guyana	10	10	10
Paraguay	17	16	17
Peru	24	17	24
Suriname	10	10	10
Uruguay	19	1	10
Venezuela	24	24	24

For Brazil, models indicate the area of highest susceptibility to occur between the midwest and Atlantic coast in a southeastward direction. Some areas of low susceptibility are noted in the northwest and northeast portions of the country. In 2005, in response to the invasive behavior of the giant African snail and its implications on health, biodiversity, and agriculture, IBAMA (Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis) forbade the sale and breeding of *Achatina fulica* (Carvalho-Junior and Nunes 2009). Remarkably, Brazil is the only South American country with *A. fulica* regulatory legislation.

We identified a larger area in South America that is climatically susceptible to be invaded by the giant African snail than that previously reported by Borrero *et al.* (2009). A comparative analysis of our models clearly indicates that several subcontinent temperate regions contain the minimal environmental conditions able to support an invasion of *Achatina fulica*. Several newly-predicted susceptible areas involve boundaries with countries where the invasive species is known to occur. This knowledge is especially useful to governmental authorities in countries where the invasive species was recently confirmed (*e.g.*, a single location in Argentina that borders Brazil and Paraguay; Gutiérrez Gregoric *et al.* 2011) or where it is yet unreported (*e.g.*, Chile and Uruguay). We expect that control and management authorities can use this information to establish priority areas for surveillance and to plan preventive actions aimed at limiting the spread of the species.



**Figure 5.** Jackknife test of training gain for the MaxEnt model. Abbreviations: **alt**, elevation, **bio1** through **bio19**, bioclimatic variables (Table 2).

## ACKNOWLEDGMENTS

The authors wish to thank Dr. Karen De Matteo for helpful comments and English editing.

## LITERATURE CITED

- Agudo-Padrón, A. I. 2008. Listagem sistemática dos moluscos continentais ocorrentes no Estado de Santa Catarina, Brasil. *Comunicaciones de la Sociedad Malacológica del Uruguay* 9: 147–179. [In Portuguese].
- Albuquerque, F. S., M. C. Peso-Aguiar, M. J. T. Assunção-Albuquerque, and L. Gálvez. 2009. Do climate variables and human density affect *Achatina fulica* (Bowditch) (Gastropoda: Pulmonata) shell length, total weight and condition factor? *Brazilian Journal of Biology* 69: 879–885.
- Anderson, R. P., D. Lew, and A. T. Peterson. 2003. Evaluating predictive models of species' distribution: Criteria for selecting optimal models. *Ecological Modelling* 162: 211–232.
- Artacho, P. and R. Nespolo. 2009. Natural selection reduces energy metabolism in the garden snail, *Helix aspersa* (*Cornu aspersum*). *Evolution* 63: 1044–1050.
- Baldwin, R. A. 2009. Use of maximum entropy modeling in wildlife research. *Entropy* 11: 854–866.
- Barçante, J. M. P., T. A. Barçante, S. R. C. Dias, and W. S. Lima. 2005. Ocorrência de *Achatina fulica* Bowdich, 1822 (Mollusca:

- Gastropoda: Achatinoidea) no Estado de Minas Gerais, Brasil. *Boletim do Museu de Biologia Mello Leitão* **18**: 65–70. [In Portuguese].
- Berto, B. and T. Bogéa. 2007. Occurrence of nematode larvae in *Achatina fulica* Bowdich, 1822 (Gastropoda: Achatinidae) snails in Vargem Pequena, Rio de Janeiro, Brazil. *Revista de Patologia Tropical* **36**: 171–177.
- Borrero, F. J., A. S. H. Breure, C. Christensen, M. Correoso, and V. Mogollón Ávila. 2009. Into the Andes: Three new introductions of *Lissachatina fulica* (Gastropoda, Achatinidae) and its potential distribution in South America. *Tentacle* **17**: 6–8.
- Cádiz, F. J. and C. S. Gallardo. 2007. *Arion intermedius* (Gastropoda: Stylommatophora); first record of this introduced slug in Chile, with notes on its anatomy and natural history. *Revista Chilena de Historia Natural* **80**: 99–108.
- Carvalho, O. S., H. M. S. Teles, E. M. Mota, C. L. G. F. Mendonça, and H. L. Lenzi. 2003. Potentiality of *Achatina fulica* Bowdich, 1822 (Mollusca: Gastropoda) as intermediate host of the *Angiostrongylus costaricensis* Morera & Céspedes 1971. *Revista da Sociedade Brasileira de Medicina Tropical* **36**: 743–745.
- Carvalho-Júnior, V. C. B. and J. R. S. Nunes. 2009. Ocorrência e distribuição do caramujo africano *Achatina fulica* Bowdich, 1822, no município de Várzea Grande, MT. *Engenharia Ambiental* **6**: 606–620. [In Portuguese].
- Colacicco-Mayhugh, M. G., P. M. Masuoka, and J. P. Grieco. 2010. Ecological niche model of *Phlebotomus alexandri* and *P. papatasi* (Diptera: Psychodidae) in the Middle East. *International Journal of Health Geographics* **9**: 2. doi:10.1186/1476-072X-9-2.
- Colley, E. 2010. Medidas de controle de *Achatina fulica*. In: M. L. Fischer and L. C. M. Costa, eds., *O Caramujo Gigante Africano Achatina fulica no Brasil*. Editora Universitária Champagnat, Curitiba, Brazil. Pp. 203–230. [In Portuguese].
- Colley, E. and M. L. Fischer. 2009. Avaliação dos problemas enfrentados no manejo do caramujo gigante africano *Achatina fulica* (Gastropoda: Pulmonata) no Brasil. *Zoologia* **26**: 674–683. [In Portuguese].
- Colley, E., M. L. Fischer, M. Simião, R. D. Zenni, D. A. T. Silva, and N. Latoski. 2005. Diagnóstico da invasão do caramujo gigante africano *Achatina fulica* Bowdich, 1822 em um fragmento de Floresta Ombrófila Densa em Morretes, Paraná, Brasil. In: *Anais do VII Congresso de Ecologia do Brasil*. (November 2011). Available at: <http://www.seb-ecologia.org.br/viiceb>. Accessed August 2012. 2 pp. [In Portuguese].
- Correoso Rodríguez, M. 2006. Estrategia preliminar para evaluar y erradicar *Achatina fulica* (Gastropoda: Achatinaceae) en Ecuador. *Boletín Técnico IASA, Serie Zoológica* **2**: 45–52. [In Spanish].
- Correoso, M. and M. Coello. 2009. Modelación y distribución de *Lissachatina fulica* (Gastropoda: Achatinidae) en Ecuador. Potenciales impactos ambientales y sanitarios. *Revista Geoespacial* **6**: 79–90. [In Spanish].
- Cowie, R. H. and D. G. Robinson. 2003. Pathways of introduction of nonindigenous land and freshwater snails and slugs. In: G. M. Ruiz and J. T. Carlton, eds., *Invasive Species: Vectors and Management Strategies*. Island Press, Washington, District of Columbia. Pp. 93–122.
- De Meyer, M., M. P. Robertson, A. T. Peterson, and M. W. Mansell. 2008. Ecological niches and potential geographical distributions of Mediterranean fruit fly (*Ceratitis capitata*) and Natal fruit fly (*Ceratitis rosa*). *Journal of Biogeography* **35**: 270–281.
- Elith, J. and J. R. Leathwick. 2009. Species distribution models: Ecological explanation and prediction across space and time. *Annual Review of Ecology, Evolution, and Systematics* **40**: 677–697.
- Eston, M. R., G. V. Menezes, A. Z. Antunes, A. S. R. Santos, and A. M. R. Santos. 2006. Espécie invasora em unidade de conservação: *Achatina fulica* (Bowdich, 1822) no Parque Estadual Carlos Botelho, Sete Barras, SP, Brasil (Nota Científica). *Revista do Instituto Florestal* **18**: 173–179. [In Portuguese].
- Fernandez, M. A., S. C. Thiengo, and L. R. L. Simone. 2003. Distribution of the introduced freshwater snail *Melanoides tuberculatus* (Gastropoda: Thiariidae) in Brazil. *The Nautilus* **117**: 78–82.
- Fielding, A. H. and J. F. Bell. 1997. A review of methods for the assessment of prediction errors in conservation presence/absence models. *Environmental Conservation* **24**: 38–49.
- Fukasawa, K., F. Koike, N. Tanaka, and K. Otsu. 2009. Predicting future invasion of an invasive alien tree in a Japanese oceanic island by process-based statistical models using recent distribution maps. *Ecological Research* **24**: 965–975.
- Giovanelli, J. G. R., C. F. B. Haddad, and J. Alexandrino. 2008. Predicting the potential distribution of the alien invasive American bullfrog (*Lithobates catesbeianus*) in Brazil. *Biological Invasions* **10**: 585–590.
- Gutiérrez Gregoric, D. E. and R. E. Vogler. 2010. Riesgo de establecimiento del gasterópodo dulceacuícola invasor *Melanoides tuberculatus* (Thiariidae) en el Río de La Plata (Argentina-Uruguay). *Revista Mexicana de Biodiversidad* **81**: 573–577. [In Spanish].
- Gutiérrez Gregoric, D. E., V. Núñez, R. E. Vogler, and A. Rumi. 2011. Invasion of the Argentinean Paranense Rainforest by the Giant African Snail *Achatina fulica*. *American Malacological Bulletin* **29**: 135–137.
- Hijmans, R. J., S. E. Cameron, J. L. Parra, P. G. Jones, and A. Jarvis. 2005. Very high resolution-interpolated climate surfaces for global land areas. *International Journal of Climatology* **25**: 1965–1978.
- Jeschke, J. M. and D. L. Strayer. 2008. Usefulness of bioclimatic models for studying climate change and invasive species. *Annals of the New York Academy of Sciences* **1134**: 1–24.
- Larson, S. R., J. P. DeGroot, L. C. Bartholomay, and R. Sugumaran. 2010. Ecological niche modeling of potential West Nile virus vector mosquito species in Iowa. *Journal of Insect Science* **10**: 1–17.
- Letelier, S., L. A. M. Ramos, and M. L. G. Huaquín. 2007. Moluscos dulceacuícolas exóticos en Chile. *Revista Mexicana de Biodiversidad* **78**: 9S–13S. [In Spanish].
- Loo, S. E., R. Mac Nally, and P. S. Lake. 2007. Forecasting New Zealand mudsnail invasion range: Model comparisons using native and invaded ranges. *Ecological Applications* **17**: 181–189.
- Lowe, S., M. Browne, S. Boudjelas, and M. De Poorter. 2000. 100 of the World's worst invasive alien species. A selection from the global invasive species database. *Aliens* **12**. Available at: [http://www.issg.org/database/species/reference\\_files/100English.pdf](http://www.issg.org/database/species/reference_files/100English.pdf). Accessed February 2012.

- Maldonado Júnior, A., R. O. Simões, A. P. M. Oliveira, E. M. Motta, M. A. Fernandez, Z. M. Pereira, S. S. Monteiro, E. J. L. Torres, and S. C. Thiengo. 2010. First report of *Angiostrongylus cantonensis* (Nematoda: Metastrongylidae) in *Achatina fulica* (Mollusca: Gastropoda) from Southeast and South Brazil. *Memórias do Instituto Oswaldo Cruz* **105**: 938–941.
- Martínez-Escarbassiere, R., E. O. Martínez, and O. Castillo. 2008. Distribución geográfica de *Achatina (Lissachatina) fulica* (Bowdich, 1882) (Gastropoda-Stylommatophora-Achatinidae) en Venezuela. *Memoria de la Fundación La Salle de Ciencias Naturales* **169**: 93–106. [In Spanish].
- Miranda, M. J. and M. G. Cuezco. 2010. Biodiversidad de gasterópodos terrestres (Mollusca) en el Parque Biológico Sierra de San Javier, Tucumán, Argentina. *Revista de Biología Tropical* **58**: 1009–1029. [In Spanish].
- Nyári, A., C. Ryall, and A. T. Peterson. 2006. Global invasive potential of the house crow *Corvus splendens* based on ecological niche modelling. *Journal of Avian Biology* **37**: 306–311.
- Ohlweiler, F. P., M. C. A. Guimarães, F. Y. Takahashi, and J. M. Eduardo. 2010. Current distribution of *Achatina fulica*, in the State of São Paulo including records of *Aelurostrongylus abstrusus* (Nematoda) larvae infestation. *Revista do Instituto de Medicina Tropical de São Paulo* **52**: 211–214.
- Oliveira, D. S., V. L. S. Jeraldo, and C. M. Melo. 2007. Estudo de populações urbanas de *Achatina fulica* no Estado de Sergipe-Brasil. In: *Anais do VIII Congresso de Ecologia do Brasil*. (November 2011). Available at: <http://www.seb-ecologia.org.br/viiiceb>. Accessed August 2012. 2 pp. [In Portuguese].
- Paraguay Biodiversidad. 2010. Gastropoda. (November 2011). Available at: <http://www.pybio.org/201/gastropoda>. Accessed November 2011. [In Spanish].
- Paredes-García, D. M., A. Ramírez-Bautista, and M. A. Martínez-Morales. 2011. Distribución y representatividad de las especies del género *Crotalus* en las áreas naturales protegidas de México. *Revista Mexicana de Biodiversidad* **82**: 689–700. [In Spanish].
- Paula, A. M. D. and V. L. D. Lopes. 2007. Caramujo africano gigante (*Achatina fulica* Bowdich, 1822) no Estado de Mato Grosso: Distribuição, influências na saúde, ambientes e estratégias de controle do molusco. In: *II Mostra de Saúde. Governo do estado de Mato Grosso, Secretaria de Estado de Saúde, Superintendência de Vigilância em Saúde. Escola de Saúde Pública*. 6 pp. [In Portuguese].
- Pearce, J. and S. Ferrier. 2000. Evaluating the predictive performance of habitat models developed using logistic regression. *Ecological Modelling* **133**: 225–245.
- Peso, J. G., D. C. Pérez, and R. E. Vogler. 2011. The invasive snail *Melanoides tuberculata* in Argentina and Paraguay. *Limnologia* **41**: 281–284.
- Phillips, S. J., R. P. Anderson, and R. E. Schapire. 2006. Maximum entropy modeling of species geographic distributions. *Ecological Modelling* **190**: 231–259.
- Phillips, S. J., M. Dudík, and R. E. Schapire. 2004. A maximum entropy approach to species distribution modelling. In: *Proceedings of the 21<sup>st</sup> International Conference on Machine Learning*. Banff, Canada. Pp. 655–662.
- Raut, S. K. and G. M. Barker. 2002. *Achatina fulica* Bowdich and other Achatinidae as pests in tropical agriculture. In: G. M. Barker, ed., *Molluscs as Crop Pests*. CABI Publishing, Wallingford, United Kingdom. Pp. 55–114.
- Robinson, T. P., R. D. van Klinken, and G. Metternicht. 2010. Comparison of alternative strategies for invasive species distribution modeling. *Ecological Modelling* **221**: 2261–2269.
- Rumi, A., J. Sánchez, and N. Ferrando. 2010. *Theba pisana* (Müller, 1774) (Gastropoda, Helicidae) and other alien land molluscs species in Argentina. *Biological Invasions* **12**: 2985–2990.
- Sacramento, J. M. C., S. A. L. Gusmão, M. T. A. Gusmão, and W. V. D. Silvestre. 2006. Ocorrência de caramujo africano (*Achatina fulica*) em hortas do entorno de Belém-PA. In: *Anais do 46<sup>o</sup> Congresso Brasileiro de Olericultura*. (November 2011). Available at: <http://www.abhorticultura.com.br/biblioteca>. Accessed August 2012. 4 pp. [In Portuguese].
- Salgado, N. C. 2011. Biodiversidade da malacofauna terrestre brasileira II: Um olhar curioso à Amazônia legal. In: C. A. Oliveira de Meirelles, C. X. Barroso, and B. B. Batista, eds., *Livro de Resumos XXII Encontro Brasileiro de Malacologia*, Fortaleza. Pp. 7–9. [In Portuguese].
- Scachetti-Pereira, R. 2002. Desktop GARP v. 1.1.6. The University of Kansas Biodiversity Research Center and Reference Center for Environmental Information. (February 2012). Available at: <http://www.nhm.ku.edu/desktopgarp/index.html>. Accessed February 2012.
- Scachetti-Pereira, R. and M. F. D. Siqueira. 2007. Algoritmo genético para produção de conjuntos de regras (GARP). *Megadiversidade* **3**: 46–55. [In Portuguese].
- Schiffler, C. L., F. F. Gomes, N. B. Ederli, and F. C. R. Oliveira. 2008. *Achatina fulica* Bowdich (1822) um novo hospedeiro de espécies de *Cryptosporidium* (Apicomplexa, Cryptosporidiidae). *Revista Brasileira de Parasitologia Veterinária* **17**: 273–276. [In Portuguese].
- Stockwell, D. R. B. and I. R. Noble. 1992. Induction of sets of rules from animal distribution data: A robust and informative method of data analysis. *Mathematics and Computers in Simulation* **33**: 385–390.
- Stockwell, D. R. B. and D. Peters. 1999. The GARP modelling system: Problems and solutions to automated spatial prediction. *International Journal of Geographical Information Science* **13**: 143–158.
- Teles, H. M. S. and L. R. Fontes. 2002. Implicações da introdução e dispersão de *Achatina fulica* Bowdich, 1822 no Brasil. *Boletim do Instituto Adolfo Lutz* **12**: 3–5. [In Portuguese].
- Teles, H. M. S., J. F. Vaz, L. R. Fontes, and M. F. Domingos. 1997. Registro de *Achatina fulica* Bowdich, 1822 (Mollusca, Gastropoda) no Brasil: Caramujo hospedeiro intermediário da angiostrongilíase. *Revista de Saúde Pública* **31**: 310–312. [In Portuguese].
- Thiengo, S. C., F. A. Faraco, N. C. Salgado, R. H. Cowie, and M. A. Fernandez. 2007. Rapid spread of an invasive snail in South America: The giant African snail, *Achatina fulica*, in Brazil. *Biological Invasions* **9**: 693–702.
- Thiengo, S. C., M. A. Fernandez, E. J. L. Torres, P. M. Coelho, and R. M. Lanfredi. 2008. First record of a nematode Metastrongyloidea (*Aelurostrongylus abstrusus* larvae) in *Achatina (Lissachatina) fulica* (Mollusca, Achatinidae) in Brazil. *Journal of Invertebrate Pathology* **98**: 34–39.

- Thiengo, S. C., A. Maldonado, E. M. Mota, E. J. L. Torres, R. Caldeira, O. S. Carvalho, A. P. M. Oliveira, R. O. Simões, M. A. Fernandez, and R. M. Lanfredi. 2010. The giant African snail *Achatina fulica* as natural intermediate host of *Angiostrongylus cantonensis* in Pernambuco, northeast Brazil. *Acta Tropica* **115**: 194–199.
- Torres, R. and J. P. Jayat. 2010. Modelos predictivos de distribución para cuatro especies de mamíferos (Cingulata, Artiodactyla y Rodentia) típicas del Chaco en Argentina. *Mastozoología Neotropical* **17**: 335–352. [In Spanish].
- Vasconcellos, M. C. and E. Pile. 2001. Ocorrência de *Achatina fulica* no Vale do Paraíba, Estado do Rio de Janeiro, Brasil. *Revista de Saúde Pública* **35**: 582–584. [In Portuguese].
- Zenni, R. D. and S. R. Ziller. 2010. Invasões biológicas: Problemas econômicos e ambientais com ênfase em *Achatina fulica*. In: M. L. Fischer and L. C. M. Costa, eds., *O Caramujo Gigante Africano Achatina fulica no Brasil*. Editora Universitária Champagnat, Curitiba. Pp. 175–188. [In Portuguese].

**Submitted:** 29 April 2012; **accepted:** 1 September 2012;  
**final revisions received:** 22 November 2012