

# Functional Perspective of Feeder Organelle from Three-dimensional Ultrastructural Characteristics in *Cryptosporidium parvum*

著者	BOCHIMOTO Hiroki, KONDOH Daisuke, ISHIHARA Yo, Mohammad Hazzaz Bin KABIR, KATO Kentaro
journal or publication title	Journal of Integrated Field Science
volume	17
page range	30-30
year	2020-03
URL	<a href="http://hdl.handle.net/10097/00128796">http://hdl.handle.net/10097/00128796</a>

## **O-1. Functional Perspective of Feeder Organelle from Three-dimensional Ultrastructural Characteristics in *Cryptosporidium parvum***

Hiroki BOCHIMOTO<sup>1,2</sup>, Daisuke KONDOH<sup>3</sup>, Yo ISHIHARA<sup>4</sup>, Mohammad Hazzaz Bin KABIR<sup>2</sup>  
and Kentaro KATO<sup>2,5</sup>

<sup>1</sup>Division of Aerospace Medicine, Department of Cell Physiology, The Jikei University School of Medicine

<sup>2</sup>National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine

<sup>3</sup>Laboratory of Veterinary Anatomy, Obihiro University of Agriculture and Veterinary Medicine

<sup>4</sup>Shonan Kamakura General Hospital

<sup>5</sup>Laboratory of Sustainable Animal Environment, Graduate School of Agricultural Science, Tohoku University

*Cryptosporidium* is a parasite causing extensive illness both in livestock and humans. Feeder organelle of *Cryptosporidium* is the multi-membranous structures localized on the parasite-host-cell interface that deprive nutrients from host cells. Although the feeder organelle has been summarized as being highly invaginated membranous structure, the three-dimensional fine structure remains unclear. Osmium-maceration procedure for scanning electron microscopy (OS-SEM) is one of the methods to enable visualization of the intracellular ultrastructure including depth direction information after removing soluble proteins. Recently, we investigated and assessed *C. parvum* possessed on the surface of ileal epithelial cells of SCID mouse by using transmission electron microscopy (TEM) and OS-SEM. By TEM observation, feeder organelles were recognized as aggregated structures of concentric-, vertically- and randomly-lined bars. Correspondingly, OS-SEM observation revealed the reticulated network of stacked flat bursiform, ring-shaped bursiform and reticulated tubular membranes. These findings of the three-dimensional ultrastructural characteristics of feeder organelle, which are more intricate than expected, may potentially reinforce the limited knowledge regarding the nature of this attachment interface and the functional mechanisms around extraction of nutrients.