

Indian J. Fish., 44(3) : 279-286, Jul.-Sep., 1997

Branchial, renal and hepatic lesions in an estuarine mullet, *Liza parsia* Hamilton - Buchanan, induced by sublethal exposure to BHC

A.K. PANDEY¹, K.C. GEORGE AND M. PEER MOHAMED

Central Marine Fisheries Research Institute, Cochin - 682 014, India

ABSTRACT

Gill of *Liza parsia* exposed to sublethal concentration (0.5 ppm) of BHC responded initially with the copious secretion of mucus followed by oedematous separation of epithelial lining cells from the basement membrane. Fusion of secondary gill lamellae, tissue hyperplasia and multiple telangiectases (aneurysms) were noticed on day 10 of the exposure. By day 15, the entire interlamellar spaces became filled with the hyperplastic epithelium. Kidney revealed an initial hypertrophy of the cells lining proximal convoluted tubules followed by shrinkage in the glomerular tufts resulting in the increase of Bowman's space and oedema. Tubulonecrosis, hyperemia and fibrosis were noticed in the renal tissue after day 19 of BHC intoxication. Exposure to BHC elicited an initial dilation of bile canaliculi, enhanced secretion of bile and cellular hypertrophy on day 4. Necrotic changes like excessive vacuolation, and karyorrhexis and karyolysis were seen on day 8. By day 10 and 15, complete vacuolation of hepatocytes, pycnosis of nuclei and focal necrosis were noticed.

Introduction

Organochlorine (OC) pesticides have become very popular because of their broad spectrum insecticidal efficiency and relatively low operational cost (Ruvio, 1972). Benzene hexachloride (BHC) or hexachlorocyclohexane (HCH) is used extensively in India for pest management in agriculture and forestry (Basak and Konar, 1977). The pesticide gets accumulated in fish and thus finds its way to food chain (Ruvio, 1972; Dikshith *et al.*, 1990; Bakre *et al.*, 1990). Mathur (1964) has recorded the patho-

logical symptoms in the liver of freshwater teleosts exposed to acute doses of BHC. However, nothing is known about the toxic effects of this pesticide on the fishes inhabiting coastal or marine environments (Sen Gupta and Qasim, 1985). Therefore, an attempt has been made to record the branchial, renal and hepatic lesions caused by sublethal exposure of BHC to the estuarine mullet, *Liza parsia*, a commercially important species inhabiting both the coasts of India.

Present address : ¹ National Bureau of Fish Genetic Resources, Post Box No. 19, Lucknow - 226 004, India.

Materials and methods

Live specimens of *Liza parsia* (average length 10.5±0.8 cm, average weight 8.6±0.4 g) were collected by operating cast net from Vypeen Island (09°57.8' N, 76° 14.4' E) near Cochin and transported to the laboratory. Fish were acclimatised in the aquaria of 120 litres capacity containing well-aerated sea water (salinity 22 ppt, pH 7.69, oxygen content 4.32 ml/litre, water temperature 29.2°C) for a period of one week prior to use. Thereafter, they were divided into two equal groups of 40 specimens each at random and were subjected to the following treatments.

Group A : Fishes were maintained in sea water and served as controls.

Group B : Experimental fishes were exposed to the sublethal (chronic-one-half of LC₆₀ value for 96 hrs) concentration (0.5 ppm) of BHC (Hindustan Insecticides Limited, Delhi). Since the pesticide is insoluble in water, it was initially dissolved in acetone and the required concentration was achieved by adding sea water.

The media were renewed every alternate day. Since fish did not accept the formulated feed under the experimental conditions, they were not fed during the entire course of investigation. About 30 % mortality was observed among the Group B fishes after day 10 of the exposure. Dead fishes were removed immediately from the aquarium to avoid oxygen depletion and discarded from the study.

Five specimens from both the groups were killed on day 1, 2, 4, 8, 10 and 15 of the treatment. Tissues (gill, kidney and liver) were surgically removed and fixed immediately in freshly prepared aqueous Bouin's solution and Zenker's

fluid. After 24 hrs, the tissues were washed thoroughly in running tap water, dehydrated in ascending series of alcohol, cleared in xylene and embedded in paraffin wax at 60°C. Serial sections were cut at 6-8 μ m and stained in hematoxylin-eosin (H&E) and periodic Acid-Schiff's reagent (PAS).

Results

BHC exposure elicited excessive secretion of mucus in the interlamellar spaces of the gill by 24 hr. Dilation of blood vessels, swelling of the tips of secondary gill lamellae and partial separation of epithelial lining cells from the basement membrane were observed on day 2. By day 4, the experimental group mullets revealed lamellar oedema and complete separation of epithelial cells from the basement membrane of secondary gill lamellae. Furthermore, the epithelial cells of secondary lamellae depicted necrotic changes too (Fig.1). The first telangiectatic secondary lamellae was observed on day 8 (Fig. 2).



Fig. 1. Gill of *Liza parsia* on day 4 of BHC exposure. Mark the oedematous separation of epithelium from the basement membrane of pillar cells (arrow). H&E x 600.

fluid. After 24 lire, **the tissues** were washed thoroughly in running tap water, dehydrated in ascending series of alcohol, **cleared in** xylene and embedded in paraffin wax at **60°C**. Serial sections were **cut at 8-8 u** and stained in **hematoxylin-eosin (H&E)** and **periodic Acid-Schiff's** reagent (PAS).

Results

BUG exposure elicited **excessive** secretion **of mucus in the interlamellar spaces of the gill** by **24 hr**. **Dilation** of blood vessels, swelling of the **tips** of secondary gill lamellae and partial separation of epithelial lining" cells from **the basement membrane** were observed **on** day 2. By clay 4, **the** experimental group **mulletts** revealed **lamellar oedema** and complete separation **of** epithelial cells **from the basement membrane** of secondary **gill** lamellae. Furthermore, **the** epithelial cells of secondary lamellae depicted **necrotic** changes **too** (Fig. 1). The first telangiectatic secondary lamellae was observed on day 8 (Fig. 2).

1 'i, , i , 'I, , , 1, (. i - F), | j , i , <
 <ii 'j|, <(> J I I ' * * 't . > H , (D L . , , >, UI,
 ' " I ' , > 1, 1" (> I I I I I I I I I I I I I I I I I I
 ' " ' < " (> I ' * I ' H, . * I j ' ' * * < L , ' I I '
 i, il' . I u ' i ' i ' < . . . , ' lic O ; (/ (. , . i . * <
 i ' ' < " > ; * i ' I < > ; , i ' I - , (> i - in i
 < > ii , - ' . | Hi K - I , . I * * il ' ; ; , (
 ' n < , ' i ' ' until i . i ' i " - I , , I * * (R < | i
 , ' > < ' * I j ij (< < r n / ifi, i' I * of
 I ' ' I ' I fr , i i l ' h j . f) . ' s I * s i f ' i k , (i l < ' :
 " . i ;) < d ; , , ' i / b , i . - " i < i * i i i , , i - i i it , .
 " I < ' ' s ' i l ' ' i - i " ' - ' I ' I ' ' i i I - m I
 I , i ' H ' ' i ' I * ' ' i ' n < i l l > i l i l ' i n . i l l ' - I
 i ' - > i . H - I ' (! ' I *) - > ; i l I i l ' , i t ' ,) * (
 il * i * ' . . j * * I , , i . > ' - ! i i , (i - n i ' M
 > > i H) < ' ' i , ' i l ' . i < I n i l , ! < . . , < ' . . , (.
 I e - I ' ' H - t , i h ' . . . , i ' , ! > ; , . . j , i I i i n <

BHC treatment for. 2 clays elicited dilation of **bile canaliculi**, increased secretion of **bile** and cellular **hypertrophy**. Most of the **hepatocytes depicted cytoplasmic vacuolation** resulting in the **displacement of nuclei towards** the

in on day 10 of BHC
 the separation d^{fr}
from basement mem-
condary gill lamellae

Fig. 9. Kidney of *Liza parsia* on day 12 of BBC exposure? exhibiting degenerative changes like **hyperemia, pycnosis** (arrow) **and** fibrosis (broken arrow), H&E x 800,

Liver is an important target organ affected by pollutants in fishes (Roberts, 1989). Several workers have reported degenerative changes in hepatic tissue subjected to pollution by various pesticides and insecticides (Kumar and Pant 1984; Gill *et al.*, 1990; Pandey *et al.*, 1993). The present observation in *Liza parsia* clearly demonstrates liver to be the organ affected most severely in response to BHC exposure.

Acknowledgment

We are grateful to Dr. P.S.B.R. James, Ex-Director, C.M.F.R.I., Cochin for encouragement and inspiration.

References

- Bakre, P.R., B. Mishra and P. Bhatnagar 1990. Residues of organochlorine insecticides in fish from Malala water reservoir, Jaipur, India. *Bull. Environ. Contam. Toxicol.*, 45 : 394-398.
- Basak, P.K. and S.K. Konar 1977. Estimation of safe concentration of insecticides : a new method tested on DDT and BHC. *J. Inland Fish. Soc. India*, 9 : 9-29.
- Dikshith, T.S.S., R.B. Raizada, S.N. Kumar, M.K. Srivastava, S.K. Kulshrestha and U.N. Andholia 1990. Residues of DDT and HCH in major sources of drinking water in Bhopal, India. *Bull. Environ. Contam. Toxicol.*, 45 : 389-393.
- Gill, T.S., J.S. Pandey and H. Tewari 1990. Hepatopathotoxicity of three pesticides in a freshwater fish, *Puntius conchoni* Ham. *J. Environ. Sci. Hlth.*, 25A : 653-663.
- Jones, T.C. and R.D. Hunt 1983. *Veterinary Pathology*. 5th Eda, Lea and Febiger, Philadelphia.
- Kumar, S. and S.C. Pant 1984. Organal damage caused by aldicarb to a freshwater teleost, *Barbus conchoni* Hamilton. *Bull. Environ. Contam. Toxicol.*, 33 : 50-55.
- Kumar, S. and S.C. Pant 1985. Renal pathology in fish (*Puntius conchoni* Hamilton) following exposure to acutely lethal and sublethal concentration of monocrotophos. *Bull. Environ. Contam. Toxicol.*, 35 : 228-233.
- Mathur, D.S. 1964. Determination of the accumulation of DDT and BHC in liver and intestine of certain fishes. *Proc. Nat. Acad. Sci. India*, 34B : 377-338.
- Pandey, A.K., M.P. Mohamed, K.C. George and Shyamlal 1993. Histopathological changes in gill, kidney and liver of an estuarine mullet, *Liza parsia*, induced by sublethal exposure to DDT. *J. Indian Fish Assoc.*, 23 : 55-63.
- Ramamurthy, R., R. Nagarathnamma, B. Jayasunderamma and P. Rama Rao 1987. Histopathological lesions in the gill of freshwater teleost, *Cyprinus carpio*, induced by methyl parathion. *Matsya*, 13 : 144-147.
- Roberts, R.J. 1989. *Fish Pathology*. 2nd Edn., Bailliere, Tindall, London.
- Ruvio, M. 1972. *Marine Pollution and Sea Life*. FAO News Ltd., London.
- Saxena, M.P., K. Gopal, W. Jones and P.K. Ray 1992. Immune response to *Aeromonas hydrophila* in catfish

- (*Heteropneustes fossilis*) exposed to cadmium and hexachlorocyclohexane. *Bull. Environ. Contam. Toxicol.*, 48 : 194-201.
- Sen Gupta, R. and S.Z. Qasim 1985. Indian Oceans and environmental overviews. In : *Oceans : Realities and Prospects*. R.C. Sharma (Ed.), p. 165-188. Rajesh Publication, New Delhi.
- Verma, S.R., S.P. Gupta and M.P. Tyagi 1975. Studies on the toxicity of lindane on *Colia fasciata*. 1. TLM measurements and histopathological changes in certain tissues. *Geg. Morphol. Jahrb. (Leipzig)*, 122 : 38-54.