

**A NOTE ON THE FOOD OF MALABAR TREVALLY,
CARANGOIDES MALABARICUS (BLOCH & SCHNEIDER)
FROM THE NORTH-WESTERN BAY OF BENGAL**

Carangoides malabaricus (Bloch & Schneider) is the most important carangid landed by the trawlers operating in the north-western Bay of Bengal. It is observed in the trawl catches at all hours of the day. Preliminary observations on the food and feeding habits of the fish collected during 1964-66 show that it is essentially a carnivore and a column feeder and that *Acetes*, prawns, *Squilla*, crabs and miscellaneous small fishes are the important items of food besides small quantities of mysids, amphipods, other crustaceans, squids and cuttle fish. However, sometimes the food exclusively consists of only one of the items mentioned above.

On 18th July, 1964 and 8th August, 1966 an unusual item of food was found in the stomachs of *C. malabaricus* and was identified as *Euphausia distinguenda* Hansen. The euphausiids were very fresh and majority mature, as shown by well developed petasma in males and spermatophores attached to the thelyca in females. Details regarding the collection of fish samples and percentage composition of the feed on the days of occurrence of the euphausiids in the stomachs of the fish are given below.

TABLE

Date.	18-7-1964.	8-8-1966.
Vessel.	M.T. Ashok.	M.V. Champa
Area.	Lat. 18° 35'N., Long. 84° 35'E.	Lat. 17° 35'N., Long. 83° 25'E.
Time of fishing.	0630-1115 hrs.	0800-1200 hrs.
Number of stomachs examined.	8	24
Length range of fish in mm.	124-172	122-190
Number of stomachs in which euphausiid is seen.	2	24
Average number of euphausiids per stomach	53	168
Length range of the euphausiid in mm.	7.0-8.05	7.5-13.5

Percentage composition of the feed in the stomachs in which euphausiids were seen.

(a) Euphausiids	75.0%	69.9%
(b) <i>Squilla</i> sp.	20.0%	—
(c) Crabs	5.0%	0.7%
(d) Prawns	—	0.9%
(e) Fish juveniles	—	1.5%

It is interesting to note that 21 out of 24 stomachs examined on 8-8-1966 contained only euphausiids. Most of the stomachs were full. The quantity of the feed indicates that the fish had fed on a thick patch of the euphausiid while the freshness of the feed shows that the fish had fed on the euphausiids just before they were caught. Examination of the stomachs of *C. malabaricus* collected a few days before and after 8-8-1966 revealed no euphausiids in them but mainly or only *Acetes*. The present observation also indicates that *C. malabaricus* is not particularly selective in regard to food.

A review of the literature shows no reports on the occurrence of *E. distinguenda* Hansen in the guts of fish from Indian waters although this species is reported to be abundant in the Arabian Sea (Illig, 1930; Tattersall, 1939; Brinton, 1963 and Pono-

mareva, 1964). However, Ganapati and Rao (1957) report the presence of euphausiids (species not given) from the stomachs of *Sardinella gibbosa* (Blkr.) off Waltair coast.

The appearance of *E. distinguenda*, which is usually a deep water form (Illig, 1930 and Tattersall, 1939) in the near-shore waters (see table) as in the present case is interesting. The fact that although *C. malabaricus* is landed in the trawl nets at all hours of the day, *E. distinguenda* was observed in the stomachs of fishes caught only in the forenoon is suggestive of a vertical movement of the latter in the Bay. The euphausiid material comprised both adults and sub-adults on the first occasion (18-7-1964) but only mature adults on the second occasion (8-8-1966). As against this the extent of the feed suggests that the euphausiid was perhaps present in a thick patch in the Bay on the second occasion, but not so on the first occasion. This supports the view of Ponomareva (1958) and others that during spawning season euphausiids congregate in swarms and ascend to deposit eggs. Sebastian (1965) states that *E. distinguenda* occurs mostly in swarms.

It is known that euphausiids are among the organisms responsible for the deep scattering layer in the sea (Marshall, 1954), but from the seas around India there is practically no information in this regard. Their importance in the economy of the sea is well known. Further information on the occurrence of euphausiids in the stomach contents of fishes, especially carangids and other column feeders, their maturity and the time of occurrence should be interesting.

The author wishes to thank Dr. K. V. Sekharan for much help and encouragement in preparing this note.

Central Marine Fisheries Research Institute,
Mandapam Camp.

S. REUBEN

REFERENCES

- BRINTON, E. 1963. *All-India Seminar on Marine Sciences*, April, 26-27, Waltair, 1-3 (Mimeo).
- GANAPATI, P. N. & SRINIVASARAO, K. 1957. *Jour. Zool. Soc. India*, 9(2) : 162-182.
- ILLIG, G. 1930. *Wiss. Ergebn. Deutsch. Tief-See Exped., 'Valdivia'* 22(6) : 400-625.
- MARSHALL, N. B. 1954. *Aspects of Deep Sea Biology*, pp. 380.
- PONOMAREVA, L. A. 1958. *Rybn. Khozy.*, 4 : 10-12.
- . 1964. *Trud. Inst. Okeanol.*, LXIV : 265-270.
- SEBASTIAN, M. J. 1965. *Symposium on Crustacea, Part I*, Mar. biol. Ass. India, Mandapam Camp : 233-254.
- TATTERSALL, W. M. 1939. *Sci. Rept.*, 5(8) : 203-246.