

CMFRI

bulletin 42

Part One

AUGUST 1988



NATIONAL SEMINAR ON SHELLFISH RESOURCES AND FARMING

TUTICORIN

19-21 January, 1987

Session - I

CENTRAL MARINE FISHERIES RESEARCH INSTITUTE
(Indian Council of Agricultural Research)
P. B. No. 2704, E. R. G. Road, Cochin-682 031, India

35. ON THE CHROMOSOMES OF A BIVALVE, *ANADARA RHOMBEA*

Sm. P. Muthu

Department of Biology, Sri Parama/oiyani College, Alwarkurichi-627 412

ABSTRACT

Study of chromosomes of bivalves is technically difficult. Spermatogonia and spermatocytes, oocytes, and cleavage stages of the fertilized eggs are more favourable than the somatic tissues of the adults for the study of chromosomes. Hypotonic treatment, prior to fixation, is an essential pre-treatment for obtaining well-spread chromosomes. The study of the chromosomes of a bivalve, *Anadara rhombea*, from the Porto Novo waters, has been the first of its kind in an Indian bivalve. Giemsa and first metaphase stages from 'squash' preparations of testis show that in male $2n = 28$ in this species; the sex chromosomes, if any, are not distinguishable.

INTRODUCTION

A survey of the literature on the chromosomal cytology of the molluscs available upto 1978 (when the study reported here was undertaken) will reveal the relative paucity of

information on the chromosomes of this group despite the fact that molluscs constitute the third large phylum in the animal kingdom (Fretter & Graham 1962); among molluscs, very few bivalves have been studied cytologically. Lillie's (1901) study on the egg of */?* is the earliest

known on the chromosomes of a bivalve, based on its maturation, fertilization and cleavage. Other works on the chromosomes of bivalves include those of; Ahmed and Sparks (1967 a) on oysters, clams and mussels, Ahmed and Sparks (1967 b) on *Ostrea* and *Crossostrea*, Ahmed and Sparks (1970) on two spp of *Mytilus*; Ieyama (1975) on three spp of Pteriomorpha; Ieyama and Inaba (1974) on ten spp of Pteriomorpha; Keyl (1956) on the mussel, *Sphaerium*; Kobayashi (1954) on two spp of *Ostrea*; Longwell et al (1967) on *Crossostrea*; Menzel (1988) on nine families of marine bivalves; Menzel and Menzel (1965) on two spp of *Mercenaria* and their hybrids, and Ropes (1972) on the surf clam, *Spisula*. It will be obvious from this list that there was no work on Indian bivalves till 1978.

MATERIAL AND METHODS

The specimens of *Anadara rhombea* were kindly provided by Dr. George John who was then working in the Centre of Advanced Study in Marine Biology, Porto Novo, on the Biology of *Anadara* spp. The specimens were obtained from Vellar estuary. Somatic tissues from the ctenidia and midgut gland, and the germinal material from the testis were used for this study. Testis proved to be the most favourable material for the study of chromosomes. Fragments of live testis were subjected, for 30 minutes, to hypotonic treatment prior to fixation; filtered estuarine water was diluted with an equal quantity of distilled water and this was used as the hypotonic fluid. The fragments were then fixed in Clarke's fluid or Newcomer's fluid, and stained in lacto-propiono-orcein or acetic-orcein. The chromosomes were studied from 'squash' preparations which were dehydrated in ethanol and mounted in euparal.

The slides were examined in an Olympus trinocular research microscope under oil immersion objective of N. A. 1.3 and with Kohler illumination; camera lucida sketches were drawn with a table top magnification of 3200 dia.

OBSERVATION

The chromosomes were studied from spermatocytes undergoing first meiotic division; cells in diakinesis and I metaphase provided

satisfactory preparations. The haploid number of chromosomes in *A. rhombea* was found to be 14 in males (Fig 1). The chromosomes are metacentric and form 14 bivalents. There is no localisation of chiasmata, there being proximal, interstitial and distal chiasmata. The meiotic behaviour of the chromosomes does not show any peculiarity. It was not possible to distinguish any sex chromosomes based on the morphology and behaviour of the chromosomes,



Fig. 1. Chromosomes of *Anadara rhombea*. Squash preparations of testes fixed, after hypotonic treatment, in Clarke's fluid and stained in lacto-propiono-orcein.

Camera lucida sketches; magnification: 3200 X.

➤ Bivalents of a cell in diakinesis; $n = 14$.

▲, ▲'▲'''''''' ○' ● '' '' ' ''«Ph"e; $n = u$,

DISCUSSION

The dearth of information on the chromosomes of bivalves is in all probability due to the technical difficulty of obtaining good preparations of well-spread chromosomes; molluscs, and the bivalves, in particular, have their chromosomes (which are in some spp large in number) in the relatively small volume of their nuclei. The conventional methods of chromosome preparations, employed in the majority of most other invertebrates, are not adequate for the investigation of the bivalve chromosomes; it is absolutely necessary to subject the tissues to hypotonic treatment in order to swell the nuclei and to disperse the chromosomes so that they are spread well while squashing, without an overlapping of the chromosomes. Besides diluted natural medium, 1% aqueous solution of sodium acetate or 0.4% aqueous solution of potassium chloride maybe used for hypotonic treatment of the material; 20 to 30 minutes in the hypotonic fluid will suffice; the tissues may then be fixed and stained; squashes can be prepared from such material and they will show a good number of well-spread chromosomes for observation.

Regarding the choice of favourable material for the study of chromosomes, in the experience of the present author, it is better to choose germinal material from the males or females during active gametogenesis. The most favourable materials for chromosomal studies are the spermatogonia and spermatocytes in the males, and the maturation stages of the oocytes in the females; cleavage stages of the fertilized eggs provide good mitotic chromosomes. Somatic tissues of the adults, such as the ctenidia or midgut glands, do provide mitotic stages, but are not as good as the gametocytes or the early stages of cleavages. For instance, Alagarwami & Sreenivasan (MS) found that the oocytes of pearl oysters were the most favourable for a study of the chromosomes; more than even the testis, the oocytes provided excellent stages of maturation in which the chromosomes were distinctly discernible. Cleavage stages in the oysters were not very useful in their studies

Patterson (1969) had reviewed the chromosome numbers of the bivalves and stated that the haploid numbers ranged from 10-23. Later work has indicated that the lower haploid number is 7. Among the bivalves cytologically investigated, the family Ostreidae have been the most investigated; except in two spp of *Ostrea*, in all other spp of *Ostrea* and *Crossostrea* which belong to this family, the haploid number has been recorded to be 10. In every family of the bivalves, the haploid number is constant for all the genera. Thus, the haploid number for the family Arcidae has been reported (Ieyama 1975) to be 14. In *Anadara rhombea*, which belongs to Arcidae, the haploid number has been found by the present author to be 14. Thus, the present observation confirms the constancy of the familial haploid number.

ACKNOWLEDGEMENTS

The author is thankful to Dr. George John for collecting and providing the specimens. He is indebted to Professor R. Natarajan, Director of the Centre of Advanced Study in Marine Biology, Porto Novo, for providing the facilities and for his constant encouragement. The award of a Teacher Fellowship under the Faculty Improvement Programme by the U. G. C. is gratefully acknowledged.

REFERENCES

AHMED, M. AND A.K. SPARKS. 1967a. Chromosomes of oysters, clams and mussels. *Proc. Nat. Shellfish Assoc.* Abstracts N S A Pacific Coast section, 58 : 10.

AHMED, M. AND A. K. SPARKS. 1967b. A preliminary study of chromosomes of two species of oysters (*Ostrea lurida* and *Crossostrea gigas*.) *J. Fish. Res. Bd Canads*, 24 : 2155-2159.

AHMED, M. AND A.K SPARKS. 1970. Chromosome number, structure and autosomal polymorphism in marine mussels, *Mytilus edulis* and *Mytilus californianus*. *fl/o. flu//*, 138 (1) : 1-13.

RETTER V. AND A. GRAHAM. 1962. Br.t.sh Prosobranch Molluscs. The Ray Society, London.

IEYAMA, H. 1975. Chromosome numbers of three species in three families of Pteriomorpha (Bivalvia). *Venus*, 34 (1-2) : 26-32.

IEYAMA, H. AND A. INABA. 1974. Chromosome numbers of ten species in four families of Pteriomorpha (Bivalvia). *Venus*, 33 (3) : 129-137.

KEYL, H, G. 1956. Beobachtungen über die Omeiose der Muschel *Spharium corneum*-Chromosome, 8*12-M.

KOBAYASHI, H. 1954. Über die Chromosomen von japanischen Arten von *Cytologia*, 19 (4) : 371-376.

LILLIE, F.R. 1901. The organization of the egg of *Unio*, based on a study of its maturation, fertilization, and cleavage, *J. Morph.* 17 : 227-292.

LONGWELL, A. C, S. S. STILES AND D. G. SMITH. 1967. Chromosome complement of the American oyster, *Crossostrea virginica*, as seen in meiotic and cleavage stages. *Canad. J. Genet. and Cytol.*, 9 : 845-856.

MENZEL, R. W. 1965. Chromosome number in nine families of marine pelecypod mollusks. *Nautilus*, 82 (2) : 45-58.

MENZEL, R. W AND M. Y. MENZEL. 1965. Chromosomes of two species of quahog. *Biol Bull.* 129-181-188

PATTERSON, C. M. 1969. Chromosomes of molluscs. *Proc. Symp. Mollusca. Marine Association of India Pt. 2* : 635-686.

ROPES, J. W. 1972. Chromosome number of surf clam, *Spisula soJidissima*. *Nautilus*, 85 (3) : 93-95.